

Limit theorems for nonlinear functionals of Volterra processes via white noise analysis

SÉBASTIEN DARSE¹, IVAN NOURDIN² and DAVID NUALART³

¹Université Aix-Marseille I, 39 rue Joliot Curie, 13453 Marseille Cedex 13, France.

E-mail: darses@cmi.univ-mrs.fr

²Laboratoire de Probabilités et Modèles Aléatoires, Université Pierre et Marie Curie (Paris VI), Boîte Courrier 188, 4 place Jussieu, 75252 Paris Cedex 05, France. E-mail: ivan.nourdin@upmc.fr

³Department of Mathematics, University of Kansas, Lawrence, KS 66045, USA.

E-mail: nualart@math.ku.edu

By means of white noise analysis, we prove some limit theorems for nonlinear functionals of a given Volterra process. In particular, our results apply to fractional Brownian motion (fBm) and should be compared with the classical convergence results of the 1980s due to Breuer, Dobrushin, Giraitis, Major, Surgailis and Taqqu, as well as the recent advances concerning the construction of a Lévy area for fBm due to Coutin, Qian and Unterberger.

Keywords: fractional Brownian motion; limit theorems; Volterra processes; white noise analysis

1. Introduction

Fix $T > 0$ and let $B = (B_t)_{t \geq 0}$ be a fractional Brownian motion with Hurst index $H \in (0, 1)$, defined on some probability space (Ω, \mathcal{B}, P) . Assume that \mathcal{B} is the completed σ -field generated by B . Fix an integer $k \geq 2$ and, for $\varepsilon > 0$, consider

$$G_\varepsilon = \varepsilon^{-k(1-H)} \int_0^T h_k \left(\frac{B_{u+\varepsilon} - B_u}{\varepsilon^H} \right) du. \quad (1.1)$$

Here, and in the rest of this paper,

$$h_k(x) = (-1)^k e^{x^2/2} \frac{d^k}{dx^k} (e^{-x^2/2}) \quad (1.2)$$

stands for the k th Hermite polynomial. We have $h_2(x) = x^2 - 1$, $h_3(x) = x^3 - 3x$ and so on.

Since the seminal works [3,6,7,19,20] by Breuer, Dobrushin, Giraitis, Major, Surgailis and Taqqu, the following three convergence results are classical:

- if $H < 1 - \frac{1}{2k}$, then

$$\left((B_t)_{t \in [0, T]}, \varepsilon^{k(1-H)-1/2} G_\varepsilon \right) \xrightarrow[\varepsilon \rightarrow 0]{\text{Law}} \left((B_t)_{t \in [0, T]}, N \right), \quad (1.3)$$

where $N \sim \mathcal{N}(0, T \times k! \int_0^T \rho^k(x) dx)$ is independent of B , with $\rho(x) = \frac{1}{2}(|x+1|^{2H} + |x-1|^{2H} - 2|x|^{2H})$;

- if $H = 1 - \frac{1}{2k}$, then

$$\left((B_t)_{t \in [0, T]}, \frac{G_\varepsilon}{\sqrt{\log(1/\varepsilon)}} \right) \xrightarrow[\varepsilon \rightarrow 0]{\text{Law}} \left((B_t)_{t \in [0, T]}, N \right), \tag{1.4}$$

where $N \sim \mathcal{N}(0, T \times 2k!(1 - \frac{1}{2k})^k (1 - \frac{1}{k})^k)$ is independent of B ;

- if $H > 1 - \frac{1}{2k}$, then

$$G_\varepsilon \xrightarrow[\varepsilon \rightarrow 0]{L^2(\Omega)} Z_T^{(k)}, \tag{1.5}$$

where $Z_T^{(k)}$ denotes the Hermite random variable of order k ; see Section 4.1 for its definition.

Combining (1.3) with the fact that $\sup_{0 < \varepsilon \leq 1} E[|\varepsilon^{k(1-H)-1/2} G_\varepsilon|^p] < \infty$ for all $p \geq 1$ (use the boundedness of $\text{Var}(\varepsilon^{k(1-H)-1/2} G_\varepsilon)$ and a classical hypercontractivity argument), we have, for all $\eta \in L^2(\Omega)$ and if $H < 1 - \frac{1}{2k}$, that

$$\varepsilon^{k(1-H)-1/2} E[\eta G_\varepsilon] \xrightarrow[\varepsilon \rightarrow 0]{} E(\eta N) = E(\eta)E(N) = 0$$

(a similar statement holds in the critical case $H = 1 - \frac{1}{2k}$). This means that $\varepsilon^{k(1-H)-1/2} G_\varepsilon$ converges *weakly* in $L^2(\Omega)$ to zero. The following question then arises. Is there a normalization of G_ε ensuring that it converges *weakly* towards a *non-zero* limit when $H \leq 1 - \frac{1}{2k}$? If so, then what can be said about the limit? The first purpose of the present paper is to provide an answer to this question in the framework of *white noise analysis*.

In [14], it is shown that for all $H \in (0, 1)$, the time derivative \dot{B} (called the *fractional white noise*) is a distribution in the sense of Hida. We also refer to Bender [1], Biagini *et al.* [2] and references therein for further works on the fractional white noise.

Since we have $E(B_{u+\varepsilon} - B_u)^2 = \varepsilon^{2H}$, observe that G_ε defined in (1.1) can be rewritten as

$$G_\varepsilon = \int_0^T \left(\frac{B_{u+\varepsilon} - B_u}{\varepsilon} \right)^{\diamond k} du, \tag{1.6}$$

where $(\dots)^{\diamond k}$ denotes the k th Wick product. In Proposition 9 below, we will show that for all $H \in (\frac{1}{2} - \frac{1}{k}, 1)$,

$$\lim_{\varepsilon \rightarrow 0} \int_0^T \left(\frac{B_{u+\varepsilon} - B_u}{\varepsilon} \right)^{\diamond k} du = \int_0^T \dot{B}_u^{\diamond k} du, \tag{1.7}$$

where the limit is in the $(S)^*$ sense.

In particular, we observe two different types of asymptotic results for G_ε when $H \in (\frac{1}{2} - \frac{1}{k}, 1 - \frac{1}{2k})$: convergence (1.7) in $(S)^*$ to a Hida distribution, and convergence (1.3) in law to a normal law, with rate $\varepsilon^{1/2-k(1-H)}$. On the other hand, when $H \in (1 - \frac{1}{2k}, 1)$, we obtain from (1.5) that the Hida distribution $\int_0^T \dot{B}_s^{\diamond k} ds$ turns out to be the square-integrable random variable $Z_T^{(k)}$, which is an interesting result in its own right.

In Proposition 4, the convergence (1.7) in $(S)^*$ is proved for a general class of Volterra processes of the form

$$\int_0^t K(t, s) dW_s, \quad t \geq 0, \tag{1.8}$$

where W stands for a standard Brownian motion, provided the kernel K satisfies some suitable conditions; see Section 3.

We also provide a new proof of the convergence (1.3) based on the recent general criterion for the convergence in distribution to a normal law of a sequence of multiple stochastic integrals established by Nualart and Peccati [15] and by Peccati and Tudor [17], which avoids the classical method of moments.

In two recent papers [9,10], Marcus and Rosen have obtained central and non-central limit theorems for a functional of the form (1.1), where B is a mean zero Gaussian process with stationary increments such that the covariance function of B , defined by $\sigma^2(|t - s|) = \text{Var}(B_t - B_s)$, is either convex (plus some additional regularity conditions), concave or given by $\sigma^2(h) = h^r$ with $1 < r < 2$. These theorems include the convergence (1.3) and, unlike our simple proof, are based on the method of moments.

In the second part of the paper, we develop a similar analysis for functionals of two independent fractional Brownian motions (or, more generally, Volterra processes) related to the Lévy area. More precisely, consider two *independent* fractional Brownian motions $B^{(1)}$ and $B^{(2)}$ with (for simplicity) the same Hurst index $H \in (0, 1)$. We are interested in the convergence, as $\varepsilon \rightarrow 0$, of

$$\tilde{G}_\varepsilon := \int_0^T B_u^{(1)} \frac{B_{u+\varepsilon}^{(2)} - B_u^{(2)}}{\varepsilon} du \tag{1.9}$$

and

$$\check{G}_\varepsilon := \int_0^T \left(\int_0^u \frac{B_{v+\varepsilon}^{(1)} - B_v^{(1)}}{\varepsilon} dv \right) \frac{B_{u+\varepsilon}^{(2)} - B_u^{(2)}}{\varepsilon} du. \tag{1.10}$$

Note that \tilde{G}_ε coincides with the ε -integral associated with the forward Russo–Vallois integral $\int_0^T B^{(1)} d^- B^{(2)}$; see, for example, [18] and references therein. Over the last decade, the convergences of \tilde{G}_ε and \check{G}_ε (or of related families of random variables) have been intensively studied. Since $\varepsilon^{-1} \int_0^u (B_{v+\varepsilon}^{(1)} - B_v^{(1)}) dv$ converges pointwise to $B_u^{(1)}$ for any u , we could think that the asymptotic behaviors of \tilde{G}_ε and \check{G}_ε are very close as $\varepsilon \rightarrow 0$. Surprisingly, this is not the case. Actually, only the result for \check{G}_ε agrees with the seminal result of Coutin and Qian [4] (that is, we have convergence of \check{G}_ε in $L^2(\Omega)$ if and only if $H > 1/4$) and with the recent result by Unterberger [21] (that is, adequately renormalized, \check{G}_ε converges in law if $H < 1/4$). More precisely:

- if $H < 1/4$, then

$$\left((B_t^{(1)}, B_t^{(2)})_{t \in [0, T]}, \varepsilon^{1/2-2H} \check{G}_\varepsilon \right) \xrightarrow[\varepsilon \rightarrow 0]{\text{Law}} \left((B_t^{(1)}, B_t^{(2)})_{t \in [0, T]}, N \right), \tag{1.11}$$

where $N \sim \mathcal{N}(0, T\check{\sigma}_H^2)$ is independent of $(B^{(1)}, B^{(2)})$ and

$$\check{\sigma}_H^2 = \frac{1}{4(2H+1)(2H+2)} \int_{\mathbb{R}} (|x+1|^{2H} + |x-1|^{2H} - 2|x|^{2H}) \times (2|x|^{2H+2} - |x+1|^{2H+2} - |x-1|^{2H+2}) dx;$$

- if $H = 1/4$, then

$$\left((B_t^{(1)}, B_t^{(2)})_{t \in [0, T]}, \frac{\check{G}_\varepsilon}{\sqrt{\log(1/\varepsilon)}} \right) \xrightarrow[\varepsilon \rightarrow 0]{\text{Law}} \left((B_t^{(1)}, B_t^{(2)})_{t \in [0, T]}, N \right), \tag{1.12}$$

where $N \sim \mathcal{N}(0, T/8)$ is independent of $(B^{(1)}, B^{(2)})$;

- if $H > 1/4$, then

$$\check{G}_\varepsilon \xrightarrow[\varepsilon \rightarrow 0]{L^2(\Omega)} \int_0^T B_u^{(1)} \diamond \dot{B}_u^{(2)} du = \int_0^T B_u^{(1)} dB_u^{(2)}; \tag{1.13}$$

- for all $H \in (0, 1)$, we have

$$\check{G}_\varepsilon \xrightarrow[\varepsilon \rightarrow 0]{(S)^*} \int_0^T B_u^{(1)} \diamond \dot{B}_u^{(2)} du. \tag{1.14}$$

However, for \tilde{G}_ε , we have, in contrast:

- if $H < 1/2$, then

$$\left((B_t^{(1)}, B_t^{(2)})_{t \in [0, T]}, \varepsilon^{1/2-H} \tilde{G}_\varepsilon \right) \xrightarrow[\varepsilon \rightarrow 0]{\text{Law}} \left((B_t^{(1)}, B_t^{(2)})_{t \in [0, T]}, N \times S \right), \tag{1.15}$$

where

$$S = \sqrt{\int_0^\infty (|x+1|^{2H} + |x-1|^{2H} - 2|x|^{2H}) dx \times \int_0^T (B_u^{(1)})^2 du}$$

and $N \sim \mathcal{N}(0, 1)$, independent of $(B^{(1)}, B^{(2)})$;

- if $H \geq 1/2$, then

$$\tilde{G}_\varepsilon \xrightarrow[\varepsilon \rightarrow 0]{L^2(\Omega)} \int_0^T B_u^{(1)} \diamond \dot{B}_u^{(2)} du = \int_0^T B_u^{(1)} dB_u^{(2)}; \tag{1.16}$$

- for all $H \in (0, 1)$, we have

$$\tilde{G}_\varepsilon \xrightarrow[\varepsilon \rightarrow 0]{(S)^*} \int_0^T B_u^{(1)} \diamond \dot{B}_u^{(2)} du. \tag{1.17}$$

Finally, we study the convergence, as $\varepsilon \rightarrow 0$, of the so-called ε -covariation (following the terminology of [18]) defined by

$$\widehat{G}_\varepsilon := \int_0^T \frac{B_{u+\varepsilon}^{(1)} - B_u^{(1)}}{\varepsilon} \times \frac{B_{u+\varepsilon}^{(2)} - B_u^{(2)}}{\varepsilon} du \tag{1.18}$$

and we get:

- if $H < 3/4$, then

$$\left((B_t^{(1)}, B_t^{(2)})_{t \in [0, T]}, \varepsilon^{3/2-2H} \widehat{G}_\varepsilon \right) \xrightarrow[\varepsilon \rightarrow 0]{\text{Law}} \left((B_t^{(1)}, B_t^{(2)})_{t \in [0, T]}, N \right) \tag{1.19}$$

with $N \sim \mathcal{N}(0, T\widehat{\sigma}_H^2)$ independent of $(B^{(1)}, B^{(2)})$ and

$$\widehat{\sigma}_H^2 = \frac{1}{4} \int_{\mathbb{R}} (|x+1|^{2H} + |x-1|^{2H} - 2|x|^{2H})^2 dx;$$

- if $H = 3/4$, then

$$\left((B_t^{(1)}, B_t^{(2)})_{t \in [0, T]}, \frac{\widehat{G}_\varepsilon}{\sqrt{\log(1/\varepsilon)}} \right) \xrightarrow[\varepsilon \rightarrow 0]{\text{Law}} \left((B_t^{(1)}, B_t^{(2)})_{t \in [0, T]}, N \right) \tag{1.20}$$

with $N \sim \mathcal{N}(0, 9T/32)$ independent of $(B^{(1)}, B^{(2)})$;

- if $H > 3/4$, then

$$\widehat{G}_\varepsilon \xrightarrow[\varepsilon \rightarrow 0]{L^2(\Omega)} \int_0^T \dot{B}_u^{(1)} \diamond \dot{B}_u^{(2)} du; \tag{1.21}$$

- for all $H \in (0, 1)$, we have

$$\widehat{G}_\varepsilon \xrightarrow[\varepsilon \rightarrow 0]{(S)^*} \int_0^T \dot{B}_u^{(1)} \diamond \dot{B}_u^{(2)} du. \tag{1.22}$$

The paper is organized as follows. In Section 2, we introduce some preliminaries on white noise analysis. Section 3 is devoted to the study, using the language and tools of the previous section, of the asymptotic behaviors of G_ε , $\widetilde{G}_\varepsilon$ and \widehat{G}_ε in the (more general) context where B is a Volterra process. Section 4 is concerned with the fractional Brownian motion case. In Section 5 (resp., Section 6), we prove (1.3) and (1.4) (resp., (1.11), (1.12), (1.15), (1.19) and (1.20)).

2. White noise functionals

In this section, we present some preliminaries on white noise analysis. The classical approach to white noise distribution theory is to endow the space of tempered distributions $\mathcal{S}'(\mathbb{R})$ with a Gaussian measure \mathbb{P} such that, for any rapidly decreasing function $\eta \in \mathcal{S}(\mathbb{R})$,

$$\int_{\mathcal{S}'(\mathbb{R})} e^{i\langle x, \eta \rangle} \mathbb{P}(dx) = e^{-|\eta|_0^2/2}.$$

Here, $|\cdot|_0$ denotes the norm in $L^2(\mathbb{R})$ and $\langle \cdot, \cdot \rangle$ the dual pairing between $\mathcal{S}'(\mathbb{R})$ and $\mathcal{S}(\mathbb{R})$. The existence of such a measure is ensured by Minlos' theorem [8].

In this way, we can consider the probability space $(\Omega, \mathcal{B}, \mathbb{P})$, where $\Omega = \mathcal{S}'(\mathbb{R})$. The pairing $\langle x, \xi \rangle$ can be extended, using the norm of $L^2(\Omega)$, to any function $\xi \in L^2(\mathbb{R})$. Then,

$W_t = \langle \cdot, \mathbf{1}_{[0,t]} \rangle$ is a two-sided Brownian motion (with the convention that $\mathbf{1}_{[0,t]} = -\mathbf{1}_{[t,0]}$ if $t < 0$) and for any $\xi \in L^2(\mathbb{R})$,

$$\langle \cdot, \xi \rangle = \int_{-\infty}^{\infty} \xi \, dW = I_1(\xi)$$

is the Wiener integral of ξ .

Let $\Phi \in L^2(\Omega)$. The classical Wiener chaos expansion of Φ says that there exists a sequence of symmetric square-integrable functions $\phi_n \in L^2(\mathbb{R}^n)$ such that

$$\Phi = \sum_{n=0}^{\infty} I_n(\phi_n), \tag{2.1}$$

where I_n denotes the multiple stochastic integral.

2.1. The space of Hida distributions

Let us recall some basic facts concerning tempered distributions. Let $(\xi_n)_{n=0}^{\infty}$ be the orthonormal basis of $L^2(\mathbb{R})$ formed by the Hermite functions given by

$$\xi_n(x) = \pi^{-1/4} (2^n n!)^{-1/2} e^{-x^2/2} h_n(x), \quad x \in \mathbb{R}, \tag{2.2}$$

where h_n are the Hermite polynomials defined in (1.2). The following two facts can immediately be checked: (a) there exists a constant $K_1 > 0$ such that $\|\xi_n\|_{\infty} \leq K_1 (n + 1)^{-1/12}$; (b) since $\xi'_n = \sqrt{\frac{n}{2}} \xi_{n-1} - \sqrt{\frac{n+1}{2}} \xi_{n+1}$, there exists a constant $K_2 > 0$ such that $\|\xi'_n\|_{\infty} \leq K_2 n^{5/12}$.

Consider the positive self-adjoint operator A (whose inverse is Hilbert–Schmidt) given by $A = -\frac{d^2}{dx^2} + (1 + x^2)$. We have $A\xi_n = (2n + 2)\xi_n$.

For any $p \geq 0$, define the space $\mathcal{S}_p(\mathbb{R})$ to be the domain of the closure of A^p . Endowed with the norm $|\xi|_p := |A^p \xi|_0$, it is a Hilbert space. Note that the norm $|\cdot|_p$ can be expressed as follows, if one uses the orthonormal basis (ξ_n) :

$$|\xi|_p^2 = \sum_{n=0}^{\infty} \langle \xi, \xi_n \rangle^2 (2n + 2)^{2p}.$$

We denote by $\mathcal{S}'_p(\mathbb{R})$ the dual of $\mathcal{S}_p(\mathbb{R})$. The norm in $\mathcal{S}'_p(\mathbb{R})$ is given by (see [16], Lemma 1.2.8)

$$|\xi|_{-p}^2 = \sum_{n=0}^{\infty} |\langle \xi, A^{-p} \xi_n \rangle|^2 = \sum_{n=0}^{\infty} \langle \xi, \xi_n \rangle^2 (2n + 2)^{-2p}$$

for any $\xi \in \mathcal{S}'_p(\mathbb{R})$. One can show that the projective limit of the spaces $\mathcal{S}_p(\mathbb{R})$, $p \geq 0$, is $\mathcal{S}(\mathbb{R})$, that the inductive limit of the spaces $\mathcal{S}_p(\mathbb{R})'$, $p \geq 0$, is $\mathcal{S}'(\mathbb{R})$ and that

$$\mathcal{S}(\mathbb{R}) \subset L^2(\mathbb{R}) \subset \mathcal{S}'(\mathbb{R})$$

is a Gel'fand triple.

We can now introduce the Gel'fand triple

$$(\mathcal{S}) \subset L^2(\Omega) \subset (\mathcal{S})^*,$$

via the second quantization operator $\Gamma(A)$. This is an unbounded and densely defined operator on $L^2(\Omega)$ given by

$$\Gamma(A)\Phi = \sum_{n=0}^{\infty} I_n(A^{\otimes n} \phi_n),$$

where Φ has the Wiener chaos expansion (2.1). If $p \geq 0$, then we denote by $(\mathcal{S})_p$ the space of random variables $\Phi \in L^2(\Omega)$ with Wiener chaos expansion (2.1) such that

$$\|\Phi\|_p^p := E[|\Gamma(A)^p \Phi|^2] = \sum_{n=0}^{\infty} n! |\phi_n|_p^2 < \infty.$$

In the above formula, $|\phi_n|_p$ denotes the norm in $\mathcal{S}_p(\mathbb{R})^{\otimes n}$. The projective limit of the spaces $(\mathcal{S})_p$, $p \geq 0$, is called the space of test functions and is denoted by (\mathcal{S}) . The inductive limit of the spaces $(\mathcal{S})_{-p}$, $p \geq 0$, is called the space of Hida distributions and is denoted by $(\mathcal{S})^*$. The elements of $(\mathcal{S})^*$ are called *Hida distributions*. The main example is the time derivative of the Brownian motion, defined as $\dot{W}_t = \langle \cdot, \delta_t \rangle$. One can show that $|\delta_t|_{-p} < \infty$ for some $p > 0$.

We denote by $\langle\langle \Phi, \Psi \rangle\rangle$ the dual pairing associated with the spaces (\mathcal{S}) and $(\mathcal{S})^*$. On the other hand (see [16], Theorem 3.1.6), for any $\Phi \in (\mathcal{S})^*$, there exist $\phi_n \in \mathcal{S}(\mathbb{R}^n)'$ such that

$$\langle\langle \Phi, \Psi \rangle\rangle = \sum_{n=0}^{\infty} n! \langle \phi_n, \psi_n \rangle,$$

where $\Psi = \sum_{n=0}^{\infty} I_n(\psi_n) \in (\mathcal{S})$. Moreover, there exists $p > 0$ such that

$$\|\Phi\|_{-p}^2 = \sum_{n=0}^{\infty} n! |\phi_n|_{-p}^2.$$

Then, with a convenient abuse of notation, we say that Φ has a generalized Wiener chaos expansion of the form (2.1).

2.2. The S-transform

A useful tool to characterize elements in $(\mathcal{S})^*$ is the \mathcal{S} -transform. The Wick exponential of a Wiener integral $I_1(\eta)$, $\eta \in L^2(\mathbb{R})$, is defined by

$$: e^{I_1(\eta)} : = e^{I_1(\eta) - |\eta|_0^2/2}.$$

The S -transform of an element $\Phi \in (\mathcal{S})^*$ is then defined by

$$S(\Phi)(\xi) = \langle \langle \Phi, : e^{I_1(\xi)} : \rangle \rangle,$$

where $\xi \in \mathcal{S}(\mathbb{R})$. One can easily see that the S -transform is injective on $(\mathcal{S})^*$.

If $\Phi \in L^2(\Omega)$, then $S(\Phi)(\xi) = E[\Phi : e^{I_1(\xi)} :]$. For instance, the S -transform of the Wick exponential is

$$S(: e^{I_1(\eta)} :)(\xi) = e^{\langle \eta, \xi \rangle}.$$

Also, $S(W_t)(\xi) = \int_0^t \xi(s) ds$ and $S(\dot{W}_t)(\xi) = \xi(t)$.

Suppose that $\Phi \in (\mathcal{S})^*$ has a generalized Wiener chaos expansion of the form (2.1). Then, for any $\xi \in \mathcal{S}(\mathbb{R})$,

$$S(\Phi)(\xi) = \sum_{n=0}^{\infty} \langle \phi_n, \xi^{\otimes n} \rangle,$$

where the series converges absolutely (see [16], Lemma 3.3.5).

The Wick product of two functionals $\Psi = \sum_{n=0}^{\infty} I_n(\psi_n)$ and $\Phi = \sum_{n=0}^{\infty} I_n(\phi_n)$ belonging to $(\mathcal{S})^*$ is defined as

$$\Psi \diamond \Phi = \sum_{n,m=0}^{\infty} I_{n+m}(\psi_n \otimes \phi_m).$$

It can be proven that $\Psi \diamond \Phi \in (\mathcal{S})^*$. The following is an important property of the S -transform:

$$S(\Phi \diamond \Psi)(\xi) = S(\Phi)(\xi)S(\Psi)(\xi). \tag{2.3}$$

If Ψ, Φ and $\Psi \diamond \Phi$ belong to $L^2(\Omega)$, then we have $E[\Psi \diamond \Phi] = E[\Psi]E[\Phi]$.

The following is a useful characterization theorem.

Theorem 1. *A function F is the S -transform of an element $\Phi \in (\mathcal{S})^*$ if and only if the following conditions are satisfied:*

- (1) *for any $\xi, \eta \in \mathcal{S}$, $z \mapsto F(z\xi + \eta)$ is holomorphic on \mathbb{C} ;*
- (2) *there exist non-negative numbers K, a and p such that for all $\xi \in \mathcal{S}$,*

$$|F(\xi)| \leq K \exp(a|\xi|_p^2).$$

Proof. See [8], Theorems 8.2 and 8.10. □

In order to study the convergence of a sequence in $(\mathcal{S})^*$, we can use its S -transform, by virtue of the following theorem.

Theorem 2. *Let $\Phi_n \in (\mathcal{S})^*$ and $S_n = S(\Phi_n)$. Then, Φ_n converges in $(\mathcal{S})^*$ if and only if the following conditions are satisfied:*

- (1) *$\lim_{n \rightarrow \infty} S_n(\xi)$ exists for each $\xi \in \mathcal{S}$;*

(2) there exist non-negative numbers K, a and p such that for all $n \in \mathbb{N}, \xi \in (\mathcal{S}),$

$$|S_n(\xi)| \leq K \exp(a|\xi|_p^2).$$

Proof. See [8], Theorem 8.6. □

3. Limit theorems for Volterra processes

3.1. One-dimensional case

Consider a Volterra process $B = (B_t)_{t \geq 0}$ of the form

$$B_t = \int_0^t K(t, s) dW_s, \tag{3.1}$$

where $K(t, s)$ satisfies $\int_0^t K(t, s)^2 ds < \infty$ for all $t > 0$ and W is the Brownian motion defined on the white noise probability space introduced in the last section. Note that the S -transform of the random variable B_t is given by

$$S(B_t)(\xi) = \int_0^t K(t, s)\xi(s) ds \tag{3.2}$$

for any $\xi \in \mathcal{S}(\mathbb{R})$. We introduce the following assumptions on the kernel K :

(H₁) K is continuously differentiable on $\{0 < s < t < \infty\}$ and for any $t > 0$, we have

$$\int_0^t \left| \frac{\partial K}{\partial t}(t, s) \right| (t - s) ds < \infty;$$

(H₂) $k(t) = \int_0^t K(t, s) ds$ is continuously differentiable on $(0, \infty)$.

Consider the operator K_+ defined by

$$K_+\xi(t) = k'(t)\xi(t) + \int_0^t \frac{\partial K}{\partial t}(t, r)(\xi(r) - \xi(t)) dr,$$

where $t > 0$ and $\xi \in \mathcal{S}(\mathbb{R})$. From Theorem 1, it follows that the linear mapping $\xi \rightarrow K_+\xi(t)$ is the S -transform of a Hida distribution. More precisely, according to [14], define the function

$$C(t) = |k'(t)| + \int_0^t \left| \frac{\partial K}{\partial t}(t, r) \right| (t - r) dr, \quad t \geq 0, \tag{3.3}$$

and observe that the following estimates hold (recall the definition (2.2) of ξ_n):

$$\begin{aligned}
 |K_+\xi(t)| &\leq C(t)(\|\xi\|_\infty + \|\xi'\|_\infty) \\
 &\leq C(t) \sum_{n=0}^\infty |\langle \xi, \xi_n \rangle| (\|\xi_n\|_\infty + \|\xi'_n\|_\infty) \\
 &\leq C(t)M \sum_{n=0}^\infty |\langle \xi, \xi_n \rangle| (n+1)^{5/12} \\
 &\leq C(t)M \sqrt{\sum_{n=0}^\infty |\langle \xi, \xi_n \rangle|^2 (2n+2)^{17/6}} \sqrt{\sum_{n=0}^\infty (n+1)^{-2}} \\
 &= C(t)M|\xi|_{17/12}
 \end{aligned}
 \tag{3.4}$$

for some constants $M > 0$ whose values are not always the same from one line to the next.

We have the following preliminary result.

Lemma 3. Fix an integer $k \geq 1$. Let B be a Volterra process with kernel K satisfying the conditions (H_1) and (H_2) . Assume, moreover, that C defined by (3.3) belongs to $L^k([0, T])$. The function $\xi \mapsto \int_0^T (K_+\xi(s))^k ds$ is then the S -transform of an element of $(S)^*$. This element is denoted by $\int_0^T \dot{B}_u^{\circ k} du$.

Proof. We use Theorem 1. Condition (1) therein is immediately checked, while for condition (2), we just write, using (3.4),

$$\left| \int_0^T (K_+\xi(s))^k ds \right| \leq \int_0^T |K_+\xi(s)|^k ds \leq M|\xi|_{17/12} \int_0^T C^k(s) ds. \quad \square$$

Fix an integer $k \geq 1$ and consider the following, additional, condition.

(H_3^k) The maximal function $D(t) = \sup_{0 < \varepsilon \leq \varepsilon_0} \frac{1}{\varepsilon} \int_t^{t+\varepsilon} C(s) ds$ belongs to $L^k([0, T])$ for any $T > 0$ and for some $\varepsilon_0 > 0$.

We can now state the main result of this section.

Proposition 4. Fix an integer $k \geq 1$. Let B be a Volterra process with kernel K satisfying the conditions (H_1) , (H_2) and (H_3^k) . The following convergence then holds:

$$\int_0^T \left(\frac{B_{u+\varepsilon} - B_u}{\varepsilon} \right)^{\circ k} du \xrightarrow[\varepsilon \rightarrow 0]{(S)^*} \int_0^T \dot{B}_u^{\circ k} du.$$

Proof. Fix $\xi \in \mathcal{S}(\mathbb{R})$ and set

$$S_\varepsilon(\xi) = S\left(\int_0^T \left(\frac{B_{u+\varepsilon} - B_u}{\varepsilon}\right)^{\diamond k} du\right)(\xi).$$

From linearity and property (2.3) of the S -transform, we obtain

$$S_\varepsilon(\xi) = \int_0^T \frac{(S(B_{u+\varepsilon} - B_u)(\xi))^k}{\varepsilon^k} du. \tag{3.5}$$

Equation (3.2) yields

$$S(B_{u+\varepsilon} - B_u)(\xi) = \int_0^{u+\varepsilon} K(u + \varepsilon, r)\xi(r) dr - \int_0^u K(u, r)\xi(r) dr. \tag{3.6}$$

We claim that

$$\int_0^{u+\varepsilon} K(u + \varepsilon, r)\xi(r) dr - \int_0^u K(u, r)\xi(r) dr = \int_u^{u+\varepsilon} K_+\xi(s) ds. \tag{3.7}$$

Indeed, we can write

$$\begin{aligned} \int_u^{u+\varepsilon} K_+\xi(s) ds &= \int_u^{u+\varepsilon} k'(s)\xi(s) ds + \int_u^{u+\varepsilon} \left(\int_0^s \frac{\partial K}{\partial s}(s, r)(\xi(r) - \xi(s)) dr\right) ds \\ &= A_u^{(1)} + A_u^{(2)}. \end{aligned} \tag{3.8}$$

We have, using Fubini's theorem, that

$$\begin{aligned} A_u^{(2)} &= - \int_u^{u+\varepsilon} ds \int_0^s dr \frac{\partial K}{\partial s}(s, r) \int_r^s d\theta \xi'(\theta) \\ &= - \int_0^{u+\varepsilon} d\theta \xi'(\theta) \int_0^\theta dr (K(u + \varepsilon, r) - K(\theta \vee u, r)). \end{aligned} \tag{3.9}$$

This can be rewritten as

$$\begin{aligned} A_u^{(2)} &= - \int_0^u (K(u + \varepsilon, r) - K(u, r))(\xi(u) - \xi(r)) dr \\ &\quad - \int_u^{u+\varepsilon} d\theta \xi'(\theta) \int_0^\theta dr (K(u + \varepsilon, r) - K(\theta, r)). \end{aligned} \tag{3.10}$$

On the other hand, integration by parts yields

$$\begin{aligned} A_u^{(1)} &= \xi(u + \varepsilon) \int_0^{u+\varepsilon} K(u + \varepsilon, r) dr \\ &\quad - \xi(u) \int_0^u K(u, r) dr - \int_u^{u+\varepsilon} ds \xi'(s) \int_0^s dr K(s, r). \end{aligned} \tag{3.11}$$

Therefore, adding (3.11) and (3.10) yields

$$\begin{aligned}
 A_u^{(1)} + A_u^{(2)} &= \xi(u + \varepsilon) \int_0^{u+\varepsilon} K(u + \varepsilon, r) \, dr - \xi(u) \int_0^u K(u, r) \, dr \\
 &\quad - \int_0^u (K(u + \varepsilon, r) - K(u, r)) (\xi(u) - \xi(r)) \, dr \\
 &\quad - \int_u^{u+\varepsilon} d\theta \xi'(\theta) \int_0^\theta K(u + \varepsilon, r) \, dr.
 \end{aligned} \tag{3.12}$$

Note that, by integrating by parts, we have

$$\begin{aligned}
 & - \int_u^{u+\varepsilon} d\theta \xi'(\theta) \int_0^\theta K(u + \varepsilon, r) \, dr \\
 &= -\xi(u + \varepsilon) \int_0^{u+\varepsilon} K(u + \varepsilon, r) \, dr + \xi(u) \int_0^u K(u + \varepsilon, r) \, dr \\
 &\quad + \int_u^{u+\varepsilon} K(u + \varepsilon, r) \xi(r) \, dr.
 \end{aligned} \tag{3.13}$$

Thus, substituting (3.13) into (3.12), we obtain

$$A_u^{(1)} + A_u^{(2)} = \int_0^{u+\varepsilon} K(u + \varepsilon, r) \xi(r) \, dr - \int_0^u K(u, r) \xi(r) \, dr,$$

which completes the proof of (3.7). As a consequence, from (3.5)–(3.7), we obtain

$$S_\varepsilon(\xi) = \int_0^T \left(\frac{1}{\varepsilon} \int_u^{u+\varepsilon} K_+ \xi(s) \, ds \right)^k \, du.$$

On the other hand, using (3.4) and the definition of the maximal function D , we get

$$\begin{aligned}
 \sup_{0 < \varepsilon \leq \varepsilon_0} \left| \frac{1}{\varepsilon} \int_u^{u+\varepsilon} K_+ \xi(s) \, ds \right|^k &\leq M^k |\xi|_{17/12}^k \sup_{0 < \varepsilon \leq \varepsilon_0} \left(\frac{1}{\varepsilon} \int_u^{u+\varepsilon} C(s) \, ds \right)^k \\
 &= M^k |\xi|_{17/12}^k D^k(u).
 \end{aligned} \tag{3.14}$$

Therefore, using hypothesis (H_3^k) and the dominated convergence theorem, we have

$$\lim_{\varepsilon \rightarrow 0} S_\varepsilon(\xi) = \int_0^T (K_+ \xi(s))^k \, ds. \tag{3.15}$$

Moreover, since $|S_\varepsilon(\xi)| \leq M^k |\xi|_{17/12}^k \int_0^T D^k(u) \, du$ for all $0 < \varepsilon \leq \varepsilon_0$ (see (3.14)), conditions (1) and (2) in Proposition 4 are fulfilled. Consequently, $\varepsilon^{-k} \int_0^T (B_{u+\varepsilon} - B_u)^{\circ k} \, du$ converges in (\mathcal{S}^*) as $\varepsilon \rightarrow 0$.

To complete the proof, it suffices to observe that the right-hand side of (3.15) is, by definition (see Lemma 3), the S -transform of $\int_0^T \dot{B}_s^{\circ k} ds$. □

In [14], it is proved that under some additional hypotheses, the mapping $t \rightarrow B_t$ is differentiable from $(0, \infty)$ to $(S)^*$ and that its derivative, denoted by \dot{B}_t , is a Hida distribution whose S -transform is $K_+\xi(t)$.

3.2. Bidimensional case

Let $W = (W_t)_{t \in \mathbb{R}}$ be a two-sided Brownian motion defined in the white noise probability space $(\mathcal{S}'(\mathbb{R}), \mathcal{B}, \mathbb{P})$. We can consider two independent standard Brownian motions as follows: for $t \geq 0$, we set $W_t^{(1)} = W_t$ and $W_t^{(2)} = W_{-t}$.

In this section, we consider a bidimensional process $B = (B_t^{(1)}, B_t^{(2)})_{t \geq 0}$, where $B^{(1)}$ and $B^{(2)}$ are independent Volterra processes of the form

$$B_t^{(i)} = \int_0^t K(t, s) dW_s^{(i)}, \quad t \geq 0, i = 1, 2. \tag{3.16}$$

For simplicity only, we work with the same kernel K for the two components.

First, using exactly the same lines of reasoning as in the proof of Lemma 3, we get the following result.

Lemma 5. *Let $B = (B_t^{(1)}, B_t^{(2)})_{t \geq 0}$ be given as above, with a kernel K satisfying the conditions (H_1) and (H_2) . Assume, moreover, that C defined by (3.3) belongs to $L^2([0, T])$ for any $T > 0$. We then have the following results:*

- (1) *the function $\xi \mapsto \int_0^T (\int_0^u K_+\xi(-y) dy) K_+\xi(u) du$ is the S -transform of an element of $(S)^*$, denoted by $\int_0^T B_u^{(1)} \diamond \dot{B}_u^{(2)} du$;*
- (2) *the function $\xi \mapsto \int_0^T K_+\xi(-u) K_+\xi(u) du$ is the S -transform of an element of $(S)^*$, denoted by $\int_0^T \dot{B}_u^{(1)} \diamond \dot{B}_u^{(2)} du$.*

We can now state the following result.

Proposition 6. *Let $B = (B_t^{(1)}, B_t^{(2)})_{t \geq 0}$ be given as above, with a kernel K satisfying the conditions (H_1) , (H_2) and (H_3^2) . The following convergences then hold:*

$$\begin{aligned} & \int_0^T B_u^{(1)} \frac{B_{u+\varepsilon}^{(2)} - B_u^{(2)}}{\varepsilon} du \xrightarrow[\varepsilon \rightarrow 0]{(S)^*} \int_0^T B_u^{(1)} \diamond \dot{B}_u^{(2)} du, \\ & \int_0^T \left(\int_0^u \frac{B_{v+\varepsilon}^{(1)} - B_v^{(1)}}{\varepsilon} dv \right) \frac{B_{u+\varepsilon}^{(2)} - B_u^{(2)}}{\varepsilon} du \xrightarrow[\varepsilon \rightarrow 0]{(S)^*} \int_0^T B_u^{(1)} \diamond \dot{B}_u^{(2)} du, \\ & \int_0^T \frac{B_{u+\varepsilon}^{(1)} - B_u^{(1)}}{\varepsilon} \times \frac{B_{u+\varepsilon}^{(2)} - B_u^{(2)}}{\varepsilon} du \xrightarrow[\varepsilon \rightarrow 0]{(S)^*} \int_0^T \dot{B}_u^{(1)} \diamond \dot{B}_u^{(2)} du. \end{aligned}$$

Proof. Set

$$\tilde{G}_\varepsilon = \int_0^T B_u^{(1)} \frac{B_{u+\varepsilon}^{(2)} - B_u^{(2)}}{\varepsilon} du = \int_0^T B_u^{(1)} \diamond \frac{B_{u+\varepsilon}^{(2)} - B_u^{(2)}}{\varepsilon} du.$$

From linearity and property (2.3) of the S -transform, we have

$$S(\tilde{G}_\varepsilon)(\xi) = \frac{1}{\varepsilon} \int_0^T S(B_u^{(1)})(\xi) S(B_{u+\varepsilon}^{(2)} - B_u^{(2)})(\xi) du$$

so that

$$S(\tilde{G}_\varepsilon)(\xi) = \int_0^T \left(\int_0^u K_{+\xi}(-y) dy \right) \left(\frac{1}{\varepsilon} \int_u^{u+\varepsilon} K_{+\xi}(x) dx \right) du.$$

Therefore, using (3.4) and (3.14), we can write

$$\begin{aligned} |S(\tilde{G}_\varepsilon)(\xi)| &\leq M^2 |\xi|_{17/12}^2 \int_0^T \left(\int_0^u C(t) dt \right) D(u) du \\ &\leq M^2 |\xi|_{17/12}^2 \int_0^T \left(\int_0^u D(t) dt \right) D(u) du \\ &= \frac{1}{2} M^2 |\xi|_{17/12}^2 \left(\int_0^T D(u) du \right)^2 \\ &\leq \frac{T}{2} M^2 |\xi|_{17/12}^2 \int_0^T D^2(u) du. \end{aligned}$$

Hence, by the dominated convergence theorem, we get

$$\lim_{\varepsilon \rightarrow 0} S(\tilde{G}_\varepsilon)(\xi) = \int_0^T \left(\int_0^u K_{+\xi}(-y) dy \right) K_{+\xi}(u) du. \tag{3.17}$$

The right-hand side of (3.17) is the S -transform of $\int_0^T B_u^{(1)} \diamond \dot{B}_u^{(2)} du$, due to Lemma 5. Therefore, by Theorem 2, we obtain the desired result in point (1).

The proofs of the other two convergences follow exactly the same lines of reasoning and are therefore left to the reader. □

4. Fractional Brownian motion case

4.1. One-dimensional case

Consider a (one-dimensional) fractional Brownian motion (fBm) $B = (B_t)_{t \geq 0}$ of Hurst index $H \in (0, 1)$. This means that B is a zero mean Gaussian process with covariance function

$$R_H(t, s) = E(B_t B_s) = \frac{1}{2}(t^{2H} + s^{2H} - |t - s|^{2H}).$$

It is well known that B is a Volterra process. More precisely (see [5]), B has the form (3.1) with the kernel $K(t, s) = K_H(t, s)$ given by

$$K_H(t, s) = c_H \left[\left(\frac{t(t-s)}{s} \right)^{H-1/2} - \left(H - \frac{1}{2} \right) s^{1/2-H} \int_s^t u^{H-3/2} (u-s)^{H-1/2} du \right].$$

Here, c_H is a constant depending only on H . Observe that

$$\frac{\partial K_H}{\partial t}(t, s) = c_H \left(H - \frac{1}{2} \right) (t-s)^{H-3/2} \left(\frac{s}{t} \right)^{1/2-H} \quad \text{for } t > s > 0. \tag{4.1}$$

Denote by \mathcal{E} the set of all \mathbb{R} -valued step functions defined on $[0, \infty)$. Consider the Hilbert space \mathfrak{H} obtained by closing \mathcal{E} with respect to the inner product

$$\langle \mathbf{1}_{[0,u]}, \mathbf{1}_{[0,v]} \rangle_{\mathfrak{H}} = E(B_u B_v).$$

The mapping $\mathbf{1}_{[0,t]} \mapsto B_t$ can be extended to an isometry $\varphi \mapsto B(\varphi)$ between \mathfrak{H} and the Gaussian space \mathcal{H}_1 associated with B . Also, write $\mathfrak{H}^{\otimes k}$ to indicate the k th tensor product of \mathfrak{H} . When $H > 1/2$, the inner product in the space \mathfrak{H} can be written as follows, for any $\varphi, \psi \in \mathcal{E}$:

$$\langle \varphi, \psi \rangle_{\mathfrak{H}} = H(2H - 1) \int_0^\infty \int_0^\infty \varphi(s) \psi(s') |s - s'|^{2H-2} ds ds'.$$

By approximation, this extends immediately to any $\varphi, \psi \in \mathcal{S}(\mathbb{R}) \cup \mathcal{E}$.

We will make use of the multiple integrals with respect to B (we refer to [13] for a detailed account on the properties of these integrals). For every $k \geq 1$, let \mathcal{H}_k be the k th Wiener chaos of B , that is, the closed linear subspace of $L^2(\Omega)$ generated by the random variables $\{h_k(B(\varphi)), \varphi \in \mathfrak{H}, \|\varphi\|_{\mathfrak{H}} = 1\}$, where h_k is the k th Hermite polynomial (1.2). For any $k \geq 1$, the mapping $I_k(\varphi^{\otimes k}) = h_k(B(\varphi))$ provides a linear isometry between the symmetric tensor product $\mathfrak{H}^{\otimes k}$ (equipped with the modified norm $\sqrt{k!} \|\cdot\|_{\mathfrak{H}^{\otimes k}}$) and the k th Wiener chaos \mathcal{H}_k .

Following [12], let us now introduce the Hermite random variable $Z_T^{(k)}$ mentioned in (1.5). Fix $T > 0$ and let $k \geq 1$ be an integer. The family $(\varphi_\varepsilon)_{\varepsilon > 0}$, defined by

$$\varphi_\varepsilon = \varepsilon^{-k} \int_0^T \mathbf{1}_{[u, u+\varepsilon]}^{\otimes k} du, \tag{4.2}$$

satisfies

$$\lim_{\varepsilon, \eta \rightarrow 0} \langle \varphi_\varepsilon, \varphi_\eta \rangle_{\mathfrak{H}^{\otimes k}} = H^k (2H - 1)^k \int_{[0, T]^2} |s - s'|^{(2H-2)k} ds ds' = c_{k,H} T^{(2H-2)k+2} \tag{4.3}$$

with $c_{k,H} = \frac{H^k (2H-1)^k}{(Hk-k+1)(2Hk-2k+1)}$. This implies that φ_ε converges, as ε tends to zero, to an element of $\mathfrak{H}^{\otimes k}$. The limit, denoted by $\pi_{\mathbf{1}_{[0,T]}}^k$, can be characterized as follows. For any $\xi_i \in \mathcal{S}(\mathbb{R})$,

$i = 1, \dots, k$, we have

$$\begin{aligned} & \langle \pi_{\mathbf{1}_{[0,T]}}^k, \xi_1 \otimes \dots \otimes \xi_k \rangle_{\mathfrak{H}^{\otimes k}} \\ &= \lim_{\varepsilon \rightarrow 0} \langle \varphi_\varepsilon, \xi_1 \otimes \dots \otimes \xi_k \rangle_{\mathfrak{H}^{\otimes k}} \\ &= \lim_{\varepsilon \rightarrow 0} \varepsilon^{-k} \int_0^T du \prod_{i=1}^k \langle \mathbf{1}_{[u, u+\varepsilon]}, \xi_i \rangle_{\mathfrak{H}} \\ &= \lim_{\varepsilon \rightarrow 0} \varepsilon^{-k} H^k (2H - 1)^k \int_0^T du \prod_{i=1}^k \int_u^{u+\varepsilon} ds \int_0^T dr |s - r|^{2H-2} \xi_i(r) \\ &= H^k (2H - 1)^k \int_0^T du \prod_{i=1}^k \int_0^T dr |u - r|^{2H-2} \xi_i(r). \end{aligned}$$

We define the k th Hermite random variable by $Z_T^{(k)} = I_k(\pi_{\mathbf{1}_{[0,T]}}^k)$. Note that, by using the isometry formula for multiple integrals and since $G_\varepsilon = I_k(\varphi_\varepsilon)$, the convergence (1.5) is just a corollary of our construction of $Z_T^{(k)}$. Moreover, by (4.3), we have

$$E[(Z_T^{(k)})^2] = c_{k,H} \times t^{(2H-2)k+2}.$$

We will need the following preliminary result.

Lemma 7.

- (1) The fBm B verifies the assumptions (H_1) , (H_2) and (H_3^k) if and only if $H \in (\frac{1}{2} - \frac{1}{k}, 1)$.
- (2) If $H \in (\frac{1}{2} - \frac{1}{k}, 1)$, then $\int_0^T \dot{B}_u^{\otimes k} du$ is a well-defined element of $(S)^*$ (in the sense of Lemma 3).
- (3) If we assume that $H > \frac{1}{2}$, then $\int_0^T \dot{B}_u^{\otimes k} du$ belongs to $L^2(\Omega)$ if and only if $H > 1 - \frac{1}{2k}$.

Proof. (1) Since

$$k'(t) = k'_H(t) = (H + \frac{1}{2})c_1(H)t^{H-1/2} \tag{4.4}$$

and

$$\int_0^t \left| \frac{\partial K_H}{\partial t}(t, s) \right| (t - s) ds = \left| \int_0^t \frac{\partial K_H}{\partial t}(t, s)(t - s) ds \right| = c_2(H)t^{H+1/2} \tag{4.5}$$

for some constants $c_1(H)$ and $c_2(H)$, we immediately see that assumptions (H_1) and (H_2) are satisfied for all $H \in (0, 1)$. It therefore remains to focus on assumption (H_3^k) . For all $H \in (0, 1)$, we have

$$\sup_{0 < \varepsilon \leq \varepsilon_0} \frac{1}{\varepsilon} \int_t^{t+\varepsilon} s^{H-1/2} ds \leq t^{H-1/2} \vee (t + \varepsilon_0)^{H-1/2} \tag{4.6}$$

and

$$\sup_{0 < \varepsilon \leq \varepsilon_0} \frac{1}{\varepsilon} \int_t^{t+\varepsilon} s^{H+1/2} ds \leq (t + \varepsilon_0)^{H+1/2}. \tag{4.7}$$

Consequently, since $\int_0^T t^{kH-k/2} dt$ is finite when $H > \frac{1}{2} - \frac{1}{k}$, we deduce from (4.4)–(4.7) that (H_3^k) holds in this case. Now, assume that $H \leq \frac{1}{2} - \frac{1}{k}$. Using the fact that $D(t) \geq C(t)$, we obtain

$$\int_0^T D^k(t) dt \geq \int_0^T C^k(t) dt = \left(H + \frac{1}{2}\right)^k c_1(H)^k \int_0^T t^{kH-k/2} dt = \infty.$$

Therefore, in this case, assumption (H_3^k) is not verified.

(2) This fact can be proven immediately: simply combine the previous point with Lemma 3.

(3) By definition of $\int_0^T \dot{B}_u^{\otimes k} du$ (see Lemma 3), it is equivalent to show that the distribution $\tau_{\mathbf{1}_{[0,T]}}^k$, defined via the identity $\int_0^t \dot{B}_s^{\otimes k} ds = I_k(\tau_{\mathbf{1}_{[0,t]}}^k)$, can be represented as a function belonging to $L^2([0, T]^k)$. We can write

$$\begin{aligned} \langle \tau_{\mathbf{1}_{[0,T]}}^k, \xi_1 \otimes \dots \otimes \xi_k \rangle &= \int_0^T K_+ \xi_1(s) \dots K_+ \xi_k(s) ds \\ &= \int_0^T ds \prod_{i=1}^k \int_0^s \frac{\partial K_H}{\partial s}(s, r) \xi_i(r) dr \end{aligned}$$

for any $\xi_1, \dots, \xi_k \in \mathcal{S}(\mathbb{R})$. Observe that $K_+ \xi(s) = \int_0^s \frac{\partial K_H}{\partial s}(s, r) \xi(r) dr$ because $K_H(s, s) = 0$ for $H > 1/2$. Using Fubini’s theorem, we deduce that the distribution $\tau_{\mathbf{1}_{[0,T]}}^k$ can be represented as the function

$$\tau_{\mathbf{1}_{[0,T]}}^k(x_1, \dots, x_k) = \mathbf{1}_{[0,T]^k}(x_1, \dots, x_k) \int_{\max(x_1, \dots, x_k)}^T \frac{\partial K_H}{\partial s}(s, x_1) \dots \frac{\partial K_H}{\partial s}(s, x_k) ds.$$

We then obtain

$$\begin{aligned} &\| \tau_{\mathbf{1}_{[0,T]}}^k \|_{L^2([0,T]^k)}^2 \\ &= \int_{[0,T]^k} \int_{\max(x_1, \dots, x_k)}^T \int_{\max(x_1, \dots, x_k)}^T \frac{\partial K_H}{\partial s}(s, x_1) \dots \frac{\partial K_H}{\partial s}(s, x_k) \\ &\quad \times \frac{\partial K_H}{\partial s}(r, x_1) \dots \frac{\partial K_H}{\partial s}(r, x_k) ds dr dx_1 \dots dx_k \\ &= \int_{[0,T]^2} \left(\int_0^{r \wedge s} \frac{\partial K_H}{\partial s}(s, x) \frac{\partial K_H}{\partial s}(r, x) dx \right)^k dr ds. \end{aligned}$$

Using the equality (4.1) and the same computations as in [13], page 278, we obtain, for $s < r$,

$$\int_0^s \frac{\partial K_H}{\partial s}(s, x) \frac{\partial K_H}{\partial r}(r, x) dx = H(2H - 1)(r - s)^{2H-2}. \tag{4.8}$$

Therefore,

$$\|\tau_{\mathbf{1}_{[0,T]}}^k\|_{L^2([0,T]^k)}^2 = (H(2H - 1))^k \int_0^T \int_0^T |r - s|^{2Hk-2k} dr ds.$$

We immediately check that $\|\tau_{\mathbf{1}_{[0,T]}}^k\|_{L^2([0,T]^k)}^2 < \infty$ if and only if $2Hk - 2k > -1$, that is, $H > 1 - \frac{1}{2k}$. Thus, in this case, the Hida distribution $\int_0^T \dot{B}_s^{\circ k} ds$ is a square-integrable random variable with

$$E \left[\left(\int_0^T \dot{B}_s^{\circ k} ds \right)^2 \right] = \|\tau_{\mathbf{1}_{[0,T]}}^k\|_{L^2([0,T]^k)}^2 = c_{k,H} \times T^{2Hk-2k+2}. \quad \square$$

Remark 8. According to our result, the two distributions $\tau_{\mathbf{1}_{[0,T]}}^k$ and $\pi_{\mathbf{1}_{[0,T]}}^k$ should coincide when $H > 1/2$. We can check this fact by means of elementary arguments. Let $\xi_i \in \mathcal{S}(\mathbb{R})$, $i = 1, \dots, k$. From (3.7), we deduce that

$$\langle \mathbf{1}_{[u, u+\varepsilon]}, \xi_i \rangle_{\mathcal{S}} = \int_u^{u+\varepsilon} K_+ \xi_i(s) ds$$

and then

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \langle \mathbf{1}_{[u, u+\varepsilon]}, \xi_i \rangle_{\mathcal{S}} = K_+ \xi_i(u).$$

Using (3.14) with $k = 1$ for each ξ_i and applying the dominated convergence theorem, since the fractional Brownian motion satisfies the assumption (H_3^k) when $H \in (\frac{1}{2} - \frac{1}{k}, 1)$, we get, for φ_ε defined in (4.2),

$$\lim_{\varepsilon \rightarrow 0} \langle \varphi_\varepsilon, \xi_1 \otimes \dots \otimes \xi_k \rangle_{\mathcal{S}^{\otimes k}} = \int_0^T K_+ \xi_1(u) \dots K_+ \xi_k(u) du,$$

which yields $\tau_{\mathbf{1}_{[0,T]}}^k = \pi_{\mathbf{1}_{[0,T]}}^k$.

We can now state the main result of this section.

Proposition 9. *Let $k \geq 2$ be an integer. If $H > \frac{1}{2} - \frac{1}{k}$ (note that this condition is immaterial for $k = 2$), the random variable*

$$G_\varepsilon = \int_0^T \left(\frac{B_{u+\varepsilon} - B_u}{\varepsilon} \right)^{\circ k} du = \varepsilon^{-k(1-H)} \int_0^T h_k \left(\frac{B_{u+\varepsilon} - B_u}{\varepsilon^H} \right) du$$

converges in (\mathcal{S}^*) , as $\varepsilon \rightarrow 0$, to the Hida distribution $\int_0^T \dot{B}_u^{\circ k} du$. Moreover, G_ε converges in $L^2(\Omega)$ if and only if $H > 1 - \frac{1}{2k}$. In this case, the limit is $\int_0^T \dot{B}_u^{\circ k} du = Z_T^{(k)}$.

Proof. The first point follows directly from Proposition 4 and Lemma 7 (point 1). On the other hand, we already know (see (1.5)) that G_ε converges in $L^2(\Omega)$ to $Z_T^{(k)}$ when $H > 1 - \frac{1}{2k}$. This

implies that when $H > 1 - \frac{1}{2k}$, $\int_0^T \dot{B}_s^{\diamond k} ds$ must be a square-integrable random variable equal to $Z_T^{(k)}$. Assume, now, that $H \leq 1 - \frac{1}{2k}$. From the proof of (1.3) and (1.4) below, it follows that $E(G_\varepsilon^2)$ tends to $+\infty$ as ε tends to zero, so G_ε does not converge in $L^2(\Omega)$. \square

4.2. Bidimensional case

Let $B^{(1)}$ and $B^{(2)}$ denote two independent fractional Brownian motions with (the same) Hurst index $H \in (0, 1)$, defined by the stochastic integral representation (3.16), as in Section 3.2.

By combining Lemma 7 (point 1 with $k = 2$) and Lemma 5, we have the following preliminary result.

Lemma 10. *For all $H \in (0, 1)$, the Hida distributions $\int_0^T B_u^{(1)} \diamond \dot{B}_u^{(2)} du$ and $\int_0^T \dot{B}_u^{(1)} \diamond \dot{B}_u^{(2)} du$ are well-defined elements of $(S)^*$ (in the sense of Lemma 5).*

We can now state the following result.

Proposition 11.

- (1) *For all $H \in (0, 1)$, \tilde{G}_ε defined by (1.9) converges in (S^*) , as $\varepsilon \rightarrow 0$, to the Hida distribution $\int_0^T B_u^{(1)} \diamond \dot{B}_u^{(2)} du$. Moreover, \tilde{G}_ε converges in $L^2(\Omega)$ if and only if $H \geq 1/2$.*
- (2) *For all $H \in (0, 1)$, \check{G}_ε defined by (1.10) converges in (S^*) , as $\varepsilon \rightarrow 0$, to the Hida distribution $\int_0^T B_u^{(1)} \diamond \dot{B}_u^{(2)} du$. Moreover, \check{G}_ε converges in $L^2(\Omega)$ if and only if $H > 1/4$.*
- (3) *For all $H \in (0, 1)$, \hat{G}_ε defined by (1.18) converges in (S^*) , as $\varepsilon \rightarrow 0$, to the Hida distribution $\int_0^T \dot{B}_u^{(1)} \diamond \dot{B}_u^{(2)} du$. Moreover, \hat{G}_ε converges in $L^2(\Omega)$ if and only if $H > 3/4$.*

Proof. (1) The first point follows directly from Proposition 6 and Lemma 7 (point 1 with $k = 2$). Assume that $H < 1/2$. From the proof of Theorem 12 below, it follows that $E(\tilde{G}_\varepsilon^2) \rightarrow \infty$ as ε tends to zero, so \tilde{G}_ε does not converge in $L^2(\Omega)$. Assume that $H = 1/2$. By a classical result of Russo and Vallois (see, for example, the survey [18]) and since we are, in this case, in a martingale setting, we have that \tilde{G}_ε converges in $L^2(\Omega)$ to the Itô integral $\int_0^T B_u^{(1)} dB_u^{(2)}$. Finally, assume that $H > 1/2$. For $\varepsilon, \eta > 0$, we have

$$E(\tilde{G}_\varepsilon \tilde{G}_\eta) = \frac{1}{\varepsilon \eta} \int_{[0, T]^2} \rho_{\varepsilon, \eta}(u - u') R_H(u, u') du du',$$

where

$$\rho_{\varepsilon, \eta}(x) = \frac{1}{2} [|x + \varepsilon|^{2H} + |x - \eta|^{2H} - |x|^{2H} - |x + \varepsilon - \eta|^{2H}]. \tag{4.9}$$

Note that as ε and η tend to zero, the quantity $(\varepsilon \eta)^{-1} \rho_{\varepsilon, \eta}(u - u')$ converges pointwise to (and is bounded by) $H(2H - 1)|u - u'|^{2H-2}$. Then, by the dominated convergence theorem, it follows that $E(\tilde{G}_\varepsilon \tilde{G}_\eta)$ converges to

$$H(2H - 1) \int_{[0, T]^2} |u - u'|^{2H-2} R_H(u, u') du du'$$

as $\varepsilon, \eta \rightarrow 0$, with $\int_{[0,T]^2} |u - u'|^{2H-2} |R_H(u, u')| du du' < \infty$, since $H > 1/2$. Hence, \check{G}_ε converges in $L^2(\Omega)$.

(2) The first point follows directly from Proposition 6 and Lemma 7 (point 1 with $k = 2$). Assume that $H \leq 1/4$. From the proof of Theorem 13 below, it follows that $E(\check{G}_\varepsilon^2) \rightarrow \infty$ as ε tends to zero, so \check{G}_ε does not converge in $L^2(\Omega)$. Assume that $H > 1/4$. For $\varepsilon, \eta > 0$, we have

$$E(\check{G}_\varepsilon \check{G}_\eta) = \frac{1}{\varepsilon^2 \eta^2} \int_{[0,T]^2} du du' \rho_{\varepsilon,\eta}(u - u') \int_0^u ds \int_0^{u'} ds' \rho_{\varepsilon,\eta}(s - s')$$

with $\rho_{\varepsilon,\eta}$ given by (4.9). Note that, as ε and η tend to zero, the quantity $(\varepsilon\eta)^{-1} \rho_{\varepsilon,\eta}(u - u')$ converges pointwise to $H(2H - 1)|u - u'|^{2H-2}$, whereas $(\varepsilon\eta)^{-1} \int_0^u ds \int_0^{u'} ds' \rho_{\varepsilon,\eta}(s - s')$ converges pointwise to $R_H(u, u')$. It then follows that $E(\check{G}_\varepsilon \check{G}_\eta)$ converges to

$$-\frac{H}{2}(2H - 1) \int_{[0,T]^2} |u - u'|^{4H-2} du du' + H \int_0^T u^{2H} (u^{2H-1} + (T - u)^{2H-1}) du$$

as $\varepsilon, \eta \rightarrow 0$ and each integral is finite since $H > 1/4$. Hence, \check{G}_ε converges in $L^2(\Omega)$.

(3) Once again, the first point follows from Proposition 6 and Lemma 7 (point 1 with $k = 2$). Assume that $H \leq 3/4$. From the proof of Theorem 14 below, it follows that $E(\widehat{G}_\varepsilon^2) \rightarrow \infty$ as ε tends to zero, so \widehat{G}_ε does not converge in $L^2(\Omega)$. Assume, now, that $H > 3/4$. For $\varepsilon, \eta > 0$, we have

$$E(\widehat{G}_\varepsilon \widehat{G}_\eta) = \frac{1}{\varepsilon^2 \eta^2} \int_{[0,T]^2} \rho_{\varepsilon,\eta}(u - u')^2 du du'$$

with $\rho_{\varepsilon,\eta}$ given by (4.9). Since the quantity $(\varepsilon\eta)^{-1} \rho_{\varepsilon,\eta}(u - u')$ converges pointwise to (and is bounded by) $H(2H - 1)|u - u'|^{2H-2}$, we have, by the dominated convergence theorem, that $E(\widehat{G}_\varepsilon \widehat{G}_\eta)$ converges to

$$H^2(2H - 1)^2 \int_{[0,T]^2} |u - u'|^{4H-4} du du'$$

as $\varepsilon, \eta \rightarrow 0$, with $\int_{[0,T]^2} |u - u'|^{4H-4} du du' < \infty$, since $H > 3/4$. Hence, \widehat{G}_ε converges in $L^2(\Omega)$. □

5. Proof of the convergences (1.3) and (1.4)

In this section, we provide a new proof of these convergences by means of a recent criterion for the weak convergence of sequences of multiple stochastic integrals established in [15] and [17]. We refer to [9] for a proof in the case of more general Gaussian processes, using different kind of tools.

Let us first recall the aforementioned criterion. We continue to use the notation introduced in Section 4.1. Also, let $\{e_i, i \geq 1\}$ denote a complete orthonormal system in \mathfrak{H} . Given $f \in \mathfrak{H}^{\otimes k}$

and $g \in \mathfrak{H}^{\odot l}$, for every $r = 0, \dots, k \wedge l$, the contraction of f and g of order r is the element of $\mathfrak{H}^{\otimes(k+l-2r)}$ defined by

$$f \otimes_r g = \sum_{i_1, \dots, i_r=1}^{\infty} \langle f, e_{i_1} \otimes \dots \otimes e_{i_r} \rangle_{\mathfrak{H}^{\otimes r}} \otimes \langle g, e_{i_1} \otimes \dots \otimes e_{i_r} \rangle_{\mathfrak{H}^{\otimes r}}.$$

(Note that $f \otimes_0 g = f \otimes g$ equals the tensor product of f and g while, for $k = l$, $f \otimes_k g = \langle f, g \rangle_{\mathfrak{H}^{\otimes k}}$.) Fix $k \geq 2$ and let $(F_\varepsilon)_{\varepsilon>0}$ be a family of the form $F_\varepsilon = I_k(\phi_\varepsilon)$ for some $\phi_\varepsilon \in \mathfrak{H}^{\odot k}$. Assume that the variance of F_ε converges as $\varepsilon \rightarrow 0$ (to σ^2 , say). The criterion of Nualart and Peccati [15] asserts that $F_\varepsilon \xrightarrow{\text{Law}} N \sim \mathcal{N}(0, \sigma^2)$ if and only if $\|\phi_\varepsilon \otimes_r \phi_\varepsilon\|_{\mathfrak{H}^{\otimes(2k-2r)}} \rightarrow 0$ for any $r = 1, \dots, k-1$. In this case, due to the result proved by Peccati and Tudor [17], we automatically have that

$$(B_{t_1}, \dots, B_{t_k}, F_\varepsilon) \xrightarrow{\text{Law}} (B_{t_1}, \dots, B_{t_k}, N)$$

for all $t_k > \dots > t_1 > 0$, with $N \sim \mathcal{N}(0, \sigma^2)$ independent of B .

For $x \in \mathbb{R}$, set

$$\rho(x) = \frac{1}{2}(|x + 1|^{2H} + |x - 1|^{2H} - 2|x|^{2H}), \tag{5.1}$$

and note that $\rho(u - v) = E[(B_{u+1} - B_u)(B_{v+1} - B_v)]$ for all $u, v \geq 0$ and that $\int_{\mathbb{R}} |\rho(x)|^k dx$ is finite if and only if $H < 1 - \frac{1}{2k}$ (since $\rho(x) \sim H(2H - 1)|x|^{2H-2}$ as $|x| \rightarrow \infty$).

We now proceed with the proof of (1.3). The proof of (1.4) would follow similar arguments.

Proof of (1.3). Because $\varepsilon^{k(1-H)-1/2}G_\varepsilon$ can be expressed as a k th multiple Wiener integral, we can use the criterion of Nualart and Peccati. By the scaling property of the fBm, it is actually equivalent to considering the family of random variables $(F_\varepsilon)_{\varepsilon>0}$, where

$$F_\varepsilon = \sqrt{\varepsilon} \int_0^{T/\varepsilon} h_k(B_{u+1} - B_u) du.$$

Step 1. Convergence of the variance. We can write

$$\begin{aligned} E(F_\varepsilon^2) &= \varepsilon k! \int_0^{T/\varepsilon} du \int_0^{T/\varepsilon} ds \rho(u - s)^k \\ &= \varepsilon k! \int_{-T/\varepsilon}^{T/\varepsilon} \rho(x)^k (T/\varepsilon - |x|) dx, \end{aligned}$$

where the function ρ is defined in (5.1). Therefore, by the dominated convergence theorem,

$$\lim_{\varepsilon \downarrow 0} E(F_\varepsilon^2) = Tk! \int_{\mathbb{R}} \rho(x)^k dx.$$

Step 2. Convergence of the contractions. Observe that the random variable $h_k(B_{u+1} - B_u)$ coincides with the multiple stochastic integral $I_k(\mathbf{1}_{[u, u+1]}^{\otimes k})$. Therefore, $F_\varepsilon = I_k(\phi_\varepsilon)$, where $\phi_\varepsilon =$

$\sqrt{\varepsilon} \int_0^{T/\varepsilon} \mathbf{1}_{[u, u+1]}^{\otimes k} du$. Let $r \in \{1, \dots, k-1\}$. We have

$$\phi_\varepsilon \otimes_r \phi_\varepsilon = \varepsilon \int_0^{T/\varepsilon} \int_0^{T/\varepsilon} (\mathbf{1}_{[u, u+1]}^{\otimes(k-r)} \otimes \mathbf{1}_{[s, s+1]}^{\otimes(k-r)}) \rho(u-s)^r du ds.$$

As a consequence, $\|\phi_\varepsilon \otimes_r \phi_\varepsilon\|_{\mathfrak{H}^{\otimes(2k-2r)}}^2$ equals

$$\varepsilon^2 \int_{[0, T/\varepsilon]^4} \rho(u-s)^r \rho(u'-s')^r \rho(u-u')^{k-r} \rho(s-s')^{k-r} ds ds' du du'.$$

Making the changes of variables $x = u - s$, $y = u' - s'$ and $z = u - u'$, we obtain that $\|\phi_\varepsilon \otimes_r \phi_\varepsilon\|_{\mathfrak{H}^{\otimes(2k-2r)}}^2$ is less than

$$A_\varepsilon = \varepsilon \int_{D_\varepsilon} |\rho(x)|^r |\rho(y)|^r |\rho(z)|^{k-r} |\rho(y+z-x)|^{k-r} dx dy dz,$$

where $D_\varepsilon = [-T/\varepsilon, T/\varepsilon]^3$. Consider the decomposition

$$\begin{aligned} A_\varepsilon &= \varepsilon \int_{D_\varepsilon \cap \{|x| \vee |y| \vee |z| \leq K\}} |\rho(x)|^r |\rho(y)|^r |\rho(z)|^{k-r} |\rho(y+z-x)|^{k-r} dx dy dz \\ &\quad + \varepsilon \int_{D_\varepsilon \cap \{|x| \vee |y| \vee |z| > K\}} |\rho(x)|^r |\rho(y)|^r |\rho(z)|^{k-r} |\rho(y+z-x)|^{k-r} dx dy dz \\ &= B_{\varepsilon, K} + C_{\varepsilon, K}. \end{aligned}$$

Clearly, for any fixed $K > 0$, the term $B_{\varepsilon, K}$ tends to zero because ρ is a bounded function. On the other hand, we have

$$D_\varepsilon \cap \{|x| \vee |y| \vee |z| > K\} \subset D_{\varepsilon, K, x} \cup D_{\varepsilon, K, y} \cup D_{\varepsilon, K, z},$$

where $D_{\varepsilon, K, x} = \{|x| > K\} \cap \{|y| \leq T/\varepsilon\} \cap \{|z| \leq T/\varepsilon\}$ ($D_{\varepsilon, K, y}$ and $D_{\varepsilon, K, z}$ being defined similarly). Set

$$C_{\varepsilon, K, x} = \varepsilon \int_{D_{\varepsilon, K, x}} |\rho(x)|^r |\rho(y)|^r |\rho(z)|^{k-r} |\rho(y+z-x)|^{k-r} dx dy dz.$$

By Hölder's inequality, we have

$$\begin{aligned} C_{\varepsilon, K, x} &\leq \varepsilon \left(\int_{D_{\varepsilon, K, x}} |\rho(x)|^k |\rho(y)|^k dx dy dz \right)^{r/k} \\ &\quad \times \left(\int_{D_{\varepsilon, K, x}} |\rho(z)|^k |\rho(y+z-x)|^k dx dy dz \right)^{1-r/k} \\ &\leq 2T \left(\int_{\mathbb{R}} |\rho(t)|^k dt \right)^{2-r/k} \left(\int_{|x|>K} |\rho(x)|^k dx \right)^{r/k} \xrightarrow{K \rightarrow \infty} 0. \end{aligned}$$

Similarly, we prove that $C_{\varepsilon, K, y} \rightarrow 0$ and $C_{\varepsilon, K, z} \rightarrow 0$ as $K \rightarrow \infty$. Finally, it suffices to choose K large enough in order to get the desired result, that is, $\|\phi_\varepsilon \otimes_r \phi_\varepsilon\|_{\mathfrak{H}^{\otimes(2k-2r)}} \rightarrow 0$ as $\varepsilon \rightarrow 0$.

Step 3. Proof of the first point. By step 1, the family

$$((B_t)_{t \in [0, T]}, \varepsilon^{1/2-2H} G_\varepsilon)$$

is tight in $C([0, T]) \times \mathbb{R}$. By step 2, we also have the convergence of the finite-dimensional distributions, as a by-product of the criteria of Nualart and Peccati [15] and Peccati and Tudor [17] (see the preliminaries at the beginning of this section). Hence, the proof of the first point is complete. □

6. Convergences in law for some functionals related to the Lévy area of the fractional Brownian motion

Let $B^{(1)}$ and $B^{(2)}$ denote two independent fractional Brownian motions with Hurst index $H \in (0, 1)$. Recall the definition (1.9) of \tilde{G}_ε :

$$\tilde{G}_\varepsilon = \int_0^T B_u^{(1)} \frac{B_{u+\varepsilon}^{(2)} - B_u^{(2)}}{\varepsilon} du.$$

Theorem 12. *Convergence in law (1.15) holds.*

Proof. We fix $H < 1/2$. The proof is divided into several steps.

Step 1. Computing the variance of $\varepsilon^{1/2-H} \tilde{G}_\varepsilon$.

By using the scaling properties of the fBm, first observe that $\varepsilon^{1/2-H} \tilde{G}_\varepsilon$ has the same law as

$$\tilde{F}_\varepsilon = \varepsilon^{1/2+H} \int_0^{T/\varepsilon} B_u^{(1)} (B_{u+1}^{(2)} - B_u^{(2)}) du. \tag{6.1}$$

For $\rho(x) = \frac{1}{2}(|x+1|^{2H} + |x-1|^{2H} - 2|x|^{2H})$, we have

$$\begin{aligned} E(\tilde{F}_\varepsilon^2) &= \varepsilon^{1+2H} \int_0^{T/\varepsilon} du \int_0^{T/\varepsilon} ds R_H(u, s) \rho(u-s) \\ &= \alpha_\varepsilon - \beta_\varepsilon, \end{aligned}$$

where

$$\begin{aligned} \alpha_\varepsilon &= \varepsilon^{1+2H} \int_0^{T/\varepsilon} du u^{2H} \int_0^{T/\varepsilon} ds \rho(u-s), \\ \beta_\varepsilon &= \varepsilon^{1+2H} \int_0^{T/\varepsilon} du \int_0^u ds (u-s)^{2H} \rho(u-s). \end{aligned}$$

Let us first study β_ε . We can write

$$\beta_\varepsilon = \varepsilon^{2H} \int_0^{T/\varepsilon} x^{2H} \rho(x)(T - \varepsilon x) dx.$$

The integral $\int_0^\infty x^{2H} \rho(x) dx$ is convergent for $H < 1/4$, while $\int_0^{T/\varepsilon} x^{2H} \rho(x) dx$ diverges as $-\frac{1}{8} \log(1/\varepsilon)$ for $H = 1/4$ and as $H(2H - 1)T^{4H-1}\varepsilon^{1-4H}$ for $1/4 < H < 1/2$. The integral $\int_0^{T/\varepsilon} x^{2H+1} \rho(x) dx$ diverges as $H(2H - 1)T^{4H}\varepsilon^{-4H}$. Therefore,

$$\lim_{\varepsilon \rightarrow 0} \beta_\varepsilon = 0.$$

Second, let us write α_ε as

$$\begin{aligned} \alpha_\varepsilon &= \varepsilon^{1+2H} \int_0^{T/\varepsilon} du u^{2H} \int_0^{T/\varepsilon} ds \rho(u - s) \\ &= \varepsilon^{1+2H} \left(\int_0^{T/\varepsilon} du u^{2H} \int_0^u ds \rho(u - s) + \int_0^{T/\varepsilon} du u^{2H} \int_u^{T/\varepsilon} ds \rho(u - s) \right) \\ &= \frac{1}{2H + 1} \int_0^{T/\varepsilon} \rho(x)(T^{2H+1} - (\varepsilon x)^{2H+1} + (T - \varepsilon x)^{2H+1}) dx. \end{aligned}$$

Hence, by the dominated convergence theorem, we have

$$\lim_{\varepsilon \rightarrow 0} \alpha_\varepsilon = \frac{2T^{2H+1}}{2H + 1} \int_0^\infty \rho(x) dx$$

so that

$$\lim_{\varepsilon \rightarrow 0} \varepsilon^{1-2H} E[\tilde{G}_\varepsilon^2] = \lim_{\varepsilon \rightarrow 0} E[\tilde{F}_\varepsilon^2] = \frac{2T^{2H+1}}{2H + 1} \int_0^\infty \rho(x) dx.$$

Step 2. Showing the convergence in law in (1.15).

By the previous step, the distributions of the family

$$\left((B_t^{(1)}, B_t^{(2)})_{t \in [0, T]}, \varepsilon^{1/2-H} \tilde{G}_\varepsilon \right)_{\varepsilon > 0}$$

are tight in $C([0, T]^2) \times \mathbb{R}$ and it suffices to show the convergence of the finite-dimensional distributions. We need to show that for any $\lambda \in \mathbb{R}$, any $0 < t_1 \leq \dots \leq t_k$, any $\theta_1, \dots, \theta_k \in \mathbb{R}$ and any $\mu_1, \dots, \mu_k \in \mathbb{R}$, we have

$$\begin{aligned} \lim_{\varepsilon \downarrow 0} E \left[e^{i \sum_{j=1}^k \theta_j B_{t_j}^{(1)}} e^{i \sum_{j=1}^k \mu_j B_{t_j}^{(2)}} e^{i \lambda \varepsilon^{1/2-H} \tilde{G}_\varepsilon} \right] \\ = E \left[e^{-(1/2) \text{Var}(\sum_{j=1}^k \mu_j B_{t_j}^{(2)})} \right] E \left[e^{i \sum_{j=1}^k \theta_j B_{t_j}^{(1)}} e^{-\lambda^2 S^2 / 2} \right], \end{aligned} \tag{6.2}$$

where $S = \sqrt{2 \int_0^\infty \rho(x) dx \int_0^T (B_u^{(1)})^2 du}$. We can write

$$\begin{aligned} & E\left[e^{i \sum_{j=1}^k \theta_j B_{t_j}^{(1)}} e^{i \sum_{j=1}^k \mu_j B_{t_j}^{(2)}} e^{i \lambda \varepsilon^{1/2-H} \tilde{G}_\varepsilon} \right] \\ &= E\left[e^{i \sum_{j=1}^k \theta_j B_{t_j}^{(1)}} E\left[e^{i \sum_{j=1}^k \mu_j B_{t_j}^{(2)}} e^{i \lambda \varepsilon^{1/2-H} \tilde{G}_\varepsilon} \mid \mathcal{B}^{(1)} \right] \right] \\ &= E\left[e^{i \sum_{j=1}^k \theta_j B_{t_j}^{(1)}} e^{-\lambda \varepsilon^{1/2-H} \sum_{j=1}^k \mu_j \int_0^T B_u^{(1)} E(B_{t_j}^{(2)} \times (B_{u+\varepsilon}^{(2)} - B_u^{(2)})/\varepsilon) du} \right. \\ &\quad \left. \times e^{-(\lambda^2/2)\varepsilon^{1-2H} \int_{[0,T]^2} B_u^{(1)} B_v^{(1)} \rho_\varepsilon(u-v) du dv} e^{-(1/2) \text{Var}(\sum_{j=1}^k \mu_j B_{t_j}^{(2)})} \right] \end{aligned}$$

with $\rho_\varepsilon(x) = \frac{1}{2}(|x + \varepsilon|^{2H} + |x - \varepsilon|^{2H} - 2|x|^{2H})$. Observe that

$$\int_{[0,T]^2} B_u^{(1)} B_v^{(1)} \rho_\varepsilon(u - v) du dv \geq 0$$

since $\rho_\varepsilon(u - v) = E[(B_{u+\varepsilon}^{(2)} - B_u^{(2)})(B_{v+\varepsilon}^{(2)} - B_v^{(2)})]$ is a covariance function. Moreover, for any fixed $t \geq 0$, we have

$$\begin{aligned} & \int_0^T B_u^{(1)} E\left(B_t^{(2)} \times \frac{B_{u+\varepsilon}^{(2)} - B_u^{(2)}}{\varepsilon} \right) du \\ &= \frac{1}{2} \int_0^T B_u^{(1)} \left(\frac{(u + \varepsilon)^{2H} - u^{2H}}{\varepsilon} + \frac{|t - u|^{2H} - |t - u - \varepsilon|^{2H}}{\varepsilon} \right) du \\ &\xrightarrow[\varepsilon \rightarrow 0]{\text{a.s.}} H \int_0^T B_u^{(1)} (u^{2H-1} - |t - u|^{2H-1}) du. \end{aligned}$$

Since $H < 1/2$, this implies that

$$e^{-\lambda \varepsilon^{1/2-H} \sum_{j=1}^k \mu_j \int_0^T B_u^{(1)} E(B_{t_j}^{(2)} \times (B_{u+\varepsilon}^{(2)} - B_u^{(2)})/\varepsilon) du} \xrightarrow[\varepsilon \rightarrow 0]{\text{a.s.}} 1.$$

Hence, to get (6.2), it suffices to show that

$$E\left[e^{i \sum_{j=1}^k \theta_j B_{t_j}^{(1)}} e^{-(\lambda^2/2)\varepsilon^{1-2H} \int_{[0,T]^2} B_u^{(1)} B_v^{(1)} \rho_\varepsilon(u-v) du dv} \right] \xrightarrow[\varepsilon \rightarrow 0]{} E\left[e^{i \sum_{j=1}^k \theta_j B_{t_j}^{(1)}} e^{-(\lambda^2/2)S^2} \right]. \tag{6.3}$$

We have

$$\begin{aligned} C_\varepsilon &:= E\left[\exp\left(i \sum_{j=1}^k \theta_j B_{t_j}^{(1)} - \frac{\lambda^2}{2} \varepsilon^{1-2H} \int_{[0,T]^2} B_u^{(1)} B_v^{(1)} \rho_\varepsilon(u - v) du dv \right) \right] \\ &= E\left[\exp\left(i \sum_{j=1}^k \theta_j B_{t_j}^{(1)} - \lambda^2 \varepsilon^{1-2H} \int_0^T B_u^{(1)} \left(\int_0^u B_{u-x}^{(1)} \rho_\varepsilon(x) dx \right) du \right) \right] \end{aligned}$$

$$\begin{aligned}
 &= E \left[\exp \left(i \sum_{j=1}^k \theta_j B_{t_j}^{(1)} - \lambda^2 \varepsilon^{1-2H} \int_0^T \rho_\varepsilon(x) \left(\int_x^T B_u^{(1)} B_{u-x}^{(1)} du \right) dx \right) \right] \\
 &= E \left[\exp \left(i \sum_{j=1}^k \theta_j B_{t_j}^{(1)} - \lambda^2 \int_0^{T/\varepsilon} \rho(x) \left(\int_{\varepsilon x}^T B_u^{(1)} B_{u-\varepsilon x}^{(1)} du \right) dx \right) \right],
 \end{aligned}$$

the last inequality following from the relation $\rho_\varepsilon(x) = \varepsilon^{2H} \rho(x/\varepsilon)$. By the dominated convergence theorem, we obtain

$$\begin{aligned}
 C_\varepsilon &\xrightarrow{\varepsilon \rightarrow 0} E \left[\exp \left(i \sum_{j=1}^k \theta_j B_{t_j}^{(1)} - \lambda^2 \int_0^\infty \rho(x) dx \times \int_0^T (B_u^{(1)})^2 du \right) \right] \\
 &= E \left[\exp \left(i \sum_{j=1}^k \theta_j B_{t_j}^{(1)} - \frac{\lambda^2}{2} S^2 \right) \right],
 \end{aligned}$$

that is, (6.3). The proof of the theorem is thus completed. □

Recall the definition (5.1) of ρ and the definition of \check{G}_ε :

$$\check{G}_\varepsilon = \int_0^T \left(\int_0^u \frac{B_{v+\varepsilon}^{(1)} - B_v^{(1)}}{\varepsilon} dv \right) \frac{B_{u+\varepsilon}^{(2)} - B_u^{(2)}}{\varepsilon} du.$$

Theorem 13. *Convergences in law (1.11) and (1.12) hold.*

Proof. We only show the first convergence, the proof of the second one being very similar. By using the scaling properties of the fBm, first observe that $\varepsilon^{1/2-2H} \check{G}_\varepsilon$ has the same law as

$$\check{F}_\varepsilon = \sqrt{\varepsilon} \int_0^{T/\varepsilon} \left(\int_0^u (B_{v+1}^{(1)} - B_v^{(1)}) dv \right) (B_{u+1}^{(2)} - B_u^{(2)}) du.$$

We now fix $H < 1/4$ and the proof is divided into several steps.

Step 1. Computing the variance of \check{F}_ε . We can write

$$E(\check{F}_\varepsilon^2) = \varepsilon \int_{[0, T/\varepsilon]^2} du du' \rho(u - u') \int_0^u dv \int_0^{u'} dv' \rho(v - v')$$

with $\rho(x) = \frac{1}{2}(|x + 1|^{2H} + |x - 1|^{2H} - 2|x|^{2H})$. We have

$$\int_0^u dv \int_0^{u'} dv' \rho(v - v') = \frac{\Psi(u - u') - \Psi(u) - \Psi(u') + 2}{2(2H + 1)(2H + 2)},$$

where

$$\Psi(x) = 2|x|^{2H+2} - |x + 1|^{2H+2} - |x - 1|^{2H+2}. \tag{6.4}$$

Consider first the contribution of the term $\Psi(u - u')$. We have

$$\lim_{\varepsilon \rightarrow 0} \varepsilon \int_{[0, T/\varepsilon]^2} \rho(u - u') \Psi(u - u') \, du \, du' = T \int_{\mathbb{R}} \rho(x) \Psi(x) \, dx.$$

Note that $\rho(x) \sim H(2H - 1)|x|^{2H-2}$ and $\Psi(x) \sim -(2H + 2)(2H + 1)|x|^{2H}$ as $|x| \rightarrow \infty$ so that $\int_{\mathbb{R}} |\rho(x) \Psi(x)| \, dx < \infty$ because $H < 1/4$. On the other hand, we have

$$\varepsilon \int_{[0, T/\varepsilon]^2} \rho(u - u') \Psi(u) \, du \, du' = \varepsilon \int_0^{T/\varepsilon} \, du \, \Psi(u) \int_{u-T/\varepsilon}^u \, dx \, \rho(x)$$

and this converges to zero as $\varepsilon \rightarrow 0$. Indeed, since $\rho(x) \sim H(2H - 1)x^{2H-2}$ as $x \rightarrow \infty$, we have $\int_u^\infty \rho(x) \, dx \sim Hu^{2H-1}$ as $u \rightarrow \infty$; hence, since $\int_{\mathbb{R}} \rho(x) \, dx = 0$, $H < 1/4$ and $\Psi(u) \sim -(2H + 2)(2H + 1)u^{2H}$ as $u \rightarrow \infty$, we have

$$\lim_{u \rightarrow \infty} \Psi(u) \int_{-\infty}^u \rho(x) \, dx = - \lim_{u \rightarrow \infty} \Psi(u) \int_u^\infty \rho(x) \, dx = 0.$$

Also, we have

$$\lim_{\varepsilon \rightarrow 0} \varepsilon \int_{[0, T/\varepsilon]^2} \rho(u - u') \, du \, du' = \int_{\mathbb{R}} \rho(x) \, dx = 0.$$

Therefore, $\lim_{\varepsilon \rightarrow 0} E(\check{F}_\varepsilon^2) = \check{\sigma}_H^2$.

Step 2. Showing the convergence in law (1.11). We first remark that by step 1, the laws of the family $((B_t^{(1)}, B_t^{(2)})_{t \in [0, T]}, \varepsilon^{1/2-2H} \check{G}_\varepsilon)_{\varepsilon > 0}$ are tight. Therefore, we only have to prove the convergence of the finite-dimensional laws. Moreover, by the main result of Peccati and Tudor [17], it suffices to prove that

$$\varepsilon^{1/2-2H} \check{G}_\varepsilon \stackrel{\text{Law}}{=} \check{F}_\varepsilon \xrightarrow{\text{Law}} \mathcal{N}(0, T\check{\sigma}_H^2) \quad \text{as } \varepsilon \rightarrow 0. \tag{6.5}$$

We have

$$E(e^{i\lambda \check{F}_\varepsilon}) = E\left(\exp\left\{-\frac{\lambda^2 \varepsilon}{2} \int_{[0, T/\varepsilon]^2} (B_{u+1}^{(2)} - B_u^{(2)})(B_{u'+1}^{(2)} - B_{u'}^{(2)}) \times \left(\int_0^u \int_0^{u'} \rho(v - v') \, dv \, dv'\right) \, du \, du'\right\}\right).$$

Since $\rho(v - v') = E[(B_{v+1}^{(1)} - B_v^{(1)})(B_{v'+1}^{(1)} - B_{v'}^{(1)})]$ is a covariance function, observe that the quantity inside the exponential in the right-hand side of the previous identity is negative. Hence, since $x \mapsto \exp(-\frac{\lambda^2}{2}x_+)$ is continuous and bounded by 1 on \mathbb{R} , (6.5) will be a consequence of the convergence

$$A_\varepsilon \stackrel{\text{law}}{\longrightarrow} T\check{\sigma}_H^2 \quad \text{as } \varepsilon \rightarrow 0 \tag{6.6}$$

with

$$A_\varepsilon := \varepsilon \int_{[0, T/\varepsilon]^2} (B_{u+1} - B_u)(B_{u'+1} - B_{u'}) \left(\int_0^u \int_0^{u'} \rho(v - v') \, dv \, dv' \right) \, du \, du',$$

B denoting a fractional Brownian motion with Hurst index H . The proof of (6.6) will be achieved by showing that the expectation (resp., the variance) of A_ε tends to $T\check{\sigma}_H^2$ (resp., zero). By step 1, observe that

$$E(A_\varepsilon) = E(\check{F}_\varepsilon^2) \rightarrow T\check{\sigma}_H^2$$

as $\varepsilon \rightarrow 0$. We now want to show that the variance of A_ε converges to zero. Performing the changes of variables $s = u\varepsilon$ and $t = u'\varepsilon$ yields

$$A_\varepsilon = \varepsilon^{-1} \int_{[0, T]^2} (B_{s/\varepsilon+1} - B_{s/\varepsilon})(B_{t/\varepsilon+1} - B_{t/\varepsilon}) \left(\int_0^{s/\varepsilon} \int_0^{t/\varepsilon} \rho(v - v') \, dv \, dv' \right) \, ds \, dt,$$

which has the same distribution as

$$\begin{aligned} C_\varepsilon &= \varepsilon^{-1-2H} \int_{[0, T]^2} (B_{s+\varepsilon} - B_s)(B_{t+\varepsilon} - B_t) \left(\int_0^{s/\varepsilon} \int_0^{t/\varepsilon} \rho(u - u') \, du \, du' \right) \, ds \, dt \\ &= \varepsilon^{-1-2H} \int_{[0, T]^2} (B_{s+\varepsilon} - B_s)(B_{t+\varepsilon} - B_t) \Lambda_\varepsilon(s, t) \, ds \, dt, \end{aligned}$$

where $\Lambda_\varepsilon(s, t) = \int_0^{s/\varepsilon} \int_0^{t/\varepsilon} \rho(u - u') \, du \, du'$. This can be written as

$$C_\varepsilon = \varepsilon^{-1-2H} \int_{\mathbb{R}^2} B_s B_t \Sigma_\varepsilon(s, t) \, ds \, dt,$$

where

$$\begin{aligned} \Sigma_\varepsilon(s, t) &= \mathbf{1}_{[\varepsilon, T+\varepsilon]}(s) \mathbf{1}_{[\varepsilon, T+\varepsilon]}(t) \Lambda_\varepsilon(s - \varepsilon, t - \varepsilon) - \mathbf{1}_{[0, T]}(s) \mathbf{1}_{[\varepsilon, T+\varepsilon]}(t) \Lambda_\varepsilon(s, t - \varepsilon) \\ &\quad - \mathbf{1}_{[\varepsilon, T+\varepsilon]}(s) \mathbf{1}_{[0, T]}(t) \Lambda_\varepsilon(s - \varepsilon, t) + \mathbf{1}_{[0, T]}(s) \mathbf{1}_{[0, T]}(t) \Lambda_\varepsilon(s, t). \end{aligned} \tag{6.7}$$

Moreover,

$$C_\varepsilon - E(C_\varepsilon) = \varepsilon^{-1-2H} I_2 \left(\int_{\mathbb{R}^2} \mathbf{1}_{[0, s]} \otimes \mathbf{1}_{[0, t]} \Sigma_\varepsilon(s, t) \, ds \, dt \right),$$

where I_2 is the double stochastic integral with respect to B . Therefore,

$$\begin{aligned} \text{Var}(C_\varepsilon) &= 2\varepsilon^{-2-4H} \left\| \int_{\mathbb{R}^2} \mathbf{1}_{[0, s]} \otimes \mathbf{1}_{[0, t]} \Sigma_\varepsilon(s, t) \, ds \, dt \right\|_{\mathfrak{H} \otimes \mathfrak{H}}^2 \\ &= 2\varepsilon^{-2-4H} \int_{\mathbb{R}^4} R_H(s, s') R_H(t, t') \Sigma_\varepsilon(s, t) \Sigma_\varepsilon(s', t') \, ds \, dt \, ds' \, dt'. \end{aligned}$$

Taking into account that the partial derivatives $\frac{\partial R_H}{\partial s}$ and $\frac{\partial R_H}{\partial t}$ are integrable, we can write

$$\begin{aligned} \text{Var}(C_\varepsilon) &= 2\varepsilon^{-2-4H} \int_{\mathbb{R}^4} \left(\int_0^s \frac{\partial R_H}{\partial \sigma}(\sigma, s') d\sigma \right) \left(\int_0^{t'} \frac{\partial R_H}{\partial \tau}(t, \tau) d\tau \right) \\ &\quad \times \Sigma_\varepsilon(s, t) \Sigma_\varepsilon(s', t') ds dt ds' dt'. \end{aligned}$$

Hence, by integrating by parts, we get

$$\begin{aligned} \text{Var}(C_\varepsilon) &= 2\varepsilon^{-2-4H} \int_{\mathbb{R}^4} \frac{\partial R_H}{\partial s}(s, s') \frac{\partial R_H}{\partial t'}(t, t') \\ &\quad \times \left(\int_0^s \Sigma_\varepsilon(\sigma, t) d\sigma \right) \left(\int_0^{t'} \Sigma_\varepsilon(s', \tau) d\tau \right) ds dt ds' dt'. \end{aligned}$$

From (6.7), we obtain

$$\int_0^s \Sigma_\varepsilon(\sigma, t) d\sigma = \mathbf{1}_{[0, T]}(s) (\mathbf{1}_{[0, \varepsilon]}(t) - \mathbf{1}_{[T, T+\varepsilon]}(t)) \int_{s-\varepsilon}^s \Lambda_\varepsilon(\sigma, t - \varepsilon) d\sigma.$$

In the same way,

$$\int_0^{t'} \Sigma_\varepsilon(s', \tau) d\tau = \mathbf{1}_{[0, T]}(t') (\mathbf{1}_{[0, \varepsilon]}(s') - \mathbf{1}_{[T, T+\varepsilon]}(s')) \int_{t'-\varepsilon}^{t'} \Lambda_\varepsilon(s' - \varepsilon, \tau) d\tau.$$

As a consequence,

$$\begin{aligned} \text{Var}(C_\varepsilon) &= 2\varepsilon^{-2-4H} \int_{\mathbb{R}^4} \frac{\partial R_H}{\partial s}(s, s') \frac{\partial R_H}{\partial t'}(t, t') \left(\int_{s-\varepsilon}^s \Lambda_\varepsilon(\sigma, t - \varepsilon) d\sigma \right) \\ &\quad \times \left(\int_{t'-\varepsilon}^{t'} \Lambda_\varepsilon(s' - \varepsilon, \tau) d\tau \right) \mathbf{1}_{[0, T]}(s) (\mathbf{1}_{[0, \varepsilon]}(t) - \mathbf{1}_{[T, T+\varepsilon]}(t)) \\ &\quad \times \mathbf{1}_{[0, T]}(t') (\mathbf{1}_{[0, \varepsilon]}(s') - \mathbf{1}_{[T, T+\varepsilon]}(s')) ds dt ds' dt' = \sum_{i=1}^4 H_\varepsilon^i, \end{aligned}$$

where

$$\begin{aligned} H_\varepsilon^1 &= \int_0^T \int_0^\varepsilon \int_0^\varepsilon \int_0^T G_\varepsilon(s, t, s', t') ds dt ds' dt', \\ H_\varepsilon^2 &= - \int_0^T \int_T^{T+\varepsilon} \int_0^\varepsilon \int_0^T G_\varepsilon(s, t, s', t') ds dt ds' dt', \\ H_\varepsilon^3 &= - \int_0^T \int_0^\varepsilon \int_T^{T+\varepsilon} \int_0^T G_\varepsilon(s, t, s', t') ds dt ds' dt', \\ H_\varepsilon^4 &= \int_0^T \int_0^{T+\varepsilon} \int_0^{T+\varepsilon} \int_0^T G_\varepsilon(s, t, s', t') ds dt ds' dt' \end{aligned}$$

and

$$G_\varepsilon(s, t, s', t') = 2\varepsilon^{-2-4H} \frac{\partial R_H}{\partial s}(s, s') \frac{\partial R_H}{\partial t'}(t, t') \times \left(\int_{s-\varepsilon}^s \Lambda_\varepsilon(\sigma, t - \varepsilon) d\sigma \right) \left(\int_{t'-\varepsilon}^{t'} \Lambda_\varepsilon(s' - \varepsilon, \tau) d\tau \right).$$

We only consider the term H_ε^1 because the others can be handled in the same way. We have, with Ψ given by (6.4),

$$\Lambda_\varepsilon(s, t) = \int_0^{s/\varepsilon} \int_0^{t/\varepsilon} \rho(u - u') du du' = \frac{\Psi((s-t)/\varepsilon) - \Psi(s/\varepsilon) - \Psi(t/\varepsilon) + 2}{2(2H+1)(2H+2)}.$$

Note that

$$\left| \Psi\left(\frac{s-t}{\varepsilon}\right) \right| \leq \varepsilon^{-2H-2} |2|s-t|^{2H+2} - |s-t+\varepsilon|^{2H+2} - |s-t-\varepsilon|^{2H+2}| \leq C\varepsilon^{-2H}$$

for any $s, t \in [0, T]$. Therefore, $|\Lambda_\varepsilon(s, t)| \leq C\varepsilon^{-2H}$ and we obtain the estimate

$$|G_\varepsilon(s, t, s', t')| \leq C\varepsilon^{-8H} (s^{2H-1} + |s-s'|^{2H-1})(t^{2H-1} + |t-t'|^{2H-1}).$$

As a consequence,

$$\begin{aligned} |H_\varepsilon^1| &\leq \int_0^T \int_0^\varepsilon \int_0^\varepsilon \int_0^T |G_\varepsilon(s, t, s', t')| ds dt ds' dt' \\ &\leq C\varepsilon^{-8H} \int_0^T \int_0^\varepsilon \int_0^\varepsilon \int_0^T (s^{2H-1} + |s-s'|^{2H-1}) \\ &\quad \times (t^{2H-1} + |t-t'|^{2H-1}) ds dt ds' dt' \\ &\leq C\varepsilon^{2-8H}, \end{aligned}$$

which converges to zero because $H < \frac{1}{4}$. □

Recall the definition (1.18) of \widehat{G}_ε ,

$$\widehat{G}_\varepsilon = \int_0^T \frac{B_{u+\varepsilon}^{(1)} - B_u^{(1)}}{\varepsilon} \times \frac{B_{u+\varepsilon}^{(2)} - B_u^{(2)}}{\varepsilon} du.$$

We have the following result.

Theorem 14. *Convergences (1.19) and (1.20) hold.*

Proof. We use the same trick as in [11], Remark 1.3, point 4. Let β and $\tilde{\beta}$ be two independent one-dimensional fractional Brownian motions with index H . Set $B^{(1)} = (\beta + \tilde{\beta})/\sqrt{2}$ and $B^{(2)} = (\beta - \tilde{\beta})/\sqrt{2}$. It is easily checked that $B^{(1)}$ and $B^{(2)}$ are also two independent fractional Brownian motions with index H . Moreover, we have

$$\begin{aligned} \varepsilon^{3/2-2H} \widehat{G}_\varepsilon &= \frac{1}{2} \varepsilon^{3/2-2H} \int_0^T \left(\frac{\beta_{u+\varepsilon} - \beta_u}{\varepsilon} \right)^2 du - \frac{1}{2} \varepsilon^{3/2-2H} \int_0^T \left(\frac{\tilde{\beta}_{u+\varepsilon} - \tilde{\beta}_u}{\varepsilon} \right)^2 du \\ &= \frac{1}{2\sqrt{\varepsilon}} \int_0^T \left(\frac{\beta_{u+\varepsilon} - \beta_u}{\varepsilon^H} \right)^2 du - \frac{1}{2\sqrt{\varepsilon}} \int_0^T \left(\frac{\tilde{\beta}_{u+\varepsilon} - \tilde{\beta}_u}{\varepsilon^H} \right)^2 du \\ &= \frac{1}{2\sqrt{\varepsilon}} \int_0^T h_2 \left(\frac{\beta_{u+\varepsilon} - \beta_u}{\varepsilon^H} \right) du - \frac{1}{2\sqrt{\varepsilon}} \int_0^T h_2 \left(\frac{\tilde{\beta}_{u+\varepsilon} - \tilde{\beta}_u}{\varepsilon^H} \right) du. \end{aligned} \quad (6.8)$$

The proofs of the desired convergences in law are now direct consequences of the convergence (1.3) with $k = 2$, taking into account that β and $\tilde{\beta}$ are independent. \square

Remark 15. As a by-product of the decomposition (6.8), and taking into account (1.5) for $k = 2$, we get that $\int_0^T \dot{B}_u^{(1)} \diamond \dot{B}_u^{(2)} du$ and $(Z_T^{(2)} - \tilde{Z}_T^{(2)})/2$ have the same law when $H > 3/4$, where $\tilde{Z}_T^{(2)}$ stands for an independent copy of the Hermite random variable $Z_T^{(2)}$.

Acknowledgements

The research of I. Nourdin was supported in part by the ANR project ‘Exploration des Chemins Rugueux’. The research of D. Nualart was supported by NSF Grant DMS-0904538.

References

- [1] Bender, C. (2003). An S -transform approach to integration with respect to a fractional Brownian motion. *Bernoulli* **9** 955–983. [MR2046814](#)
- [2] Biagini, F., Øksendal, B., Sulem, A. and Wallner, N. (2004). An introduction to white noise theory and Malliavin calculus for fractional Brownian motion. *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.* **460** 347–372. [MR2052267](#)
- [3] Breuer, P. and Major, P. (1983). Central limit theorems for nonlinear functionals of Gaussian fields. *J. Multivariate Anal.* **13** 425–441. [MR0716933](#)
- [4] Coutin, L. and Qian, Z. (2002). Stochastic rough path analysis and fractional Brownian motion. *Probab. Theory Related Fields* **122** 108–140. [MR1883719](#)
- [5] Decreusefond, L. and Üstünel, A.S. (1998). Stochastic analysis of the fractional Brownian motion. *Potential Anal.* **10** 177–214. [MR1677455](#)
- [6] Dobrushin, R.L. and Major, P. (1979). Non-central limit theorems for nonlinear functionals of Gaussian fields. *Z. Wahrsch. Verw. Gebiete* **50** 27–52. [MR0550122](#)
- [7] Giraitis, L. and Surgailis, D. (1985). CLT and other limit theorems for functionals of Gaussian processes. *Z. Wahrsch. Verw. Gebiete* **70** 191–212. [MR0799146](#)
- [8] Kuo, H.-H. (1996). *White Noise Distribution Theory*. Boca Raton, FL: CRC Press. [MR1387829](#)

- [9] Marcus, M. and Rosen, J. (2008). CLT for L^p moduli of continuity of Gaussian processes. *Stochastic Process. Appl.* **118** 1107–1135. [MR2428711](#)
- [10] Marcus, M. and Rosen, J. (2008). Non-normal CLTs for functions of the increments of Gaussian processes with convex increment's variance. Available at [ArXiv: 0707.3928](#).
- [11] Nourdin, I. (2009). A change of variable formula for the 2D fractional Brownian motion of Hurst index bigger or equal to $1/4$. *J. Funct. Anal.* **256** 2304–2320. [MR2498767](#)
- [12] Nourdin, I., Nualart, D. and Tudor, C.A. (2010). Central and non-central limit theorems for weighted power variations of fractional Brownian motion. *Ann. Inst. H. Poincaré Probab. Statist.* To appear.
- [13] Nualart, D. (2006). *The Malliavin Calculus and Related Topics*, 2nd ed. Berlin: Springer. [MR2200233](#)
- [14] Nualart, D. (2005). A white noise approach to fractional Brownian motion. In *Stochastic Analysis: Classical and Quantum* 112–126. Hackensack, NJ: World Sci. Publ. [MR2233155](#)
- [15] Nualart, D. and Peccati, G. (2005). Central limit theorems for sequences of multiple stochastic integrals. *Ann. Probab.* **33** 177–193. [MR2118863](#)
- [16] Obata, N. (1994). *White Noise Calculus and Fock Space. Lecture Notes in Math.* **1577** 1–183. Berlin: Springer. [MR1301775](#)
- [17] Peccati, G. and Tudor, C.A. (2005). Gaussian limits for vector-valued multiple stochastic integrals. In *Séminaire de Probabilités XXXVIII. Lecture Notes in Math.* **1857** 247–262. Berlin: Springer. [MR2126978](#)
- [18] Russo, F. and Vallois, P. (2007). Elements of stochastic calculus via regularization. In *Séminaire de Probabilités XL. Lecture Notes in Math.* **1899** 147–185. Berlin: Springer. [MR2409004](#)
- [19] Taqqu, M.S. (1975). Weak convergence to fractional Brownian motion and to Rosenblatt process. *Z. Wahrsch. Verw. Gebiete* **31** 287–302. [MR0400329](#)
- [20] Taqqu, M.S. (1979). Convergence of integrated processes of arbitrary Hermite rank. *Z. Wahrsch. Verw. Gebiete* **50** 53–83. [MR0550123](#)
- [21] Unterberger, J. (2008). A central limit theorem for the rescaled Lévy area of two-dimensional fractional Brownian motion with Hurst index $H < 1/4$. Available at [ArXiv: 0808.3458](#).

Received April 2009