

Strongly barycentrically associative and preassociative functions

Bruno Teheux

(joint work with Jean-Luc Marichal)

Let X be a nonempty set and X^* be the free monoid generated by X . Recall that a function $F: X^* \rightarrow X \cup \{\varepsilon\}$ is barycentrically associative if the function value of a string does not change when replacing every letter of a substring of consecutive letters with the value of this substring.

In this talk, we investigate the weaker property of *strong barycentric associativity* which stipulates that the function value of a string does not change when replacing every letter of *any* substring with the value of this substring. Equivalently, a function $F: X^* \rightarrow X \cup \{\varepsilon\}$ is strongly barycentrically associative if and only if it satisfies the equation

$$F(\mathbf{xyz}) = F(F(\mathbf{xz})^{|\mathbf{x}|} \mathbf{y} F(\mathbf{xz})^{|\mathbf{z}|}), \quad \mathbf{xyz} \in X^*.$$

We also investigate a variant of strong barycentric associativity called *strong barycentric preassociativity* which does not involve composition of functions in its definition. We establish links between strong barycentric associativity and strong barycentric preassociativity. We recall a variant of Kolmogoroff-Nagumo's characterization of the class of quasi-arithmetic means based on the strong barycentric associativity property, and we generalize this characterization to strongly barycentrically-preassociative functions.