

EQUIVALENCES OF COISOTROPIC SUBMANIFOLDS

FLORIAN SCHÄTZ AND MARCO ZAMBON

ABSTRACT. We study the role that Hamiltonian and symplectic diffeomorphisms play in the deformation problem of coisotropic submanifolds. We prove that the action by Hamiltonian diffeomorphisms corresponds to the gauge-action of the L_∞ -algebra of Oh and Park. Moreover we introduce the notion of extended gauge-equivalence and show that in the case of Oh and Park's L_∞ -algebra one recovers the action of symplectic isotopies on coisotropic submanifolds. Finally, we consider the transversally integrable case in detail.

CONTENTS

Introduction	1
1. (Pre-)Symplectic Geometry	3
2. Deformations of coisotropic submanifolds	5
3. Hamiltonian diffeomorphisms	8
4. Symplectomorphisms	15
5. The transversally integrable case	22
Appendix A. The extended gauge-action	28
References	29

INTRODUCTION

Coisotropic submanifolds form an important class of sub-objects in symplectic and Poisson geometry. They naturally generalize Lagrangian submanifolds, play an important role in the theory of constraints and also appear in theoretical physics in the form of “branes”, i.e. boundary conditions of sigma models [5, 2].

In this note we consider coisotropic deformations inside a symplectic manifold. The nearby deformations of a Lagrangian submanifold L are well-understood: by Weinstein's normal form theorem, one can replace the ambient symplectic manifold by the cotangent bundle T^*L . The graph of a 1-form α is Lagrangian if and only if α is closed. If one identifies closed 1-forms which are related through an Hamiltonian isotopy, one arrives at the first de Rham cohomology group $H^1(L, \mathbb{R})$ of L as the appropriate moduli space of nearby Lagrangian deformations.

The generalization of these statements to coisotropic submanifolds is not obvious, since the space of coisotropic deformations is not linear and not even modelled on a topological vector space, see [18, 11]. However, the general pattern of deformation theory teaches us that

every deformation problem¹ should be captured by differential graded Lie algebras or their homotopical cousins, known as L_∞ -algebras. That this is indeed the case was established by Oh and Park in [11]. To be more precise, Oh and Park constructed an L_∞ -algebra that controls the formal deformation problem for coisotropic submanifolds. In the special case of a Lagrangian submanifold L , their construction recovers the de Rham complex of L .

In [15], we studied convergence issues arising in the framework of [11]. One finds that the *Maurer-Cartan equation*, which replaces the condition of being closed from the Lagrangian case, is always convergent, and that it converges to zero if and only if one is dealing with a coisotropic deformation.² Having established a firmer link to actual geometric deformations, it is natural to turn attention to the geometric symmetries that are present in the problem. In particular, one might wonder how the actions of Hamiltonian and symplectic isotopies on the space of coisotropic deformations can be understood. A natural symmetry acting on Maurer-Cartan elements of Oh and Park's L_∞ -algebra are the inner automorphisms, known as *gauge-transformations*. Our main result is that these agree with the action by Hamiltonian isotopies, while the action by symplectic isotopies agrees with certain extended gauge-equivalences, that we specify below.

In Section 3 we deal with Hamiltonian isotopies. It turns out that the gauge-transformations of Oh and Park's L_∞ -algebra correspond to certain special Hamiltonian isotopies. The remaining problem is to show that any Hamiltonian isotopy can be reduced to such a special one. This is parallel to the Lagrangian situation: there the main task is also to show that arbitrary Hamiltonian isotopies can be reduced to a function f on the Lagrangian submanifold, which acts on the space of closed 1-forms (whose graphs we are interested in) simply by $\alpha \mapsto \alpha + df$. We establish the appropriate generalization in Theorem 3.21, Subsection 3.5. As a consequence, we identify

$$\frac{\{\text{coisotropic submanifolds}\}}{\text{Hamiltonian isotopies}} \simeq \frac{\{\text{Maurer-Cartan elements}\}}{\text{gauge-equivalences}},$$

which is the content of Theorem 3.22. For an alternative treatment within the BFV-formalism we refer to the article [14] by the first named author.

Section 4 is concerned with symplectic isotopies. Given a Lagrangian submanifold, any of its Lagrangian deformations is related to the original submanifold by a symplectomorphism, so we do not obtain an interesting moduli space. In the general coisotropic case the situation is much more complicated and we do obtain another reasonable equivalence relation on the space of deformations by considering symplectic isotopies. In order to fit this into the algebraic framework, we have to revise a construction of Oh and Park's L_∞ -algebra due to Cattaneo and Felder [3], which relies on Voronov's derived bracket construction [16, 17]. We show that every L_∞ -algebra which arises through Voronov's construction comes along with additional automorphisms. As a consequence, we obtain more ways to identify Maurer-Cartan elements. We refer to this extended equivalence relation as *extended gauge-equivalence*. The content of Theorem 4.17, Subsection 4.4 is that if one applies this to Oh and Park's L_∞ -algebra, one precisely recovers the action of symplectic isotopies on

¹... in characteristic zero...

²For an alternative treatment of the coisotropic deformation problem in terms of a Maurer-Cartan equation, see [13].

the space of coisotropic deformations. As a consequence, we identify

$$\frac{\{\text{coisotropic submanifolds}\}}{\text{symplectic isotopies}} \cong \frac{\{\text{Maurer-Cartan elements}\}}{\text{extended gauge-equivalences}},$$

see Theorem 4.18.

In Section 5, we consider coisotropic submanifolds which are transversally integrable. This regularity condition allows one to make some of the previous constructions more explicit. In particular, one can give formula for nearby coisotropic deformations which are obtained by an Hamiltonian or symplectic isotopy from the original coisotropic submanifold, see Proposition 5.12.

Organization of the paper: In Section 1 we recall background material on coisotropic submanifolds. In Section 2 we review the results about deformations of coisotropic submanifolds which are relevant in the subsequent discussion. In particular, we introduce Oh and Park's L_∞ -algebra and review the relation between its Maurer-Cartan elements and the deformation problem. In Section 3 we discuss Hamiltonian isotopies, while in Section 4 we deal with symplectic isotopies. In the final Section 5, we consider the case of transversally integrable submanifolds.

Comparison with the literature: While we were completing this note, the preprint [8] by Lê, Oh, Tortorella and Vitagliano appeared. It considers coisotropic deformations in the very general setting of Jacobi manifolds, which include Poisson and symplectic manifolds as special cases. There is an overlap between the results presented there in [8, Subsection 4.4] - once specialized to the symplectic case - and one of the main sections of the present note, namely Section 3. In particular, Thm. 3.21 (i.e. the equivalence of Hamiltonian equivalence and gauge equivalence, under a compactness assumption) corresponds to [8, Corollary 4.24]. Notice that in the latter the assumption on the compactness of the coisotropic submanifold is omitted.

Acknowledgements: M.Z. thanks Luca Vitagliano and Alberto Martín Zamora for useful conversations. In particular, the proof of Lemma 3.14 was communicated to us by Luca Vitagliano. Moreover we thank Hồng Vân Lê for useful comments on a draft-version of this note.

F.S. was supported by the Center of Excellence Grant "Centre for Quantum Geometry of Moduli Spaces" from the Danish National Research Foundation (DNRF95). M.Z. was partially supported by grants MTM2011-22612 and ICMAT Severo Ochoa SEV-2011-0087 (Spain), and Pesquisador Visitante Especial grant 88881.030367/2013-01 (CAPES/Brazil).

1. (PRE-)SYMPLECTIC GEOMETRY

We summarize background information about coisotropic submanifolds and associated structures.

Remark 1.1. Throughout this paper, (M, ω) will denote a symplectic manifold. Let C be a submanifold of M and $E \rightarrow C$ a vector subbundle of $TM|_C$. The symplectic orthogonal

E^\perp to E is the vector bundle whose fibre over $x \in C$ is

$$E_x^\perp := \{e \in T_x M \text{ such that } \forall v \in E_x \text{ we have } \omega_x(e, v) = 0\}.$$

Another way to characterize E^\perp is as the pre-image of the annihilator E° of E under the sharp-map

$$\omega^\sharp : TM \rightarrow T^*M, \quad v \mapsto \omega(v, -).$$

Definition 1.2. A submanifold C of (M, ω) is **coisotropic** if the symplectic orthogonal TC^\perp to TC is contained in TC .

Remark 1.3. An alternative way to express the coisotropy of C is in terms of the Poisson bivector-field Π associated to ω , defined by the requirement that $\Pi^\sharp : T^*M \rightarrow TM$, $\xi \mapsto \Pi(\xi, -)$ equals $-(\omega^\sharp)^{-1}$. Let $\mathcal{X}^\bullet(M)$ denote the space of multivector-fields on M , i.e. sections of $\wedge TM$. There is a natural projection map

$$P : \mathcal{X}^\bullet(M) \rightarrow \Gamma(\wedge(TM|_C/TC)),$$

which is given by restricting multivector-fields to C , followed by composition with the natural projection $\wedge TM|_C \rightarrow \wedge(TM|_C/TC)$. The submanifold C is coisotropic if and only if the Poisson bivector field Π lies in the kernel of P .

Definition 1.4. A two-form η on C that is closed and whose rank is constant is called a **pre-symplectic structure**.

Lemma 1.5. Let C be a coisotropic submanifold of (M, ω) . The pull back of ω to C along the inclusion $\iota : C \hookrightarrow M$ is a closed two-form of constant rank $2 \dim C - \dim M$. We denote this form by ω_C .

Remark 1.6.

- (1) Let η be any pre-symplectic structure on C . The closedness of η implies that the kernel of $\eta^\sharp : TC \rightarrow T^*C$ is an involutive subbundle of TC . Hence C carries a foliation, called the characteristic foliation of η .
- (2) We now consider the case of the pre-symplectic structure ω_C associated to a coisotropic submanifold C of (M, ω) . We always denote the kernel of ω_C by $K(= TC^\perp)$ and the corresponding characteristic foliation by \mathcal{F} in this situation. Moreover, observe that, in this situation, the vector bundle morphism

$$TM|_C \xrightarrow{\omega^\sharp} T^*M|_C \longrightarrow (TC^\perp)^*$$

is surjective and has kernel TC . Hence we obtain an isomorphism between the normal bundle $TM|_C/TC$ and K^* .

Remark 1.7. We saw that every coisotropic submanifold comes along with a pre-symplectic structure. An important observation is that this can be reversed: *every* pre-symplectic structure can be realized as the pre-symplectic structure associated to a coisotropic submanifold. Moreover, this realization is essentially unique. We start with a pre-symplectic structure η on a manifold C . Let K be the kernel of η^\sharp and G a complement to K . The choice of G yields an inclusion $j : K^* \hookrightarrow T^*C$. Recall that T^*C carries a canonical symplectic structure ω_{T^*C} . We now combine η and ω_{T^*C} into the two-form

$$\Omega := \pi^* \omega_C + j^* \omega_{T^*C}.$$

on K^* , where π denotes the projection map $K^* \rightarrow C$.

The two-form Ω restricts to η on C and is symplectic on a tubular neighborhood U of the zero section $C \subset K^*$. We refer to (U, Ω) as the *local symplectic model* associated to the the pre-symplectic manifold (C, η) .

The local symplectic model depends on the choice of complement G to K , but choosing different complements will lead to local symplectic models which are symplectomorphic in neighborhoods of C , and one can choose a symplectomorphism that restricts to the identity on C . Hence we will speak of *the* local symplectic model of (C, η) .

The following theorem of Gotay [4] asserts that actually *every* symplectic manifold (M, ω) into which C embeds as a coisotropic submanifold, such that $\omega_C = \eta$, looks like the local symplectic model in a neighborhood of C :

Theorem 1.8 (Gotay [4]). *Let C be a coisotropic submanifold of a symplectic manifold (M, ω) . There is a symplectomorphism ψ between a tubular neighborhood of C inside M and a tubular neighborhood of C inside its local symplectic model (U, Ω) . Moreover, ψ can be chosen such that the restriction of ψ to C is the identity.*

Throughout the rest of the paper we fix a local symplectic model (U, Ω) of the coisotropic submanifold C . Since the local symplectic model is a neighborhood of the zero section in a vector bundle $E \rightarrow C$, it comes equipped with an embedding of the zero section C in U , with coisotropic image, as well as with a surjective submersion $\pi: U \rightarrow C$. Recall that E is isomorphic to K^* , the dual to the kernel of the pre-symplectic structure ω_C . To avoid unnecessary confusion about signs, we also assume that U was chosen invariant with respect to fibrewise multiplication by -1 .

Summarizing, the setting we assume in the rest of the paper is:

(M, ω) is a symplectic manifold,
 C is a coisotropic submanifolds with induced presymplectic form ω_C ,
 (U, Ω) is the local symplectic model,
 where U is a neighborhood of the zero section in a vector bundle $E \rightarrow C$.

2. DEFORMATIONS OF COISOTROPIC SUBMANIFOLDS

We set up the problem of deforming a given coisotropic submanifold and discuss some relevant results, setting the stage for the subsequent development. In particular, the precise relationship between the deformation problem and the $L_\infty[1]$ -algebra of Oh and Park [11, 15] is recalled.

2.1. The deformation problem. It is natural to wonder how the “space of coisotropic submanifolds close to C ” looks like, i.e. we ask

Which deformations of C are coisotropic submanifolds of (U, Ω) ?

Definition 2.1. *The space of coisotropic sections of U is*

$$\text{Def}_U(C) := \{s \in \Gamma(U) : \text{the graph of } s \text{ is coisotropic inside } (U, \Omega)\}.$$

We now translate the above question into:

How can one describe the set $\text{Def}_U(C)$?

Theorem 2.9 in Subsection 2.3 provides an answer to this question.

2.2. Infinitesimal deformations. We discuss the infinitesimal version of the space $\text{Def}_U(C)$, which turns out to be closely related to the foliated de Rham complex.

Remark 2.2. Recall from Section 1 that the kernel K of the pre-symplectic structure ω_C on C is involutive, and the associated foliation \mathcal{F} of C is called the characteristic foliation. One has the following foliated version of the de Rham complex:

$$\begin{aligned}\Omega_{\mathcal{F}}(C) &:= \Gamma(\wedge K^*), \\ (d_{\mathcal{F}}\omega)(s_0, \dots, s_k) &:= \sum_{i=0}^k (-1)^i s_i(\omega(s_0, \dots, s_{i-1}, \widehat{s}_i, s_{i+1}, \dots, s_k)) \\ &\quad + \sum_{i < j} (-1)^{i+j} \omega([s_i, s_j], s_1, \dots, \widehat{s}_i, \dots, \widehat{s}_j, \dots, s_k).\end{aligned}$$

In Remark 1.6, we obtained a vector bundle isomorphism

$$E = TM|_C/TC \rightarrow K^*,$$

by restricting ω^\sharp . This yields an isomorphism $\Gamma(\wedge E) \cong \Gamma(\wedge K^*) = \Omega_{\mathcal{F}}(C)$. The foliated de Rham operator $d_{\mathcal{F}}$ then corresponds to the operator

$$\xi \mapsto P([\Pi, \xi]),$$

where $\xi \in \Gamma(\wedge E)$ is interpreted as a vertical multivector-field that is constant along the fibres of E , and $[\cdot, \cdot]$ is the Schouten-Nijenhuis bracket, see [15, Proof of Prop 3.5] for more details.

Remark 2.3. We will show that the formal tangent space to $\text{Def}_U(C)$ can be identified with the space of $d_{\mathcal{F}}$ -closed foliated one-forms on C . To this end, we rewrite the condition for a section s of U to be coisotropic in a more algebraic way. First, every section $s \in \Gamma(U)$ yields a diffeomorphism

$$\psi_{-s} : E \rightarrow E, \quad (x, e) \mapsto (x, e - s_x),$$

which maps $\text{graph}(s)$ of to the zero section $C \subset E$.

The graph of s is coisotropic with respect to Ω if and only if the zero section is coisotropic with respect to $(\psi_{-s})_*\Pi$, where Π denotes the Poisson bivector-field corresponding to Ω . As discussed in Section 1, the latter statement can be expressed by saying that $(\psi_{-s})_*\Pi$ lies in the kernel of the projection map

$$(1) \quad P : \mathcal{X}^\bullet(E) \rightarrow \Gamma(\wedge E),$$

given by restriction to C , composed with the projection $\wedge TE|_C \rightarrow \wedge E$.

Hence, if we define μ to be the map

$$\mu : \Gamma(U) \rightarrow \Gamma(\wedge^2 E), \quad s \mapsto P((\psi_{-s})_*\Pi),$$

a section s will be coisotropic if and only if it is mapped to zero under μ .

The map μ seems non-local since it involves the symplectic form away from C . However, the symplectic structure Ω_C of the local symplectic model (U, Ω) is determined by ω_C . We will return to this point in Subsection 2.3, where we see that the equation $\mu(-s) = 0$ can in fact be recovered as the Maurer-Cartan equation of an L_∞ -algebra whose structure maps are multi-differential operators on C .

Proposition 2.4. *Let s_t be a smooth one-parameter family of sections of U which starts at the zero section $s_0 = 0$. Then*

$$\frac{\partial}{\partial t}\Big|_{t=0}\mu(s_t) = -d_{\mathfrak{F}}\left(\frac{\partial}{\partial t}\Big|_{t=0}s_t\right)$$

under the identification $E \cong K^*$.

Proof. Consider the one-parameter family of diffeomorphisms $\psi_{-s_t}: E \rightarrow E$. The corresponding time-dependent vector field is $Y_t := -\frac{\partial}{\partial t}s_t$, a vertical vector field which is constant on each fiber of E . Using this and the definition of μ , we see that $\frac{\partial}{\partial t}\Big|_{t=0}\mu(s_t)$ equals the image under the projection $P: \chi^\bullet(E) \rightarrow \Gamma(\wedge E)$ of $\mathcal{L}_{\frac{\partial}{\partial t}\Big|_{t=0}s_t}\Pi = -[\Pi, \frac{\partial}{\partial t}\Big|_{t=0}s_t]$. By Remark 2.2 this is exactly the formula for the image of $\frac{\partial}{\partial t}\Big|_{t=0}s_t$ under $d_{\mathfrak{F}}$, if we apply the identification $E \cong K^*$. \square

Corollary 2.5. *Let s_t be a smooth one-parameter family of coisotropic sections of E with $s_0 = 0$. Then $\frac{\partial}{\partial t}\Big|_{t=0}s_t$ is closed with respect to $d_{\mathfrak{F}}$.*

Proof. We have $\mu(s_t) = 0$ for all t by Remark 2.3, hence the statement follows from Proposition 2.4. \square

Remark 2.6. The proof of the corollary shows that the space of closed elements of $\Omega_{\mathcal{F}}^1(C)$ is the formal tangent space to $\text{Def}_U(C)$ at C , where the formal tangent space is defined as the space of solutions to the linearized equation. We point out that it is known that not all cohomology classes of $H_{\mathcal{F}}^1(C)$ can be realized through one-parameter families of deformations, see [18, 11].

2.3. Oh and Park's $L_\infty[1]$ -algebra. We recall the $L_\infty[1]$ -algebra associated to C [11, 3].³

Definition 2.7. *An $L_\infty[1]$ -algebra is a \mathbb{Z} -graded vector space W , equipped with a collection of graded symmetric brackets $(\lambda_k: W^{\otimes k} \rightarrow W)_{k \geq 1}$ of degree 1 which satisfy a collection of quadratic relations [6], called higher Jacobi identities.*

The Maurer-Cartan series of a degree zero element $\beta \in W$ is the infinite sum

$$\text{MC}(\beta) := \sum_{k \geq 1} \frac{1}{k!} \lambda_k(\beta^{\otimes k}).$$

We say that β is a Maurer-Cartan element if its Maurer-Cartan series converges to zero⁴. We denote the set of all Maurer-Cartan elements of W by $\text{MC}(W)$.

Remark 2.8. In order to describe the $L_\infty[1]$ -algebra associated to the coisotropic submanifold C of (U, Ω) as explicitly as possible, we consider the Poisson structure Π associated to Ω . As explained in Section 1, the coisotropy of C is equivalent to $P(\Pi) = 0$, where

$$P: \chi^\bullet(E) \rightarrow \Gamma(\wedge E)$$

is as in Equation (1).

³The reader is referred to [8, Appendix D] for a proof that the construction from [11] coincides with the one from [3], specialized to the symplectic case.

⁴...with respect to a suitable topology. For the specific examples of $L_\infty[1]$ -algebras with which we will be concerned later on, we will make this precise.

As shown in [11] and [3], the space $\Gamma(\wedge E)[1]$ is equipped with a canonical $L_\infty[1]$ -algebra structure. We denote the structure maps of this $L_\infty[1]$ -algebra by

$$\lambda_k : \Gamma(\wedge E)[1]^{\otimes k} \rightarrow \Gamma(\wedge E)[1].$$

The evaluation of λ_k on $s \otimes \cdots \otimes s$ for $s \in \Gamma(E)$ yields

$$(2) \quad \lambda_k(s, \dots, s) := P([\dots [\Pi, s], s] \dots], s]),$$

where s is interpreted as a fibrewise constant vertical vector-field on E . Hence the Maurer-Cartan series of s reads $\text{MC}(s) = P(e^{[\cdot, s]}\Pi)$.

The following result, which is – partly in an implicit manner – contained in [11], is essentially [15, Thm. 2.8]. It relies on the fact that the Poisson bivector field associated to Ω is analytic in the fibre direction⁵, which is true thanks to [15, Cor. 2.7].

Theorem 2.9. *Consider the $L_\infty[1]$ -algebra $\Gamma(\wedge E)[1]$ associated to the coisotropic submanifold C . For any $s \in \Gamma(E)$ such that $\text{graph}(s)$ is contained in (U, Ω) , the Maurer-Cartan series $\text{MC}(-s)$ is convergent. Furthermore, for any such s the following two statements are equivalent:*

- (1) $\text{graph}(s)$ is a coisotropic submanifold of (U, Ω) .
- (2) The Maurer-Cartan series $\text{MC}(-s)$ converges to zero in the sense of pointwise convergence.

Remark 2.10. In other words, if we restrict attention to those sections whose graphs lie inside U , the map $s \mapsto -s$ restricts to a bijection between the set of coisotropic sections

$$\text{Def}_U(C) := \{s \in \Gamma(U) : \text{the graph of } s \text{ is coisotropic inside } (U, \Omega)\}$$

from Subsection 2.1, and

$$\text{MC}_U(\Gamma(\wedge E)[1]) := \{ \text{Maurer-Cartan elements of } \Gamma(\wedge E)[1] \text{ whose graphs lie in } U \}.$$

Notice that the first structure map λ_1 of the $L_\infty[1]$ -algebra $\Gamma(\wedge E)[1]$ coincides with the foliated de Rham differential $d_{\mathfrak{F}}$ under the isomorphism $\Gamma(\wedge E) \cong \Omega_{\mathcal{F}}(C)$. We could – a posteriori – use this fact to recover the infinitesimal description of $\text{Def}_U(C)$ which we obtained in Subsection 2.2.

3. HAMILTONIAN DIFFEOMORPHISMS

In this section we investigate the action of Hamiltonian diffeomorphisms on the space of coisotropic submanifolds. More precisely, we provide a description of the induced equivalence relation on the space of coisotropic sections. As the main result, we show that for compact coisotropic submanifolds this equivalence relation coincides with the gauge-equivalence in Oh and Park's $L_\infty[1]$ -algebra. This result was obtained independently by Lê, Oh, Tortorella and Vitagliano in [8, Corollary 4.24].

⁵In [15], such bivector-fields are called fibrewise entire.

3.1. The deformation problem. Recall that by Definition 2.1 a section s of π is called coisotropic if $\text{graph}(s)$ is a coisotropic submanifold of (U, Ω) , and that we denote the set of all such sections by $\text{Def}_U(C)$.

Definition 3.1. *Two coisotropic sections s_0 and s_1 are called **Hamiltonian equivalent** if there is a family of coisotropic sections s_t , agreeing with the given ones at $t = 0$ and $t = 1$, and an isotopy of Hamiltonian diffeomorphisms ϕ_t such that ϕ_t maps the graph of s_0 to the graph of s_t for all $t \in [0, 1]$.*

Remark 3.2. To be more precise, we assume that we are given a locally defined Hamiltonian isotopy, i.e. a family of diffeomorphisms between open subsets of U , generated by a family of locally defined Hamiltonian vector fields, which maps $\text{graph}(s_0)$ onto $\text{graph}(s_t)$.

It is straight-forward to check that Hamiltonian equivalence actually defines an equivalence relations on the set $\text{Def}_E(C)$, which we denote by \sim_{Ham} . It is natural to wonder about the equivalence classes, so we define:

Definition 3.3. *The Hamiltonian moduli space of coisotropic sections is the set*

$$\mathcal{M}_U^{\text{Ham}}(C) := \text{Def}_U(C) / \sim_{\text{Ham}} .$$

We ask:

How can one describe the set $\mathcal{M}_U^{\text{Ham}}(C)$?

Theorem 3.22 of Subsection 3.5 provides an answer in terms of the $L_\infty[1]$ -algebra of Oh and Park.

3.2. Infinitesimal moduli. We discuss the infinitesimal version of $\mathcal{M}_U^{\text{Ham}}(C)$. In particular, we argue that the formal tangent space to $\mathcal{M}_U^{\text{Ham}}(C)$ at the equivalence class of the zero-section C is given by the first foliated cohomology $H_{\mathfrak{F}}^1(C)$, with \mathfrak{F} the characteristic foliation of the pre-symplectic structure on C . The results of this subsection can be recovered – via specialization to the symplectic case – from the results obtained by Lê and Oh, [7, Subsection 6.3], who studied deformations of coisotropic submanifolds in locally conformal symplectic manifolds.

Remark 3.4. Let $(s_t)_{t \in [0,1]}$ be a family of coisotropic sections that starts at the zero-section. In Subsection 2.2 we saw that $\frac{\partial s_t}{\partial t}|_{t=0} \in \Gamma(E)$ lies in the kernel of the complex $(\Gamma(\wedge E), P([\Pi, -]))$ and that the latter is isomorphic to the foliated de Rham complex $(\Omega_{\mathfrak{F}}(C), d_{\mathfrak{F}})$.

Proposition 3.5. *Suppose that $(s_t)_{t \in [0,1]}$ is a family of coisotropic sections that starts at the zero-section and is trivial under Hamiltonian equivalence, i.e. there is an Hamiltonian isotopy ϕ_t such that the graph of s_t coincides with the image of the zero section under ϕ_t .*

Then the cohomology class of $\frac{\partial s_t}{\partial t}|_{t=0}$ in $H_{\mathfrak{F}}^1(C)$ is trivial.

Proof. Suppose that ϕ_t is generated by the family of Hamiltonian vector fields X_{H_t} . We can write $\frac{\partial s_t}{\partial t}|_{t=0}$ as $P(X_{H_0}) = P([\Pi, H_0])$ (see Lemma 3.13 later on). We observe that the latter expression equals $P([\Pi, H_0|_C])$, because $\Pi^\sharp|_C$ maps the co-normal bundle to the tangent bundle TC , whose sections lie in the kernel of P . As a consequence, the cohomology class of $\frac{\partial s_t}{\partial t}|_{t=0}$ equals the cohomology class of $P([\Pi, H_0|_C])$, which is trivial. Now apply the isomorphism between $\Gamma(\wedge E)$ and the foliated de Rham complex from Remark 2.3. \square

Remark 3.6. For every $f \in C^\infty(C)$, let ϕ_t be the flow of the Hamiltonian vector field $X_{\pi^* f}$, and $(s_t)_{t \in [0, \epsilon]}$ the family of coisotropic sections determined by $\text{graph}(s_t) = \phi_t(C)$. Then the proof of Proposition 3.5 shows that $\frac{\partial s_t}{\partial t}|_{t=0}$ corresponds to $d_{\mathfrak{F}} f$ under the isomorphism $\Gamma(E) \cong \Omega_{\mathfrak{F}}^1(C)$. Hence we can refine Proposition 3.5 as follows: the formal tangent space of the set of coisotropic sections which are trivial under Hamiltonian equivalence is precisely $\Omega_{\mathfrak{F}, \text{exact}}^1(C)$.

This and Remark 2.6 imply that the formal tangent space at zero to $\mathcal{M}_U^{\text{Ham}}(C)$ is $H_{\mathfrak{F}}^1(C)$. In the special case of C Lagrangian, this reduces to the first deRham cohomology $H^1(C)$ of C , as expected.

3.3. Gauge-equivalence.

Remark 3.7. Ignoring convergence issues, every $L_\infty[1]$ -algebra W comes along with its group of inner automorphisms, called the gauge-group, which acts on W_0 , the elements of degree 0. It is generated by elements γ of degree -1 . The vector field on W corresponding to γ reads

$$\lambda_1(\gamma) + \lambda_2(\gamma, \beta) + \frac{1}{2!} \lambda_3(\gamma, \beta, \beta) + \frac{1}{3!} \lambda_4(\gamma, \beta, \beta, \beta) + \dots$$

at the point $\beta \in W_0$. This action preserves the set of Maurer-Cartan elements of W and induces a canonical equivalence relations there:

Definition 3.8. *Two Maurer-Cartan elements β_0 and β_1 of an $L_\infty[1]$ -algebra W are **gauge-equivalent** if there is a one-parameter family γ_t of degree -1 elements of W and a one-parameter family β_t of degree zero elements of W , agreeing with the given ones at $t = 0$ and $t = 1$, such that*

$$\frac{\partial}{\partial t} \beta_t = \lambda_1(\gamma_t) + \lambda_2(\gamma_t, \beta_t) + \frac{1}{2!} \lambda_3(\gamma_t, \beta_t, \beta_t) + \frac{1}{3!} \lambda_4(\gamma_t, \beta_t, \beta_t, \beta_t) + \dots$$

We presuppose that W is equipped with a suitable topology and that the right-hand side of the above equation converges.

We apply this to the $L_\infty[1]$ -algebra structure on $\Gamma(\wedge E)[1]$ from Subsection 2.3. We are interested in $\text{MC}_U(\Gamma(\wedge E)[1])$, the Maurer-Cartan elements of $\Gamma(\wedge E)[1]$ whose graphs lie in U (see Remark 2.10). We define an equivalence relation on $\text{MC}_U(\Gamma(\wedge E)[1])$ as in Def. 3.8, but additionally requiring that the one-parameter family of degree zero elements β_t consists of sections of U (rather than E). We use the bijection $\text{Def}_U(C) \cong \text{MC}_U(\Gamma(\wedge E)[1])$, $s \mapsto -s$ described in Remark 2.10 to transport the above equivalence relation to $\text{Def}_U(C)$:

Definition 3.9. *Two coisotropic sections s_0 and s_1 are called **gauge equivalent**, $s_0 \sim_{\text{gauge}} s_1$, if $-s_0$ and $-s_1$ are equivalent elements (in the sense above) of $\text{MC}_U(\Gamma(\wedge E)[1])$.*

Remark 3.10. We make the equivalence relation \sim_{gauge} more explicit. Two elements s_0 and s_1 in $\text{Def}_U(C)$ are declared gauge-equivalent if there is a smooth one-parameter family s_t in $\Gamma(U)$, coinciding with s_0 and s_1 at the endpoints, such that

$$\begin{aligned} \frac{\partial}{\partial t}(-s_t) &= P([\Pi, \pi^* f_t]) + P([\Pi, \pi^* f_t], -s_t) + \frac{1}{2!} P([[[\Pi, \pi^* f_t], -s_t], -s_t]) + \dots \\ &= P(e^{[\cdot, -s_t]} X_{\pi^* f_t}). \end{aligned}$$

Here $-s_t$ is interpreted as a family of fibrewise constant vertical vector field and f_t is a one-parameter family of smooth functions on C . Observe that the latter can be seen as a one-parameter family of degree -1 elements of the $L_\infty[1]$ -algebra $\Gamma(\wedge E)[1]$. To rewrite the condition in more geometric terms, recall that for $s \in \Gamma(E)$, ψ_s is the diffeomorphism of E that consists of fibrewise addition with s . Moreover, let \mathfrak{p}_s^v be the projection of $TE|_{\text{graph}(s)}$ onto the vertical part of TE along $T\text{graph}(s)$.

We now compute

$$\begin{aligned} P(e^{[\cdot, -s_t]} X_{\pi^* f_t}) &= P((\psi_{-s_t})_* X_{\pi^* f_t}) = \mathfrak{p}_0^v((\psi_{-s_t})_*(X_{\pi^* f_t}|_{\text{graph}(s_t)})) \\ &= (\psi_{-s_t})_*(\mathfrak{p}_{s_t}^v(X_{\pi^* f_t}|_{\text{graph}(s_t)})) = \mathfrak{p}_{s_t}^v(X_{\pi^* f_t}|_{\text{graph}(s_t)}). \end{aligned}$$

We use [15, Prop. 1.15] in the first equality⁶, which applies since the vector field $X_{\pi^* f_t}$ is fiberwise entire in the terminology of [15]. In the last equality we used the fact that ψ_{-s_t} maps $\text{graph}(s_t)$ to the zero section C and preserves the fibers of the projection $\pi : U \rightarrow C$.

After reversing the signs in front of f_t , this shows:

Proposition 3.11. *Elements s_0 and s_1 of $\text{Def}_U(C)$ are gauge equivalent if and only if there is a one-parameter family $s_t \in \Gamma(U)$, agreeing with s_0 and s_1 at the endpoints, and a one-parameter family $f_t \in \mathcal{C}^\infty(C)$ such that*

$$(3) \quad \frac{\partial}{\partial t} s_t = \mathfrak{p}_{s_t}^v(X_{\pi^* f_t}|_{\text{graph}(s_t)})$$

holds for all $t \in [0, 1]$.

3.4. Technical Lemmata. We establish some technical lemmata that we use subsequently to relate various notions of equivalence between coisotropic sections.

Remark 3.12. Throughout this subsection, A denotes a vector bundle over a smooth manifold M . Given a section s of A , we have a splitting $T_y A = V_y \oplus T_y \text{graph}(s)$ of the tangent space to A at y , where $V := \ker(d\pi)$ is the vertical bundle. We will denote by \mathfrak{p}_s^v the projection $T_y A \rightarrow V_y$ with kernel $T_y \text{graph}(s)$. It is defined for all $y \in \text{graph}(s)$.

Lemma 3.13. *Let X_t be a one-parameter family of vector fields on A , and ϕ_t its flow. Moreover, let s_t be a one-parameter family of sections of A such that*

$$\text{graph}(s_t) = \phi_t(\text{graph}(s_0))$$

holds for all $t \in [0, 1]$.

Then s_t satisfies the equation

$$\frac{\partial}{\partial t} s_t = \mathfrak{p}_{s_t}^v X_t, \quad \forall t \in [0, 1],$$

which we see as an equality of sections of $V|_{\text{graph}(s_t)}$.

Proof. If we define ψ_t to be the isotopy of M given by $\pi \circ \phi_t \circ s_0$, we have

$$s_t = \phi_t \circ s_0 \circ (\psi_t)^{-1} : M \rightarrow A.$$

⁶[15, Prop. 1.15] is stated for bivector fields, but it carries over immediately to the case of vector fields.

Evaluating at $x \in M$ and taking the time derivative we obtain

$$\frac{\partial}{\partial t}(s_t(x)) = X_t|_{s_t(x)} + (\phi_t)_*(s_0)_* \frac{\partial}{\partial t}((\psi_t)^{-1}(x)).$$

We finish noticing that the last summand is tangent to $\phi_t(\text{graph}(s_0)) = \text{graph}(s_t)$, and that $\frac{\partial}{\partial t}(s_t(x))$ lies in $V_{s_t(x)}$. \square

The following Lemma, whose (geometric) proof was communicated to us by Luca Vitagliano, is a converse to Lemma 3.13.

Lemma 3.14. *Let X_t be a one-parameter family of vector fields on A , and ϕ_t its flow, assumed to exist for all $t \in [0, 1]$. Suppose s_t is a one-parameter family of sections of A that satisfies*

$$(4) \quad \frac{\partial}{\partial t}s_t = p_{s_t}^v X_t, \quad \forall t \in [0, 1].$$

Then the family of submanifolds $\text{graph}(s_t)$ coincides with $\phi_t(\text{graph}(s_0))$ for all $t \in [0, 1]$.

Proof. We work on the vector bundle $A \times [0, 1] \rightarrow M \times [0, 1]$, and denote by t the standard coordinate on the $[0, 1]$ -factor. Define $\widehat{s} \in \Gamma(A \times [0, 1])$ by

$$\widehat{s}(x, t) = (s_t(x), t)$$

and the vector field \widehat{X} on $A \times [0, 1]$ by

$$\widehat{X}|_{(y,t)} = (X_t)|_y + \frac{\partial}{\partial t}.$$

Notice that the flow φ_t of \widehat{X} takes $(y, 0)$ to $(\phi_t(y), t)$ for all $y \in A$.

The key observation is that the vector field \widehat{X} is tangent to the submanifold $\text{graph}(\widehat{s})$. To this end we compute

$$\frac{d}{dt}\widehat{s}(x, t) = \frac{d}{dt}s_t(x) + \frac{\partial}{\partial t} = (X_t)|_{s_t(x)} - v + \frac{\partial}{\partial t} = \widehat{X}|_{(s_t(x), t)} - v$$

for some vector $v \in T_{s_t(x)}(\text{graph}(s_t))$, making use of equation (4) in the second equality. This implies that $\widehat{X}|_{(s_t(x), t)} = \frac{d}{dt}\widehat{s}(x, t) + v$ is the sum of two vectors tangent to $\text{graph}(\widehat{s})$.

Hence the flow φ_t of \widehat{X} maps $\text{graph}(\widehat{s}|_{M \times \{0\}}) = \text{graph}(s_0) \times \{0\}$ to $\text{graph}(\widehat{s}|_{M \times \{t\}}) = \text{graph}(s_t) \times \{t\}$. On the other hand, we saw above that φ_t maps $\text{graph}(s_0) \times \{0\}$ to $\phi_t(\text{graph}(s_0)) \times \{t\}$. \square

In Lemma 3.14 we assume that the flow of X_t is defined on the interval $[0, 1]$. We now show that this assumption can be replaced by asking that the base M of the vector bundle be compact.

Lemma 3.15. *Let $\pi: A \rightarrow M$ be a vector bundle over a compact base M . Let X_t be a one-parameter family of vector fields on A and s_t be a one-parameter family of sections of A that satisfies*

$$\frac{\partial}{\partial t}s_t = p_{s_t}^v X_t, \quad \forall t \in [0, 1].$$

Then the flow lines of X_t starting at $\text{graph}(s_0)$ exist for $t \in [0, 1]$ and the equality

$$\text{graph}(s_t) = \phi_t(\text{graph}(s_0))$$

holds.

Proof. Fix an auxiliary fibre metric on A . We let $K \subset A$ be the compact subset given by all vectors of length less than or equal to $l + \delta$ for some $\delta > 0$, where

$$l := \max_{x \in M, t \in [0,1]} (||s_t(x)||).$$

Let φ be a function on A with compact support, and so that $\varphi|_K \equiv 1$. Then $(\varphi X_t)_{t \in [0,1]}$ is a time-dependent vector field whose integral curves are defined for all times. Let T be the maximal element of $[0, 1]$ such that $\text{graph}(s_t) = \phi_t(\text{graph}(s_0))$ holds for all $t \in [0, T]$. Suppose $T < 1$. There is $\epsilon > 0$ such that $\phi_t(\text{graph}(s_0)) \subset K$ for all $t \in [0, T + \epsilon]$. But since the one-parameter families X_t and φX_t agree on K , we see as in Lemma 3.14 that $\text{graph}(s_t) = \phi_t(\text{graph}(s_0))$ actually holds for all $t \in [0, \min\{1, T + \epsilon\}]$, which is a contradiction. \square

Remark 3.16. The compactness assumption in Lemma 3.15 can not be omitted, as the following counter-example shows. Take a non-compact manifold M , a vector field X on M whose flow is not defined on the whole of $[0, 1]$. Take the trivial bundle $A := M \times [0, 1]$ and let X_t be the horizontal lift of X to A . Moreover, let s_t be $M \times \{0\}$. Notice that $\frac{\partial}{\partial t} s_t$ and $p_{s_t}^\vee X_t$ agree, since they both vanish identically.

3.5. Hamiltonian equivalence = gauge-equivalence. Our aim is to compare the two equivalence relations \sim_{gauge} and \sim_{Ham} on $\text{Def}_U(E)$. As an intermediate notion we introduce:

Definition 3.17. One can restrict Hamiltonian equivalence \sim_{Ham} by only allowing Hamiltonian flows generated by functions of the type $\pi^* f$, with $f \in C^\infty(C)$. We call the resulting equivalence relation **base Hamiltonian equivalence** and denote it by $\sim_{b\text{Ham}}$.

Proposition 3.18. The following chain of implications holds between the three equivalence relations on $\text{Def}_U(C)$:

$$\begin{array}{c} \text{base Hamiltonian equivalence } \sim_{b\text{Ham}} \\ \Downarrow (1) \\ \text{Hamiltonian equivalence } \sim_{\text{Ham}} \\ \Downarrow (2) \\ \text{gauge-equivalence } \sim_{\text{gauge}} . \end{array}$$

Proof. Implication (1) is clear, so we pass on to implication (2). Let s_t be a smooth family of coisotropic sections of U and suppose that H_t is a smooth family of functions on U such that the Hamiltonian flow $\phi_t^{H_t}$ of H_t maps $\text{graph}(s_0)$ to $\text{graph}(s_t)$. By Lemma 3.13, this implies that the equation

$$\frac{\partial}{\partial t} s_t = p_{s_t}^\vee X_{H_t},$$

holds for all $t \in [0, 1]$. Define $f_t \in C^\infty(C)$ to be $H_t \circ s_t$.

Observe that $p_{s_t}^\vee (X_{H_t} - X_{\pi^* f_t})$ is zero since $H_t - \pi^* f_t$ vanishes on $\text{graph}(s_t)$ and consequently $X_{H_t - \pi^* f_t} = X_{H_t} - X_{\pi^* f_t}$ gets mapped to $T\text{graph}(s_t)$ under Π^\sharp since $\text{graph}(s_t)$ is

coisotropic. We conclude that the equation

$$\frac{\partial}{\partial t} s_t = p_{s_t}^v X_{H_t} = p_{s_t}^v (X_{\pi^* f_t})$$

holds. By Proposition 3.11 we have that s_0 and s_1 are gauge-equivalent as claimed. \square

Under the assumption that C is compact, we can “close the circle” of the implications of Proposition 3.18:

Proposition 3.19. *Suppose C is compact coisotropic submanifold. Then the following implication holds for the local symplectic model of C :*

$$\begin{array}{c} \text{gauge-equivalence} \sim_{\text{gauge}} \\ \Downarrow \\ \text{base Hamiltonian equivalence} \sim_{\text{bHam}} . \end{array}$$

Proof. Suppose that s_0 and s_1 of $\text{Def}_U(C)$ are gauge-equivalent. This means that there is a one-parameter family s_t in $\text{Def}_U(C)$ and a one-parameter family of functions f_t on C such that

$$\frac{\partial}{\partial t} s_t = p_{s_t}^v X_{\pi^* f_t}$$

holds for all $t \in [0, 1]$.

The compactness of C allows us to apply Lemma 3.15, which states that the flow ϕ_t of $X_{\pi^* f_t}$ exists for all $t \in [0, 1]$ and indeed maps $\text{graph}(s_0)$ to $\text{graph}(s_t)$. \square

Remark 3.20. When C is a Lagrangian submanifold, Hamiltonian equivalence implies base Hamiltonian equivalence without any compactness assumption: this follows from Proposition 3.18 and Proposition 3.19, noticing that in the latter in the Lagrangian case no compactness is necessary, for $X_{\pi^* f_t}$ is a vertical vector field on $U \subset T^*C$. In particular, if $(\phi_t)_{t \in [0, 1]}$ is an isotopy by Hamiltonian diffeomorphisms mapping the zero section C to sections of U for all $t \in [0, 1]$, then $\phi_1(C)$ is the graph of an exact 1-form on C . This is in agreement with [9, Proposition 9.33].

Combining Proposition 3.18 and Proposition 3.19 we arrive at the main result of this section:

Theorem 3.21. *Let C be a compact coisotropic submanifold with local symplectic model (U, Ω) . The equivalence relations on*

$$\text{Def}_U(C) := \{s \in \Gamma(U) : s \text{ is coisotropic}\}$$

given by

- Hamiltonian equivalence \sim_{Ham} (Definition 3.1) and
- gauge-equivalence \sim_{gauge} (Definition 3.9, see also Proposition 3.11)

coincide.

As a consequence we obtain the following result:

Theorem 3.22. *Let C be a compact coisotropic submanifold with local symplectic model (U, Ω) . The bijection*

$$\mathrm{Def}_U(C) \cong \mathrm{MC}_U(\Gamma(\wedge E)[1])$$

descends to a bijection

$$\mathcal{M}_U^{\mathrm{Ham}}(C) := \mathrm{Def}_U(C) / \sim_{\mathrm{Ham}} \cong \mathrm{MC}_U(\Gamma(\wedge E)[1]) / \sim_{\mathrm{gauge}} .$$

Remark 3.23.

- (1) One could use Theorem 3.22 to rederive the infinitesimal description of $\mathcal{M}_U^{\mathrm{Ham}}(C)$ from Subsection 3.2 by linearizing the Maurer-Cartan equation and the gauge-equivalence.
- (2) A description of $\mathcal{M}_U^{\mathrm{Ham}}(C)$ similar to Theorem 3.22 was obtained in [14]. There the differential graded Lie algebra associated to the BFV-complex was used to encode deformations of C and the action of Hamiltonian diffeomorphisms. The BFV-complex has the advantage that it works also in the Poisson case, unlike the $L_\infty[1]$ -algebra from [11] and [3]. The drawbacks of the approach via the BFV-complex is that one needs to single out the geometrically relevant Maurer-Cartan elements by hand and is forced to deal with symmetries of symmetries.

4. SYMPLECTOMORPHISMS

Next we consider the action of symplectomorphisms on the space of coisotropic sections, which we encode by an equivalence relation \sim_{Sym} on the space of coisotropic sections $\mathrm{Def}_U(C)$. In the search for an interpretation of \sim_{Sym} in terms of Oh and Park's $L_\infty[1]$ -algebra, we are led to reconsider Voronov's derived bracket construction [16, 17].

4.1. The deformation problem. Let C be a coisotropic submanifold with local symplectic model (U, Ω) .

Definition 4.1. *Two coisotropic sections s_0 and s_1 of U are called **symplectic equivalent**, $s_0 \sim_{\mathrm{Sym}} s_1$ if there is a family of coisotropic sections $s_t \in \Gamma(U)$, agreeing with the given ones at $t = 0$ and $t = 1$, and an isotopy of local symplectomorphisms ϕ_t such that ϕ_t maps $\mathrm{graph}(s_0)$ to $\mathrm{graph}(s_t)$ for all $t \in [0, 1]$.*

Remark 4.2. As for Hamiltonian equivalence, it is straight-forward to check that \sim_{Sym} is in fact an equivalence relation. We define the **symplectic moduli space of coisotropic sections** to be the set

$$\mathcal{M}_U^{\mathrm{Sym}}(C) := \mathrm{Def}_U(C) / \sim_{\mathrm{Sym}} .$$

Our aim is to answer

How can one describe the set $\mathcal{M}_U^{\mathrm{Sym}}(C)$?

which we will achieve in Theorem 4.18 of Subsection 4.4.

4.2. Infinitesimal moduli. We first consider the infinitesimal counterpart of $\mathcal{M}_U^{\mathrm{Sym}}(C)$. We argue – see Remark 4.6 – that the formal tangent space to $\mathcal{M}_U^{\mathrm{Sym}}(C)$ at the equivalence class of the zero-section C is given by the cokernel of a certain map $r : H^1(C) \rightarrow H_{\mathfrak{F}}^1(C)$.

Remark 4.3.

- (1) Recall that every coisotropic submanifold C comes along with a pre-symplectic structure ω_C , whose kernel K is an involutive distribution, whose corresponding foliation on C is denoted by \mathfrak{F} . Restriction to K yields a chain map

$$r : \Omega(C) \rightarrow \Omega_{\mathcal{F}}(C)$$

between the ordinary and the foliated de Rham complex of C .

- (2) As we observed in Subsection 2.2, $\Omega_{\mathcal{F}}(C)$ is isomorphic to $\Gamma(\wedge E)$, equipped with the differential $P([\Pi, \cdot])$, where P is the projection from multivector-fields on E onto $\Gamma(\wedge E)$.

Lemma 4.4. *Let C be a coisotropic submanifold of (E, ω) with inclusion map ι . Given $\beta \in \Omega^1(E)$, denote by X_β the unique vector field on E which satisfies*

$$i_{X_\beta} \omega = \beta.$$

Then the triangle

$$\begin{array}{ccccc}
 & & \beta \in & \Omega^1(E) & \\
 & \swarrow & & \searrow & \\
 P(X_\beta) \in & & & & \Omega_{\mathcal{F}}^1(C). \\
 & \swarrow & & \searrow & \\
 & \Gamma(E) & \xrightarrow{\cong} & &
 \end{array}$$

commutes.

Proof. The identification $E \cong K^*$ from Section 1, which is used in the bottom map of the above diagram, maps $e \in E_x$ to $\omega^\sharp(e)|_{K_x}$. We have

$$\omega^\sharp(P(X_\beta))|_K = \omega^\sharp(X_\beta)|_K = \beta|_K,$$

where in the first equality we used that $\omega(v, -)$ vanishes on K for all $v \in TC$. This proves the desired commutativity. \square

The following proposition is a special instance of Lemma 6.7 in [7, Subsection 6.3.], where the more general case of locally conformal symplectic manifolds is treated. In its formulation we make use of the above isomorphism in order to view $\frac{\partial s_t}{\partial t}|_{t=0} \in \Gamma(E)$ as an element of $\Omega_{\mathcal{F}}^1(C)$.

Proposition 4.5. *Suppose that $(s_t)_{t \in [0,1]}$ is a family of coisotropic sections that starts at the zero-section and is trivial under symplectic equivalence, i.e. there is a symplectic isotopy ϕ_t such that the image of the zero section under ϕ_t coincides with the graph of s_t .*

Then the cohomology class of $\frac{\partial s_t}{\partial t}|_{t=0}$ in $H_{\mathfrak{F}}^1(C)$ lies in the image of $r : H^1(C) \rightarrow H_{\mathfrak{F}}^1(C)$.

Proof. Suppose that ϕ_t is the symplectic isotopy generated by the family of vector fields X_t . Since ϕ_t is symplectic, $\beta_t := i_{X_t} \omega$ is a family of closed one-forms. By Lemma 3.13, we can write $\frac{\partial s_t}{\partial t}|_{t=0}$ as $P(X_0)$. By the previous lemma, this equals the image of β_0 under $r \circ \iota^*$. In particular, the cohomology class of $\frac{\partial s_t}{\partial t}|_{t=0}$ coincides with the cohomology class $(r \circ \iota^*)[\beta_0]$, hence lies in the image of $r : H^1(C) \rightarrow H_{\mathfrak{F}}^1(C)$. \square

Remark 4.6. Proposition 4.5 is an analogue of Proposition 3.5, where we showed that if a family $(s_t)_{t \in [0,1]}$ is trivial under Hamiltonian equivalence then the cohomology class of $\frac{\partial s_t}{\partial t}|_{t=0}$ is zero.

One can strengthen Proposition 4.5 by observing that, by the same proof, every element in the image of the map $r: \Omega_{\text{closed}}^1(C) \rightarrow \Omega_{\mathfrak{F}, \text{closed}}^1(C)$ is of the form $\frac{\partial s_t}{\partial t}|_{t=0}$, where $(s_t)_{t \in [0,\epsilon]}$ arises through the action of a symplectic isotopy on the zero-section. Indeed, for every $\gamma \in \Omega_{\text{closed}}^1(C)$ one considers the symplectic isotopy generated by the vector field $(\omega^\sharp)^{-1}(\pi^*\gamma)$.

In full analogy to Remark 3.6, this together with Remark 2.6 shows that the formal tangent space at zero to $\mathcal{M}_U^{\text{Sym}}(C)$ is

$$\Omega_{\mathfrak{F}, \text{closed}}^1 / r(\Omega_{\text{closed}}^1(C)) \cong H_{\mathfrak{F}}^1(C) / r(H^1(C)),$$

that is, the cokernel of $r: H^1(C) \rightarrow H_{\mathfrak{F}}^1(C)$. The isomorphism is obtained by quotienting both terms on the left-hand side by $\Omega_{\mathfrak{F}, \text{exact}}^1$ and by using the following linear algebra statement for the denominator: if $f: V_1 \rightarrow V_2$ is a linear map and W_1, W_2 are subspaces such that $f(W_1) = W_2$, then $f(V_1)/W_2 = \text{Im}([f]: V_1/W_1 \rightarrow V_2/W_2)$.

We note that if C is Lagrangian we have $H_{\mathfrak{F}}^1(C) = H^1(C)$ and r is the identity, so its cokernel is trivial, as expected.

Notice also, by the above and Remark 3.6, that the formal tangent space at zero of $\mathcal{M}_U^{\text{Sym}}(C)$ is a quotient of the formal tangent space to $\mathcal{M}_U^{\text{Ham}}(C)$, and that they agree iff $r: H^1(C) \rightarrow H_{\mathfrak{F}}^1(C)$ is the zero map. This happens for instance if $H^1(C) = 0$, in which cases it is clear a priori that $\mathcal{M}_U^{\text{Sym}}(C) = \mathcal{M}_U^{\text{Ham}}(C)$, for all symplectic vector fields on U are Hamiltonian. In Example 4.23 below we display an example in which r is not the zero map.

4.3. The extended formal picture. We explain now how to interpret the equivalence relation \sim_{Sym} from the point of view of Oh and Park's $L_\infty[1]$ -algebra structure on $\Gamma(\wedge E)[1]$.

To this aim, we first need to prove an algebraic statements about Voronov's derived brackets construction, see Lemma 4.9.

Remark 4.7 (on coalgebras and $L_\infty[1]$ -algebras).

- (1) Let \mathfrak{a} be a graded vector space. Consider $S\mathfrak{a} := \bigoplus_{i \geq 0} S^i \mathfrak{a}$, the graded symmetric coalgebra of \mathfrak{a} . It is obtained from the tensor algebra on \mathfrak{a} , modulo the homogeneous ideal generated by

$$x \otimes y - (-1)^{|x||y|} y \otimes x.$$

Deconcatenation

$$\begin{aligned} \Delta(x_1 \otimes \cdots \otimes x_n) &:= 1 \otimes (x_1 \otimes \cdots \otimes x_n) + \sum_{i=1}^{n-1} (x_1 \otimes \cdots \otimes x_i) \otimes (x_{i+1} \otimes \cdots \otimes x_n) + \\ &\quad (x_1 \otimes \cdots \otimes x_n) \otimes 1 \end{aligned}$$

defines a cocommutative coassociative coproduct on $S\mathfrak{a}$. Sometimes we work with the completion $\hat{S}\mathfrak{a}$ of $S\mathfrak{a}$, which consists of formal power series in \mathfrak{a} . The coproduct makes $\hat{S}\mathfrak{a}$ into a topological coalgebra. We notice that $S\mathfrak{a}$ and its completion $\hat{S}\mathfrak{a}$ come equipped with an augmentation map ϵ to \mathbb{R} , which just projects onto the constant term.

- (2) Recall that an element ξ of an augmented coalgebra of degree 0 is called primitive if $\epsilon(\xi) = 0$ and $\Delta(\xi) = \xi \otimes 1 + 1 \otimes \xi$. Similarly, an element g is called group-like if its image under the augmentation map is 1 and $\Delta(g) = g \otimes g$. These conditions still make sense for a topological coalgebra and one can prove that the primitive elements of $\hat{S}\mathfrak{a}$ are exactly those sitting in $\mathfrak{a} \subset \hat{S}\mathfrak{a}$, see [10, Theorem 5.18]. From this one can deduce that the group-like elements of $\hat{S}\mathfrak{a}$ are all of the form

$$e^\beta = 1 + \beta + \frac{1}{2!}\beta \otimes \beta + \frac{1}{3!}\beta \otimes \beta \otimes \beta + \dots$$

for some $\beta \in \mathfrak{a}$.

- (3) The language of coalgebras can be used to encode $L_\infty[1]$ -algebra quite efficiently: An $L_\infty[1]$ -algebra structure on \mathfrak{a} is the same as a degree 1 coderivation \tilde{D} of the coalgebra $S\mathfrak{a}$ that annihilates $1 \in \mathbb{R} \subset S\mathfrak{a}$ and squares to zero, i.e. an endomorphism \tilde{D} of $S\mathfrak{a}$ that satisfied

$$\Delta \circ \tilde{D} = (\tilde{D} \otimes \text{id} + \text{id} \otimes \tilde{D}) \circ \Delta, \quad \tilde{D}(1) = 0, \quad \text{and} \quad \tilde{D} \circ \tilde{D} = 0.$$

Observe that this means that an $L_\infty[1]$ -algebra structure on \mathfrak{a} corresponds to a Maurer-Cartan element \tilde{D} in the graded Lie algebra of coderivations $\text{Coder}(S\mathfrak{a})$, equipped with the commutator bracket. We note that the Maurer-Cartan equation for β in \mathfrak{a}_0 is equivalent to

$$e^{-\beta} \tilde{D}(e^\beta) = 0.$$

This follows from the fact that $e^{-\beta} \tilde{D}(e^\beta)$ is primitive, hence lives in \mathfrak{a} and equals the image of $\tilde{D}e^\beta$ under the projection $S\mathfrak{a} \rightarrow \mathfrak{a}$. The latter expression coincides with the Maurer-Cartan series of β .

- (4) We will also consider coderivations of $S\mathfrak{a}$ of degree 0. Observe that if D is such a coderivation and $\beta \in \mathfrak{a}_0$, then

$$e^{-\beta} D(e^\beta)$$

will be automatically primitive, i.e. an element of \mathfrak{a}_0 . We denote this element by $\hat{D}(\beta)$.

Remark 4.8 (on Voronov's derived brackets).

- (1) We briefly review the framework of Voronov's derived brackets [16, 17] and the main result of [1]. Let L be a graded Lie algebra, \mathfrak{a} an abelian subalgebra of L , $P : L \rightarrow \mathfrak{a}$ a projection whose kernel is a subalgebra of L . In [1] it is shown that these data induce a morphism of graded Lie algebras

$$\Phi : L \rightarrow \text{Coder}(S\mathfrak{a}).$$

- (2) Given X be a Maurer-Cartan element of L , i.e. $X \in L_1$ and $[X, X] = 0$, its image under Φ is a Maurer-Cartan element $\Phi(X)$ of $\text{Coder}(S\mathfrak{a})$. As explained above, a Maurer-Cartan element of $\text{Coder}(S\mathfrak{a})$ which maps 1 to 0 is nothing but an $L_\infty[1]$ -algebra structure on \mathfrak{a} . To ensure that the coderivation $\Phi(X)$ maps 1 to 0, one has to impose the additional condition $P(X) = 0$.

- (3) We now consider the gauge-action introduced in Subsection 3.3 in the present setting. So suppose γ is an element of degree 0 of L . As explained in Subsection 3.3, such an element generates an automorphism of L_1 – since L is just a graded Lie algebra, this reduces to the adjoint action.

The importance of the following lemma is that it implies that the set of Maurer-Cartan elements of an $L_\infty[1]$ -algebra which arises through Voronov’s derived bracket construction comes along with a *second natural group of automorphisms*, which we call the *extended gauge-automorphisms*. The interested reader can find a proof of the lemma in Appendix A.

Lemma 4.9. *Let L be a graded Lie algebra with abelian subalgebra \mathfrak{a} and a projection $L \rightarrow \mathfrak{a}$ whose kernel is a Lie subalgebra. Denote the morphism of graded Lie algebras from [1] by*

$$\Phi : L \rightarrow \text{Coder}(S\mathfrak{a})$$

and fix a Maurer-Cartan element X of L such $P(X) = 0$.

- (1) The set

$$Z(X) := \{\sigma \in L_0 : [\gamma, X] = 0\}$$

is a Lie subalgebra of L_0 which generates an involutive distribution \mathcal{E} on \mathfrak{a}_0 , whose value at $\beta \in \mathfrak{a}_0$ is the image of

$$Z(X) \rightarrow T_\beta \mathfrak{a} = \mathfrak{a}, \quad \sigma \mapsto \widehat{\Phi(\sigma)}(\beta),$$

where $\widehat{\Phi(\sigma)}(\beta)$ is defined to be $e^{-\beta}\Phi(\sigma)(e^\beta)$ and equals $P(e^{[\cdot, \beta]}\sigma)$.

- (2) The distribution \mathcal{E} is tangent to the set of Maurer-Cartan elements $\text{MC}(\mathfrak{a}, \Phi(X))$ of the $L_\infty[1]$ -algebra structure corresponding to $\Phi(X)$.

Remark 4.10. Using Lemma 4.9 we can define an equivalence relation $\sim_{\text{ext-gauge}}$ on the set of Maurer-Cartan elements $\text{MC}(\mathfrak{a}, \Phi(X))$ as follows: two Maurer-Cartan elements β_0 and β_1 are called **extended gauge-equivalent** if there is a one-parameter family σ_t of degree 0 elements of L which commute with X and a one-parameter family β_t of elements of \mathfrak{a}_0 , agreeing with the given ones at $t = 0$ and $t = 1$, such that

$$\frac{\partial}{\partial t}\beta_t = P(\sigma_t) + P([\sigma_t, \beta_t]) + \frac{1}{2!}P([[\sigma_t, \beta_t], \beta_t]) + \frac{1}{3!}P([[[\sigma_t, \beta_t], \beta_t], \beta_t]) + \dots$$

holds for all $t \in [0, 1]$.

We note that families of elements of the form $[X, \gamma_t]$, for $\gamma_t \in \mathfrak{a}_{-1}$, automatically commute with X and hence give rise to extended gauge-equivalences. If we substitute such a family $[X, \gamma_t]$ for σ_t in the above formula, we obtain

$$\frac{\partial}{\partial t}\beta_t = P([X, \gamma_t]) + P([[X, \gamma_t], \beta_t]) + \frac{1}{2!}P([[[X, \gamma_t], \beta_t], \beta_t]) + \frac{1}{3!}P([[[[X, \gamma_t], \beta_t], \beta_t], \beta_t]) + \dots$$

This expression coincides with the defining formula of an (ordinary) gauge-equivalence between the Maurer-Cartan elements β_0 and β_1 , see Definition 3.8 in Subsection 3.3, hence gauge-equivalence implies extended gauge-equivalence.

Now we return to the equivalence relation \sim_{Sym} on the space of coisotropic deformations. If we apply Voronov’s derived bracket construction (see Remark 4.8) to the data

- $L = (\chi^\bullet(E)[1], [-, -])$,
- $\mathfrak{a} = \Gamma(\wedge E)[1]$

- $P : L \rightarrow \mathfrak{a}$ the projection as before,
- $X = \Pi \in \chi^2(E)$ the Poisson bivector field corresponding to ω ,

one recovers Oh and Park's $L_\infty[1]$ -algebra structure on $\Gamma(\wedge E)[1]$ from Subsection 2.3.

By Lemma 4.9 its Maurer-Cartan elements are endowed with a second equivalence relation, arising from the degree 0-elements of $\chi^\bullet(E)[1]$ that commute with the Poisson bivector field. These are exactly the symplectic vector fields. Lemma 4.9 prompts us to repeat the definition of gauge-equivalence from Subsection 3.3, with the Hamiltonian vector fields $X_{\pi^* f_t}$ replaced with any family of symplectic vector fields. However, in order to maintain the link to geometry, we restrict ourselves to symplectic vector fields on E which are fibrewise entire.

Definition 4.11. *Let (U, Ω) be a local symplectic model for the coisotropic submanifold C .*

*Two elements s_0 and s_1 of $\text{Def}_U(C)$ are **extended gauge-equivalent**, $s_0 \sim_{\text{ext-gauge}} s_1$, if there is a one-parameter family $s_t \in \Gamma(U)$, agreeing with s_0 and s_1 at the endpoints, and a family of symplectic, fibrewise entire vector fields X_t on U such that*

$$\frac{\partial}{\partial t}(-s_t) = P(e^{[\cdot, -s_t]} X_t)$$

holds for all $t \in [0, 1]$.

Remark 4.12. We denote the induced equivalence relation on $\text{Def}_U(C)$ by $\sim_{\text{ext-gauge}}$. The proof of Proposition 3.11 goes through mutatis mutandis and we obtain:

Proposition 4.13. *Elements s_0 and s_1 of $\text{Def}_U(C)$ are extended gauge-equivalent if and only if there is a one-parameter family $s_t \in \Gamma(U)$, agreeing with s_0 and s_1 at the endpoints, and a one-parameter family X_t of symplectic and fibrewise entire vector fields on U such that*

$$\frac{\partial}{\partial t} s_t = p_{s_t}^v(X_t|_{\text{graph}(s_t)})$$

holds for all $t \in [0, 1]$.

4.4. Symplectic equivalence = extended gauge-equivalence. Our aim is to compare the two equivalence relations $\sim_{\text{ext-gauge}}$ and \sim_{Sym} on $\text{Def}_U(E)$.

Remark 4.14. The following two results are proved in parallel to Proposition 3.18 and Proposition 3.19. The key point is the following: if we are given a section s of U whose graph is coisotropic, and a closed 1-form β on E , the vector fields $(\omega^\sharp)^{-1}(\pi^* s^* \beta)$ and $(\omega^\sharp)^{-1}(\beta)$ have the same vertical projection onto $E|_{\text{graph}(s)}$ along $T\text{graph}(s)$. As in the proofs of Proposition 3.18 and Propositions 3.19, this fact allows one to replace any family of symplectic isotopies by a family of symplectic isotopies generated by fibrewise entire symplectic vector fields.

Proposition 4.15. *The following implication holds between the equivalence relations on $\text{Def}_U(C)$:*

$$\begin{array}{c} \text{symplectic equivalence} \sim_{\text{Sym}} \\ \Downarrow \\ \text{extended gauge-equivalence} \sim_{\text{ext-gauge}} . \end{array}$$

Under the assumption that C is compact, we can reverse the implications of Proposition 4.15:

Proposition 4.16. *Suppose C is compact coisotropic submanifold. Then the following implication holds for the local symplectic model of C :*

$$\begin{array}{c} \text{extended gauge-equivalence} \sim_{\text{ext-gauge}} \\ \Downarrow \\ \text{symplectic equivalence} \sim_{\text{Sym}} . \end{array}$$

Combining the two previous propositions, we obtain the main result of this section:

Theorem 4.17. *Let C be a compact coisotropic submanifold with local symplectic model (U, Ω) . The equivalence relations on*

$$\text{Def}_U(C) := \{s \in \Gamma(E) : s \text{ is coisotropic and } \text{graph}(s) \subset U\}$$

given by

- symplectic equivalence \sim_{Sym} (Definition 4.1) and
- extended gauge-equivalence $\sim_{\text{ext-gauge}}$ (Definition 4.11, see also Proposition 4.13)

coincide.

As a consequence we have:

Theorem 4.18. *Let C be a compact coisotropic submanifold with local symplectic model (U, Ω) . The bijection*

$$\text{Def}_U(C) \cong \text{MC}_U(\Gamma(\wedge E)[1])$$

descends to a bijection

$$\mathcal{M}_U^{\text{Sym}}(C) := \text{Def}_U(C) / \sim_{\text{Sym}} \cong \text{MC}_U(\Gamma(\wedge E)[1]) / \sim_{\text{ext-gauge}} .$$

4.5. Comparison with Hamiltonian equivalence. In this note we considered both Hamiltonian equivalence (Definition 3.1) and symplectic equivalence (Definition 4.1) of coisotropic submanifolds. Here we summarize some results of Ruan [12] about the relation between these two kinds of equivalence. Ruan considers a restricted class of coisotropic submanifolds, which he calls integral.

Definition 4.19. *A coisotropic submanifold C is **integral** if the leaves of its characteristic foliation \mathfrak{F} are all compact and the set of leaves S admits a smooth structure such that the natural map is a submersion. (In other words: if $C \rightarrow S$ is a smooth fiber bundle with compact fibers.)*

Remark 4.20.

- (1) As Ruan noticed in [12], being integral is not preserved under small deformations inside the space of coisotropic submanifolds. In the following, we restrict attention to the space of coisotropic sections which are integral, and denote them by $\text{Def}_U^{\text{int}}(C)$.
- (2) Recall that every fibre bundle $p : M \rightarrow S$ with compact fibres S inherits a local system \mathbf{H} , given by the fibrewise cohomology, i.e.

$$\mathbf{H}_s := H^\bullet(p^{-1}(s), \mathbb{R}),$$

equipped with the Gauss-Manin connection. The cohomology $H^\bullet(S, \mathbf{H})$ is the second sheet of the Leray-Serre spectral sequence associated to $p : C \rightarrow S$, which converges

to the cohomology of C . We will focus on $\mathbf{H}_s^1 := H^1(p^{-1}(s), \mathbb{R})$. Observe that the differential d_2 of the second sheet gives a natural linear map

$$d_2 : H^0(S, \mathbf{H}^1) \rightarrow H^2(S, \mathbf{H}^0).$$

Notice that the former group is the space of global, flat sections of the vector bundle \mathbf{H}^1 over S . Since the fibres of p are connected, the latter group is just $H^2(S, \mathbb{R})$.

In [12, Theorem 1] Ruan establishes the following result:

Theorem 4.21. *Let C be an integral coisotropic submanifold.*

(1) *There is an open embedding*

$$\text{Def}_U^{\text{int}}(C) / \sim_{\text{Ham}} \hookrightarrow H^0(S, \mathbf{H}^1).$$

(2) *The image of the equivalence class of C with respect to symplectic equivalence \sim_{Sym} under the map $\text{Def}_U^{\text{int}}(C) \rightarrow \text{Def}_U^{\text{int}}(C) / \sim_{\text{gauge}}$ is given nearby C as the kernel of $d_2 : H^0(S, \mathbf{H}^1) \rightarrow H^2(S, \mathbb{R})$.*

Below we reproduce an example from [12]:

Example 4.22. Consider the unit sphere $C = S^3$ in \mathbb{R}^4 , with the canonical symplectic form. The characteristic leaves of S^3 are circles, and $p: S^3 \rightarrow S = S^2$ is the Hopf fibration. \mathbf{H}^1 is a trivial rank 1 vector bundle over S^2 , so $H^0(S, \mathbf{H}^1) \cong \mathbb{R}$, one generator being represented by a connection 1-form on the Hopf fibration. The map $H^0(S, \mathbf{H}^1) \rightarrow H^2(S, \mathbb{R}) \cong \mathbb{R}$ is an isomorphism, reflecting the fact that the connection is not flat.

Hence, by Theorem 4.21, not all nearby integral coisotropic deformations of S^3 are related to C by a symplectomorphism, for instance all spheres of radius r for $r \neq 1$ are not. But those which are, are actually equivalent to C by a Hamiltonian diffeomorphism. The latter statement follows, since $H^1(C) = 0$ implies that all symplectic vector fields in a tubular neighborhood of C are Hamiltonian.

Another example is:

Example 4.23. Consider the 3-torus $C = \mathbb{T}^3$, which “coordinates” $\theta_1, \theta_2, \theta_3$, as the zero section of $(\mathbb{T}^3 \times \mathbb{R}, d\theta_1 \wedge d\theta_2 + d\theta_3 \wedge dx_4)$, where x_4 is the standard coordinate on \mathbb{R} . The characteristic leaves are again circles, and $p: \mathbb{T}^3 \rightarrow S = \mathbb{T}^2$ is the trivial fibration. Again, \mathbf{H}^1 is a trivial rank 1 vector bundle, so $H^0(S, \mathbf{H}^1) \cong \mathbb{R}$, one generator being represented by $d\theta_3$. The map $H^0(S, \mathbf{H}^1) \cong \mathbb{R} \rightarrow H^2(S, \mathbb{R}) \cong \mathbb{R}$ is the zero map, reflecting the fact that $d\theta_3$ is a closed 1-form.

We conclude that all nearby integral coisotropic deformations of C are related to C by a symplectomorphism, but not all of them are related to C by a Hamiltonian diffeomorphism. For instance, the 3-tori given by $\{x_4 = c\}$ for constants $c \neq 0$ are not. Notice that the latter statement is in accordance with the fact that $\mathcal{M}_U^{\text{Sym}}(C) \neq \mathcal{M}_U^{\text{Ham}}(C)$, which is a consequence of Remark 4.6 since the map $r : H^1(C) \rightarrow H_{\mathfrak{F}}^1(C)$ has one-dimensional image.

5. THE TRANSVERSALLY INTEGRABLE CASE

In this section we consider coisotropic submanifolds C that admit a foliation that is complementary to the characteristic foliation:

Definition 5.1. A coisotropic submanifold C of (M, ω) is called **transversally integrable** if the kernel K of the pre-symplectic structure ω_C admits a complementary subbundle G which is involutive.

Remark 5.2. A transversally integrable coisotropic submanifold C comes equipped with two foliations: the characteristic foliation \mathfrak{F} , given by the maximal leaves of K , and another foliation, given by the maximal leaves of G . Since K is the kernel of the pre-symplectic structure on C , the leaves of $G \cong TC/K$ inherit a symplectic structure.

The assumption of transversal integrability leads to many simplifications. We recover a result by Oh and Park [11] that says that the $L_\infty[1]$ -algebra associated to a transversally integrable C is a differential graded Lie algebra (Prop. 5.3). Moreover, we give a formula for the coisotropic section generated by moving the zero section by a basic Hamiltonian flow (Prop. 5.12).

5.1. Oh and Park's $L_\infty[1]$ -algebra. Let C be a coisotropic submanifold and (U, Ω) be a local symplectic model of C as in Section 1. As seen there, the normal model is a neighborhood of the zero section in a vector bundle $E \rightarrow C$, so it comes equipped with a surjective submersion $\pi: U \rightarrow C$.

The following proposition was already proven in [11, Equation (9.17)] (see also Theorem 9.3 there). We provide an alternative proof here.

Proposition 5.3. *Let C be a coisotropic submanifold, and assume there exists an involutive complement G to $K = \ker(\omega_C)$. Then the $L_\infty[1]$ -algebra structure on $\Gamma(\wedge E)[1]$, $E = K^*$, associated to C as in Subsection 2.3 corresponds⁷ to a differential graded Lie algebra.*

Proof. The structure maps λ_r of the $L_\infty[1]$ -algebra from Subsection 2.3 are derivations in each argument. Consequently they can be evaluated locally. Moreover, the derivation property and a degree-count using the fact that Π is a bivector field show that it suffices to evaluate them on tuples of the form

$$(f, g, s_1, \dots, s_{r-2}), \quad (f, s_1, \dots, s_{r-1}) \quad \text{and} \quad (s_1, \dots, s_r)$$

with $f, g \in \mathcal{C}^\infty(C)$ and $s_i \in \Gamma(E)$, seen as vertical vector fields on E , in order to determine them completely.

We now compute the multibracket λ_k of Oh and Park's $L_\infty[1]$ -algebra structure on $\Gamma(\wedge E)[1]$ in local coordinates. As we already noticed, the leaves of the involutive subbundle G complementary to K are symplectic. Choose coordinates $q_1, \dots, q_{n-k}, y_1, \dots, y_{2k}$ on C adapted to the foliations integrating K and G , respectively. That is, K is spanned by the $\frac{\partial}{\partial q}$'s and G is spanned by the $\frac{\partial}{\partial y}$'s. Add conjugate coordinates $p_1, \dots, p_{n-k}, u_1, \dots, u_{2k}$ to obtain a coordinate system on T^*C . The subbundle $G^\circ \subset T^*C$ is locally given by $\{u_1 = \dots = u_{2k} = 0\}$. Hence the symplectic form on $E = K^* \cong G^\circ$ (see Section 1) reads

$$(5) \quad \Omega = \sum_{i=1}^{n-k} dq_i \wedge dp_i + \pi^* \omega_C.$$

⁷That is, the L_∞ -algebra obtained after applying the degree shift operator $[-1]$ is a differential graded Lie algebra, i.e. the structure maps λ_k vanish for $k > 2$.

Notice that, in coordinates, ω_C has the form $\sum h_{jl} dy_j \wedge dy_l$ for some functions h_{jl} on C . Notice further that Ω (and therefore the Poisson bivector field Π obtained by inverting Ω) are invariant under all of the vertical vector fields $\frac{\partial}{\partial p_1}, \dots, \frac{\partial}{\partial p_{n-k}}$. The structure maps λ_r are determined by their evaluation on tuples of the form

$$(f, g, \frac{\partial}{\partial q_{i_1}}, \dots, \frac{\partial}{\partial q_{i_{r-2}}}), \quad (f, \frac{\partial}{\partial q_{i_1}}, \dots, \frac{\partial}{\partial q_{i_{r-1}}}) \quad \text{and} \quad (\frac{\partial}{\partial q_{i_1}}, \dots, \frac{\partial}{\partial q_{i_r}}).$$

Consider the term on the right-hand side of Equation (2) in Subsection 2.3 before applying the projection P , that is,

$$[[\dots [\Pi, -], -] \dots], -].$$

As we argued above, it suffices to evaluate this expression on tuples consisting of functions on C and vertical vector fields $\frac{\partial}{\partial q_i}$. Since π^*f and Ω is invariant under any of the vertical vector fields, $\frac{\partial}{\partial q_i}$, the structure map λ_r vanish whenever we evaluate it on a tuple that contains a $\frac{\partial}{\partial q_i}$. Hence, only λ_1 and λ_2 can be non-zero. \square

Remark 5.4. The non-trivial structure maps of the differential graded Lie algebra associated to a transversally integrable coisotropic submanifold are given by

$$\lambda_1(f) = \pm P(X_{\pi^*f}) \quad \text{and} \quad \lambda_2(f, g) = \pm \{f, g\}^G,$$

the fact that λ_1 and λ_2 annihilate the coordinate vector fields $\frac{\partial}{\partial q}$ associated to adapted coordinates on C , and the derivation rule. Here, $\{\cdot, \cdot\}^G$ denotes the leafwise Poisson structure associated to the symplectic foliation integrating G .

5.2. Hamiltonian equivalences. We want to be more explicit about lifting constructions from a coisotropic submanifold C to its local symplectic model (U, Ω) . To this end, the concept of a partial Ehresmann connection will be of great importance.

Definition 5.5. Let G be an involutive distribution on C . Suppose $\pi : U \rightarrow C$ is a surjective submersion. A **partial Ehresmann connection** on U is a choice of a complementary subbundle G to K and a subbundle G^\sharp of TU such that the differential of $d_x\pi$ at $x \in U$ maps G_x^\sharp isomorphically onto $G_{\pi(x)}$.

Remark 5.6. We notice that the last condition implies that G^\sharp is complementary to $(d\pi)^{-1}(K)$.

Lemma 5.7. Let C be a coisotropic submanifold with local symplectic model (U, Ω) . Suppose G is the subbundle complementary to the kernel K of the pre-symplectic form ω_C which was chosen in the construction of (U, Ω) .

- (1) The subbundles $(d\pi)^{-1}G$ and $V = \ker(d\pi)$ of TU are symplectically orthogonal to each other.
- (2) The bundle

$$G^\sharp := ((d\pi)^{-1}(K))^\perp$$

defines a partial Ehresmann connection on U .

Proof. We take $\xi \in \ker(d_x\pi)$ and $v \in (d_x\pi)^{-1}(G)$. Plugging the two vectors into the symplectic form ω yields

$$\Omega_x(\xi, v) = \omega_C(d\pi(\xi), d\pi(v)) + \omega_{T^*C}(d_xj(\xi), d_xj(v)) = 0.$$

Since the ranks of the two subbundles add up to the rank of TU , the first claim follows.

Concerning (2), the inclusion $\ker(d_x\pi) \subset (d_x\pi)^{-1}(K)$ implies

$$((d_x\pi)^{-1}(K))^\perp \subset (\ker(d_x\pi))^\perp = (d_x\pi)^{-1}(G),$$

i.e. G^\sharp maps indeed onto G under $d\pi$. To check that the map is an isomorphism, it suffices to check that the dimensions match, which is straight-forward. \square

Remark 5.8. The partial Ehresmann connection was first considered in [11], see Equation (6.3) there. Observe that G^\sharp is usually not linear, i.e. compatible with the linear structure on $E \supset U$.

Notice further that a partial Ehresmann connection allows one to define the horizontal lift X^{hor} of a section X of G : Just define X^{hor} at a point $x \in U$ to be the pre-image of $X_{\pi(x)}$ under the isomorphism $d_x\pi : G_x^\sharp \rightarrow G_x$. If G is involutive, it makes sense to ask whether the assignment

$$\Gamma(G) \rightarrow \mathcal{X}(U), \quad X \mapsto X^{\text{hor}}$$

is compatible with the Lie bracket of vector fields. If this is the case one calls G^\sharp flat.

Proposition 5.9. *Let C be a coisotropic submanifold that is transversally integrable, with G an involutive transversal. Let (U, Ω) be the corresponding local symplectic model.*

- (1) *The partial connection G^\sharp on $U \subset E$ is linear and flat.*
- (2) *For all $f \in C^\infty(C)$, we have*

$$X_{\pi^*f} = P(X_{\pi^*f}) + (X_f^G)^{\text{hor}}$$

where:

- (i) *$P(X_{\pi^*f}) \in \Gamma(E)$ is seen as a vertical vector field on $U \subset E$, constant along the fibers,*
- (ii) *X_f^G denotes the leafwise Hamiltonian vector field of f with respect to the symplectic foliation integrating G and*
- (iii) *$(X_f^G)^{\text{hor}}$ denotes the horizontal lift of X_f^G with respect to the partial Ehresmann connection G^\sharp .*

Proof. Choose coordinates $y_1, \dots, y_{2k}, q_1, \dots, q_{n-k}, p_1, \dots, p_{n-k}$ on U as in the proof of Proposition 5.3.

(1) Equation (5) shows that at every point $x \in U$, Ω_x is the sum of two symplectic forms, one defined on the subspace spanned by the $\frac{\partial}{\partial p}$ and $\frac{\partial}{\partial q}$'s, the other one defined on the subspace spanned by the $\frac{\partial}{\partial y}$'s. As $((d\pi)^{-1}E)^\circ$ is spanned by the dy 's, we obtain

$$(6) \quad G^\sharp = \text{span}\left\{\frac{\partial}{\partial y_1}, \dots, \frac{\partial}{\partial y_{2k}}\right\}.$$

In other words, in the trivialization of the vector bundle $E = K^*$ given by the chosen coordinates, G^\sharp is a trivial partial connection.

From this we deduce that the parallel transport with respect to G^\sharp along paths contained in a leaf of G is by linear isomorphisms between the fibers of E , showing that the partial connection G^\sharp is linear. Second, the linear partial connection G^\sharp is flat, since the distribution G^\sharp is clearly involutive.

(2) In the above coordinates, by Equation (5), we have

$$X_{\pi^*f} = [\Pi, \pi^*f] = \sum_i \frac{\partial f}{\partial q_i} \frac{\partial}{\partial p_i} + (X_f^G)^{\text{hor}},$$

where for the horizontal component we used (6). Its vertical component is invariant under each of the vertical vector fields $\frac{\partial}{\partial p_1}, \dots, \frac{\partial}{\partial p_{n-k}}$, therefore it agrees with the vertical component at $\pi(x) \in C$, which is $P(X_{\pi^*f})$. \square

Remark 5.10. Our next aim is to explicitly describe the sections of E which are Hamiltonian equivalent to the zero section $\iota : C \rightarrow E$. If C is compact, we can replace Hamiltonian equivalence by base Hamiltonian equivalence, see Definition 3.17 and Propositions 3.18 and 3.19. Recall that this means that we have to consider the time one flow of a time-dependent vector field $X_{\pi^*f_t}$ where $f_t \in C^\infty(C)$. Such vector fields are not vertical in general, hence solving explicitly the ODE to find their flow is not easy. We are able to do so when G is involutive, making use of the following result:

Lemma 5.11. *Let $A \rightarrow M$ be a vector bundle with a linear connection ∇ . Let $(X_t)_{t \in [0,1]}$ be a one-parameter family of vector fields on M , and $(\alpha_t)_{t \in [0,1]}$ a one-parameter family of sections of A . Consider the one-parameter family of vector fields on A given by*

$$\alpha_t + (X_t)^{\text{hor}}$$

where α_t is viewed as a vertical vector field which is constant along the fibers of A , and $(X_t)^{\text{hor}}$ is the horizontal lift of X_t with respect to the connection ∇ . The integral curve of $\alpha_t + (X_t)^{\text{hor}}$ starting at $q \in C$ is given by

$$s(t) = \int_0^t \begin{array}{c} \gamma(t) \\ \gamma(\tau) \end{array} \parallel [\alpha_\tau |_{\gamma(\tau)}] d\tau \in A_{\gamma(t)},$$

where \parallel denotes the parallel transport with respect to ∇ along the curve $\gamma(t) := \psi_t(q)$, and $\psi_t : C \rightarrow C$ the flow of X_t .

Proof. We have $s = A \circ \Delta$, where $\Delta : [0, 1] \rightarrow [0, 1]^2$ is the diagonal map and

$$A(r, t) := \int_0^r \begin{array}{c} \gamma(t) \\ \gamma(\tau) \end{array} \parallel [\alpha_\tau |_{\gamma(\tau)}] d\tau.$$

Hence $\frac{\partial}{\partial t}|_{t_0} s(t) = \frac{\partial}{\partial t}|_{t_0} A(t, t_0) + \frac{\partial}{\partial t}|_{t_0} A(t_0, t)$ can be written as the sum of two terms, for all $t_0 \in [0, 1]$. The first one is the vertical vector

$$\frac{\partial}{\partial t}|_{t_0} \int_0^{t_0} \begin{array}{c} \gamma(t_0) \\ \gamma(\tau) \end{array} \parallel [\alpha_\tau |_{\gamma(\tau)}] d\tau = \alpha_{t_0} |_{\gamma(t_0)},$$

as can be seen noticing that the integrand is a curve in $A_{\gamma(t_0)}$, parametrized by s , and applying the fundamental theorem of calculus.

For the second term, we claim that

$$(7) \quad \frac{\partial}{\partial t}|_{t_0} \int_0^{t_0} \begin{array}{c} \gamma(t) \\ \gamma(\tau) \end{array} \parallel [\alpha_\tau |_{\gamma(\tau)}] d\tau = (X_{t_0})^{\text{hor}} |_{s(t_0)}.$$

Notice that the integral on the left-hand side of Equation (7) is an element of the fiber of V over $\gamma(t)$, hence applying $\frac{\partial}{\partial t}|_{t_0}$ we obtain an element of $T_{s(t_0)}A$ that projects to $\frac{\partial}{\partial t}|_{t_0} \gamma(t) =$

$X_{t_0}|_{\gamma(t_0)}$ under π . We now argue that the left-hand side of Equation (7) is a horizontal lift, which would conclude the statement. Let $r, t \in [0, 1]$. Under the identification $A_{\gamma(r)} \cong A_{\gamma(t)}$ given by the parallel transport $\gamma(r) \parallel \gamma(t)$, the elements $\gamma(r) \parallel [\alpha_r|_{\gamma(r)}]$ and $\gamma(t) \parallel [\alpha_t|_{\gamma(t)}]$ agree for every τ . The same holds for the integral from $\tau = 0$ to $\tau = t_0$ of these elements, since parallel transport is a linear isomorphism. Hence the integral on the left-hand side of (7), as t varies, defines a parallel section of A over γ . Hence applying $\frac{\partial}{\partial t}|_{t_0}$ we obtain an horizontal element of $T_{s(t_0)}A$. \square

Proposition 5.12. *Let C be a coisotropic submanifold that is transversally integrable, with G an involutive transversal. Let (U, Ω) be the corresponding local symplectic model. Take a one-parameter family $(f_t)_{t \in [0,1]} \in C^\infty(C)$, and denote by Φ the time-1 flow of the time-dependent vector field $(X_{\pi^* f_t})_{t \in [0,1]}$. Then $\Phi(C)$ is the graph of the following section of $U \subset E$:*

$$p \mapsto \int_0^1 \frac{\sigma(1)}{\sigma(t)} \parallel [P(X_{\pi^* f_t})|_{\sigma(t)}] dt$$

where \parallel denotes the parallel transport with respect to the partial connection G^\sharp along the curve $\sigma(t) := \psi_t((\psi_1)^{-1}p)$, for $\psi_t: C \rightarrow C$ the flow of $X_{f_t}^G$.

Proof. By Proposition 5.9, G^\sharp is a partial linear connection on $U \subset E$, and $X_{\pi^* f_t} = P(X_{\pi^* f_t}) + (X_{f_t}^G)^{\text{hor}}$. We note that, in particular, this vector field covers $X_{f_t}^G$, which is tangent to the leaves of G . Fix $p \in C$, and let $L \subset C$ be the leaf of G through p . Consider the vector bundle $E|_L \rightarrow L$, equipped with the linear connection obtained by restricting G^\sharp . We apply Lemma 5.11 to the one-parameter family of vector fields $(X_{f_t}^G)|_L$ and to the one-parameter family of sections $P(X_{\pi^* f_t})|_L$. Choosing the point q so that $\psi_1(q) = p$ and setting $t = 1$ finishes the proof. \square

Remark 5.13.

- (1) We observe that Propositions 5.9 and 5.12 continue to hold for symplectomorphisms, i.e. one obtains explicit formulae for the symplectic vector field associated to a closed 1-form obtained via pull-back from the base C , as well as for the image of C under the flow of such a vector field.
- (2) When C is Lagrangian, U is open in the cotangent bundle T^*C , hence $X_{\pi^* f_t}$ is a (constant) vertical vector field and $P(X_{\pi^* f_t}) = df_t$. Further $G = \{0\}$, so the curve σ through p is constant. Therefore we recover the well-known result that $\Phi(C)$ is the graph of the exact one-form $d(\int_0^1 f_t dt)$.

We exemplify the above discussion in the case of C hypersurface, i.e. of co-dimension 1. While all smooth deformations of a co-dimension 1 submanifold are automatically coisotropic, it turns out that the equivalence problem is non-trivial.

Example 5.14. Fix a codimension 1 compact submanifold C of (M, ω) , which we assume to be oriented. The annihilator $TC^\circ \cong K$ is a trivial line bundle, so there is $\alpha \in \Omega^1(C)$ such that $G := \ker(\alpha)$ satisfies $G \oplus E = TC$. As usual K is the characteristic distribution of C , i.e., $K := \ker(\omega_C)$. We assume that $d\alpha = 0$, which in particular implies that G is

involutive. By [9, Exercise 3.36] a tubular neighborhood of C in M is symplectomorphic to

$$(U, \Omega) := (C \times I, \pi^* \omega_C - du \wedge \pi^* \alpha),$$

where I is an open interval containing 0, u the standard coordinate on I , and $\pi: C \times I \rightarrow C$ is the projection. In the following we denote by $\hat{\xi}$ the unique vector field on C lying in K such that $\alpha(\hat{\xi}) = 1$.

Take a one-parameter family $f_t \in C^\infty(C)$, and denote by Φ the time-1 flow of the vector field $(X_{\pi^* f_t})_{t \in [0,1]}$. Then $\Phi(C)$ is the graph of

$$s: C \rightarrow \mathbb{R}, \quad s(p) = - \int_0^1 \hat{\xi}(f_t)|_{\psi_t((\psi_1)^{-1}p)} dt$$

where $\psi_\tau: C \rightarrow C$ is the flow of $(X_{f_t}^G)_{t \in [0,1]}$. This follows from Prop. 5.12, since $X_{\pi^* f} = -\hat{\xi}(f_t) \frac{\partial}{\partial u} + X_f^G$ at points of C , and G^\sharp is the trivial partial connection by Equation (6).

APPENDIX A. THE EXTENDED GAUGE-ACTION

We prove Lemma 4.9.

We first observe that since Φ is a morphism of graded Lie algebras, and since we assume $[\sigma, X] = 0$, $\Phi(\sigma)$ and $\Phi(X)$ will commute as well. In particular, the adjoint action by $\Phi(\sigma)$ leaves $\Phi(X)$ invariant.

(1) It is immediate that $Z(X)$ is a Lie subalgebra of L_0 .

We now show that $\widehat{\Phi(\sigma)}(\beta)$ equals $P(e^{[\cdot, \beta]}\sigma)$. We already know that $\widehat{\Phi(\sigma)}(\beta) = e^{-\beta} \Phi(\sigma)(e^\beta)$ is an element of \mathfrak{a} . To compute it, it suffices to compute the component of $\Phi(\sigma)(e^\beta)$ in \mathfrak{a} . By construction of Φ , $\Phi(\sigma)$ is the coderivation corresponding to the derived bracket construction with σ . Hence the component of $\Phi(\sigma)(e^\beta)$ in \mathfrak{a} is

$$P(\sigma) + P([\sigma, \beta]) + \frac{1}{2!} P([[\sigma, \beta], \beta]) + \frac{1}{3!} P([[[\sigma, \beta], \beta], \beta]) + \dots = P(e^{[\cdot, \beta]}\sigma).$$

To check the involutivity of \mathcal{E} , recall that a vector field X on a vector space V can be seen as a map from V to itself. In the case at hand $V = \mathfrak{a}_0$ and the vector fields we are interested in correspond to the maps $f(\beta) := \widehat{\Phi(\sigma)}(\beta)$ and $g(\beta) := \widehat{\Phi(\sigma')}\beta$, respectively.

The Lie bracket of two such vector fields is given by

$$(8) \quad (d_\beta g)(f(\beta)) - (d_\beta f)(g(\beta)).$$

A formal computation yields

$$\begin{aligned} (d_\beta g)(\xi) &= -\xi e^{-\beta} \Phi(\sigma')(e^\beta) + e^{-\beta} \Phi(\sigma')(\xi e^\beta) \\ &= -\xi g(\beta) + e^{-\beta} \Phi(\sigma')(\xi e^\beta) \end{aligned}$$

and we arrive at

$$(d_\beta g)(f(\beta)) = -f(\beta)g(\beta) + e^{-\beta} \Phi(\sigma')(f(\beta)e^\beta).$$

For the Lie bracket (8) of the vector fields corresponding to f and g we obtain

$$e^{-\beta} \left(\Phi(\sigma')(f(\beta)e^\beta) - \Phi(\sigma)(g(\beta)e^\beta) \right),$$

which equals $e^{-\beta}([\Phi(\sigma'), \Phi(\sigma)])(e^\beta) = \widehat{\Phi([\sigma', \sigma])}(\beta)$.

- (2) We have to prove that for $\beta \in \text{MC}(\mathfrak{a}, \Phi(X))$, the element $\widehat{\Phi(\sigma)}(\beta) \in \mathfrak{a}_0 = T_\beta \mathfrak{a}_0$ is contained in the formal tangent space

$$T_\beta \text{MC}(\mathfrak{a}, \Phi(X))$$

at β to the set of Maurer-Cartan elements of \mathfrak{a} , equipped with the $L_\infty[1]$ -algebra structure corresponding to $\Phi(X)$. This formal tangent space is given by

$$T_\beta \text{MC}(\mathfrak{a}, \Phi(X)) = \{\xi \in \mathfrak{a}_0 : \Phi(X)(\xi e^\beta) = 0\}$$

and can be written more explicitly as the set of those $\xi \in \mathfrak{a}_0$ such that

$$P(e^{[-, \beta]}([X, \xi])) = P([X, \xi]) + P([[X, \xi], \beta]) + \frac{1}{2}P([[[X, \xi], \beta], \beta]) + \dots$$

vanishes. Hence it is sufficient to compute

$$\begin{aligned} \Phi(X) \left(\widehat{\Phi(\sigma)}(\beta) e^\beta \right) &= \Phi(X) \left(e^{-\beta} \Phi(\sigma)(e^\beta) e^\beta \right) \\ &= \Phi(X) \left(\Phi(\sigma)(e^\beta) \right) = \Phi(\sigma) \left(\Phi(X)(e^\beta) \right) = 0, \end{aligned}$$

where in the first equality we used the definition of $\widehat{\Phi(\sigma)}$, in the second equality we used the commutativity of the product in $S\mathfrak{a}$ and in the last equality we used that $\Phi(X)$ and $\Phi(\sigma)$ commute.

REFERENCES

- [1] R. Bandiera. Non-abelian higher derived brackets. 04 2013, ArXiv:1304.4097.
- [2] A. S. Cattaneo and G. Felder. Coisotropic submanifolds in Poisson geometry and branes in the Poisson sigma model. *Lett. Math. Phys.*, 69:157–175, 2004.
- [3] A. S. Cattaneo and G. Felder. Relative formality theorem and quantisation of coisotropic submanifolds. *Adv. in Math.*, 208:521–548, 2007.
- [4] M. J. Gotay. On coisotropic imbeddings of presymplectic manifolds. *Proc. Amer. Math. Soc.*, 84(1):111–114, 1982.
- [5] A. Kapustin and D. Orlov. Remarks on A-branes, mirror symmetry, and the Fukaya category. *J. Geom. Phys.*, 48(1):84–99, 2003.
- [6] T. Lada and J. Stasheff. Introduction to SH Lie algebras for physicists. *Internat. J. Theoret. Phys.*, 32(7):1087–1103, 1993.
- [7] H. V. Lê and Y.-G. Oh. Deformations of coisotropic submanifolds in locally conformal symplectic manifolds. 08 2012, ArXiv:1208.3590.
- [8] H. V. Lê, Y.-G. Oh, A. G. Tortorella, and L. Vitagliano. Deformations of Coisotropic Submanifolds in Abstract Jacobi Manifolds. 10 2014, ArXiv:1410.8446.
- [9] D. McDuff and D. Salamon. *Introduction to symplectic topology*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, second edition, 1998.
- [10] J. W. Milnor and J. C. Moore. On the structure of Hopf algebras. *Ann. of Math. (2)*, 81:211–264, 1965.
- [11] Y.-G. Oh and J.-S. Park. Deformations of coisotropic submanifolds and strong homotopy Lie algebroids. *Invent. Math.*, 161(2):287–360, 2005.
- [12] W.-D. Ruan. Deformation of integral coisotropic submanifolds in symplectic manifolds. *J. Symplectic Geom.*, 3(2):161–169, 2005.
- [13] F. Schätz. BFV-complex and higher homotopy structures. *Comm. Math. Phys.*, 286(2):399–443, 2009.
- [14] F. Schätz. Moduli of coisotropic sections and the BFV-complex. *Asian J. Math.*, 15(1):71–100, 2011.
- [15] F. Schätz and M. Zambon. Deformations of coisotropic submanifolds for fibrewise entire Poisson structures. *Lett. Math. Phys.*, 103(7):777–791, 2013.

- [16] T. Voronov. Higher derived brackets and homotopy algebras. *J. Pure Appl. Algebra*, 202(1-3):133–153, 2005.
- [17] T. T. Voronov. Higher derived brackets for arbitrary derivations. In *Travaux mathématiques. Fasc. XVI*, Trav. Math., XVI, pages 163–186. Univ. Luxemb., Luxembourg, 2005.
- [18] M. Zambon. An example of coisotropic submanifolds C^1 -close to a given coisotropic submanifold. *Differential Geom. Appl.*, 26(6):635–637, 2008.
E-mail address: `florian.schaetz@gmail.com`

CENTRE FOR QUANTUM GEOMETRY OF MODULI SPACES, AARHUS UNIVERSITY, NY MUNKEGADE 118, DK-8000 AARHUS C, DENMARK.

E-mail address: `marco.zambon@uam.es`, `marco.zambon@icmat.es`, `marco.zambon@wis.kuleuven.be`

UNIVERSIDAD AUTÓNOMA DE MADRID (DEPARTAMENTO DE MATEMÁTICAS) AND ICMAT(CSIC-UAM-UC3M-UCM), CAMPUS DE CANTOBLANCO, 28049 - MADRID, SPAIN. CURRENT ADDRESS: KU LEUVEN, DEPARTMENT OF MATHEMATICS, CELESTIJNENLAAN 200B BOX 2400, BE-3001 LEUVEN, BELGIUM.