

Products in the category of \mathbb{Z}_2^n -manifolds

Andrew Bruce and Norbert Poncin*

Abstract

We prove that the category of \mathbb{Z}_2^n -manifolds has all finite products. Further, we show that a \mathbb{Z}_2^n -manifold (resp., a \mathbb{Z}_2^n -morphism) can be reconstructed from its algebra of global \mathbb{Z}_2^n -functions (resp., from its algebra morphism between global \mathbb{Z}_2^n -function algebras). These results are of importance in the study of \mathbb{Z}_2^n Lie groups. The investigation is all the more challenging, since the completed tensor product of the structure sheafs of two \mathbb{Z}_2^n -manifolds is not a sheaf. We rely on a number of results on (pre)sheaves of topological algebras, which we establish in the appendix.

Contents

1	Introduction	2
2	\mathbb{Z}_2^n-manifolds and their morphisms	3
3	Reconstructions of \mathbb{Z}_2^n-manifolds and \mathbb{Z}_2^n-morphisms	5
3.1	Reconstruction of the topological base space	5
3.2	Reconstruction of the structure sheaf	7
3.3	Reconstruction of a \mathbb{Z}_2^n -morphism	11
4	Finite products in the category of \mathbb{Z}_2^n-manifolds	13
4.1	Cartesian product of \mathbb{Z}_2^n -manifolds	13
4.2	Fundamental isomorphisms	15
4.3	Categorical products of \mathbb{Z}_2^n -manifolds	23
4.4	Products of \mathbb{Z}_2^n -morphisms	27
5	Appendix	29
5.1	Topological vector subspaces	29
5.2	Completions of topological vector spaces	30
5.3	Locally convex spaces	30
5.4	Presheaves of topological algebras and sheafification	31
5.5	Direct image and sheafification	35

*University of Luxembourg, Mathematics Research Unit, 4364 Esch-sur-Alzette, Luxembourg, andrew.bruce@uni.lu, norbert.poncin@uni.lu

1 Introduction

\mathbb{Z}_2^n -Geometry is an emerging framework in mathematics and mathematical physics, which has been introduced in the foundational papers [CGP16a] and [COP12]. This non-trivial extension of standard Supergeometry allows for \mathbb{Z}_2^n -gradings, where

$$\mathbb{Z}_2^n = \mathbb{Z}_2^{\times n} = \mathbb{Z}_2 \times \dots \times \mathbb{Z}_2 \quad \text{and} \quad n \in \mathbb{N}.$$

The corresponding \mathbb{Z}_2^n -commutation rule for coordinates $(u^A)_A$ with degrees $\deg u^A \in \mathbb{Z}_2^n$ does not use the product of the (obviously defined) parities, but the scalar product $\langle -, - \rangle$ of \mathbb{Z}_2^n :

$$u^A u^B = (-1)^{\langle \deg u^A, \deg u^B \rangle} u^B u^A. \quad (1)$$

A brief description of the category $\mathbb{Z}_2^n \mathbf{Man}$ of \mathbb{Z}_2^n -manifolds $\mathcal{M} = (M, \mathcal{O}_M)$ and morphisms $\Phi = (\phi, \phi^*)$ between them can be found in Section 2. For $n = 1$, one recovers the category \mathbf{SMan} of supermanifolds. A survey on \mathbb{Z}_2^n -Geometry is available in [Pon16]. The differential calculus and the splitting theorem for \mathbb{Z}_2^n -manifolds have been investigated in [CKP16] and [CGP16b], respectively. In the introduction of [BP18], the reader finds motivations for the study of \mathbb{Z}_2^n -Geometry. The present paper uses the main results of [BP18].

Applications of \mathbb{Z}_2^n -Geometry are based in particular on \mathbb{Z}_2^n Lie groups and their actions on \mathbb{Z}_2^n -manifolds (supergravity), on \mathbb{Z}_2^n vector bundles and their sections (\mathbb{Z}_2^n Lie algebroids), on the internal Hom functor in $\mathbb{Z}_2^n \mathbf{Man}$ (\mathbb{Z}_2^n -gradings and \mathbb{Z}_2^n -parities in field theory), ... All these notions rely themselves on *products* in the category $\mathbb{Z}_2^n \mathbf{Man}$. On the other hand, a comparison of different approaches to \mathbb{Z}_2^n vector bundles is more challenging than in the supercase [BCC11]. A generalization to \mathbb{Z}_2^n -manifolds of the Schwarz-Voronov-Molotkov embedding is needed. This extension, which embeds $\mathbb{Z}_2^n \mathbf{Man}$ into the category of contravariant functors from \mathbb{Z}_2^n -points to a specific category of Fréchet manifolds, uses the *reconstructions* of \mathbb{Z}_2^n -manifolds and \mathbb{Z}_2^n -morphisms from the \mathbb{Z}_2^n -commutative associative unital \mathbb{R} -algebras of global \mathbb{Z}_2^n -functions and the \mathbb{Z}_2^n -graded unital algebra morphisms between them, respectively.

The existence of categorical products and the mentioned reconstruction theorems are the main topics of the present paper. The text is organized as follows. Section 3 contains the proofs of the above-mentioned \mathbb{Z}_2^n reconstruction results. The definition of a product \mathbb{Z}_2^n -manifold and the proof of its meaningfulness are rather obvious, see Definition 13. However, the proof of the existence of categorical products in $\mathbb{Z}_2^n \mathbf{Man}$ is quite tough. It relies on the generalization of the well-known isomorphism of topological vector spaces

$$C^\infty(\Omega') \widehat{\otimes} C^\infty(\Omega'') \simeq C^\infty(\Omega' \times \Omega'')$$

(for open subsets $\Omega' \subset \mathbb{R}^p, \Omega'' \subset \mathbb{R}^r$) to an isomorphism of locally convex topological algebras of formal power series

$$C^\infty(\Omega')[[\xi]] \widehat{\otimes} C^\infty(\Omega'')[[\eta]] \simeq C^\infty(\Omega' \times \Omega'')[[\xi, \eta]] \quad (2)$$

(for \mathbb{Z}_2^n -domains $\mathcal{U}^{p|\mathbf{q}} = (\Omega', C_{\Omega'}^\infty[[\xi]])$ and $\mathcal{V}^{r|\mathbf{s}} = (\Omega'', C_{\Omega''}^\infty[[\eta]])$, with

$$\xi = (\xi_1, \dots, \xi_{N(\mathbf{q})}) \quad \text{and} \quad \eta = (\eta_1, \dots, \eta_{N(\mathbf{s})}),$$

see Theorem 14. The issue here is the formal power series, which replace the polynomials of standard Supergeometry. Moreover, if $\mathcal{M} = (M, \mathcal{O}_M)$ and $\mathcal{N} = (N, \mathcal{O}_N)$ are two \mathbb{Z}_2^n -manifolds, and $\mathcal{M} \times \mathcal{N} = (M \times N, \mathcal{O}_{M \times N})$ is their product \mathbb{Z}_2^n -manifold, one gets from (2) that, for an open subset $u \times v \subset M \times N$ of the basis \mathfrak{B} made of products of \mathbb{Z}_2^n -chart domains, we have

$$\mathcal{E}(u \times v) \simeq \mathcal{O}_{M \times N}(u \times v) ,$$

where \mathcal{E} is the \mathfrak{B} -presheaf

$$\mathcal{E}(u \times v) = \mathcal{O}_M(u) \widehat{\otimes} \mathcal{O}_N(v) .$$

Let now $\bar{\mathcal{F}}$ be the standard extension of the \mathfrak{B} -presheaf \mathcal{F} that assigns to any open subset $U \times V \subset M \times N$ (where $U \subset M$ and $V \subset N$ are not necessarily chart domains) the algebra

$$\mathcal{F}(U \times V) = \mathcal{O}_M(U) \widehat{\otimes} \mathcal{O}_N(V) .$$

The presheaf $\bar{\mathcal{F}}$ and the sheaf $\mathcal{O}_{M \times N}$ are thus two extensions of the \mathfrak{B} -presheaf \mathcal{E} . However, this does not mean that $\bar{\mathcal{F}} \simeq \mathcal{O}_{M \times N}$ and that $\bar{\mathcal{F}}$ is a sheaf. Indeed, \mathfrak{B} -sheaves have unique extensions, but \mathfrak{B} -presheaves do not. Also the reconstruction results mentioned above do not allow us to prove that $\bar{\mathcal{F}}$ is a sheaf. Hence, we prove the next best result, i.e., the existence of an isomorphism of sheaves of algebras

$$\mathcal{O}_{M \times N} \simeq \bar{\mathcal{F}}^+ \tag{3}$$

between the structure sheaf of the product \mathbb{Z}_2^n -manifold and the sheafification of the presheaf $\bar{\mathcal{F}}$, see Theorem 16. In the case $n = 1$, we thus recover the definition of a product supermanifold used in [BBH91]. The isomorphism (3) allows us to prove the existence of all finite categorical products in $\mathbb{Z}_2^n \mathbf{Man}$, see Theorem 19. The proof uses the results on sheafification and presheaves of locally convex topological algebras proven in Subsections 5.4 and 5.5 of the Appendix. Products of \mathbb{Z}_2^n -morphisms are obtained from the universality property of categorical products. They are explicitly described in Proposition 20.

2 \mathbb{Z}_2^n -manifolds and their morphisms

We denote by \mathbb{Z}_2^n the cartesian product of n copies of \mathbb{Z}_2 . Further, we use systematically the following *standard order* of the 2^n elements of \mathbb{Z}_2^n : first the even degrees are ordered lexicographically, then the odd ones are also ordered lexicographically. For example,

$$\mathbb{Z}_2^3 = \{(0, 0, 0), (0, 1, 1), (1, 0, 1), (1, 1, 0), (0, 0, 1), (0, 1, 0), (1, 0, 0), (1, 1, 1)\} .$$

A \mathbb{Z}_2^n -domain has, aside from the usual coordinates $x = (x^1, \dots, x^p)$ of degree $\deg x^i = 0 \in \mathbb{Z}_2^n$, also formal coordinates or parameters $\xi = (\xi^1, \dots, \xi^Q)$ of non-zero degrees $\deg \xi^a \in \mathbb{Z}_2^n$. These coordinates $u = (x, \xi)$ commute according to the generalized sign rule

$$u^A u^B = (-1)^{\langle \deg u^A, \deg u^B \rangle} u^B u^A , \tag{4}$$

where $\langle -, - \rangle$ denotes the standard scalar product. For instance,

$$\langle (0, 1, 1), (1, 0, 1) \rangle = 1 .$$

Observe that, in contrast with ordinary \mathbb{Z}_2 - or super-domains, even coordinates may anticommute, odd coordinates may commute, and even nonzero degree coordinates are not nilpotent. Of course, for $n = 1$, we recover the classical situation. We denote by p the number of coordinates x^i of degree 0, by q_1 the number of coordinates ξ^a which have the first non-zero degree of \mathbb{Z}_2^n , and so on. We get that way a tuple $\mathbf{q} = (q_1, \dots, q_N) \in \mathbb{N}^N$ with $N := 2^n - 1$. The dimension of the considered \mathbb{Z}_2^n -domain is then given by $p|\mathbf{q}|$. Clearly the Q above is the sum $|\mathbf{q}| = \sum_{i=1}^N q_i$.

We recall the definition of a \mathbb{Z}_2^n -manifold.

Definition 1. A locally \mathbb{Z}_2^n -ringed space is a pair (M, \mathcal{O}_M) made of a topological space M and a sheaf of \mathbb{Z}_2^n -graded \mathbb{Z}_2^n -commutative (in the sense of (4)) associative unital \mathbb{R} -algebras over it, such that at every point $m \in M$ the stalk $\mathcal{O}_{M,m}$ is a local graded ring.

A smooth \mathbb{Z}_2^n -manifold of dimension $p|\mathbf{q}|$ is a locally \mathbb{Z}_2^n -ringed space $\mathcal{M} = (M, \mathcal{O}_M)$, which is locally isomorphic to the smooth \mathbb{Z}_2^n -domain $\mathbb{R}^{p|\mathbf{q}} := (\mathbb{R}^p, C_{\mathbb{R}^p}^\infty[[\xi]])$, and whose underlying topological space M is second-countable and Hausdorff. Sections of the structure sheaf $C_{\mathbb{R}^p}^\infty[[\xi]]$ are formal power series in the \mathbb{Z}_2^n -commutative parameters ξ , with coefficients in smooth functions:

$$C_{\mathbb{R}^p}^\infty(U)[[\xi]] := \left\{ \sum_{\alpha \in \mathbb{N}^{\times |\mathbf{q}|}} f_\alpha(x) \xi^\alpha \mid f_\alpha \in C^\infty(U) \right\} \quad (U \text{ open in } \mathbb{R}^p) .$$

\mathbb{Z}_2^n -morphisms between \mathbb{Z}_2^n -manifolds are just morphisms of \mathbb{Z}_2^n -ringed spaces, i.e., pairs $\Phi = (\phi, \phi^*) : (M, \mathcal{O}_M) \rightarrow (N, \mathcal{O}_N)$ made of a continuous map $\phi : M \rightarrow N$ and a sheaf morphism $\phi^* : \mathcal{O}_N \rightarrow \phi_* \mathcal{O}_M$, i.e., a family of \mathbb{Z}_2^n -graded unital \mathbb{R} -algebra morphisms, which commute with restrictions and are defined, for any open $V \subset N$, by

$$\phi_V^* : \mathcal{O}_N(V) \rightarrow \mathcal{O}_M(\phi^{-1}(V)) .$$

We denote the category of \mathbb{Z}_2^n -manifolds and \mathbb{Z}_2^n -morphisms between them by $\mathbb{Z}_2^n\text{Man}$.

Remark 2. Let us stress that the base space M corresponds to the degree zero coordinates (and not to the even degree coordinates), and let us mention that it can be proven that the topological base space M carries a natural smooth manifold structure of dimension p , that the continuous base map $\phi : M \rightarrow N$ is in fact smooth, and that the algebra morphisms

$$\phi_m^* : \mathcal{O}_{\phi(m)} \rightarrow \mathcal{O}_m \quad (m \in M)$$

between stalks, which are induced by the considered \mathbb{Z}_2^n -morphism $\Phi : \mathcal{M} \rightarrow \mathcal{N}$, respect the unique homogeneous maximal ideals of the local graded rings $\mathcal{O}_{\phi(m)}$ and \mathcal{O}_m .

3 Reconstructions of \mathbb{Z}_2^n -manifolds and \mathbb{Z}_2^n -morphisms

In this section, we reconstruct a \mathbb{Z}_2^n -manifold (M, \mathcal{O}_M) from the \mathbb{Z}_2^n -commutative unital algebra $\mathcal{O}_M(M)$ of global sections of its function sheaf. We also reconstruct a \mathbb{Z}_2^n -morphism

$$\Phi = (\phi, \phi^*) : (M, \mathcal{O}_M) \rightarrow (N, \mathcal{O}_N)$$

from its pullback \mathbb{Z}_2^n -graded unital algebra morphism

$$\phi_N^* : \mathcal{O}_N(N) \rightarrow \mathcal{O}_M(M)$$

between global sections.

3.1 Reconstruction of the topological base space

Algebraic characterizations of spaces can be traced back to I. Gel'fand and A. Kolmogoroff [GK39]. In that paper, compact topological spaces K are characterized by the algebras $C^0(K)$ of continuous functions on them. In particular, the points m of these spaces are identified with the maximal ideals

$$I_m = \{f \in C^0(K) : f(m) = 0\}$$

of these algebras. A similar characterization holds for the points of second countable Hausdorff smooth manifolds.

Let $\mathcal{M} = (M, \mathcal{O}_M)$ be a \mathbb{Z}_2^n -manifold. We denote the maximal spectrum of $\mathcal{O}(M)$ (subscript omitted) by $\text{Spm}(\mathcal{O}(M))$ (we actually consider here the real maximal spectrum, in the sense that the quotient $\mathcal{O}(M)/\mu$ by an ideal μ in the spectrum is isomorphic to the field \mathbb{R} of real numbers). Note that any $m \in M$ induces a map

$$\varepsilon_m : \mathcal{O}(M) \ni f \mapsto (\varepsilon_M f)(m) \in \mathbb{R} ,$$

which is referred to as the *evaluation map* at m and is a \mathbb{Z}_2^n -graded unital \mathbb{R} -algebra morphism

$$\varepsilon_m \in \text{Hom}_{\mathbb{Z}_2^n \text{UAlg}}(\mathcal{O}(M), \mathbb{R}) .$$

The kernel

$$\mu_m = \ker \varepsilon_m = \{f \in \mathcal{O}(M) : (\varepsilon_M f)(m) = 0\} \in \text{Spm}(\mathcal{O}(M))$$

is a maximal ideal. More generally, the kernel of an arbitrary algebra morphism

$$\psi \in \text{Hom}_{\mathbb{Z}_2^n \text{UAlg}}(\mathcal{O}(M), \mathbb{R})$$

is a maximal ideal, since $\mathcal{O}(M)/\ker \psi \simeq \mathbb{R}$. Indeed, to any class $[f]$ in the quotient we can associate the real number $\psi(f)$. This map is well-defined and injective. It is also surjective, since, for any $r \in \mathbb{R}$, the image of $[r \cdot 1_{\mathcal{O}(M)}]$ is r . It follows in particular that any class in the quotient is of type $[r \cdot 1_{\mathcal{O}(M)}]$ for a unique $r \in \mathbb{R}$. We have the following

Proposition 3. *The maps*

$$\mathfrak{b} : M \ni m \mapsto \mu_m \in \text{Spm}(\mathcal{O}(M))$$

and

$$\flat : \text{Hom}_{\mathbb{Z}_2^n \text{UAlg}}(\mathcal{O}(M), \mathbb{R}) \ni \psi \mapsto \ker \psi \in \text{Spm}(\mathcal{O}(M))$$

are 1:1 correspondences.

Proof. To prove that \mathfrak{b} is bijective, consider a maximal ideal $\mu \in \text{Spm}(\mathcal{O}(M))$. The image $\varepsilon_M(\mu) \subset C^\infty(M)$ is a maximal ideal. Indeed, it is an ideal, since the map ε_M is surjective (the short sequence of sheafs [BP18, Equation (3)] is exact for the good reason that it is exact for any open subset of M). To see that it is maximal, assume there is an ideal ν , such that $\varepsilon_M(\mu) \subset \nu \subset C^\infty(M)$, so that $\mu \subset \varepsilon_M^{-1}(\nu) \subset \mathcal{O}_M(M)$. It follows that $\varepsilon_M^{-1}(\nu) = \mu$ or $\varepsilon_M^{-1}(\nu) = \mathcal{O}_M(M)$, and that $\nu = \varepsilon_M(\mu)$ or $\nu = C^\infty(M)$. Hence,

$$\varepsilon_M(\mu) = I_m = \{f \in C^\infty(M) : f(m) = 0\},$$

since any maximal ideal of $C^\infty(M)$ is known to be of type I_m for a unique $m \in M$. Finally, we get

$$\mu \subset \varepsilon_M^{-1}(I_m) = \{f \in \mathcal{O}(M) : (\varepsilon_M f)(m) = 0\} = \mu_m \subset \mathcal{O}(M).$$

Since $\mu_m \neq \mathcal{O}(M)$, we have $\mu = \mu_m$, which proves the bijectivity of \mathfrak{b} . Indeed, if $\mu = \mu_n$, we obtain $\varepsilon_M(\mu_n) \subset I_n \subset C^\infty(M)$, so that

$$I_n = \varepsilon_M(\mu_n) = \varepsilon_M(\mu_m) = I_m,$$

and $m = n$.

Since any $\mu \in \text{Spm}(\mathcal{O}(M))$ reads $\mu = \mu_m = \ker \varepsilon_m = \flat(\varepsilon_m)$, the map \flat is surjective. Let ψ, ϕ be unital algebra morphisms, such that $\ker \psi = \ker \phi = \mu$. For any $f \in \mathcal{O}(M)$, there exists a unique $r \in \mathbb{R}$, such that $[f] = [r \cdot 1_{\mathcal{O}(M)}]$. Thus $\psi(f) = r = \phi(f)$ and $\psi = \phi$, so that \flat is also injective. \square

Proposition 4. *The map*

$$\varepsilon_M^{-1} : \text{Spm}(C^\infty(M)) \ni I_m \mapsto \varepsilon_M^{-1}(I_m) = \mu_m \in \text{Spm}(\mathcal{O}(M))$$

is a homeomorphism with inverse ε_M , both, if the maximal spectra are endowed with their Zariski topology and if they are endowed with their Gel'fand topology. Hence, the Zariski and Gel'fand topologies coincide on $\text{Spm}(\mathcal{O}(M))$. Further, the bijection

$$\mathfrak{b} : M \ni m \mapsto \mu_m \in \text{Spm}(\mathcal{O}(M))$$

is a homeomorphism.

Proof. The maps

$$\varepsilon_M^{-1} : \text{Spm}(C^\infty(M)) \ni I_m \mapsto \varepsilon_M^{-1}(I_m) = \mu_m \in \text{Spm}(\mathcal{O}(M))$$

and

$$\varepsilon_M : \text{Spm}(\mathcal{O}(M)) \ni \mu_m \mapsto \varepsilon_M(\mu_m) = I_m \in \text{Spm}(C^\infty(M))$$

are inverses of each other.

We first equip the spectrum $\text{Spm}(C^\infty(M))$ as usual with the Zariski topology, which is defined by its basis of open subsets $V_{C^\infty}(f)$, $f \in C^\infty(M)$, given by

$$V_{C^\infty}(f) = \{I_m \in \text{Spm}(C^\infty(M)) : f \notin I_m\},$$

and we proceed similarly for $\text{Spm}(\mathcal{O}(M))$. It is straightforwardly checked that, if $f = \varepsilon_M(F)$, we have

$$\varepsilon_M^{-1}(V_{C^\infty}(f)) = V_{\mathcal{O}}(F) \quad \text{and} \quad \varepsilon_M(V_{\mathcal{O}}(F)) = V_{C^\infty}(f).$$

Hence the announced homeomorphism result for the Zariski topologies.

The Gel'fand topology of $\text{Spm}(C^\infty(M))$ is defined by the basis of open subsets $B_{C^\infty}(m, \epsilon; f_1, \dots, f_n)$, indexed by $m \in M$, $\epsilon > 0$, $n \in \mathbb{N}$, and $f_1, \dots, f_n \in C^\infty(M)$, and defined by

$$B_{C^\infty}(m, \epsilon; f_1, \dots, f_n) = \{I_n \in \text{Spm}(C^\infty(M)) : |f_i(n) - f_i(m)| < \epsilon, \forall i\}.$$

The Gel'fand topology of $\text{Spm}(\mathcal{O}(M))$ is defined analogously by

$$B_{\mathcal{O}}(m, \epsilon; F_1, \dots, F_n) = \{\mu_n \in \text{Spm}(\mathcal{O}(M)) : |(\varepsilon_M F_i)(n) - (\varepsilon_M F_i)(m)| < \epsilon, \forall i\},$$

where $F_i \in \mathcal{O}(M)$. If $f_i = \varepsilon_M(F_i)$, we have obviously

$$\varepsilon_M^{-1}(B_{C^\infty}(m, \epsilon; f_1, \dots, f_n)) = B_{\mathcal{O}}(m, \epsilon; F_1, \dots, F_n),$$

and similarly for ε_M , so that the homeomorphism result holds also for the Gel'fand topologies.

Since the Zariski and Gel'fand topologies coincide on $\text{Spm}(C^\infty(M))$, it follows from the above that there is a homeomorphism from $\text{Spm}(\mathcal{O}(M))$ endowed with the Zariski topology to itself endowed with the Gel'fand topology.

It is well-known that the map $\mathfrak{b}_{C^\infty} : M \ni m \mapsto I_m \in \text{Spm}(C^\infty(M))$ is a homeomorphism. Hence, the bijection $\mathfrak{b} = \varepsilon_M^{-1} \circ \mathfrak{b}_{C^\infty}$ is a homeomorphism as well. \square

3.2 Reconstruction of the structure sheaf

Proposition 5. *Let (M, \mathcal{O}_M) be a \mathbb{Z}_2^n -manifold and let $U \subset M$ be open. A \mathbb{Z}_2^n -function $F \in \mathcal{O}_M(U)$ is invertible if and only if its base projection $f = \varepsilon_U(F) \in C_M^\infty(U)$ is invertible.*

Proof. It is obvious that f is invertible if F is. Assume now that there exists $f^{-1} \in C^\infty(U)$ and consider a cover of U by \mathbb{Z}_2^n -chart domains V_i . For any i , we have $f^{-1}|_{V_i} = (f|_{V_i})^{-1}$, i.e., the base function $\varepsilon_{V_i}(F|_{V_i}) = \varepsilon_U(F)|_{V_i}$ is invertible in $C^\infty(V_i)$, so that $F|_{V_i} \in \mathcal{O}(V_i) \simeq C^\infty(V_i)[[\xi]]$ has an inverse $G_{V_i} \in \mathcal{O}(V_i)$. It follows that, for any V_i and V_j with intersection V_{ij} ,

$$G_{V_i}|_{V_{ij}} = (F|_{V_{ij}})^{-1} = G_{V_j}|_{V_{ij}} .$$

Hence, there is a unique \mathbb{Z}_2^n -function $G \in \mathcal{O}(U)$, such that $G|_{V_i} = G_{V_i}$. It is clear that G is the inverse of F . \square

Reconstructions of a sheaf from its global sections have been thoroughly studied in algebraic and differential geometry. A survey on such results can be found in [BPP18] and [BPP15]. The probably best known example is the construction of the structure sheaf \mathcal{O}_X of an affine scheme $X = \text{Spec } R$ from its global sections commutative unital ring $\mathcal{O}_X(X) = R$. In this case, the ring $\mathcal{O}_X(V_f)$ of functions on a Zariski open subset V_f , $f \in R$, is defined as a localization of $\mathcal{O}_X(X)$. In the case of a \mathbb{Z}_2^n -manifold (M, \mathcal{O}_M) , we reconstruct $\mathcal{O}_M(U)$ as the localization of $\mathcal{O}_M(M)$ with respect to the multiplicative subset $S_U = \{F \in \mathcal{O}_M(M) : (\varepsilon_M F)|_U \text{ is invertible}\}$. The chosen localization comes with a morphism that sends global sections with invertible projection in $C_M^\infty(U)$ to invertible sections in $\mathcal{O}_M(U)$, see Proposition 5.

We briefly address localization in the \mathbb{Z}_2^n -commutative setting. Let R be a \mathbb{Z}_2^n -commutative associative unital \mathbb{R} -algebra and let S be a multiplicative subset of R , whose elements are homogeneous even vectors. In the following, we consider right fractions $rs^{-1} \in RS^{-1}$ (left ones would work as well) and denote the degree of any element by the same symbol as the element itself. We define the equivalence $rs^{-1} \sim r's'^{-1}$ of two fractions by requiring the existence of some $\sigma \in S$ such that

$$(rs' - (-1)^{\langle s, s' \rangle} r's) \sigma = 0 . \quad (5)$$

It can be checked that the relation \sim is an equivalence. The addition of fractions is defined by

$$rs^{-1} + r's'^{-1} = \left(rs' + (-1)^{\langle s, s' \rangle} r's \right) (ss')^{-1} . \quad (6)$$

The definition is independent of the chosen representatives. The multiplication of fractions is given by

$$rs^{-1} \cdot r's'^{-1} = (-1)^{\langle r'+s', s \rangle} rr'(ss')^{-1} , \quad (7)$$

provided r' is homogeneous. Again the result is invariant under a change of representatives. If r' is not homogeneous, it uniquely reads $r' = \sum_\gamma r'^\gamma$ ($\gamma \in \mathbb{Z}_2^n$). In view of (6), we get $r's'^{-1} = \sum_\gamma (r'^\gamma s'^{-1})$, so that we can extend the definition (7) by linearity. Finally, the scalar multiplication by $\mathfrak{r} \in \mathbb{R}$ is

$$\mathfrak{r}(rs^{-1}) = (\mathfrak{r}r)s^{-1} . \quad (8)$$

For our purpose, it will actually be sufficient to consider the multiplicative subset

$$S_U = \{F \in \mathcal{O}_M^0(M) : (\varepsilon_M F)|_U \text{ is invertible}\} \subset \mathcal{O}_M(M)$$

of the \mathbb{Z}_2^n -commutative associative unital \mathbb{R} -algebra $\mathcal{O}_M(M)$. Since the \mathbb{Z}_2^n -functions of this subset are not only even, but of degree 0, the signs in the equivalence (5) and the operations (6) and (7) disappear.

Proposition 6. *The operations (6), (7), and (8) (without signs) endow the localization $\mathcal{O}_M(M) \cdot S_U^{-1}$ with a \mathbb{Z}_2^n -commutative associative unital \mathbb{R} -algebra structure, whose grading is naturally induced by the grading of $\mathcal{O}_M(M)$ (for a homogeneous r , the degree of rs^{-1} is the degree of r), and whose zero (resp., unit) is represented by $0 \cdot 1^{-1}$ (resp., $1 \cdot 1^{-1}$).*

Proof. Straightforward verification. □

We thus get a presheaf

$$\mathcal{L}_M : \text{Open}(M) \ni U \mapsto \mathcal{O}_M(M) \cdot S_U^{-1} \in \mathbb{Z}_2^n \mathbf{UAlg}$$

on M valued in the category of \mathbb{Z}_2^n -commutative associative unital \mathbb{R} -algebras. Indeed, if $V \subset U$ is open, the obvious inclusion $\iota_V^U : S_U \hookrightarrow S_V$ provides a natural well-defined restriction

$$r_V^U : \mathcal{L}_M(U) \ni Fs^{-1} \mapsto F(\iota_V^U s)^{-1} \in \mathcal{L}_M(V),$$

and these restrictions satisfy the usual cocycle condition.

As indicated above, we will show (in several steps) that the presheaf \mathcal{L}_M coincides with the structure sheaf \mathcal{O}_M .

First, since it follows from Proposition 5 that, for any $s \in S_U$, the restriction $s|_U$ is invertible in $\mathcal{O}_M(U)$, we have a map

$$\lambda_U : \mathcal{L}_M(U) \ni Fs^{-1} \mapsto F|_U(s|_U)^{-1} \in \mathcal{O}_M(U).$$

This map is well-defined. Indeed, if $Fs^{-1} = F's'^{-1}$, there is $\sigma \in S_U$, such that $(F|_U s'|_U - F'|_U s|_U) \sigma|_U = 0$. Since the restrictions $s|_U$, $s'|_U$, and $\sigma|_U$ are invertible in $\mathcal{O}_M(U)$, the claim follows. Further, it can be straightforwardly checked that λ_U is a morphism of \mathbb{Z}_2^n -graded unital \mathbb{R} -algebras.

In fact:

Proposition 7. *For any open $U \subset M$, the localisation map $\lambda_U : \mathcal{L}_M(U) \rightarrow \mathcal{O}_M(U)$ is a \mathbb{Z}_2^n -graded unital \mathbb{R} -algebra isomorphism.*

The proof of this result uses a method that can be found in various works, see for instance [BBH91], [CCF11], and [NS03]. We give this proof for completeness, as well as to show that it goes through in our \mathbb{Z}_2^n -graded setting.

Proof. It suffices to explain why λ_U is bijective.

1. *Injectivity:* Assume that $F|_U(s|_U)^{-1} = 0$, i.e., that $F|_U = 0$, and show that $Fs^{-1} \sim 01^{-1}$, i.e., that there is $\sigma \in S_U$, such that $F\sigma = 0$. Let (V_i, ψ_i) be a partition of unity of M , such that the V_i are \mathbb{Z}_2^n -chart domains, so that $\mathcal{O}_M|_{V_i} \simeq C_M^\infty|_{V_i}[[\xi]]$. For any i , we have

$$F|_{U \cap V_i}(x, \xi) = \sum_{\alpha} F_{\alpha}|_{U \cap V_i}(x) \xi^{\alpha} = 0, \quad \text{i.e.,} \quad F_{\alpha}|_{U \cap V_i} = 0, \quad \forall \alpha.$$

Let

$$\sigma_i \in C_M^\infty(V_i) \subset \mathcal{O}_M^0(V_i), \quad \text{such that} \quad \sigma_i|_{U \cap V_i} > 0 \quad \text{and} \quad \sigma_i|_{V_i \setminus (U \cap V_i)} = 0.$$

It follows that $F|_{V_i} \sigma_i = 0$. The \mathbb{Z}_2^n -function $\sigma = \sum_i \sigma_i \psi_i \in \mathcal{O}_M^0(M)$ has the required properties. Indeed, the open subsets V_i and $\Omega_i = M \setminus \text{supp } \psi_i$ cover M and $F\sigma_i \psi_i$ vanishes on both, V_i and Ω_i , so that $F\sigma = \sum_i F\sigma_i \psi_i = 0$. In addition, for any $m \in U$, we have $(\varepsilon \psi_i)(m) \geq 0$, for all i , and there is at least one j , such that $(\varepsilon \psi_j)(m) > 0$. Since $(\varepsilon \psi_j)|_{\Omega_j} = 0$, we get $m \in U \cap V_j$, so that $(\sigma_j \varepsilon \psi_j)(m) > 0$,

$$(\varepsilon \sigma)(m) = \sum_i (\sigma_i \varepsilon \psi_i)(m) > 0,$$

and $\sigma \in S_U$.

2. *Surjectivity:* We must express an arbitrary $f \in \mathcal{O}_M(U)$ as a product $f = F|_U(s|_U)^{-1}$, with $F \in \mathcal{O}_M(M)$ and $s \in S_U$. To construct the global sections F and s , consider an increasing countable family of seminorms p_n that implements the locally convex topology of the Fréchet space $\mathcal{O}_M(M)$. Take also a countable open cover U_n of U , such that $\bar{U}_n \subset U$, as well as bump functions $\gamma_n \in \mathcal{O}_M^0(M)$, which satisfy $\gamma_n|_{U_n} = 1$ (which implies that $(\varepsilon_M \gamma_n)|_{U_n} = 1$), $\text{supp } \gamma_n \subset U$, and $\varepsilon_M \gamma_n \geq 0$. The following series converge in $\mathcal{O}_M(M)$ and provide us with the required global sections:

$$F := \sum_{n=0}^{\infty} \frac{1}{2^n} \frac{\gamma_n f}{1 + p_n(\gamma_n) + p_n(\gamma_n f)} \quad \text{and} \quad s := \sum_{n=0}^{\infty} \frac{1}{2^n} \frac{\gamma_n}{1 + p_n(\gamma_n) + p_n(\gamma_n f)}.$$

Indeed, convergence follows, if we can show that the series are Cauchy, i.e., if they are

Cauchy with respect to each p_m . If $r, s \rightarrow \infty$, we get $r \geq m$, and, since the seminorms are increasing, we have

$$p_m \left(\sum_{n=r}^s \frac{1}{2^n} \frac{\gamma_n f}{1 + p_n(\gamma_n) + p_n(\gamma_n f)} \right) \leq \sum_{n=r}^s \frac{1}{2^n} \frac{p_m(\gamma_n f)}{1 + p_n(\gamma_n) + p_n(\gamma_n f)} < \sum_{n=r}^s \frac{1}{2^n} \rightarrow 0,$$

whether the factor f is present or not. On the other hand, as restrictions are continuous, it is clear that $F|_U = fs|_U$, so that $f = F|_U(s|_U)^{-1}$, provided we show that $s \in \mathcal{O}_M^0(M)$ belongs to S_U , i.e., that $(\varepsilon_M s)(m) \neq 0$, for all $m \in U$. To see this, remark that $(\text{id}, \varepsilon) :$

$(M, C_M^\infty) \rightarrow (M, \mathcal{O}_M)$ is a morphism of \mathbb{Z}_2^n -manifolds, so that $\varepsilon_M : \mathcal{O}_M(M) \rightarrow C_M^\infty(M)$ is continuous, see [BP18, Theorem 19]. For any U_m of the cover of U , we thus get

$$(\varepsilon_M s)|_{U_m} := \sum_{n=0}^{\infty} \frac{1}{2^n} \frac{(\varepsilon_M \gamma_n)|_{U_m}}{1 + p_n(\gamma_n) + p_n(\gamma_n f)} > 0,$$

in view of the properties of γ_n . □

Theorem 8. *The \mathbb{Z}_2^n -commutative associative unital \mathbb{R} -algebra $\mathcal{O}_M(M)$ of global sections of the structure sheaf of a \mathbb{Z}_2^n -manifold (M, \mathcal{O}_M) fully determines this sheaf. More precisely, there is a presheaf isomorphism $\lambda : \mathcal{L}_M \rightarrow \mathcal{O}_M$, so that the presheaf \mathcal{L}_M , which is obtained from $\mathcal{O}_M(M)$, is actually a sheaf, which is isomorphic to the structure sheaf \mathcal{O}_M .*

Proof. It suffices to check that the family $\lambda_U : \mathcal{L}_M(U) \rightarrow \mathcal{O}_M(U)$, $U \in \mathbf{Open}(M)$, of \mathbb{Z}_2^n -graded unital \mathbb{R} -algebra isomorphisms, commutes with the restrictions r_V^U in \mathcal{L}_M and ρ_V^U in \mathcal{O}_M ($V \subset U$, $V \in \mathbf{Open}(M)$). This is actually obvious:

$$\lambda_V(r_V^U(Fs^{-1})) = F|_V(s|_V)^{-1} = \rho_V^U(\lambda_U(Fs^{-1})).$$

□

3.3 Reconstruction of a \mathbb{Z}_2^n -morphism

In algebraic geometry, any commutative unital ring morphism $\psi : S \rightarrow R$ defines a morphism of affine schemes $\Phi = (\phi, \phi^*) : (\mathrm{Spec} R, \mathcal{O}_{\mathrm{Spec} R}) \rightarrow (\mathrm{Spec} S, \mathcal{O}_{\mathrm{Spec} S})$, whose continuous base map ϕ associates to each prime ideal \mathfrak{p} the prime ideal $\psi^{-1}(\mathfrak{p})$. A similar result exists in the category of \mathbb{Z}_2^n -manifolds and \mathbb{Z}_2^n -morphisms, with the same definition of the continuous base map.

Theorem 9. *Let $\mathcal{M} = (M, \mathcal{O}_M)$ and $\mathcal{N} = (N, \mathcal{O}_N)$ be \mathbb{Z}_2^n -manifolds. The map*

$$\beta : \mathrm{Hom}_{\mathbb{Z}_2^n \mathbf{Man}}(\mathcal{M}, \mathcal{N}) \ni \Phi = (\phi, \phi^*) \mapsto \phi_N^* \in \mathrm{Hom}_{\mathbb{Z}_2^n \mathbf{UAlg}}(\mathcal{O}_N(N), \mathcal{O}_M(M))$$

is a bijection.

Proof. To show that β is surjective, we consider $\psi \in \mathrm{Hom}_{\mathbb{Z}_2^n \mathbf{UAlg}}(\mathcal{O}_N(N), \mathcal{O}_M(M))$ and construct $\Phi \in \mathrm{Hom}_{\mathbb{Z}_2^n \mathbf{Man}}(\mathcal{M}, \mathcal{N})$, such that $\phi_N^* = \psi$.

Since M (resp., N) endowed with its base space topology is homeomorphic to $\mathrm{Spm}(\mathcal{O}_M(M))$ (resp., $\mathrm{Spm}(\mathcal{O}_N(N))$) endowed with the Zariski topology, we define ϕ by

$$\phi : \mathrm{Spm}(\mathcal{O}_M(M)) \ni \ker \varepsilon_m \mapsto \ker(\varepsilon_m \circ \psi) = \ker \varepsilon_n \in \mathrm{Spm}(\mathcal{O}_N(N)),$$

see Propositions 4 and 3. This map is continuous. Indeed, for any $F \in \mathcal{O}_N(N)$, the preimage by ϕ of the open subset

$$V(F) = \{\ker \varepsilon_n \in \mathrm{Spm}(\mathcal{O}_N(N)) : \varepsilon_n(F) \neq 0\}$$

is the subset

$$\phi^{-1}(V(F)) = \{\ker \varepsilon_m \in \text{Spn}(\mathcal{O}_M(M)) : \varepsilon_m(\psi(F)) \neq 0\} = V(\psi(F)) .$$

To define, for any open $V \subset N$, a \mathbb{Z}_2^n -graded unital \mathbb{R} -algebra morphism

$$\phi_V^* : \mathcal{O}_N(V) \rightarrow (\phi_* \mathcal{O}_M)(V) ,$$

we rely on the isomorphism of \mathbb{Z}_2^n -graded unital \mathbb{R} -algebras $\mathcal{O}_N(V) \simeq \mathcal{L}_N(V)$ and the similar isomorphism in M . Hence, we define ϕ_V^* by

$$\phi_V^* : \mathcal{L}_N(V) \ni Fs^{-1} \mapsto \psi(F)\psi(s)^{-1} \in \mathcal{L}_M(\phi^{-1}(V)) .$$

This map is actually well-defined. Since $s \notin \ker \varepsilon_n$, for all $n \in V$, we have $s \notin \ker(\varepsilon_m \circ \psi)$, for all $m \in \phi^{-1}(V)$, what means that $\psi(s) \in S_{\phi^{-1}(V)}$. In view of this, it is easy to see that the image is independent of the representative. The map ϕ_V^* is a \mathbb{Z}_2^n -graded unital \mathbb{R} -algebra morphism, because ψ is.

As the family ϕ_V^* , $V \in \mathbf{Open}(N)$, commutes obviously with restrictions, the continuous base map ϕ and the family of algebra morphisms ϕ_V^* , $V \in \mathbf{Open}(N)$, define a \mathbb{Z}_2^n -morphism $\Phi : \mathcal{M} \rightarrow \mathcal{N}$. To see that $\beta(\Phi) = \phi_N^* = \psi$, it suffices to note that $\lambda_N : \mathcal{L}_N(N) \rightarrow \mathcal{O}_N(N)$ sends the fraction Fs^{-1} to the section Fs^{-1} (and similarly for M), so that ϕ_N^* and ψ coincide.

It remains to prove that β is injective. Let thus $\Phi = (\phi, \phi^*)$ and $\Psi = (\psi, \psi^*)$ be two \mathbb{Z}_2^n -morphisms from \mathcal{M} to \mathcal{N} , such that $\phi_N^* = \psi_N^*$. Since the pullbacks by a \mathbb{Z}_2^n -morphism commute with the base projections, we get, for any $m \in M$,

$$\phi(m) \simeq \ker \varepsilon_{\phi(m)} = \{F \in \mathcal{O}(N) : ((\varepsilon_N F) \circ \phi)(m) = 0\} = \{F \in \mathcal{O}(N) : (\varepsilon_M(\phi_N^* F))(m) = 0\} .$$

Hence, the continuous base maps ϕ and ψ coincide. Similarly, for any open $V \subset N$, each $F_V \in \mathcal{O}_N(V)$ reads uniquely $F_V = \lambda_V(Fs^{-1}) = F|_V(s|_V)^{-1}$, with $F \in \mathcal{O}_N(N)$ and $s \in S_V \subset \mathcal{O}_N(N)$, see Proposition 7. As the family of pullbacks ϕ^* commutes with restrictions, we obtain

$$\phi_V^*(F_V) = (\phi_V^* F|_V) (\phi_V^* s|_V)^{-1} = (\phi_N^* F)|_{\phi^{-1}(V)} (\phi_N^* s)|_{\phi^{-1}(V)}^{-1} .$$

Hence $\phi_V^* = \psi_V^*$. □

The preceding theorem, which allows us to characterize \mathcal{M} -points $\text{Hom}_{\mathbb{Z}_2^n \mathbf{Man}}(\mathcal{M}, \mathcal{N})$ of a \mathbb{Z}_2^n -manifold \mathcal{N} by algebra morphisms, has some noteworthy corollaries.

Corollary 10. *The covariant functor*

$$\mathcal{F} : \mathbb{Z}_2^n \mathbf{Man} \rightarrow \mathbb{Z}_2^n \mathbf{UAlg}^{\text{op}} ,$$

which is defined on objects by $\mathcal{F}(\mathcal{M}) = \mathcal{O}_M(M)$ and on morphisms by $\mathcal{F}(\Phi) = \phi_N^*$, is fully faithful, so that $\mathbb{Z}_2^n \mathbf{Man}$ can be viewed as full subcategory of $\mathbb{Z}_2^n \mathbf{UAlg}^{\text{op}}$.

The statement regarding the full subcategory is based on the well-known fact that any fully faithful functor is injective up to isomorphism on objects. This means that the existence of an isomorphism $\mathcal{O}_M(M) \simeq \mathcal{O}_N(N)$ of \mathbb{Z}_2^n -graded unital \mathbb{R} -algebras implies the existence of an isomorphism $\mathcal{M} \simeq \mathcal{N}$ of \mathbb{Z}_2^n -manifolds.

Corollary 11. *Let $\mathcal{M} = (M, \mathcal{O}_M)$ and $\mathcal{N} = (N, \mathcal{O}_N)$ be \mathbb{Z}_2^n -manifolds. The \mathbb{Z}_2^n -manifolds \mathcal{M} and \mathcal{N} are diffeomorphic if and only if their \mathbb{Z}_2^n -commutative associative unital \mathbb{R} -algebras $\mathcal{O}_M(M)$ and $\mathcal{O}_N(N)$ of global \mathbb{Z}_2^n -functions are isomorphic.*

Such Pursell-Shanks type results have been studied extensively by one of the authors of this paper. Algebraic characterizations similar to Corollary 11 exist for instance for the Lie algebras of first order differential operators, of differential operators, and symbols of differential operators on a smooth manifold, for the super Lie algebras of vector fields and first order differential operators on a smooth supermanifold, as well as for the Lie algebra of sections of an Atiyah algebroid, see [GP04], [GKP10], [GKP11], [GKP13].

Corollary 12. *The \mathbb{Z}_2^n -manifold $\mathcal{E} = (\emptyset, 0)$ (resp., $\mathbb{R}^{0\mathbf{0}} = (\{\text{pt}\}, \mathbb{R})$) is the initial (resp., terminal) object of the category of \mathbb{Z}_2^n -manifolds.*

Proof. For any \mathbb{Z}_2^n -manifold $\mathcal{M} = (M, \mathcal{O}_M)$, we have bijections

$$\text{Hom}_{\mathbb{Z}_2^n\text{Man}}(\mathcal{E}, \mathcal{M}) \simeq \text{Hom}_{\mathbb{Z}_2^n\text{UAlg}}(\mathcal{O}_M(M), 0) \simeq \{F \mapsto 0\}$$

and

$$\text{Hom}_{\mathbb{Z}_2^n\text{Man}}(\mathcal{M}, \mathbb{R}^{0\mathbf{0}}) \simeq \text{Hom}_{\mathbb{Z}_2^n\text{UAlg}}(\mathbb{R}, \mathcal{O}_M(M)) \simeq \{r \mapsto r \cdot 1\}.$$

□

4 Finite products in the category of \mathbb{Z}_2^n -manifolds

4.1 Cartesian product of \mathbb{Z}_2^n -manifolds

Let $\mathcal{M} = (M, \mathcal{O}_M)$ and $\mathcal{N} = (N, \mathcal{O}_N)$ be two \mathbb{Z}_2^n -manifolds of dimension $p|\mathbf{q}$ and $r|\mathbf{s}$, respectively. The products $U \times V$, $U \subset M$ and $V \subset N$ open, form a basis \mathcal{B} of the (second-countable, Hausdorff) product topology of $M \times N$. Better, since the \mathbb{Z}_2^n -chart domains U_i in M (resp., V_j in N) are a basis of the topology of M (resp., of N), the products $U_i \times V_j$ form a basis \mathfrak{B} of the product topology of $M \times N$. As \mathbb{Z}_2^n -chart domains are diffeomorphic to open subsets of some coordinate space \mathbb{R}^n , we identify the U_i and the V_j with the diffeomorphic $U_i \subset \mathbb{R}^p$ and $V_j \subset \mathbb{R}^r$. Further, we denote the coordinates of the charts with domains U_i (resp., V_j) by (x_i, ξ_i) (resp., (y_j, η_j)), or, in case we use only two domains U_i (resp., V_j), we write also (x, ξ) and (x', ξ') (resp., (y, η) and (y', η')).

Definition 13. *Let $\mathcal{M} = (M, \mathcal{O}_M)$ and $\mathcal{N} = (N, \mathcal{O}_N)$ be two \mathbb{Z}_2^n -manifolds of dimension $p|\mathbf{q}$ and $r|\mathbf{s}$, respectively. The product \mathbb{Z}_2^n -manifold $\mathcal{M} \times \mathcal{N}$, of dimension $p+r|\mathbf{q}+\mathbf{s}$, is the locally \mathbb{Z}_2^n -ringed space $(M \times N, \mathcal{O}_{M \times N})$, where $M \times N$ is the product topological space and where the sheaf $\mathcal{O}_{M \times N}$ is glued from the sheaves $C_{U_i \times V_j}^\infty(x_i, y_j)[[\xi_i, \eta_j]]$ associated to the basis \mathfrak{B} :*

$$\mathcal{O}_{M \times N}|_{U_i \times V_j} \simeq C_{U_i \times V_j}^\infty(x_i, y_j)[[\xi_i, \eta_j]]. \quad (9)$$

Recall that sheaves can be glued. More precisely, if $(U_i)_i$ is an open cover of a topological space M , if \mathcal{F}_i is a sheaf on U_i , and if $\varphi_{ji} : \mathcal{F}_i|_{U_i \cap U_j} \rightarrow \mathcal{F}_j|_{U_i \cap U_j}$ is a sheaf isomorphism such that the usual cocycle condition $\varphi_{kj} \varphi_{ji} = \varphi_{ki}$ holds, then there is a unique sheaf \mathcal{F} on M such that $\mathcal{F}|_{U_i} \simeq \mathcal{F}_i$. In the following, we set $U_{ij} = U_i \cap U_j$.

Let now $C_{U_i \times V_j}^\infty[[\xi, \eta]]$ be the standard sheaf of \mathbb{Z}_2^n -graded \mathbb{Z}_2^n -commutative associative unital \mathbb{R} -algebras of formal power series in (ξ, η) with coefficients in sections of the sheaf $C_{U_i \times V_j}^\infty$. The isomorphisms $\varphi_{ij, ij}$ between the appropriate restrictions of the sheaves of algebras $C_{U_i \times V_j}^\infty[[\xi, \eta]]$ on the open cover $(U_i \times V_j)_{i,j}$ of $M \times N$ are induced as follows. Since \mathcal{M} is a \mathbb{Z}_2^n -manifold, there are \mathbb{Z}_2^n -isomorphisms

$$\Phi_i = (\phi_i, \phi_i^*) : (U_i, \mathcal{O}_M|_{U_i}) \rightarrow (U_i, C_{U_i}^\infty[[\xi]]) ,$$

which induce \mathbb{Z}_2^n -isomorphisms or coordinate transformations

$$\Psi_{ii} = \Phi_i \Phi_i^{-1} : (U_{ii}, C_{U_{ii}}^\infty|_{U_{ii}}[[\xi]]) \rightarrow (U_{ii}, C_{U_{ii}}^\infty|_{U_{ii}}[[\xi']]) .$$

As we view U_i as both, an open subset of M and an open subset of \mathbb{R}^p , we implicitly identify U_i with its diffeomorphic image $\phi_i(U_i)$, so that $\phi_i = \text{id}_{U_i}$. Hence, the coordinate transformations reduce to the isomorphisms

$$\psi_{ii}^* = (\phi_i^*)^{-1} \phi_i^* \tag{10}$$

of sheaves of \mathbb{Z}_2^n -commutative \mathbb{R} -algebras:

$$\psi_{ii}^* : C_{U_{ii}}^\infty|_{U_{ii}}[[\xi']] \rightarrow C_{U_{ii}}^\infty|_{U_{ii}}[[\xi]] . \tag{11}$$

Similar coordinate transformations exist for \mathcal{N} :

$$\psi_{jj}^* : C_{V_{jj}}^\infty|_{V_{jj}}[[\eta']] \rightarrow C_{V_{jj}}^\infty|_{V_{jj}}[[\eta]] . \tag{12}$$

We denote the base coordinates in U_i (resp., U_i) by x (resp., x') and those in V_j (resp., V_j) by y (resp., y'). The coordinate transformations (11), $x = x(x', \xi')$, $\xi = \xi(x', \xi')$, and (12), $y = y(y', \eta')$, $\eta = \eta(y', \eta')$, implement coordinate transformations or isomorphisms of sheaves of \mathbb{Z}_2^n -commutative \mathbb{R} -algebras

$$\varphi_{ij, ij} = \psi_{ii}^* \times \psi_{jj}^* : C_{U_i \times V_j}^\infty|_{U_{ii} \times V_{jj}}[[\xi', \eta']] \rightarrow C_{U_i \times V_j}^\infty|_{U_{ii} \times V_{jj}}[[\xi, \eta]] .$$

In view of (10), the $\varphi_{ij, ij}$ satisfy the cocycle condition. We thus get a unique glued sheaf $\mathcal{O}_{M \times N}$ of \mathbb{Z}_2^n -commutative \mathbb{R} -algebras over $M \times N$ which restricts on $U_i \times V_j$ to

$$\mathcal{O}_{M \times N}|_{U_i \times V_j} \simeq C_{U_i \times V_j}^\infty[[\xi, \eta]] ,$$

i.e., we obtain a \mathbb{Z}_2^n -manifold, which we refer to as the product $\mathcal{M} \times \mathcal{N}$ of \mathcal{M} and \mathcal{N} .

4.2 Fundamental isomorphisms

Theorem 14. *Let $\mathbb{R}^{p|\mathfrak{q}}$ (resp., $\mathbb{R}^{r|\mathfrak{s}}$) be the usual \mathbb{Z}_2^n -domain $(\mathbb{R}^p, C_{\mathbb{R}^p}^\infty[[\xi]])$ (resp., $(\mathbb{R}^r, C_{\mathbb{R}^r}^\infty[[\eta]])$), and let $\Omega' \subset \mathbb{R}^p$ and $\Omega'' \subset \mathbb{R}^r$ be open. There is an isomorphism of topological algebras*

$$C_{\mathbb{R}^p}^\infty(\Omega')[[\xi]] \widehat{\otimes} C_{\mathbb{R}^r}^\infty(\Omega'')[[\eta]] \simeq C_{\mathbb{R}^p \times \mathbb{R}^r}^\infty(\Omega' \times \Omega'')[[\xi, \eta]], \quad (13)$$

where the completion is taken with respect to any locally convex topology on the algebraic tensor product $C_{\mathbb{R}^p}^\infty(\Omega')[[\xi]] \otimes C_{\mathbb{R}^r}^\infty(\Omega'')[[\eta]]$.

Proof. Let R be a commutative von Neumann regular ring. For any families $(M_\alpha)_\alpha$ and $(N_\beta)_\beta$ of free R -modules, the natural R -linear map

$$\left(\prod_{\alpha} M_{\alpha} \right) \otimes_R \left(\prod_{\beta} N_{\beta} \right) \rightarrow \prod_{\alpha\beta} (M_{\alpha} \otimes_R N_{\beta})$$

is injective, if and only if R is injective as a module over itself [Goo72]. Since any field is von Neumann regular, the regularity and injective module conditions are satisfied for $R = \mathbb{R}$. Hence, the linear map

$$\left(\prod_{\alpha} C^\infty(\Omega') \right) \otimes \left(\prod_{\beta} C^\infty(\Omega'') \right) \rightarrow \prod_{\alpha\beta} (C^\infty(\Omega') \otimes C^\infty(\Omega''))$$

is injective. Further, in view of [BP18, Corollary 17], the map

$$C^\infty(\Omega')[[\xi]] \ni \sum_{\alpha \in \mathcal{A}} f_{\alpha}(x) \xi^{\alpha} \mapsto (f_{\alpha})_{\alpha \in \mathcal{A}} \in \prod_{\alpha \in \mathcal{A}} C^\infty(\Omega') \quad (14)$$

is a TVS-isomorphism between the source and the target equipped with the standard topology and the product topology of the standard topologies, respectively. In the sequence of canonical maps

$$\begin{aligned} C^\infty(\Omega')[[\xi]] \otimes C^\infty(\Omega'')[[\eta]] &\simeq \left(\prod_{\alpha \in \mathcal{A}} C^\infty(\Omega') \right) \otimes \left(\prod_{\beta \in \mathcal{B}} C^\infty(\Omega'') \right) \rightarrow \prod_{\alpha \in \mathcal{A}, \beta \in \mathcal{B}} (C^\infty(\Omega') \otimes C^\infty(\Omega'')) \rightarrow \\ &\prod_{\alpha \in \mathcal{A}, \beta \in \mathcal{B}} C^\infty(\Omega' \times \Omega'') \simeq C^\infty(\Omega' \times \Omega'')[[\xi, \eta]], \end{aligned} \quad (15)$$

the first \simeq is a linear bijection, the first \rightarrow is a linear injection, and the second \simeq is a TVS-isomorphism for the topologies used in (14). In

$$C^\infty(\Omega') \otimes C^\infty(\Omega'') \rightarrow C^\infty(\Omega') \widehat{\otimes} C^\infty(\Omega'') \simeq C^\infty(\Omega' \times \Omega''), \quad (16)$$

the isomorphism \simeq is the well-known TVS-isomorphism [Gro52] (the target is endowed with its standard topology and the source with the topology of the completion with respect to any ($C^\infty(\Omega')$ is nuclear) locally convex topology on $C^\infty(\Omega') \otimes C^\infty(\Omega'')$ – we will not specify the latter topology), and the arrow \rightarrow is the continuous linear inclusion (any TVS is a topological vector subspace (TVSS) of its completion, see Proposition 26). This \rightarrow induces the second \rightarrow in (15), which is the inclusion of the source vector subspace into the target vector space. The

source becomes a TVSS of the target when endowed with the induced topology (the induced topology is coarser than the product topology of the induced topologies). Finally, we equip the first space in (15) with the initial topology with respect to the first \rightarrow , so that the first space gets promoted to a TVSS of the second, see Proposition 24, and the first \rightarrow becomes the continuous linear inclusion. The composite

$$\iota : C^\infty(\Omega')[[\xi]] \otimes C^\infty(\Omega''')[[\eta]] \ni \sum_{\alpha} f_{\alpha} \xi^{\alpha} \otimes \sum_{\beta} g_{\beta} \eta^{\beta} \mapsto \sum_{\alpha\beta} f_{\alpha} \otimes g_{\beta} \xi^{\alpha} \eta^{\beta} \in C^\infty(\Omega' \times \Omega''')[[\xi, \eta]]$$

of the maps of (15) is now the inclusion of a the source TVSS into the target TVS.

Note that, since the target is a LCTVS, see [BP18, Lemma 16], the source TVSS is also a LCTVS, see Proposition 29 (since $C^\infty(\Omega')[[\xi]]$ is nuclear, see [BP18, Lemma 16], the completion of the source is independent of the chosen locally convex topology). In view of Proposition 28, the completion of the source is a TVSS of the completion of the target, which, as the target is complete, see [BP18, Lemma 16], can be identified with the target due to Remark 27. In other words, the continuous extension

$$\hat{\iota} : C^\infty(\Omega')[[\xi]] \widehat{\otimes} C^\infty(\Omega''')[[\eta]] \rightarrow C^\infty(\Omega' \times \Omega''')[[\xi, \eta]] \quad (17)$$

of the inclusion ι is an injective continuous linear map, see text above Proposition 28. We will now prove that this map is surjective.

Let

$$S = \sum_{\alpha\beta} F_{\alpha\beta} \xi^{\alpha} \eta^{\beta} \in C^\infty(\Omega' \times \Omega''')[[\xi, \eta]]$$

be a formal series in the target space. In view of (16), we have [Gro52], for any $(\alpha, \beta) \in \mathcal{A} \times \mathcal{B}$,

$$F_{\alpha\beta} = \lim_{N \rightarrow +\infty} \sum_{j=0}^N f_{\alpha\beta}^j \otimes g_{\alpha\beta}^j,$$

where $f_{\alpha\beta}^j \in C^\infty(\Omega')$ and $g_{\alpha\beta}^j \in C^\infty(\Omega''')$, and where the limit is taken in $C^\infty(\Omega' \times \Omega''')$. Recall that $\mathcal{A} = \mathbb{N}^{|\mathbf{q}'|} \times \mathbb{Z}_2^{|\mathbf{q}''|}$, and similarly for \mathcal{B} . The product $\mathcal{A} \times \mathcal{B}$ is countable, since it is a finite product of countable sets. Let $I : \mathcal{A} \times \mathcal{B} \rightarrow \mathbb{N}$ be an injective map valued in \mathbb{N} . The map $J : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{I}$, with $\mathcal{I} = I(\mathcal{A} \times \mathcal{B})$, is thus a 1:1 correspondence. We identify $\mathcal{A} \times \mathcal{B}$ with \mathcal{I} via J . For any $j \in \mathbb{N}$, we set,

- for any $\alpha \in \mathcal{A}$ and any $i \in \mathcal{I}$,

$$C^\infty(\Omega') \ni \phi_{\alpha i}^j = \begin{cases} 0, & \text{if } i \simeq (\gamma, \delta) \neq (\alpha, \delta), \\ f_{\alpha\delta}^j, & \text{if } i \simeq (\gamma, \delta) = (\alpha, \delta), \end{cases} \quad \text{and,}$$

- for any $\beta \in \mathcal{B}$ and any $i \in \mathcal{I}$,

$$C^\infty(\Omega''') \ni \psi_{i\beta}^j = \begin{cases} 0, & \text{if } i \simeq (\gamma, \delta) \neq (\gamma, \beta), \\ g_{\gamma\beta}^j, & \text{if } i \simeq (\gamma, \delta) = (\gamma, \beta). \end{cases}$$

Note that \mathcal{I} is a finite set $\{0, 1, \dots, L\}$, $L \in \mathbb{N}$, (resp., is \mathbb{N}), if $\mathcal{A} \times \mathcal{B}$ is finite (resp., if $\mathcal{A} \times \mathcal{B}$ is countably infinite). For all $j \in \mathbb{N}$ and all $(\alpha, \beta) \in \mathcal{A} \times \mathcal{B}$, we get

$$\sum_{i=0}^M \phi_{\alpha i}^j \otimes \psi_{i\beta}^j = f_{\alpha\beta}^j \otimes g_{\alpha\beta}^j,$$

when $M \in \mathcal{I} \cap [J(\alpha, \beta), +\infty[$. Indeed, if $i \simeq (\gamma, \delta) \neq (\alpha, \beta)$, then, either $\gamma \neq \alpha$ and $\phi_{\alpha i}^j = 0$, or $\delta \neq \beta$ and $\psi_{i\beta}^j = 0$. However, if $i \simeq (\gamma, \delta) = (\alpha, \beta)$, then $\phi_{\alpha i}^j = f_{\alpha\beta}^j$ and $\psi_{i\beta}^j = g_{\alpha\beta}^j$, so that the announced result follows. Hence, for any $j \in \mathbb{N}$ and any $(\alpha, \beta) \in \mathcal{A} \times \mathcal{B}$, we have

$$\lim_{M \rightarrow +\infty} \sum_{i=0}^M \phi_{\alpha i}^j \otimes \psi_{i\beta}^j = f_{\alpha\beta}^j \otimes g_{\alpha\beta}^j,$$

where the sequence is constant for $M \geq J(\alpha, \beta)$ and where the limit is computed in the topology of $C^\infty(\Omega' \times \Omega'')$. If a finite number of sequences of a TVS do converge, then their sum converges to the sum of the limits. It follows that, for any $(\alpha, \beta) \in \mathcal{A} \times \mathcal{B}$ and any $N \in \mathbb{N}$,

$$\lim_{M \rightarrow +\infty} \sum_{j=0}^N \sum_{i=0}^M \phi_{\alpha i}^j \otimes \psi_{i\beta}^j = \sum_{j=0}^N f_{\alpha\beta}^j \otimes g_{\alpha\beta}^j,$$

so that, for all $(\alpha, \beta) \in \mathcal{A} \times \mathcal{B}$,

$$\lim_{N \rightarrow +\infty} \lim_{M \rightarrow +\infty} \sum_{j=0}^N \sum_{i=0}^M \phi_{\alpha i}^j \otimes \psi_{i\beta}^j = \lim_{N \rightarrow +\infty} \sum_{j=0}^N f_{\alpha\beta}^j \otimes g_{\alpha\beta}^j = F_{\alpha\beta}$$

in $C^\infty(\Omega' \times \Omega'')$, and

$$\lim_{N \rightarrow +\infty} \lim_{M \rightarrow +\infty} \left(\sum_{j=0}^N \sum_{i=0}^M \phi_{\alpha i}^j \otimes \psi_{i\beta}^j \right)_{(\alpha, \beta) \in \mathcal{A} \times \mathcal{B}} = (F_{\alpha\beta})_{(\alpha, \beta) \in \mathcal{A} \times \mathcal{B}} \quad (18)$$

in the product topology of $\prod_{\alpha\beta} C^\infty(\Omega' \times \Omega'')$, i.e., in the topology of the TVS $C^\infty(\Omega' \times \Omega'')[[\xi, \eta]]$. Therefore, the sequence

$$\left(\sum_{j=0}^N \sum_{i=0}^M \phi_{\alpha i}^j \otimes \psi_{i\beta}^j \right)_{(\alpha, \beta)} = \sum_{j=0}^N \sum_{i=0}^M \left(\sum_{\alpha} \phi_{\alpha i}^j \xi^\alpha \otimes \sum_{\beta} \psi_{i\beta}^j \eta^\beta \right) \in C^\infty(\Omega')[[\xi]] \otimes C^\infty(\Omega'')[[\eta]]$$

is a Cauchy sequence in $C^\infty(\Omega' \times \Omega'')[[\xi, \eta]]$, so a Cauchy sequence in the TVSS $C^\infty(\Omega')[[\xi]] \otimes C^\infty(\Omega'')[[\eta]]$, and also in the topological vector supspace $C^\infty(\Omega')[[\xi]] \widehat{\otimes} C^\infty(\Omega'')[[\eta]]$. Since this completion is sequentially complete, the Cauchy sequence considered converges in this space:

$$\lim_{N \rightarrow +\infty} \lim_{M \rightarrow +\infty} \sum_{j=0}^N \sum_{i=0}^M \left(\sum_{\alpha} \phi_{\alpha i}^j \xi^\alpha \otimes \sum_{\beta} \psi_{i\beta}^j \eta^\beta \right) \in C^\infty(\Omega')[[\xi]] \widehat{\otimes} C^\infty(\Omega'')[[\eta]],$$

where the limit is taken in the topology of $C^\infty(\Omega')[[\xi]] \widehat{\otimes} C^\infty(\Omega'')[[\eta]]$. Since the inclusion \hat{i} , see Equation (17), is sequentially continuous, we get

$$\begin{aligned} & \hat{i} \lim_{N \rightarrow +\infty} \lim_{M \rightarrow +\infty} \sum_{j=0}^N \sum_{i=0}^M \left(\sum_{\alpha} \phi_{\alpha i}^j \xi^\alpha \otimes \sum_{\beta} \psi_{i\beta}^j \eta^\beta \right) = \\ & \lim_{N \rightarrow +\infty} \lim_{M \rightarrow +\infty} \hat{i} \sum_{j=0}^N \sum_{i=0}^M \left(\sum_{\alpha} \phi_{\alpha i}^j \xi^\alpha \otimes \sum_{\beta} \psi_{i\beta}^j \eta^\beta \right) = \\ & \lim_{N \rightarrow +\infty} \lim_{M \rightarrow +\infty} \sum_{\alpha\beta} \sum_{j=0}^N \sum_{i=0}^M \phi_{\alpha i}^j \otimes \psi_{i\beta}^j \xi^\alpha \eta^\beta = \sum_{\alpha\beta} F_{\alpha\beta} \xi^\alpha \eta^\beta = S, \end{aligned}$$

in view of (18). This shows that the continuous linear inclusion

$$\hat{i} : C^\infty(\Omega')[[\xi]] \widehat{\otimes} C^\infty(\Omega'')[[\eta]] \rightarrow C^\infty(\Omega' \times \Omega'')[[\xi, \eta]]$$

is bijective, so that the source TVSS of the target coincides with the target as TVS.

Since the completed tensor product of two nuclear Fréchet algebras is again a nuclear Fréchet algebra [Eme17, Lemma 1.2.13], the source and target are actually topological algebras. We leave it to the reader to check that the preceding identification respects the multiplications. \square

Remark 15. If $p|\mathbf{q} = p|\mathbf{0}$ and $r|\mathbf{s} = 0|\mathbf{s}$, it follows from Theorem 14 that

$$C^\infty(\Omega) \widehat{\otimes}_{\mathbb{R}} \mathbb{R}[[\xi]] \simeq C^\infty(\Omega)[[\xi]],$$

and, if $p|\mathbf{q} = 0|\mathbf{q}$ and $r|\mathbf{s} = 0|\mathbf{s}$, we get

$$\mathbb{R}[[\xi]] \widehat{\otimes}_{\mathbb{R}} \mathbb{R}[[\eta]] \simeq \mathbb{R}[[\xi, \eta]].$$

Conversely, the general isomorphism of Theorem 14 is a consequence of the preceding particular cases and the fact that the category of complete nuclear spaces is a symmetric monoidal category with respect to the completed tensor product [Cos11].

Theorem 16. There is an isomorphism of sheaves of \mathbb{Z}_2^n -commutative \mathbb{R} -algebras

$$\mathcal{O}_{M \times N} \simeq (\mathcal{O}_M \widehat{\otimes} \mathcal{O}_N)^{-+}$$

between the structure sheaf of a product \mathbb{Z}_2^n -manifold and the sheafification of the standard extension of the \mathcal{B} -presheaf

$$\mathcal{O}_M \widehat{\otimes} \mathcal{O}_N : U \times V \mapsto \mathcal{O}_M(U) \widehat{\otimes} \mathcal{O}_N(V).$$

Proof. Recall that \mathcal{B} (resp., \mathfrak{B}) is the basis of the product topology of $M \times N$ made of the rectangular subsets $U \times V$, where $U \subset M$ and $V \subset N$ are open (resp., of the rectangular subsets $U_i \times V_j$, where $U_i \subset M$ and $V_j \subset N$ are \mathbb{Z}_2^n -chart domains). Let

$$\mathcal{F}(U \times V) := \mathcal{O}_M(U) \widehat{\otimes} \mathcal{O}_N(V) \quad (19)$$

be the completed tensor product of the nuclear \mathbb{Z}_2^n -graded Fréchet algebras $\mathcal{O}_M(U)$ and $\mathcal{O}_N(V)$ (with respect to any (reasonable) locally convex topology, e.g., the projective one). If $U' \times V' \subset U \times V$, the restrictions $\rho_{U'}^U : \mathcal{O}_M(U) \rightarrow \mathcal{O}_M(U')$ and $\rho_{V'}^V : \mathcal{O}_N(V) \rightarrow \mathcal{O}_N(V')$ of the Fréchet sheaves \mathcal{O}_M and \mathcal{O}_N are continuous linear maps. The continuous extension of the continuous linear map $\rho_{U'}^U \otimes \rho_{V'}^V$ is a continuous linear map [Gro52]

$$\rho_{U'}^U \widehat{\otimes} \rho_{V'}^V : \mathcal{O}_M(U) \widehat{\otimes} \mathcal{O}_N(V) \rightarrow \mathcal{O}_M(U') \widehat{\otimes} \mathcal{O}_N(V'),$$

which we denote by

$$\rho_{U' \times V'}^{U \times V} : \mathcal{F}(U \times V) \rightarrow \mathcal{F}(U' \times V').$$

Since the $\rho_{U'}^U$ and the $\rho_{V'}^V$ satisfy the standard presheaf conditions and the linear maps $\rho_{U' \times V'}^{U \times V}$ are continuous, it is clear that the latter satisfy these conditions as well. Hence, the pair (\mathcal{F}, ρ) is a **Set**-valued \mathcal{B} -presheaf.

This \mathcal{B} -presheaf can be extended to a **Set**-valued presheaf $(\bar{\mathcal{F}}, \bar{\rho})$. Indeed, set, for any open $\Omega \subset M \times N$,

$$\bar{\mathcal{F}}(\Omega) := \{(f_{ab})_{ab} : f_{ab} \in \mathcal{F}(U_a \times V_b), U_a \times V_b \subset \Omega, \text{ such that } \rho_{U_{aa} \times V_{bb}}^{U_a \times V_b}(f_{ab}) = \rho_{U_{aa} \times V_{bb}}^{U_a \times V_b}(f_{ab})\}, \quad (20)$$

and consider, for any $\Omega' \subset \Omega$, the map

$$\bar{\rho}_{\Omega'}^{\Omega} : \bar{\mathcal{F}}(\Omega) \rightarrow \bar{\mathcal{F}}(\Omega')$$

which sends any element of $\bar{\mathcal{F}}(\Omega)$ to the element of $\bar{\mathcal{F}}(\Omega')$ that we obtain by suppressing the f_{ab} for which $U_a \times V_b$ is not a subset of Ω' . The $\bar{\rho}_{\Omega'}^{\Omega}$ satisfy of course the standard presheaf conditions. Further, the presheaf $(\bar{\mathcal{F}}, \bar{\rho})$ extends the \mathcal{B} -presheaf (\mathcal{F}, ρ) . Indeed, for $\Omega = U \times V$, any $f \in \mathcal{F}(\Omega)$ provides a unique family $f_{ab} = \rho_{U_a \times V_b}^{U \times V}(f)$ in $\bar{\mathcal{F}}(\Omega)$, thus defining a map $\flat_{\Omega} : \mathcal{F}(\Omega) \rightarrow \bar{\mathcal{F}}(\Omega)$. If $\flat_{\Omega}(f) = \flat_{\Omega}(g)$, then, in particular,

$$f = \rho_{U \times V}^{U \times V}(f) = \rho_{U \times V}^{U \times V}(g) = g.$$

In fact \flat_{Ω} is a 1:1 correspondence. Indeed, any family f_{ab} in $\bar{\mathcal{F}}(\Omega)$ contains $f \in \mathcal{F}(\Omega)$, and $f_{ab} = \rho_{U_a \times V_b}^{U \times V}(f)$, so that $\flat_{\Omega}(f) = (f_{ab})_{ab}$. Hence,

$$\flat_{U \times V} : \mathcal{F}(U \times V) \xrightarrow{\sim} \bar{\mathcal{F}}(U \times V). \quad (21)$$

Moreover, if $\Omega' = U' \times V' \subset \Omega = U \times V$, and if $f \in \mathcal{F}(U \times V)$, then

$$\bar{\rho}_{U' \times V'}^{U \times V}(\flat_{U \times V}(f)) = \flat_{U' \times V'}(\rho_{U' \times V'}^{U \times V}(f)),$$

since both sides are made of the family of restrictions $\rho_{U_a \times V_b}^{U \times V}(f)$, for all $U_a \times V_b \subset U' \times V'$.

Let now $U_i \times V_j \in \mathfrak{B}$ be a Cartesian product of \mathbb{Z}_2^n -chart domains, and let $\Omega \subset U_i \times V_j$ be any open subset. Recall Definition (20). Since $U_a \subset U_i$ and $V_b \subset V_j$ are \mathbb{Z}_2^n -chart domains, we get in view of (13) and of (9),

$$\begin{aligned} \mathcal{F}(U_a \times V_b) &= \mathcal{O}_M(U_a) \widehat{\otimes} \mathcal{O}_N(V_b) = C^\infty(U_a)[[\xi_i]] \widehat{\otimes} C^\infty(V_b)[[\eta_j]] \simeq \\ &C^\infty(U_a \times V_b)[[\xi_i, \eta_j]] = \mathcal{O}_{M \times N}(U_a \times V_b). \end{aligned} \quad (22)$$

Due to the continuity and linearity of the restrictions $\rho_{U' \times V'}^{U \times V}$ of \mathcal{F} and the restrictions P_Ω^Ω of $\mathcal{O}_{M \times N}$, the restrictions in (20) coincide with the corresponding restrictions P of the structure sheaf of the product \mathbb{Z}_2^n -manifold, as both reduce to the same restrictions of classical functions. It follows from (20) and (22) that any family $f_{ab} \in \bar{\mathcal{F}}(\Omega)$ is made of \mathbb{Z}_2^n -functions $f_{ab} \in \mathcal{O}_{M \times N}(U_a \times V_b)$, which are defined on the cover of Ω by all the $U_a \times V_b \subset \Omega$, and whose P -restrictions coincide on all intersections. Hence, any family $f_{ab} \in \bar{\mathcal{F}}(\Omega)$ can be glued in the sheaf $\mathcal{O}_{M \times N}$ and thus provides a unique \mathbb{Z}_2^n -function $f \in \mathcal{O}_{M \times N}(\Omega)$ such that $P_{U_a \times V_b}^\Omega(f) = f_{ab}$. The resulting map $\mathfrak{b}_\Omega : \bar{\mathcal{F}}(\Omega) \rightarrow \mathcal{O}_{M \times N}(\Omega)$ is clearly injective. It is also surjective, since the restrictions $f_{ab} := P_{U_a \times V_b}^\Omega(f)$ of any $f \in \mathcal{O}_{M \times N}(\Omega)$ define a family $f_{ab} \in \bar{\mathcal{F}}(\Omega)$ whose image by \mathfrak{b}_Ω is f . Therefore, if $\Omega \subset U_i \times V_j$, we have a 1:1 correspondence

$$\mathfrak{b}_\Omega : \bar{\mathcal{F}}(\Omega) \xrightarrow{\sim} \mathcal{O}_{M \times N}(\Omega). \quad (23)$$

Moreover,

$$P_{\Omega'}^\Omega \circ \mathfrak{b}_\Omega = \mathfrak{b}_{\Omega'} \circ \bar{\rho}_{\Omega'}^\Omega. \quad (24)$$

Indeed, if $(f_{ab})_{ab} \in \bar{\mathcal{F}}(\Omega)$, the LHS map sends this family f_{ab} , $U_a \times V_b \subset \Omega$, first to the unique $f \in \mathcal{O}_{M \times N}(\Omega)$ such that $P_{U_a \times V_b}^\Omega(f) = f_{ab}$, and then to the restriction $P_{\Omega'}^\Omega(f)$. The RHS map sends this family first to the subfamily $f_{\alpha\beta}$, $U_\alpha \times V_\beta \subset \Omega'$, then to the unique $g \in \mathcal{O}_{M \times N}(\Omega')$ such that $P_{U_\alpha \times V_\beta}^{\Omega'}(g) = f_{\alpha\beta}$. It is clear that $g = P_{\Omega'}^\Omega(f)$. Hence,

$$\mathfrak{b} : \bar{\mathcal{F}}|_{U_i \times V_j} \xrightarrow{\sim} \mathcal{O}_{M \times N}|_{U_i \times V_j}$$

is a presheaf isomorphism, so that $\bar{\mathcal{F}}|_{U_i \times V_j}$ is a sheaf.

We denote the sheafification of the presheaf $\bar{\mathcal{F}}$ by $\varphi : \bar{\mathcal{F}} \rightarrow \bar{\mathcal{F}}^+$ ($\bar{\rho}^+$ refers to the restrictions of $\bar{\mathcal{F}}^+$). Recall that any presheaf and its sheafification have the same stalks, i.e., that the maps

$$\varphi_{m,n} : \bar{\mathcal{F}}_{m,n} \xrightarrow{\sim} \bar{\mathcal{F}}_{m,n}^+,$$

$(m, n) \in M \times N$, induced on stalks by the presheaf morphism φ are isomorphisms. Therefore, the sheaf morphism $\varphi|_{U_i \times V_j} : \bar{\mathcal{F}}|_{U_i \times V_j} \rightarrow \bar{\mathcal{F}}^+|_{U_i \times V_j}$ is a sheaf isomorphism. This means that

$$\varphi_\Omega : \bar{\mathcal{F}}(\Omega) \xrightarrow{\sim} \bar{\mathcal{F}}^+(\Omega)$$

is an isomorphism, or, here, a 1:1 correspondence, for any open $\Omega \subset U_i \times V_j$, so that

$$\iota_\Omega := \mathfrak{b}_\Omega \circ \varphi_\Omega^{-1} : \bar{\mathcal{F}}^+(\Omega) \xrightarrow{\sim} \mathcal{O}_{M \times N}(\Omega)$$

is also 1:1. In particular, for any $U_i \times V_j \in \mathfrak{B}$, we have

$$\iota_{U_i \times V_j} : \bar{\mathcal{F}}^+(U_i \times V_j) \xrightarrow{\sim} \mathcal{O}_{M \times N}(U_i \times V_j) . \quad (25)$$

Since φ commutes with restrictions, we get, for any $U_i \times V_j \subset U_i \times U_j$,

$$\bar{\rho}_{U_i \times V_j}^{U_i \times V_j} \circ \varphi_{U_i \times U_j}^{-1} = \varphi_{U_i \times V_j}^{-1} \circ \bar{\rho}_{U_i \times V_j}^{U_i \times U_j} ,$$

so that, when taking also (24) into account, we obtain

$$\mathbf{P}_{U_i \times V_j}^{U_i \times V_j} \circ \iota_{U_i \times V_j} = \iota_{U_i \times V_j} \circ \bar{\rho}_{U_i \times V_j}^{U_i \times U_j} . \quad (26)$$

Due to (25) and (26), the map ι is a \mathfrak{B} -sheaf isomorphism between $\bar{\mathcal{F}}^+$ and $\mathcal{O}_{M \times N}$ viewed as \mathfrak{B} -sheaves. Since a \mathfrak{B} -sheaf morphism extends to a unique sheaf morphism, there exists a sheaf isomorphism

$$\mathbf{I} : \bar{\mathcal{F}}^+ \xrightarrow{\sim} \mathcal{O}_{M \times N} .$$

The morphism \mathbf{I} is actually an isomorphism of sheaves of \mathbb{Z}_2^n -commutative \mathbb{R} -algebras. It suffices to show that ι is an isomorphism of \mathfrak{B} -sheaves of such algebras, i.e., that $\iota_{U_i \times V_j}$ is a morphism of \mathbb{Z}_2^n -graded unital \mathbb{R} -algebras. We will prove that ι_Ω , $\Omega \subset U_i \times V_j$, is an algebra morphism, leaving the remaining checks to the reader. The space $\mathcal{F}(U \times V)$ is a nuclear Fréchet algebra, because it is the completed tensor product of nuclear Fréchet algebras [Eme17, Lemma 1.2.13]. Its multiplication $\bullet_{\mathcal{F}}$ is continuous. It is given by

$$\sum_{i=0}^{\infty} f_i \otimes g_i \bullet_{\mathcal{F}} \sum_{j=0}^{\infty} h_j \otimes k_j = \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} (-1)^{\langle g_i, h_j \rangle} (f_i h_j) \otimes (g_i k_j) .$$

The multiplication $\bullet_{\mathcal{F}}$ induces a multiplication $\bullet_{\bar{\mathcal{F}}}$ on $\bar{\mathcal{F}}(\Omega)$, $\Omega \subset M \times N$, which is defined by

$$(f_{ab})_{ab} \bullet_{\bar{\mathcal{F}}} (g_{ab})_{ab} = (f_{ab} \bullet_{\mathcal{F}} g_{ab})_{ab} .$$

Addition and scalar multiplication on $\bar{\mathcal{F}}(\Omega)$ are defined similarly. As $\bar{\mathcal{F}}$ is thus a presheaf of algebras, its sheafification $\bar{\mathcal{F}}^+$ is a sheaf of algebras and the $\varphi_\Omega : \bar{\mathcal{F}}(\Omega) \rightarrow \bar{\mathcal{F}}^+(\Omega)$ are algebra morphisms, see Subsection 5.4 of the Appendix. For $\Omega \subset U_i \times V_j$, this morphism φ_Ω is an algebra isomorphism and so is φ_Ω^{-1} . The map $\mathbf{b}_\Omega : \bar{\mathcal{F}}(\Omega) \rightarrow \mathcal{O}_{M \times N}(\Omega)$ associates to each $(f_{ab})_{ab}$ the f such that $\mathbf{P}_{U_a \times V_b}^\Omega(f) = f_{ab}$. Hence, the image by \mathbf{b}_Ω of a product $(f_{ab})_{ab} \bullet_{\bar{\mathcal{F}}} (g_{ab})_{ab}$ is the function h such that $\mathbf{P}_{U_a \times V_b}^\Omega(h) = f_{ab} \bullet_{\mathcal{F}} g_{ab}$. On the other hand, the product $f \cdot g$ in $\mathcal{O}_{M \times N}(\Omega)$ of the images by \mathbf{b}_Ω satisfies

$$\mathbf{P}_{U_a \times V_b}^\Omega(f \cdot g) = \mathbf{P}_{U_a \times V_b}^\Omega(f) \cdot \mathbf{P}_{U_a \times V_b}^\Omega(g) = f_{ab} \cdot g_{ab} = f_{ab} \bullet_{\mathcal{F}} g_{ab} ,$$

see Theorem 14. It follows that $h = f \cdot g$. The map \mathbf{b}_Ω is in fact an algebra morphism. Finally $\iota_\Omega = \mathbf{b}_\Omega \circ \varphi_\Omega^{-1}$ is a morphism of algebras, as needed. \square

Remark 17. *In view of (23), (21), and (19) (as well as in view of (9) and (13)), it is clear that, for any $U_i \times V_j \in \mathfrak{B}$, we have*

$$\mathcal{O}_{M \times N}(U_i \times V_j) = \mathcal{O}_M(U_i) \widehat{\otimes} \mathcal{O}_N(V_j) ,$$

but we were unable to convince ourselves that the same holds true for any $U \times V \in \mathfrak{B}$.

Indeed, it is well-known that tensor products of sheaves (and in particular completed tensor products of function sheaves) require a sheafification (see [Sar08, Section 3]). However, section spaces of the sheafification of a presheaf do not agree with the corresponding section spaces of the presheaf.

On the other hand, attempts to get rid of the problem in Remark 17 using the reconstruction results from Section 3 below, are not really promising.

Further, although $\mathcal{O}_{M \times N}$ and $\bar{\mathcal{F}}$ are two presheaves that extend the \mathfrak{B} -presheaf $\mathcal{O}_M \hat{\otimes} \mathcal{O}_N$, they do not necessarily coincide: \mathfrak{B} -sheaves have unique extensions, but \mathfrak{B} -presheaves do not. Indeed, to show that $\mathcal{O}_{M \times N} \simeq \bar{\mathcal{F}}$, we would have to decompose sections of $\mathcal{O}_{M \times N}$ into sections of the \mathfrak{B} -presheaf and then reglue them in $\bar{\mathcal{F}}$, which is impossible, since $\bar{\mathcal{F}}$ is only a presheaf.

There is actually a condition for the presheaf $\bar{\mathcal{F}}$ that extends the \mathfrak{B} -presheaf $\mathcal{F} = \mathcal{O}_M \hat{\otimes} \mathcal{O}_N$ to be a sheaf.

The explanation of this result needs some preparation.

For any open $U \times V \subset M \times N$, we set $\mathcal{O}_{M \times N}^M(U \times V) := \mathcal{O}_M(U)$. Similarly, for any open $U' \times V' \subset U \times V$, we define

$$r_{U' \times V'}^{U \times V} : \mathcal{O}_{M \times N}^M(U \times V) \rightarrow \mathcal{O}_{M \times N}^M(U' \times V')$$

to be $\rho_{U'}^U : \mathcal{O}_M(U) \rightarrow \mathcal{O}_M(U')$. It is straightforwardly checked that $\mathcal{O}_{M \times N}^M$ is a nuclear Fréchet sheaf of algebras, hence, in particular a nuclear locally convex topological sheaf of algebras. The assignment

$$\mathcal{F} : U \times V \mapsto \mathcal{O}_{M \times N}^M(U \times V) \hat{\otimes} \mathcal{O}_{M \times N}^N(U \times V) = \mathcal{O}_M(U) \hat{\otimes} \mathcal{O}_N(V)$$

defines a presheaf $\bar{\mathcal{F}}$ on $M \times N$. Applying [Mal70, Equation (2.2)], we *would* get

$$\mathcal{O}_{M \times N}(U \times V) \simeq \bar{\mathcal{F}}^+(U \times V) \simeq \mathcal{O}_M(U) \hat{\otimes} \mathcal{O}_N(V), \quad (27)$$

for any open $U \times V \subset M \times N$, if $\mathcal{O}_{M \times N}^M$ or $\mathcal{O}_{M \times N}^N$ determined a topologically dual weakly flabby precosheaf.

Just as a presheaf \mathcal{G} on a topological space T with values in a concrete category \mathbf{C} is a contravariant functor $\mathcal{G} : \mathbf{Open}(T)^{\text{op}} \rightarrow \mathbf{C}$, a *precosheaf* \mathcal{H} on T with values in \mathbf{C} is a covariant functor $\mathcal{H} : \mathbf{Open}(T) \rightarrow \mathbf{C}$. The point is that, for open subsets $V \subset U$, there is a \mathbf{C} -morphism $e_V^U : \mathcal{H}(V) \rightarrow \mathcal{H}(U)$, which we refer to as *extension morphism*. The relevant example for our purpose is the topologically dual precosheaf \mathcal{V}' of vector spaces of a topological sheaf \mathcal{V} of vector spaces on a Hausdorff space T . This precosheaf is defined, for any open $U \subset T$, by

$$\mathcal{V}'(U) = \text{Hom}_{\text{TVS}}(\mathcal{V}(U), \mathbb{R}),$$

and, for any open subsets $V \subset U$, by

$$e_V^U = {}^t R_V^U : \mathcal{V}'(V) \ni \ell \mapsto [({}^t R_V^U)\ell : \mathcal{V}(U) \ni v \mapsto \ell(R_V^U v) \in \mathbb{R}] \in \mathcal{V}'(U),$$

where R_V^U denotes the restriction in \mathcal{V} . The precosheaf \mathcal{V}' is *weakly flabby* if, for any open $U \subset T$, the morphism $e_T^U : \mathcal{V}'(U) \rightarrow \mathcal{V}'(T)$ is surjective.

We should prove that the topologically dual precosheaf of $\mathcal{V} = \mathcal{O}_{M \times N}^M$ is weakly flabby, i.e., that, for any open $U \subset M$ and for any $L \in \text{Hom}_{\text{TVS}}(\mathcal{O}_M(M), \mathbb{R})$, there exists $\ell \in \text{Hom}_{\text{TVS}}(\mathcal{O}_M(U), \mathbb{R})$, such that ${}^t\rho_U^M \ell = L$. It turns out that this condition is not satisfied, so that we cannot conclude that (27) holds. Indeed, assume that the condition is satisfied, so that it is in particular valid for $\mathcal{M} = (\mathbb{R}, C_{\mathbb{R}}^\infty)$. Choose now any $x \in \mathbb{R}$ and any open interval $I \subset \mathbb{R}$ that does not contain x . The evaluation map

$$\varepsilon_x : C_{\mathbb{R}}^\infty(\mathbb{R}) \ni f \mapsto f(x) \in \mathbb{R}$$

is linear. It is also continuous, since there exists a compact $C \subset \mathbb{R}$ that contains x and since, for any $f \in C_{\mathbb{R}}^\infty(\mathbb{R})$, we have $|f(x)| \leq \sup_C |f|$. In view of our assumption, there exists $\ell \in \text{Hom}_{\text{TVS}}(C_{\mathbb{R}}^\infty(I), \mathbb{R})$, such that, for any $f \in C_{\mathbb{R}}^\infty(\mathbb{R})$, we have $\ell(f|_I) = f(x)$. If we take now two functions $f, g \in C_{\mathbb{R}}^\infty(\mathbb{R})$ that coincide in I and have different values at x , we get the contradiction

$$f(x) = \ell(f|_I) = \ell(g|_I) = g(x) .$$

4.3 Categorical products of \mathbb{Z}_2^n -manifolds

We recommend to first read Subsections 5.4 and 5.5 of the Appendix.

Lemma 18. *Let $\mathcal{M}_1, \mathcal{M}_2 \in \mathbb{Z}_2^n\text{Man}$. The presheaf $\bar{\mathcal{F}}$ considered in Theorem 16 is an object of the category $\text{PSh}(M_1 \times M_2, \text{LCTAlg})$ of presheaves of locally convex topological algebras over $M_1 \times M_2$.*

Proof. In the proof of Theorem 16 we showed that $\bar{\mathcal{F}} \in \text{PSh}(M_1 \times M_2, \text{Alg})$ for the obvious restrictions and algebra operations.

Recall that, for any open $\Omega \subset M_1 \times M_2$, the algebra $\bar{\mathcal{F}}(\Omega)$ is given by

$$\bar{\mathcal{F}}(\Omega) := \{(f_{ab})_{ab} : f_{ab} \in \mathcal{F}(U_a \times V_b), U_a \times V_b \subset \Omega, \text{ such that } \rho_{U_{aa} \times V_{bb}}^{U_a \times V_b}(f_{ab}) = \rho_{U_{aa} \times V_{bb}}^{U_a \times V_b}(f_{ab})\}$$

and is thus a subalgebra of

$$\prod_{U_a \times V_b \subset \Omega} \mathcal{F}(U_a \times V_b) = \prod_{U_a \times V_b \subset \Omega} \mathcal{O}_{M_1}(U_a) \hat{\otimes} \mathcal{O}_{M_2}(V_b) .$$

We equip $\bar{\mathcal{F}}(\Omega)$ with the topology induced by the product of the topologies of the locally convex topological algebras (LCTA-s) $\mathcal{O}_{M_1}(U_a) \hat{\otimes} \mathcal{O}_{M_2}(V_b)$. Since a product of LCTVS-s and a subspace of a LCTVS are themselves LCTVS-s, the algebra $\bar{\mathcal{F}}(\Omega)$ is a LCTVS. Its multiplication

$$(f_{ab})_{ab} \bullet_{\bar{\mathcal{F}}} (g_{ab})_{ab} = (f_{ab} \bullet_{\mathcal{F}} g_{ab})_{ab}$$

is continuous, since the multiplication $\bullet_{\mathcal{F}}$ is, see Proof of Theorem 16 and Lemma 32. Hence, the space $\bar{\mathcal{F}}(\Omega)$ is a LCTA.

A restriction $\bar{\rho}_{\Omega'}^{\Omega}$ ($\Omega' \subset \Omega$) – it sends any family $(f_{ab})_{ab}$ of $\bar{\mathcal{F}}(\Omega)$ indexed by the $U_a \times V_b \subset \Omega$ to the family $(f_{ab})_{ab}$ of $\bar{\mathcal{F}}(\Omega')$ indexed by the $U_a \times V_b \subset \Omega'$ – is known to be an algebra morphism. It is continuous, since it is continuous as a map

$$\bar{\rho}_{\Omega'}^{\Omega} : \prod_{U_a \times V_b \subset \Omega} \mathcal{F}(U_a \times V_b) \rightarrow \prod_{U_a \times V_b \subset \Omega'} \mathcal{F}(U_a \times V_b),$$

in view of the definition of the product topology. \square

Theorem 19. *The category $\mathbb{Z}_2^n\mathbf{Man}$ has all finite products.*

Proof. Since $\mathbb{Z}_2^n\mathbf{Man}$ has a terminal object (see Corollary 12), it suffices to prove that it has binary products. Let $\mathcal{M}_1, \mathcal{M}_2 \in \mathbb{Z}_2^n\mathbf{Man}$. We will show that the product \mathbb{Z}_2^n -manifold $\mathcal{M}_1 \times \mathcal{M}_2$ (see Definition 13) is the categorical binary product of \mathcal{M}_1 and \mathcal{M}_2 .

We first define \mathbb{Z}_2^n -morphisms

$$\Pi_i = (\pi_i, \pi_i^*) : \mathcal{M}_1 \times \mathcal{M}_2 \rightarrow \mathcal{M}_i, \quad i \in \{1, 2\}.$$

The base maps $\pi_i : \mathcal{M}_1 \times \mathcal{M}_2 \rightarrow \mathcal{M}_i$ are the canonical smooth projections. In the following, we consider the case $i = 1$ and use the notation introduced above.

The maps

$$T_U : \mathcal{O}_{M_1}(U) \ni f \mapsto f \otimes 1 \in (\pi_{1,*}\bar{\mathcal{F}})(U),$$

$U \in \mathbf{Open}(M_1)$, define a morphism $T : \mathcal{O}_{M_1} \rightarrow \pi_{1,*}\bar{\mathcal{F}}$ in $\mathbf{PSh}(M_1, \mathbf{LCTAlg})$. Indeed, we have $\mathcal{O}_{M_1}, \pi_{1,*}\bar{\mathcal{F}} \in \mathbf{PSh}(M_1, \mathbf{LCTAlg})$, see [BP18, Theorem 14] and Proof of Proposition 39. It is easy to see that the maps T_U commute with restrictions and are algebra morphisms. To show that T_U is continuous, it suffices to check that the linear map

$$T_U : \mathcal{O}_{M_1}(U) \ni f \mapsto f \otimes 1 \in \mathcal{O}_{M_1}(U) \otimes \mathcal{O}_{M_2}(M_2) \quad (28)$$

is continuous for the projective tensor topology on the target. We apply Theorem 30 and Proposition 31. Let $p_{C,D} \otimes p_{K,\Delta}$ be any of the seminorms that induce the projective tensor topology. Recall that $C \subset U$ and $K \subset M_2$ are compact subsets, and that D and Δ are differential operators acting on $\mathcal{O}_{M_1}(U)$ and $\mathcal{O}_{M_2}(M_2)$, respectively. We must prove that there is a finite number of seminorms p_{C_k, D_k} on the source and a constant $C > 0$, such that

$$p_{C,D}(f) \cdot p_{K,\Delta}(1) \leq C \max_k p_{C_k, D_k}(f),$$

for any $f \in \mathcal{O}_{M_1}(U)$. It suffices to use a single source seminorm p_{C_k, D_k} , namely $p_{C,D}$. Indeed, if $\mathfrak{C} = p_{K,\Delta}(1) = 0$, the previous condition is satisfied with $C = 1$, and if $\mathfrak{C} > 0$, it is satisfied with $C = \mathfrak{C}$.

It follows from Proposition 35 that $T^+ : \mathcal{O}_{M_1}^+ \rightarrow (\pi_{1,*}\bar{\mathcal{F}})^+$ is a morphism in $\mathbf{Sh}(M_1, \mathbf{LCTAlg})$. Moreover, in view of Proposition 39, there is an $\mathbf{Sh}(M_1, \mathbf{LCTAlg})$ -morphism $\iota : (\pi_{1,*}\bar{\mathcal{F}})^+ \rightarrow$

$\pi_{1,*}\bar{\mathcal{F}}^+$. The composite $\iota \circ T^+ : \mathcal{O}_{M_1}^+ \rightarrow \pi_{1,*}\bar{\mathcal{F}}^+$ is a morphism in $\mathbf{Sh}(M_1, \mathbf{LCTAlg})$. Proposition 35 and Theorem 16 allow us to interpret it as a morphism

$$\pi_1^* = \iota \circ T^+ : \mathcal{O}_{M_1} \rightarrow \pi_{1,*}\mathcal{O}_{M_1 \times M_2} \quad (29)$$

in $\mathbf{Sh}(M_1, \mathbf{Alg})$. The map π_1^* is actually a morphism of sheaves of \mathbb{Z}_2^n -commutative associative unital algebras, so that the map $\Pi_1 = (\pi_1, \pi_1^*) : \mathcal{M}_1 \times \mathcal{M}_2 \rightarrow \mathcal{M}_1$ is a morphism in $\mathbb{Z}_2^n\mathbf{Man}$ (the results of the appendix we use in this proof extend obviously to the graded unital setting: when speaking in the rest of the proof about algebras, we actually mean \mathbb{Z}_2^n -commutative associative unital algebras).

It remains to check the universality of our construction. Let $\mathcal{N} \in \mathbb{Z}_2^n\mathbf{Man}$ and let $\Phi_i = (\phi_i, \phi_i^*) : \mathcal{N} \rightarrow \mathcal{M}_i$ be morphisms in $\mathbb{Z}_2^n\mathbf{Man}$. We will prove that there exists a unique $\mathbb{Z}_2^n\mathbf{Man}$ -morphism $\Psi = (\psi, \psi^*) : \mathcal{N} \rightarrow \mathcal{M}_1 \times \mathcal{M}_2$, such that $\Pi_i \circ \Psi = \Phi_i$, for both i .

We set $\psi = (\phi_1, \phi_2) : \mathcal{N} \rightarrow M_1 \times M_2$. As for ψ^* , observe that, for any open $U_i \subset M_i$, the map $\phi_{i,U_i}^* : \mathcal{O}_{M_i}(U_i) \rightarrow \mathcal{O}_{\mathcal{N}}(\phi_i^{-1}(U_i))$ is a continuous algebra (\mathbb{Z}_2^n -graded unital algebra) morphism, see [BP18, Theorem 19]. Denote now by $V = V_1 \cap V_2$ the open subset

$$V = V_1 \cap V_2 = \phi_1^{-1}(U_1) \cap \phi_2^{-1}(U_2) = \psi^{-1}(U_1 \times U_2) \subset \mathcal{N},$$

and denote by $m_V = - \cdot -$ the multiplication of $\mathcal{O}_{\mathcal{N}}(V)$. In view of [BP18, Theorem 14] and Propositions 37 and 38, the map

$$\mathfrak{p}_{U_1 \times U_2} = \widehat{m}_V \circ (\rho_V^{V_1} \widehat{\otimes} \rho_V^{V_2}) \circ (\phi_{1,U_1}^* \widehat{\otimes} \phi_{2,U_2}^*) : \mathcal{O}_{M_1}(U_1) \widehat{\otimes} \mathcal{O}_{M_2}(U_2) \rightarrow \mathcal{O}_{\mathcal{N}}(V),$$

where $\rho_V^{V_1}$ and $\rho_V^{V_2}$ are restrictions in $\mathcal{O}_{\mathcal{N}}$, is a continuous algebra morphism between nuclear Fréchet algebras. The maps

$$\mathfrak{p}_{U_1 \times U_2} : \mathcal{F}(U_1 \times U_2) \ni \sum_{j=0}^{\infty} f_j \otimes g_j \mapsto \sum_{j=0}^{\infty} \rho_V^{V_1} \phi_{1,U_1}^* f_j \cdot \rho_V^{V_2} \phi_{2,U_2}^* g_j \in (\psi_*\mathcal{O}_{\mathcal{N}})(U_1 \times U_2) \quad (30)$$

define a morphism of \mathcal{B} -presheaves of locally convex topological algebras. The latter extends to a morphism $\bar{\mathfrak{p}} : \bar{\mathcal{F}} \rightarrow \psi_*\mathcal{O}_{\mathcal{N}}$ of presheaves of locally convex topological algebras over $M_1 \times M_2$.

In view of Propositions 35 and 39, there are morphisms

$$\bar{\mathfrak{p}}^+ : \bar{\mathcal{F}}^+ \rightarrow (\psi_*\mathcal{O}_{\mathcal{N}})^+ \quad \text{and} \quad \iota : (\psi_*\mathcal{O}_{\mathcal{N}})^+ \rightarrow \psi_*\mathcal{O}_{\mathcal{N}}^+$$

in $\mathbf{Sh}(M_1 \times M_2, \mathbf{LCTAlg})$. As above, we can view their composite as a morphism

$$\psi^* = \iota \circ \bar{\mathfrak{p}}^+ : \mathcal{O}_{M_1 \times M_2} \rightarrow \psi_*\mathcal{O}_{\mathcal{N}} \quad (31)$$

of sheaves of algebras, so that $\Psi = (\psi, \psi^*) : \mathcal{N} \rightarrow \mathcal{M}_1 \times \mathcal{M}_2$ is a morphism in $\mathbb{Z}_2^n\mathbf{Man}$.

To prove that $\Pi_i \circ \Psi = \Phi_i$, it suffices to show that

$$\psi_{M_1 \times M_2}^* \circ \pi_{i,M_i}^* = \beta(\Pi_i \circ \Psi) = \beta(\Phi_i) = \phi_{i,M_i}^*,$$

see Theorem 9. The latter is a straightforward consequence of Equations (28), (29), (30), (31), (38), and (40).

Let now $X = (\chi, \chi^*) : \mathcal{N} \rightarrow \mathcal{M}_1 \times \mathcal{M}_2$ be another $\mathbb{Z}_2^n \mathbf{Man}$ -morphism that satisfies $\Pi_i \circ X = \Phi_i$. As the category of smooth manifolds has finite products, we get $\chi = \psi$. We will check that $\beta(X) = \beta(\Psi)$, i.e., that, for all $\sigma \in \bar{\mathcal{F}}^+(M_1 \times M_2)$, we have

$$\chi_{M_1 \times M_2}^* \sigma = \psi_{M_1 \times M_2}^* \sigma \in \mathcal{O}_N(N) .$$

It suffices to show that these sections coincide in a neighborhood of an arbitrary point $n_0 \in N$. We use the compact notation $m \in M$ instead of $(m_1, m_2) \in M_1 \times M_2$. Recall that

$$\sigma = ([s]_m)_{m \in M} ,$$

where $s \in \mathcal{F}(U)$ ($U = U_1 \times U_2 \ni m$) reads $s = \sum_{j=0}^{\infty} f_j \otimes g_j$ ($f_j \in \mathcal{O}_{M_1}(U_1)$, $g_j \in \mathcal{O}_{M_2}(U_2)$).

In view of (30), (31), (38), and (40),

$$\psi_{M_1 \times M_2}^* \sigma = ([\sum_{j=0}^{\infty} \rho_V^{V_1} \phi_{1,U_1}^* f_j \cdot \rho_V^{V_2} \phi_{2,U_2}^* g_j]_n)_{n \in N} , \quad (32)$$

where we take the germ at n of the section induced by the representative s of the germ at $\psi(n) \in M$. Let $m_0 = \psi(n_0) \in M$. Since s is constant in a neighborhood $U_0 = U_{1,0} \times U_{2,0}$ of m_0 , the representative in the RHS of the preceding equation is constant in the neighborhood

$$V_0 = V_{1,0} \cap V_{2,0} = \phi_1^{-1}(U_{1,0}) \cap \phi_2^{-1}(U_{2,0}) = \psi^{-1}(U_0)$$

of n_0 . Hence,

$$\rho_{V_0}^N \psi_{M_1 \times M_2}^* \sigma = \sum_{j=0}^{\infty} \rho_{V_0}^{V_{1,0}} \phi_{1,U_{1,0}}^* f_j \cdot \rho_{V_0}^{V_{2,0}} \phi_{2,U_{2,0}}^* g_j \in \mathcal{O}_N(V_0) .$$

On the other hand, since $\Pi_i \circ X = \Phi_i$, we have, for any open $U_1 \subset M_1$ and any $f \in \mathcal{O}_{M_1}(U_1)$,

$$\chi_{U_1 \times M_2}^* ([f \otimes 1]_m)_{m \in U_1 \times M_2} = \phi_{1,U_1}^* f ,$$

due to (28), (29), (38), and (40). For any open $U_2 \subset M_2$, we get

$$\chi_U^* ([f \otimes 1]_m)_{m \in U} = \rho_{\psi^{-1}(U)}^{\psi^{-1}(U_1 \times M_2)} \chi_{U_1 \times M_2}^* ([f \otimes 1]_m)_{m \in U_1 \times M_2} = \rho_V^{V_1} \phi_{1,U_1}^* f . \quad (33)$$

An analogous result holds for $i = 2$. Observe now that

$$\begin{aligned} \sigma &= ([\sum_{j=0}^{\infty} f_j \otimes g_j]_m)_{m \in M} = (\pi_U^m \lim_N \sum_{j=0}^N f_j \otimes g_j)_{m \in M} = \\ &= (\lim_N \sum_{j=0}^N [f_j \otimes g_j]_m)_{m \in M} = \sum_{j=0}^{\infty} ([f_j \otimes g_j]_m)_{m \in M} , \end{aligned} \quad (34)$$

see Proof of Proposition 35. As the pullbacks χ^* and the restriction $\rho_{V_0}^N$ are continuous algebra morphisms, we get

$$\begin{aligned} \rho_{V_0}^N \chi_{M_1 \times M_2}^* \sigma &= \sum_{j=0}^{\infty} \rho_{\psi^{-1}(U_0)}^{\psi^{-1}(M)} \chi_M^* ([f_j \otimes g_j]_m)_{m \in M} = \\ &= \sum_{j=0}^{\infty} \chi_{U_0}^* ([f_j \otimes 1]_m)_{m \in U_0} \cdot \chi_{U_0}^* ([1 \otimes g_j]_m)_{m \in U_0} = \sum_{j=0}^{\infty} \rho_{V_0}^{V_{1,0}} \phi_{1,U_{1,0}}^* f_j \cdot \rho_{V_0}^{V_{2,0}} \phi_{2,U_{2,0}}^* g_j, \end{aligned}$$

due to (33). \square

4.4 Products of \mathbb{Z}_2^n -morphisms

We use again abbreviations of the type $n = (n_1, n_2) \in N = N_1 \times N_2$ (which we introduced in the proof of Theorem 19).

Proposition 20. *Let $\Psi_i : \mathcal{M}_i \rightarrow \mathcal{N}_i$, $i \in \{1, 2\}$, be a \mathbb{Z}_2^n -morphism with base map ψ_i and pullback sheaf morphism ψ_i^* . Due to the universality of the product of \mathbb{Z}_2^n -manifolds, there is a canonical \mathbb{Z}_2^n -morphism*

$$\Psi = \Psi_1 \times \Psi_2 : \mathcal{M}_1 \times \mathcal{M}_2 \rightarrow \mathcal{N}_1 \times \mathcal{N}_2.$$

Its base map is $\psi = \psi_1 \times \psi_2$ and its pullback sheaf morphism $\psi^* = (\psi_1 \times \psi_2)^*$ is given, for each open subset $\Omega \subset N_1 \times N_2$, by

$$\psi_{\Omega}^* = \left(\prod_{m \in \psi^{-1}(\Omega)} (\psi_1^* \widehat{\otimes} \psi_2^*)_{\psi(m)} \right) \circ (p_{\psi(m)})_{m \in \psi^{-1}(\Omega)}.$$

The first map in the RHS is the product of the morphisms between stalks induced by the morphism

$$\psi_1^* \widehat{\otimes} \psi_2^* : \bar{\mathcal{F}}_N \rightarrow \psi_* \bar{\mathcal{F}}_M$$

(of presheaves of locally convex topological algebras), where $\bar{\mathcal{F}}_N$ is the presheaf defined by $\mathcal{O}_{N_1} \widehat{\otimes} \mathcal{O}_{N_2}$, and similarly for $\bar{\mathcal{F}}_M$. The second map in the RHS is the tuple of morphisms

$$p_{\psi(m)} : \prod_{n \in \Omega} \bar{\mathcal{F}}_{N,n} \rightarrow \bar{\mathcal{F}}_{N,\psi(m)}.$$

To understand this claim, recall that

$$\psi_{\Omega}^* : \mathcal{O}_N(\Omega) \simeq \bar{\mathcal{F}}_N^+(\Omega) \subset \prod_{n \in \Omega} \bar{\mathcal{F}}_{N,n} \rightarrow \mathcal{O}_M(\psi^{-1}(\Omega)) \simeq \bar{\mathcal{F}}_M^+(\psi^{-1}(\Omega)) \subset \prod_{m \in \psi^{-1}(\Omega)} \bar{\mathcal{F}}_{M,m}.$$

Note now that

$$(p_{\psi(m)})_{m \in \psi^{-1}(\Omega)} : \mathcal{O}_N(\Omega) \rightarrow \prod_{m \in \psi^{-1}(\Omega)} \bar{\mathcal{F}}_{N,\psi(m)}$$

and that

$$\prod_{m \in \psi^{-1}(\Omega)} (\psi_1^* \widehat{\otimes} \psi_2^*)_{\psi(m)} : \prod_{m \in \psi^{-1}(\Omega)} \bar{\mathcal{F}}_{N,\psi(m)} \rightarrow \prod_{m \in \psi^{-1}(\Omega)} \bar{\mathcal{F}}_{M,m},$$

so that the composite of these maps may coincide with ψ_{Ω}^* .

Proof. For $i \in \{1, 2\}$, we denote by Π_i^S (resp., Π_i^T) the \mathbb{Z}_2^n -morphism $\Pi_i^S : \mathcal{M}_1 \times \mathcal{M}_2 \rightarrow \mathcal{M}_i$ (resp., $\Pi_i^T : \mathcal{N}_1 \times \mathcal{N}_2 \rightarrow \mathcal{N}_i$). The composite $\Phi_i = \Psi_i \circ \Pi_i^S$ is a \mathbb{Z}_2^n -morphism $\Phi_i : \mathcal{M}_1 \times \mathcal{M}_2 \rightarrow \mathcal{N}_i$. In view of the universality of the product $\mathcal{N}_1 \times \mathcal{N}_2$, there is a unique \mathbb{Z}_2^n -morphism $\Psi : \mathcal{M}_1 \times \mathcal{M}_2 \rightarrow \mathcal{N}_1 \times \mathcal{N}_2$, such that $\Pi_i^T \circ \Psi = \Phi_i$. We denote this morphism Ψ by $\Psi_1 \times \Psi_2$ and refer to it as the product of the \mathbb{Z}_2^n -morphisms Ψ_i . We showed in the proof of Theorem 19 that the base map of Ψ is

$$\psi = (\psi_1 \circ \pi_1^S, \psi_2 \circ \pi_2^S) = \psi_1 \times \psi_2 .$$

We investigate now the pullback morphisms ψ^* . Let $\Omega \subset N$ be open, so that $\psi_\Omega^* : \mathcal{O}_N(\Omega) \rightarrow \mathcal{O}_M(\psi^{-1}(\Omega))$, and let $\sigma \in \mathcal{O}_N(\Omega) \simeq \bar{\mathcal{F}}_N^+(\Omega)$. Recall once again that $\sigma = ([s]_n)_{n \in \Omega}$, where $s \in \mathcal{F}_N(U)$ ($U = U_1 \times U_2 \ni n$) reads

$$s = \sum_{j=0}^{\infty} f_j \otimes g_j \quad (f_j \in \mathcal{O}_{N_1}(U_1), g_j \in \mathcal{O}_{N_2}(U_2)) .$$

It follows from Equation (32) that

$$\psi_\Omega^* \sigma = \left(\left[\sum_{j=0}^{\infty} \rho_V^{V_1} (\pi_1^S)_{W_1}^* \psi_{1,U_1}^* f_j \cdot \rho_V^{V_2} (\pi_2^S)_{W_2}^* \psi_{2,U_2}^* g_j \right]_m \right)_{m \in \psi^{-1}(\Omega)} , \quad (35)$$

where

$$V = V_1 \cap V_2, \quad V_i = (\pi_i^S)^{-1}(W_i), \quad \text{and} \quad W_i = \psi_i^{-1}(U_i) \quad (\text{obviously } V = W_1 \times W_2) .$$

We interpret $\psi_{1,U_1}^* f_j \in \mathcal{O}_{M_1}(W_1)$ as

$$([\psi_{1,U_1}^* f_j]_{m_1})_{m_1 \in W_1} \in \mathcal{O}_{M_1}^+(W_1) ,$$

so that

$$\rho_V^{V_1} (\pi_1^S)_{W_1}^* \psi_{1,U_1}^* f_j = ([\psi_{1,U_1}^* f_j \otimes 1]_\mu)_{\mu \in V} ,$$

due to Equations (28) and (29), as well as to the observation that $m_1 \in W_1$ is equivalent to $m_1 = \pi_1^S(m)$ ($m \in V_1$). A similar result holds for the second factor in the RHS of (35), so that, in view of (34), we obtain

$$\begin{aligned} \psi_\Omega^* \sigma &= \left(\left[\sum_{j=0}^{\infty} ([\psi_{1,U_1}^* f_j \otimes \psi_{2,U_2}^* g_j]_\mu)_{\mu \in V} \right]_m \right)_{m \in \psi^{-1}(\Omega)} = \\ &= \left(\left[([\psi_{1,U_1}^* \hat{\otimes} \psi_{2,U_2}^* \sum_{j=0}^{\infty} f_j \otimes g_j]_\mu)_{\mu \in V} \right]_m \right)_{m \in \psi^{-1}(\Omega)} . \end{aligned}$$

This image is a family (indexed by m) of elements of $\bar{\mathcal{F}}_{M,m}^+ \simeq \bar{\mathcal{F}}_{M,m}$. The isomorphism between a stalk of a presheaf and the corresponding stalk of its sheafification is described in the proof of Lemma 6.17.2 of the Stacks Project. The description shows that

$$\psi_\Omega^* \sigma = \left(\left[\psi_{1,U_1}^* \hat{\otimes} \psi_{2,U_2}^* \sum_{j=0}^{\infty} f_j \otimes g_j \right]_m \right)_{m \in \psi^{-1}(\Omega)} =$$

$$\left(\prod_{m \in \psi^{-1}(\Omega)} (\psi_1^* \widehat{\otimes} \psi_2^*)_{\psi(m)} \right) \left((p_{\psi(m)})_{m \in \psi^{-1}(\Omega)} \sigma \right) .$$

□

5 Appendix

We prove and recall results on topological vector spaces and on topological algebras.

5.1 Topological vector subspaces

Definition 21. A topological vector subspace S of a TVS V (TVSS for short) is a subset $S \subset V$ which is a TVS for the linear operations and the topology of V .

Proposition 22. A subset $S \subset V$ of a TVS V is a TVSS of V if and only if S is a linear subspace of V and is endowed with the topology induced by the topology of V .

Proof. The restrictions to S of the continuous addition and scalar multiplication in V are continuous in the topology induced on S by V . □

Note also that, if V is a TVS, and if $S \subset V$ is a TVSS, then S is a TVS and the inclusion $\iota : S \ni s \mapsto s \in V$ is an injective continuous linear map. Conversely, if $S \subset V$ is a TVS and the inclusion ι is an injective continuous linear map, then the linear structure on $S \subset V$ is the same as in V , but S can have a topology that is finer than the induced one, so that S is not necessarily a TVSS.

Proposition 23. Let $\iota : V \rightarrow W$ be an injective linear map between TVS-s. When equipped with the induced topology, the linear subspace $\iota(V)$ is a TVSS of W . The bijective linear map $\tilde{\iota} : V \rightarrow \iota(V)$ is a TVS-isomorphism, i.e., a linear homeomorphism, if and only if the topology of V is the initial topology of $\iota : V \rightarrow W$. In this case, the space V can be viewed as a TVSS $V \simeq \iota(V)$ of W .

Proof. It suffices to prove the second claim. If $\tilde{\iota}$ is an isomorphism, then the topology $\mathcal{T}(V)$ of V is given by

$$\mathcal{T}(V) = \{ \tilde{\iota}^{-1}(\iota(V) \cap U_W) = \iota^{-1}(\iota(V) \cap U_W) = \iota^{-1}(\iota(V)) \cap \iota^{-1}(U_W) = \iota^{-1}(U_W) : U_W \in \mathcal{T}(W) \} ,$$

hence, it is the initial topology of $\iota : V \rightarrow W$. Conversely, if $\mathcal{T}(V)$ is the initial topology of ι , the maps $\tilde{\iota}$ and $\tilde{\iota}^{-1}$ are continuous. Indeed, as just explained, we have $\tilde{\iota}^{-1}(\iota(V) \cap U_W) = \iota^{-1}(U_W) \in \mathcal{T}(V)$, so that $\tilde{\iota}$ is continuous (but it would still be continuous for a finer topology on V). For $\tilde{\iota}^{-1}$ we have $\tilde{\iota}(\iota^{-1}(U_W)) = \iota(V) \cap U_W \in \mathcal{T}(\iota(V))$. □

Proposition 24. Let $\iota : V \rightarrow W$ be an injective linear map from a vector space V to a TVS W . When equipped with the initial topology of ι , the linear space V is a TVS and can be viewed as a TVSS of W .

Proof. The linear operations on V are continuous when V has the initial topology. Indeed, denote by $+_V$ (resp., $+_W$) the addition in V (resp., W), and let $\iota^{-1}(U_W)$, $U_W \in \mathcal{T}(W)$, be an arbitrary open subset in $\mathcal{T}(V)$. Then,

$$\begin{aligned} (+_V)^{-1}(\iota^{-1}(U_W)) &= \{(v, v') \in V \times V : \iota(v) +_W \iota(v') \in U_W\} = \\ &= \{(v, v') \in V \times V : (\iota(v), \iota(v')) \in (+_W)^{-1}(U_W)\} = (\iota \times \iota)^{-1}((+_W)^{-1}(U_W)) \end{aligned}$$

is open in $V \times V$, since ι and $+_W$ are continuous. The case of the scalar multiplication is similar. The second claim follows now from Proposition 23. \square

Remark 25. *When passing above from topological vector subspace structures on included subsets to topological vector subspace structures on injected subsets, we replaced the induced topology by the initial topology. Of course, if the injection is the inclusion, the initial topology with respect to it coincides with the induced topology.*

5.2 Completions of topological vector spaces

We recall now well-known properties of the completion of a TVS [Nag07].

For any TVS V , there is a complete TVS \widehat{V} , and an injective linear map $\iota : V \rightarrow \widehat{V}$, such that the linear subspace $\iota(V) \subset \widehat{V}$ is dense in \widehat{V} . Moreover, when endowed with the induced topology, the image $\iota(V) \subset \widehat{V}$ becomes a TVSS of \widehat{V} , and the map $\tilde{\iota} : V \rightarrow \iota(V)$ is promoted to a linear homeomorphism, or, equivalently, to a TVS-isomorphism. It follows from Proposition 23 that the topology of V is the initial topology of $\iota : V \rightarrow \widehat{V}$, and that $V \simeq \iota(V)$ is a TVSS of \widehat{V} . In short, the complete TVS \widehat{V} , which we refer to as the *completion* of the TVS V , contains V as a dense TVSS:

Proposition 26. *The completion \widehat{V} of V contains V as a dense subset, and it induces on V the original topology and original linear structure.*

Remark 27. *If V is already a complete TVS, then $\iota : V \rightarrow \widehat{V}$ is a TVS-isomorphism.*

Further, let $S \subset V$ be a TVSS of V and let $\iota : S \rightarrow V$ be the injective continuous linear inclusion. The continuous extension of ι , which we denote by $\hat{\iota} : \widehat{S} \rightarrow \widehat{V}$, is an injective continuous linear map. If $\hat{\iota}(\widehat{S})$ carries the induced topology, the map $\tilde{\hat{\iota}} : \widehat{S} \rightarrow \hat{\iota}(\widehat{S})$ is a TVS-isomorphism. In view of Proposition 23 this means that the topology of \widehat{S} is the initial topology of $\hat{\iota} : \widehat{S} \rightarrow \widehat{V}$, and that $\widehat{S} \simeq \hat{\iota}(\widehat{S})$ is a TVSS of \widehat{V} . In short:

Proposition 28. *The completion of a TVSS $S \subset V$ is a TVSS $\widehat{S} \subset \widehat{V}$ of the completion.*

5.3 Locally convex spaces

Proposition 29. *The initial topology of a linear map $\ell : V \rightarrow W$ from a vector space V to a LCTVS W endows V with a LCTVS structure. In particular, the induced topology on a vector subspace $S \subset W$ of a LCTVS endows S with a LCTVS structure.*

Proof. The topology of W has a convex basis \mathfrak{B} . As the preimage of a convex set by a linear map is convex, the family $\ell^{-1}(\mathfrak{B})$ is a convex basis of the initial topology. In view of the proof of Proposition 24, the initial topology endows V with a LCTVS structure. The second claim is a special case of the first one. \square

We close this subsection recalling two results.

Theorem 30. *Let V and W be two LCTVS-s and let $(p_i)_{i \in I}$ and $(q_j)_{j \in J}$ be two families of seminorms that induce the topologies of V and W , respectively. The projective tensor topology π on $V \otimes W$ is induced by $(p_i \otimes q_j)_{ij}$. Moreover, for any $t \in V \otimes W$, we have*

$$(p_i \otimes q_j)(t) = \inf \left\{ \sum_{k=1}^N p_i(v_k) q_j(w_k) : t = \sum_{k=1}^N v_k \otimes w_k, v_k \in V, w_k \in W, N \in \mathbb{N} \right\},$$

and, for any $v \in V$ and $w \in W$, we have

$$(p_i \otimes q_j)(v \otimes w) = p_i(v) q_j(w).$$

Proposition 31. *Let V and W be two LCTVS-s and let $(p_i)_{i \in I}$ and $(q_j)_{j \in J}$ be two families of seminorms that induce the topologies of V and W , respectively. A linear map $\ell : V \rightarrow W$ is continuous if and only if, for any $j \in J$, there exist $i_1, \dots, i_N \in I$ and $C > 0$, such that, for all $v \in V$, one has*

$$q_j(\ell(v)) \leq C \max_k p_{i_k}(v).$$

5.4 Presheaves of topological algebras and sheafification

In the following, we need two lemmas.

Unless otherwise stated, all cartesian products $\prod_{\alpha} T_{\alpha}$ of topological spaces T_{α} are endowed with the product topology, i.e., the weakest topology for which all projections $p_{\beta} : \prod_{\alpha} T_{\alpha} \rightarrow T_{\beta}$ are continuous.

Lemma 32. *Let $(T_{\alpha}^i)_{\alpha}$ ($i \in \{1, 2, 3\}$) be families of topological spaces, and let $(m_{\alpha})_{\alpha}$ be a family $m_{\alpha} : T_{\alpha}^1 \times T_{\alpha}^2 \rightarrow T_{\alpha}^3$ of continuous maps. Then the map*

$$m : \prod_{\alpha} T_{\alpha}^1 \times \prod_{\alpha} T_{\alpha}^2 \ni ((\tau_{\alpha})_{\alpha}, (\mathbf{t}_{\alpha})_{\alpha}) \mapsto (m_{\alpha}(\tau_{\alpha}, \mathbf{t}_{\alpha}))_{\alpha} \in \prod_{\alpha} T_{\alpha}^3$$

is continuous.

Proof. Since the cartesian product $\prod_{\alpha} T_{\alpha}$ of topological spaces T_{α} equipped with the product topology, is the product in the category of topological spaces, it follows from the universal property that a map $f : T \rightarrow \prod_{\alpha} T_{\alpha}$ from a topological space to a product space is continuous if and only if all the $p_{\beta} \circ f : T \rightarrow T_{\beta}$ are continuous. However, the map $p_{\beta}^3 \circ m : ((\tau_{\alpha})_{\alpha}, (\mathbf{t}_{\alpha})_{\alpha}) \mapsto m_{\beta}(\tau_{\beta}, \mathbf{t}_{\beta})$ is the composite $m_{\beta} \circ (p_{\beta}^1 \times p_{\beta}^2)$, which is of course continuous. \square

Lemma 33. *Let V be a vector space, let $(V_\alpha)_\alpha$ be a family of LCTVS, let $\ell_\alpha : V_\alpha \rightarrow V$ be a family of linear maps, and let V be equipped with the finest locally convex vector space topology, for which all the ℓ_α are continuous (V is then a LCTVS). If W is a LCTVS, a linear map $\ell : V \rightarrow W$ is continuous if and only if all the maps $\ell \circ \ell_\alpha : V_\alpha \rightarrow W$ are continuous.*

Proof. See [BS17, Proposition 2.3.5] □

In the present text:

Definition 34. *A topological algebra (TA for short) (resp., a locally convex topological algebra (LCTA for short)) is a (real) topological vector space (resp., a locally convex (real) topological vector space), with an associative, bilinear, and (jointly) continuous multiplication. A morphism of topological algebras (resp., a morphism of locally convex topological algebras) is a continuous algebra morphism. We denote the category of topological algebras (resp., of locally convex topological algebras) and morphisms between them by \mathbf{TAAlg} (resp., by $\mathbf{LCTAAlg}$).*

The category of Fréchet algebras is a full subcategory of the category of locally convex topological algebras, itself a full subcategory of the category of topological algebras.

We denote by $\mathbf{PSh}(T, \mathbf{TAAlg})$ (resp., $\mathbf{Sh}(T, \mathbf{TAAlg})$) the category of presheaves (resp., sheaves) of TA-s over a topological space T . Similarly, the category $\mathbf{PSh}(T, \mathbf{LCTAAlg})$ (resp., $\mathbf{Sh}(T, \mathbf{LCTAAlg})$) is the category of presheaves (resp., sheaves) of LCTA-s over T , and the category $\mathbf{PSh}(T, \mathbf{Alg})$ (resp., $\mathbf{Sh}(T, \mathbf{Alg})$) is the category of presheaves (resp., sheaves) of algebras over T .

Proposition 35. *Denote by $+$ the sheafification functor*

$$+ : \mathbf{PSh}(T, \mathbf{Alg}) \rightarrow \mathbf{Sh}(T, \mathbf{Alg}) : \text{For} ,$$

i.e., the left adjoint of the forgetful functor For. If $\mathcal{F} \in \mathbf{PSh}(T, \mathbf{TAAlg})$, we have $\mathcal{F}^+ \in \mathbf{Sh}(T, \mathbf{TAAlg})$, and the morphism $i : \mathcal{F} \rightarrow \mathcal{F}^+$ of presheaves of algebras is a morphism of presheaves of TA-s. Moreover, if $\varphi : \mathcal{F} \rightarrow \mathcal{F}'$ is a morphism in $\mathbf{PSh}(T, \mathbf{TAAlg})$, then $\varphi^+ : \mathcal{F}^+ \rightarrow \mathcal{F}'^+$ is a morphism in $\mathbf{Sh}(T, \mathbf{TAAlg})$. The same results hold, if we replace TA-s by LCTA-s.

Proof. We study the case $\mathcal{F} \in \mathbf{PSh}(T, \mathbf{LCTAAlg})$ (in particular $\mathcal{F} \in \mathbf{PSh}(T, \mathbf{Alg})$).

Since sheafification is based on stalks \mathcal{F}_x ($x \in T$), i.e., on inductive limits, and since the inductive limit of a directed system of sets endowed with a same algebraic structure has as underlying set the inductive limit of the directed system of underlying sets, it is natural that the same result holds for sheafification functors. Let $+$ be the sheafification functor

$$+ : \mathbf{PSh}(T, \mathbf{Alg}) \ni \mathcal{F} \mapsto \mathcal{F}^+ \in \mathbf{Sh}(T, \mathbf{Alg}) : \text{For} .$$

Recall that $\mathcal{F}^+(U)$ ($U \in \mathbf{Open}(T)$) is defined as the subset of $(\Pi \mathcal{F})(U) := \prod_{x \in U} \mathcal{F}_x$, which is made of those elements $\sigma = (\sigma_x)_{x \in U} = ([s]_x)_{x \in U}$, for whom the section s is constant in a neighborhood of any point of U . The algebra operations on $\mathcal{F}^+(U)$ are naturally induced by

those of the stalks. Restrictions are the obvious algebra morphisms. The morphism $i : \mathcal{F} \rightarrow \mathcal{F}^+$ of presheaves of algebras is defined by

$$i_U : \mathcal{F}(U) \ni s \mapsto ([s]_x)_{x \in U} \in \mathcal{F}^+(U), \quad (36)$$

and i_U is injective, if \mathcal{F} is separated. If $\mathcal{G} \in \mathbf{Sh}(T, \mathbf{Alg})$, the morphism $i : \mathcal{G} \rightarrow \mathcal{G}^+$ is an isomorphism of sheaves of algebras. Further, if $j : \mathcal{F} \rightarrow \mathcal{G}$ is a morphism of presheaves of algebras, the unique morphism $\bar{j} : \mathcal{F}^+ \rightarrow \mathcal{G}$ of sheaves of algebras, such that $\bar{j} \circ i = j$, is given by $\bar{j}_U = \prod_{x \in U} j_x$, where $j_x : \mathcal{F}_x \rightarrow \mathcal{G}_x$ is the morphism of algebras induced by j :

$$\bar{j}_U : \mathcal{F}^+(U) \ni ([s]_x)_{x \in U} \mapsto (j_x[s]_x)_{x \in U} \in \mathcal{G}^+(U) \simeq \mathcal{G}(U). \quad (37)$$

Similarly, if $\mathcal{F}, \mathcal{F}' \in \mathbf{PSh}(T, \mathbf{Alg})$ and $\varphi : \mathcal{F} \rightarrow \mathcal{F}'$ is a presheaf morphism, the components of the sheaf morphism $\varphi^+ : \mathcal{F}^+ \rightarrow \mathcal{F}'^+$ are

$$\varphi_U^+ : \mathcal{F}^+(U) \ni ([s]_x)_{x \in U} \mapsto (\varphi_x[s]_x)_{x \in U} \in \mathcal{F}'^+(U).$$

The stalk \mathcal{F}_x ($x \in T$) is the inductive limit algebra $\mathcal{F}_x = \varinjlim_{U \ni x} \mathcal{F}(U)$ of the directed system of algebras $(\mathcal{F}(U), \rho_V^U)$, where U is an open neighborhood of x and ρ_V^U the restriction from U to $V \subset U$. This system is actually a directed system of LCTA-s, in the sense that the $\mathcal{F}(U)$ are LCTA-s and the ρ_V^U are continuous algebra morphisms. If we endow the inductive limit algebra \mathcal{F}_x with the final locally convex vector space topology with respect to the canonical algebra morphisms $\pi_U^x : \mathcal{F}(U) \rightarrow \mathcal{F}_x$ (i.e., with the finest locally convex vector space topology, for which the π_U^x are all continuous), the limit \mathcal{F}_x is a LCTVS, whose multiplication is (jointly) continuous [Mal86, Lemma 2.2.], i.e., the stalk \mathcal{F}_x is a LCTA.

In the following, the algebra $\mathcal{F}^+(U) \subset (\Pi\mathcal{F})(U)$ carries the induced topology of the product topology. Since any product of LCTVS-s and any subspace of a LCTVS are LCTVS-s, the algebra $\mathcal{F}^+(U)$ is a LCTVS. The multiplication on $\mathcal{F}^+(U)$ is continuous in view of Lemma 32, so that $\mathcal{F}^+(U)$ is a LCTA.

To show that $\mathcal{F}^+ \in \mathbf{Sh}(T, \mathbf{LCTAlg})$, it suffices to prove that any restriction

$$r_V^U : (\Pi\mathcal{F})(U) \ni (\sigma_x)_{x \in U} \mapsto (\sigma_x)_{x \in V} \in (\Pi\mathcal{F})(V)$$

is continuous, i.e., that, for all $y \in V$, the map

$$p_y^V \circ r_V^U = p_y^U : (\Pi\mathcal{F})(U) \ni (\sigma_x)_{x \in U} \mapsto \sigma_y \in \mathcal{F}_y$$

is continuous – which is a consequence of the definition of the product topology.

The next step consists in proving that the morphism $i : \mathcal{F} \rightarrow \mathcal{F}^+$ of presheaves of algebras defined by (36), is a morphism of presheaves of LCTA-s, i.e., in proving that, for any $y \in U$, the map

$$p_y^U \circ i_U = \pi_U^y : \mathcal{F}(U) \ni s \mapsto [s]_y \in \mathcal{F}_y$$

is continuous. This holds true by definition of the final topology on \mathcal{F}_y .

Let $\varphi : \mathcal{F} \rightarrow \mathcal{F}'$ be a morphism in $\mathbf{PSh}(T, \mathbf{LCTAlg})$. To show that $\varphi^+ : \mathcal{F}^+ \rightarrow \mathcal{F}'^+$ is a morphism in $\mathbf{Sh}(T, \mathbf{LCTAlg})$, it suffices to show that

$$\prod_{x \in U} \varphi_x : (\Pi \mathcal{F})(U) \ni ([s]_x)_{x \in U} \mapsto (\varphi_x [s]_x)_{x \in U} \in (\Pi \mathcal{F}')(U) \quad (38)$$

is continuous. This is the case if and only if $\varphi_x : \mathcal{F}_x \rightarrow \mathcal{F}'_x$, $x \in U$, is continuous. In view of Lemma 33, the algebra morphism φ_x is continuous if and only if

$$\varphi_x \circ \pi_V^x = \pi_V'^x \circ \varphi_V : \mathcal{F}(V) \ni s \mapsto \varphi_x [s]_x = [\varphi_V s]_x \in \mathcal{F}'_x,$$

$V \ni x$, is continuous. This condition is obviously fulfilled. \square

Corollary 36. *When equipped with the final locally convex vector space topology with respect to the canonical algebra morphisms $\pi_U^x : \mathcal{F}(U) \rightarrow \mathcal{F}_x$, a stalk \mathcal{F}_x , $x \in T$, of a presheaf $\mathcal{F} \in \mathbf{PSh}(T, \mathbf{TAAlg})$ is the inductive limit in \mathbf{TAAlg} of the directed system $(\mathcal{F}(U), \rho_V^U)$. The same statement holds in \mathbf{LCTAlg} .*

Proof. Clearly, the $\pi_U^x : \mathcal{F}(U) \rightarrow \mathcal{F}_x$ are morphisms in $(\mathbf{LC})\mathbf{TAAlg}$. Let (F, p_U) be made of $F \in (\mathbf{LC})\mathbf{TAAlg}$ and $(\mathbf{LC})\mathbf{TAAlg}$ -morphisms $p_U : \mathcal{F}(U) \rightarrow F$, such that $p_V \circ \rho_V^U = p_U$. Due to Lemma 33, the unique \mathbf{Alg} -morphism $u : \mathcal{F}_x \rightarrow F$, such that $u \circ \pi_U^x = p_U$, is continuous, since the $u \circ \pi_U^x = p_U$ are all continuous. Hence, the claim. \square

Proposition 37. *Let $\alpha : A \rightarrow C$ and $\beta : B \rightarrow D$ be two continuous algebra morphisms between nuclear Fréchet algebras. Then $\alpha \widehat{\otimes} \beta : A \widehat{\otimes} B \rightarrow C \widehat{\otimes} D$ is a continuous algebra morphism between nuclear Fréchet algebras.*

Proof. Since α and β are continuous linear maps between locally convex spaces, the map $\alpha \otimes \beta : A \otimes B \rightarrow C \otimes D$ is continuous linear, and its continuous extension $\alpha \widehat{\otimes} \beta : A \widehat{\otimes} B \rightarrow C \widehat{\otimes} D$ is continuous linear as well.

The source and the target of $\alpha \widehat{\otimes} \beta$ are nuclear Fréchet algebras [Eme17, Lemma 1.2.13]. It remains to show that $\alpha \widehat{\otimes} \beta$ respects their (continuous) multiplications. Let $\mathbf{a} = \sum_{i=0}^{\infty} a_i \otimes b_i$ and $\mathbf{b} = \sum_{j=0}^{\infty} a'_j \otimes b'_j$ be two elements of $A \widehat{\otimes} B$. Using continuity, we get

$$(\alpha \widehat{\otimes} \beta)(\mathbf{a} \cdot \mathbf{b}) = \lim_n \lim_m \sum_{i=0}^n \sum_{j=0}^m (\alpha(a_i) \alpha(a'_j)) \otimes (\beta(b_i) \beta(b'_j)).$$

It is straightforwardly seen that $(\alpha \widehat{\otimes} \beta)\mathbf{a} \cdot (\alpha \widehat{\otimes} \beta)\mathbf{b}$ is given by the same limit of sums of tensor products. \square

Proposition 38. *The multiplication $m : A \times A \rightarrow A$ of a nuclear Fréchet algebra A extends to a continuous algebra morphism $\widehat{m} : A \widehat{\otimes} A \rightarrow A$ between nuclear Fréchet algebras.*

Proof. We equip $A \otimes A$ with the projective tensor topology. The continuous bilinear maps from $A \times A$ to A correspond exactly to the continuous linear maps from $A \otimes A$ to A . Hence, the multiplication m can be viewed as a continuous linear map $m : A \otimes A \rightarrow A$. The latter extends to a continuous linear map $\widehat{m} : A \widehat{\otimes} A \rightarrow A$. This extension respects the (continuous) multiplications. The proof is similar to the one of Proposition 37. \square

5.5 Direct image and sheafification

Proposition 39. *Let $\mathcal{F} \in \text{PSh}(T, \text{LCTAlg})$ and let $f \in C^0(T, T')$. There is a morphism*

$$\iota : (f_*\mathcal{F})^+ \rightarrow f_*\mathcal{F}^+$$

in $\text{Sh}(T', \text{LCTAlg})$.

Proof. The assignment

$$f_*\mathcal{F} : \text{Open}(T') \ni V \mapsto \mathcal{F}(f^{-1}(V)) \in \text{LCTAlg}$$

together with the restrictions

$$\rho_{f^{-1}(V')}^{f^{-1}(V)} : (f_*\mathcal{F})(V) \rightarrow (f_*\mathcal{F})(V'),$$

where ρ are the restrictions of \mathcal{F} in LCTAlg , is a presheaf $f_*\mathcal{F} \in \text{PSh}(T', \text{LCTAlg})$. In view of Proposition 35, we get $(f_*\mathcal{F})^+ \in \text{Sh}(T', \text{LCTAlg})$. Similarly, we have $f_*\mathcal{F}^+ \in \text{Sh}(T', \text{LCTAlg})$.

Further, for $x \in T$ and $U \supset U' \ni x$, the LCTAlg -morphisms $\pi_U^x : \mathcal{F}(U) \rightarrow \mathcal{F}_x$ and $\pi_{U'}^x : \mathcal{F}(U') \rightarrow \mathcal{F}_x$ satisfy $\pi_{U'}^x \circ \rho_{U'}^U = \pi_U^x$. As, due to Corollary 36, the stalk $(f_*\mathcal{F})_{f(x)} \in \text{LCTAlg}$ is the inductive limit in LCTAlg of the directed system

$$(\mathcal{F}(f^{-1}(V)), \rho_{f^{-1}(V')}^{f^{-1}(V)}) \quad (V \supset V' \ni f(x)),$$

there exists a unique LCTAlg -morphism

$$u_x : (f_*\mathcal{F})_{f(x)} \rightarrow \mathcal{F}_x, \quad (39)$$

such that $u_x \circ \pi_V^{f(x)} = \pi_{f^{-1}(V)}^x$, i.e., such that $u_x [s]_{f(x)} = [s]_x$, for all $s \in \mathcal{F}(f^{-1}(V))$ and all $V \ni f(x)$.

For any $V \in \text{Open}(T')$, the map

$$\iota_V : (\Pi(f_*\mathcal{F}))(V) \ni ([s]_y)_{y \in V} \mapsto (u_x [s]_{f(x)})_{x \in f^{-1}(V)} = ([s]_x)_{x \in f^{-1}(V)} \in (\Pi\mathcal{F})(f^{-1}(V)) \quad (40)$$

is a continuous algebra morphism. Indeed, the algebraic operations are defined component-wise. For instance, the image of a product $([s]_y \cdot [s']_y)_{y \in V}$ is the product $(u_x [s]_{f(x)} \cdot u_x [s']_{f(x)})_{x \in f^{-1}(V)}$ of the images. Moreover, the map ι_V is continuous if and only if the maps

$$p_x^{f^{-1}(V)} \circ \iota_V : (\Pi(f_*\mathcal{F}))(V) \ni ([s]_y)_{y \in V} \mapsto u_x [s]_{f(x)} \in \mathcal{F}_x \quad (x \in f^{-1}(V))$$

are. The preimage by $p_x^{f^{-1}(V)} \circ \iota_V$ of any open $\omega \subset \mathcal{F}_x$ is the product over $y \in V$, whose factors indexed by $y \neq f(x)$ are $(f_*\mathcal{F})_y$ and whose factor indexed by $f(x)$ is the open subset $u_x^{-1}(\omega)$ of $(f_*\mathcal{F})_{f(x)}$. The preimage by $p_x^{f^{-1}(V)} \circ \iota_V$ is thus open in $(\Pi(f_*\mathcal{F}))(V)$.

The restriction of ι_V to $(f_*\mathcal{F})^+(V)$, still denoted by ι_V , arrives in $(f_*\mathcal{F}^+)(V)$. Assume that $([s]_y)_{y \in V}$ is implemented in a neighborhood V_{y_0} of an arbitrary point $y_0 \in V$ by a same section $t \in (f_*\mathcal{F})(V_{y_0})$, let $x_0 \in f^{-1}(V)$, and set $U_{x_0} = f^{-1}(V_{f(x_0)})$. For any $x \in U_{x_0}$, we have

$$u_x [s]_{f(x)} = u_x [t]_{f(x)} = [t]_x,$$

with $t \in \mathcal{F}(U_{x_0})$.

The restriction $\iota_V : (f_*\mathcal{F})^+(V) \rightarrow (f_*\mathcal{F}^+)(V)$ is obviously a morphism in \mathbf{LCTAlg} . Since these ι_V commute with restrictions, they define the sheaf morphism ι announced in Proposition 39. \square

References

- [BCC11] L. Balduzzi, C. Carmeli, and G. Cassinelli, *Super Vector Bundles*, J. Phys.: Conf. Ser. 284 (2011), 1–10.
- [BBH91] C. Bartocci, U. Bruzzo, and D. Hernández-Ruipérez, *The Geometry of Supermanifolds*, Math. and Appl., 71 (1991).
- [BS17] V. I. Bogachev and O. G. Smolyanov, *Methods of constructing topological vector spaces*, Top. Vect. Spaces & Appl. (2017), 101–152.
- [BPP18] G. di Brino, D. Pištalo, and N. Poncin, *Koszul-Tate resolutions as cofibrant replacements of algebras over differential operators*, JHRS, <http://rdcu.be/JQDP> (2018).
- [BPP15] G. di Brino, D. Pištalo, and N. Poncin, *Model structure on differential graded commutative algebras over the ring of differential operators*, [arXiv:1505.07720](https://arxiv.org/abs/1505.07720) (2015).
- [BP18] A. Bruce and N. Poncin, *Functional analytic issues in \mathbb{Z}_2^n -Geometry*, to appear (2018).
- [CCF11] C. Carmeli, L. Caston, and R. Fiorese, *Mathematical foundations of Supersymmetry*, EMS series of lectures in math., 15 (2011).
- [Cos11] K. Costello, *Renormalization and Effective Field Theory*, Mathematical Surveys and Monographs, 170, AMS (2011).
- [CGP14] T. Covolo, J. Grabowski, and N. Poncin, *\mathbb{Z}_2^n -Supergeometry I: Manifolds and Morphisms*, [arXiv:1408.2939](https://arxiv.org/abs/1408.2939) (2014).
- [CGP16a] T. Covolo, J. Grabowski, and N. Poncin, *The category of \mathbb{Z}_2^n -supermanifolds*, J. Math. Phys., 57(7) (2016).
- [CGP16b] T. Covolo, J. Grabowski, and N. Poncin, *Splitting theorem for \mathbb{Z}_2^n -supermanifolds*, J. Geo. Phys., 110 (2016), 393–401.
- [CKP16] T. Covolo, S. Kwok, and N. Poncin, *Differential Calculus on \mathbb{Z}_2^n -supermanifolds*, [arXiv:1608.00949](https://arxiv.org/abs/1608.00949) (2016).
- [COP12] T. Covolo, V. Ovsienko, and N. Poncin, *Higher trace and Berezinian of matrices over a Clifford algebra*, J. Geo. Phys., 62 (2012), 2294–2319.

- [Eme17] M. Emerton, *Locally analytic vectors in representations of locally p -adic analytic groups*, Memoirs AMS, 248(1175) (2017).
- [GK39] I. Gel'fand and A. Kolmogoroff, *On rings of continuous functions on topological spaces*, C. R. (Dokl.) Acad. Sci. URSS, 22 (1939), 11–15.
- [Goo72] K. R. Goodearl, *Distributing tensor product over direct product*, Pacific J. Math., 43(1) (1972), 107–110.
- [GP04] J. Grabowski and N. Poncin, *Automorphisms of quantum and classical Poisson algebras*, Compositio Math., 140(2) (2004), 511–527.
- [GKP10] J. Grabowski, A. Kotov, and N. Poncin, *The Lie superalgebra of a supermanifold*, J. Lie Theo., 20(4) (2010), 739–749.
- [GKP11] J. Grabowski, A. Kotov, and N. Poncin, *Geometric structures encoded in the Lie structure of an Atiyah algebroid*, Trans. Groups, 16(1) (2011), 137–160.
- [GKP13] J. Grabowski, A. Kotov, and N. Poncin, *Lie superalgebras of differential operators*, J. Lie Theo., 23(1) (2013), 35–54.
- [Gro52] A. Grothendieck, *Résumé des résultats essentiels dans la théorie des produits tensoriels topologiques et des espaces nucléaires*, Ann. Inst. Fourier, 4, 73–112.
- [Ham82] R. S. Hamilton, *The inverse function theorem of Nash and Moser*, Bull. AMS, 7(1) (1982).
- [Lei80] D. Leites, *Introduction to the theory of supermanifolds*, Russ. Math. Surv., 34(1) (1980), 1–64.
- [Mal70] A. Mallios, *On topological algebra sheaves of nuclear type*, Studia Math., 38 (1970), 215–220.
- [Mal86] A. Mallios, *Topological Algebras, Selected Topics*, Math. Studies, 124, Notas de Matemática, 109, Elsevier (1986).
- [MH05] A. Mallios and M. Haralampidou (Editors), *Topological Algebras and Applications*, Fifth International Conference on Topological Algebras and Applications, June 27 - July 1, 2005, Athens, Greece, in: Contemporary Mathematics, 427, AMS, Providence, Rhode Island.
- [Nag07] G. Nagy, *Topological Vector Spaces IV: Completeness and Metrizability*, www.math.ksu.edu/~nagy/func-an-2007-2008/top-vs-4.pdf.
- [NS03] J. Navarro González and J. Sancho de Salas, *C^∞ -Differentiable Spaces*, Lecture notes in Mathematics, Springer, Berlin, Heidelberg, New York, 1824 (2003).
- [Pon16] N. Poncin, *Towards Integration on Colored Supermanifolds*, Banach Center Publications, 110 (2016), 201–217.

- [Sar08] G. Sardanashvily, *Supermetrics on Supermanifolds*, Int. J. Geo. Meth. Mod. Phys., 5(2) (2008), 271-286.
- [Tre67] F. Trèves, *Topological vector spaces, distributions and kernels*, Academic Press, New York, 75, 273 (1967).
- [Var04] V.S. Varadarajan, *Supersymmetry for Mathematicians: An Introduction*, AMS (2004).