

The functional Breuer-Major theorem

Ivan Nourdin¹ and David Nualart^{2,3}

Université du Luxembourg and University of Kansas

Abstract: Let $X = \{X_n\}_{n \in \mathbb{Z}}$ be zero-mean stationary Gaussian sequence of random variables with covariance function ρ satisfying $\rho(0) = 1$. Let $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ be a function such that $\mathbb{E}[\varphi(X_0)^2] < \infty$ and assume that φ has Hermite rank $d \geq 1$. The celebrated Breuer-Major theorem asserts that, if $\sum_{r \in \mathbb{Z}} |\rho(r)|^d < \infty$ then the finite dimensional distributions of $\frac{1}{\sqrt{n}} \sum_{i=0}^{\lfloor nt \rfloor - 1} \varphi(X_i)$ converge to those of σW , where W is a standard Brownian motion and σ is some (explicit) constant. Surprisingly, and despite the fact this theorem has become over the years a prominent tool in a bunch of different areas, a necessary and sufficient condition implying the weak convergence in the space $\mathbf{D}([0, 1])$ of càdlàg functions endowed with the Skorohod topology is still missing. Our main goal in this paper is to fill this gap. More precisely, by using suitable boundedness properties satisfied by the generator of the Ornstein-Uhlenbeck semigroup, we show that tightness holds under the sufficient (and almost necessary) natural condition that $\mathbb{E}[|\varphi(X_0)|^p] < \infty$ for some $p > 2$.

1 Introduction

Consider a zero-mean stationary Gaussian sequence of random variables $X = \{X_n\}_{n \in \mathbb{Z}}$ with covariance function $\mathbb{E}[X_n X_m] = \rho(|n - m|)$ such that $\rho(0) = 1$. Let $\gamma = N(0, 1)$ be the standard Gaussian measure on \mathbb{R} . Consider a function $\varphi \in L^2(\mathbb{R}, \gamma)$ of Hermite rank $d \geq 1$, that is, φ has a series expansion given by

$$\varphi(x) = \sum_{q=d}^{\infty} c_q H_q(x), \quad c_d \neq 0, \quad (1.1)$$

where $H_q(x)$ is the q th Hermite polynomial with leading coefficient 1.

A classical central limit theorem, proved by Breuer and Major in [3], asserts that under the condition

$$\sum_{k \in \mathbb{Z}} |\rho(k)|^d < \infty, \quad (1.2)$$

the *finite-dimensional distributions* of the process

$$Y_n(t) := \frac{1}{\sqrt{n}} \sum_{i=0}^{\lfloor nt \rfloor - 1} \varphi(X_i), \quad t \in [0, 1] \quad (1.3)$$

converge to those of σW as n tends to infinity, where $W = \{W_t\}_{t \in [0, 1]}$ is a standard Brownian motion and

$$\sigma^2 = \sum_{q=d}^{\infty} q! c_q^2 \sum_{k \in \mathbb{Z}} \rho(k)^q. \quad (1.4)$$

¹Université du Luxembourg, Maison du Nombre, 6 avenue de la Fonte, L-4364 Esch-sur-Alzette, Grand Duchy of Luxembourg. ivan.nourdin@uni.lu

²Department of Mathematics, University of Kansas, Lawrence, KS 66045, USA. nualart@ku.edu

³David Nualart was supported by the NSF grant DMS 1811181

Observe that $|\rho(k)| = |\mathbb{E}[X_k X_0]| \leq \rho(0) = 1$ by Cauchy-Schwarz, and thus σ^2 is well defined under the integrability assumption (1.2) imposed on ρ . We also refer the reader to [7, Chapter 7], where a modern proof of the Breuer-Major theorem is given, by means of the recent Malliavin-Stein approach.

What about the *functional convergence*, that is, convergence in law of Y_n to σW in the space $\mathbf{D}([0, 1])$ endowed with the Skorohod topology? The best-to-date available criterion ensuring tightness for Y_n is due to Ben Hariz [1] and Chambers and Slud [4] (the former being only a slight improvement with respect to the latter⁴), in the simpler situation where sums are replaced by integrals and convergences are understood in the space $\mathbf{C}([0, 1])$ of continuous functions endowed with the uniform topology. Transformed into our setting, the criterion in [1, 4] reads as follows⁵: tightness holds provided there exists $R > 1$ such that

$$\sum_{q=d}^{\infty} \sqrt{q!} |c_q| \left(\sum_{k \in \mathbb{Z}} |\rho(k)|^q \right)^{\frac{1}{2}} R^q < \infty. \quad (1.5)$$

But, in our opinion, condition (1.5) is not meaningful, for at least three reasons: (i) it is not very natural, (ii) it is far from being optimal, and (iii) it may be difficult to check it in practice, especially when the computation of the Hermite coefficients c_q appears to be tricky or even impossible. Moreover, the proof given in [1, 4] of the fact that (1.5) implies tightness can be simplified a lot, by proceeding as follows. Let us first recall that tightness in $\mathbf{D}([0, 1])$ holds if there exist $p > 2$ and $c > 0$ such that, for all n ,

$$\|Y_n(t) - Y_n(s)\|_{L^p(\Omega)} \leq c \left(\frac{\lfloor nt \rfloor - \lfloor ns \rfloor}{n} \right)^{1/2}, \quad 0 \leq s \leq t \leq 1 \quad (1.6)$$

(see Lemma 3.1 below). Here, we have

$$\begin{aligned} \|Y_n(t) - Y_n(s)\|_{L^p(\Omega)} &= \left\| \sum_{q=d}^{\infty} c_q \frac{1}{\sqrt{n}} \sum_{i=\lfloor ns \rfloor}^{\lfloor nt \rfloor - 1} H_q(X_i) \right\|_{L^p(\Omega)} \\ &\leq \sum_{q=d}^{\infty} |c_q| \left\| \frac{1}{\sqrt{n}} \sum_{i=\lfloor ns \rfloor}^{\lfloor nt \rfloor - 1} H_q(X_i) \right\|_{L^p(\Omega)}. \end{aligned} \quad (1.7)$$

At this stage, a crucial observation is that $\sum_{i=\lfloor ns \rfloor}^{\lfloor nt \rfloor - 1} H_q(X_i)$ belongs to the q th Wiener chaos, where all $L^p(\Omega)$ -norms are equivalent by hypercontractivity. More precisely,

$$\left\| \frac{1}{\sqrt{n}} \sum_{i=\lfloor ns \rfloor}^{\lfloor nt \rfloor - 1} H_q(X_i) \right\|_{L^p(\Omega)} \leq (p-1)^{\frac{q}{2}} \left\| \frac{1}{\sqrt{n}} \sum_{i=\lfloor ns \rfloor}^{\lfloor nt \rfloor - 1} H_q(X_i) \right\|_{L^2(\Omega)}, \quad (1.8)$$

⁴Chambers and Slud criterion corresponds to Ben Hariz criterion (1.5) with $R = \frac{3}{2}$ and without the terms $\sum_{k \in \mathbb{Z}} |\rho(k)|^q$ all bounded by (1.2).

⁵Compared to [1], condition (1.5) is stated here with $\sqrt{q!}$ instead of $(\sqrt{q!})^{-1}$ (since we work here with Hermite polynomials with leading coefficient 1) and with sums replacing integrals (since we work here in a discrete framework).

see, e.g., [7, Corollary 2.8.14]. The interest of the right-hand side of (1.8) with respect to the left-hand side is that the former is straightforward to calculate and to estimate, as follows:

$$\left\| \frac{1}{\sqrt{n}} \sum_{i=\lfloor ns \rfloor}^{\lfloor nt \rfloor - 1} H_q(X_i) \right\|_{L^2(\Omega)}^2 \leq \frac{\lfloor nt \rfloor - \lfloor ns \rfloor}{n} q! \sum_{k \in \mathbb{Z}} |\rho(k)|^q.$$

By plugging this into (1.8) and then into (1.7), we obtain

$$\|Y_n(t) - Y_n(s)\|_{L^p(\Omega)} \leq \left(\frac{\lfloor nt \rfloor - \lfloor ns \rfloor}{n} \right)^{1/2} \sum_{q=d}^{\infty} |c_q| (p-1)^{\frac{q}{2}} \sqrt{q!} \left(\sum_{k \in \mathbb{Z}} |\rho(k)|^q \right)^{\frac{1}{2}},$$

implying in turn that (1.6) is satisfied (and then tightness) under (1.5) with $R = \sqrt{p-1} > 1$.

As we have just seen, the criterion (1.5) of [1, 4] for tightness is actually not so difficult to prove. But on the other hand it is neither natural, nor easy to check in practice. The main objective of this note is thus to provide a simpler sufficient condition for the convergence $Y_n \Rightarrow \sigma W$ to hold in law in $\mathbf{D}([0, 1])$ endowed with the Skorohod topology. Actually, our finding is that only a little more integrability of the function φ is needed.

Theorem 1.1. *Let $X = \{X_n\}_{n \in \mathbb{Z}}$ be a zero-mean Gaussian stationary sequence with covariance function $\mathbb{E}[X_n X_m] = \rho(|n - m|)$ such that $\rho(0) = 1$. Consider a function $\varphi \in L^2(\mathbb{R}, \gamma)$ with expansion (1.1) and Hermite rank $d \geq 1$, and suppose that $\sum_{k \in \mathbb{Z}} |\rho(k)|^d < \infty$. Finally, recall Y_n from (1.3), let $W = \{W_t\}_{t \in [0, 1]}$ be a Brownian motion and let σ^2 be defined in (1.4). Then, as $n \rightarrow \infty$,*

1. *The finite-dimensional distributions of Y_n converge to those of σW ;*
2. *If $\varphi \in L^p(\mathbb{R}, \gamma)$ for some $p > 2$, then Y_n converges in law to σW in $\mathbf{D}([0, 1])$ endowed with the Skorohod topology.*

We can prove a similar result in the space $\mathbf{C}([0, 1])$ of continuous functions endowed with the uniform topology. Of course, in this case we have to consider the linear interpolation Z_n instead of Y_n , defined as follows:

$$Z_n(t) = \frac{nt - \lfloor nt \rfloor}{\sqrt{n}} \varphi(X_{\lfloor nt \rfloor}) + \frac{1}{\sqrt{n}} \sum_{i=0}^{\lfloor nt \rfloor - 1} \varphi(X_i), \quad t \in [0, 1]. \quad (1.9)$$

Theorem 1.2. *Let $X = \{X_n\}_{n \in \mathbb{Z}}$ be a zero-mean Gaussian stationary sequence with covariance function $\mathbb{E}[X_n X_m] = \rho(|n - m|)$ such that $\rho(0) = 1$. Consider a function $\varphi \in L^2(\mathbb{R}, \gamma)$ with expansion (1.1) and Hermite rank $d \geq 1$, and suppose that $\sum_{k \in \mathbb{Z}} |\rho(k)|^d < \infty$. Finally, recall Z_n from (1.9), let $W = \{W_t\}_{t \in [0, 1]}$ be a Brownian motion and let σ^2 be defined in (1.4). Then, as $n \rightarrow \infty$,*

1. *The finite-dimensional distributions of Z_n converge to those of σW ;*
2. *If $\varphi \in L^p(\mathbb{R}, \gamma)$ for some $p > 2$, then Z_n converges in law to σW in $\mathbf{C}([0, 1])$ endowed with the uniform topology.*

The proof of Theorems 1.1 and 1.2 are based on the application of the techniques of Malliavin calculus, especially Meyer inequalities from [6] (in the modern form taken from [8]). The method we employ is based on the representation $\varphi(X_i) = \delta^d((D(-L)^{-1})^d(\varphi(X_i)))$ where δ , D and L are the usual Malliavin operators (see Section 2). It is robust enough to be used for other families of interest than Y_n and Z_n , see indeed [5] for an application to the self-intersection local time of the fractional Brownian motion, or Section 4 in the present paper for an extension of Theorem 1.1 in a critical situation where $\sum_{|k| \leq n} |\rho(k)|^d$ diverges slowly enough when $n \rightarrow \infty$.

The rest of the paper is organized as follows. Section 2 contains some useful preliminaries on Malliavin calculus, as well as some boundedness properties of the so-called shift operator, which is our main tool in this paper. The proof of Theorem 1.1 (resp. 1.2) is given in Section 3 (resp. 4). Finally, in Section 4 we provide an extension of Theorem 1.1 in the case where $\sum_{k \in \mathbb{Z}} |\rho(k)|^d$ explodes slowly.

2 Preliminaries

In this section, we gather several preliminary results that are needed for the proofs of the main results of this paper.

2.1 Elements of Malliavin calculus with respect to the Wiener process

We refer the reader to the references [7, 8, 9] for a detailed account on the Malliavin calculus. In this paper we will make use of the following notation and results.

First, let us introduce a specific realization of the sequence $\{X_k\}_{k \in \mathbb{Z}}$. The space

$$\mathcal{H} := \overline{\text{span}\{X_k, k \in \mathbb{Z}\}}^{L^2(\Omega)}$$

being a real separable Hilbert space, it is isometrically isomorphic to either \mathbb{R}^N (for some $N \geq 1$) or $L^2(\mathbb{R}_+)$. In both cases, there exists an isometry $\Phi : \mathcal{H} \rightarrow L^2(\mathbb{R}_+)$. Set $e_k = \Phi(X_k)$ for each $k \in \mathbb{Z}$. We have

$$\rho(k-l) = \mathbb{E}[X_k X_l] = \int_0^\infty e_k(x) e_l(x) dx, \quad k, l \geq 1. \quad (2.1)$$

Let $W = \{W(h), h \in L^2(\mathbb{R}_+)\}$ be the standard Wiener process, that is, a centered Gaussian family satisfying $\mathbb{E}[W(h)W(g)] = \langle h, g \rangle_{L^2(\mathbb{R}_+)}$ for all $h, g \in L^2(\mathbb{R}_+)$. We deduce immediately from (2.1) that

$$\{X_k\}_{k \in \mathbb{Z}} \stackrel{\text{law}}{=} \{W(e_k)\}_{k \in \mathbb{Z}}.$$

Since, in this paper, the quantities we are interested in only depend on the law, starting from now and without loss of generality, we set

$$X_k := W(e_k), \quad k \in \mathbb{Z}. \quad (2.2)$$

For integers $q \geq 1$, the q th Wiener chaos is the closed linear subspace of $L^2(\Omega)$ that is generated by the random variables $\{H_q(W(h)), h \in L^2(\mathbb{R}_+), \|h\|_{L^2(\mathbb{R}_+)} = 1\}$, where H_q stands for the q th Hermite polynomial defined by

$$H_q(x) = (-1)^q e^{\frac{x^2}{2}} \frac{d^q}{dx^q} e^{-\frac{x^2}{2}}, \quad q \geq 1,$$

and $H_0(x) = 1$. For $q \geq 1$, it is known that the map

$$I_q(h^{\otimes q}) = H_q(W(h)), \quad h \in L^2(\mathbb{R}_+), \quad \|h\|_{L^2(\mathbb{R}_+)} = 1, \quad (2.3)$$

provides a linear isometry between the set of symmetric square integrable functions $L_s^2(\mathbb{R}_+^q)$ (equipped with the modified norm $\sqrt{q!} \|\cdot\|_{L^2(\mathbb{R}_+^q)}$) and the q th Wiener chaos. By convention, $I_0(x) = x$ for all $x \in \mathbb{R}$.

It is well-known that any $F \in L^2(\Omega)$ measurable with respect to W can be decomposed into Wiener chaos as follows:

$$F = \mathbb{E}[F] + \sum_{q=1}^{\infty} I_q(f_q), \quad (2.4)$$

where the kernels $f_q \in L_s^2(\mathbb{R}_+^q)$ are uniquely determined by F .

For a smooth and cylindrical random variable $F = f(W(h_1), \dots, W(h_n))$, with $h_i \in L^2(\mathbb{R}_+)$ and $f \in C_b^\infty(\mathbb{R}^n)$ (f and its partial derivatives are bounded), we define its Malliavin derivative D as the $L^2(\mathbb{R}_+)$ -valued random variable given by

$$DF = \sum_{i=1}^n \frac{\partial f}{\partial x_i}(W(h_1), \dots, W(h_n)) h_i.$$

By iteration, one can define the k -th derivative $D^k F$ as an element of $L^2(\Omega; L^2(\mathbb{R}_+^k))$. For any natural number k and any real number $p \geq 1$, we define the Sobolev space $\mathbb{D}^{k,p}$ as the closure of the space of smooth and cylindrical random variables with respect to the norm $\|\cdot\|_{k,p}$ defined by

$$\|F\|_{k,p}^p = \mathbb{E}(|F|^p) + \sum_{l=1}^k \mathbb{E}(\|D^l F\|_{L^2(\mathbb{R}_+^l)}^p).$$

For any Hilbert space V we denote by $\mathbb{D}^{k,p}(V)$ the corresponding space of V -valued random variables.

The divergence operator δ is defined as the adjoint of the derivative operator D . An element $u \in L^2(\Omega; L^2(\mathbb{R}_+))$ belongs to the domain of δ , denoted by $\text{Dom } \delta$, if there is a constant c_u depending on u such that

$$|\mathbb{E}(\langle DF, u \rangle_{L^2(\mathbb{R}_+)})| \leq c_u \|F\|_{L^2(\Omega)}$$

for any $F \in \mathbb{D}^{1,2}$. If $u \in \text{Dom } \delta$, then the random variable $\delta(u)$ is defined by the duality relationship

$$\mathbb{E}[F \delta(u)] = \mathbb{E}[\langle DF, u \rangle_{L^2(\mathbb{R}_+)}], \quad (2.5)$$

which holds for any $F \in \mathbb{D}^{1,2}$. In a similar way we can introduce the iterated divergence operator δ^k for each integer $k \geq 2$, defined by the duality relationship

$$\mathbb{E}[F \delta^k(u)] = \mathbb{E}[\langle D^k F, u \rangle_{L^2(\mathbb{R}_+^k)}], \quad (2.6)$$

for any $F \in \mathbb{D}^{k,2}$, where $u \in \text{Dom } \delta^k \subset L^2(\Omega; L^2(\mathbb{R}_+^k))$. If $u \in L_s^2(\mathbb{R}_+^k)$ is deterministic, then

$$\delta^k(u) = I_k(u). \quad (2.7)$$

For any $p > 1$ and any integer $k \geq 1$, the operator δ^k is continuous from $\mathbb{D}^{k,p}(L^2(\mathbb{R}_+^k))$ into $L^p(\Omega)$, and we have the inequality (see, for instance, [8, Proposition 1.5.4])

$$\|\delta^k(v)\|_{L^p(\Omega)} \leq c_p \sum_{j=0}^k \|D^j v\|_{L^p(\Omega; L^2(\mathbb{R}_+^j))}, \quad (2.8)$$

for any $v \in \mathbb{D}^{k,p}(L^2(\mathbb{R}_+^k))$. This inequality is a consequence of Meyer inequalities (from [6]), which states the equivalence in $L^p(\Omega)$, for any $p > 1$, of the operators D and $(-L)^{1/2}$, where L is the infinitesimal generator of the Ornstein-Uhlenbeck semigroup $(P_t)_{t \geq 0}$ in $L^2(\Omega)$ defined as

$$P_t F = \sum_{q=0}^{\infty} e^{-qt} I_q(f_q), \quad t \geq 0, \quad \text{and} \quad (-L)^r F = \sum_{q=1}^{\infty} q^r I_q(f_q), \quad r \in \mathbb{R},$$

if F is given by (2.4). More precisely, there exist two constants $c_{i,p}$, $i = 1, 2$, such that, for any $F \in \mathbb{D}^{1,p}$,

$$c_{1,p} \|DF\|_{L^p(\Omega, L^2(\mathbb{R}_+))} \leq \|(-L)^{1/2} F\|_{L^p(\Omega)} \leq c_{2,p} \|DF\|_{L^p(\Omega, L^2(\mathbb{R}_+))}. \quad (2.9)$$

More generally, we can state Meyer's inequalities in the general case (see [8, Theorem 1.5.1]): for any $p > 1$ and any integer $k \geq 1$, there exist two constants $c_{i,p,k}$, $i = 1, 2$, such that, for any $F \in \mathbb{D}^{1,p}$,

$$c_{1,k,p} \|D^k F\|_{L^p(\Omega, L^2(\mathbb{R}_+^k))} \leq \|(-L)^{k/2} F\|_{L^p(\Omega)} \leq c_{2,k,p} (\|D^k F\|_{L^p(\Omega, L^2(\mathbb{R}_+^k))} + \|F\|_{L^p(\Omega)}). \quad (2.10)$$

2.2 The shift operator

Let $\varphi \in L^2(\mathbb{R}, \gamma)$ be a function of Hermite rank $d \geq 1$ and expansion (1.1). Consider the function φ_d defined by a shift of d units in the coefficients, that is,

$$\varphi_d = \sum_{q=d}^{\infty} c_q H_{q-d}. \quad (2.11)$$

It is immediately checked that $\varphi_d \in L^2(\mathbb{R}, \gamma)$.

Given (2.11) and the relation (2.3) between Hermite polynomials and multiple stochastic integrals, the random variable $\varphi_d(W(h))$ admits the following chaotic decomposition when $h \in L^2(\mathbb{R}_+)$ has norm 1 :

$$\varphi_d(W(h)) = \sum_{q=d}^{\infty} c_q I_{q-d}(h^{\otimes(q-d)}).$$

Moreover, we claim that $\varphi_d(W(h))$ belongs to $\mathbb{D}^{2,d}$. Indeed, for any $k = 1, \dots, d$ we have that

$$D^k(\varphi_d(W(h))) = \sum_{q=d}^{\infty} c_q (q-d)(q-d-1) \cdots (q-d-k+1) I_{q-d-k}(h^{\otimes(q-d-k)}) h^{\otimes k},$$

and this series converges in $L^2(\Omega, L^2(\mathbb{R}_+^k))$ since

$$\begin{aligned}\mathbb{E}\|D^k(\varphi_d(W(h)))\|_{L^2(\mathbb{R}_+^k)}^2 &= \sum_{q=d}^{\infty} c_q^2 (q-d)^2 (q-d-1)^2 \cdots (q-d-k+1)^2 (q-d-k)! \\ &\leq \sum_{q=d}^{\infty} c_q^2 q! < \infty.\end{aligned}$$

The following two lemmas will play a crucial role in the sequel.

Lemma 2.1. *Suppose that $\varphi \in L^2(\mathbb{R}, \gamma)$ given by (1.1) has Hermite rank $d \geq 1$. We have, for any $h \in L^2(\mathbb{R}_+)$ of norm 1,*

$$\varphi(W(h)) = \delta^d(\varphi_d(W(h))h^{\otimes d}) \quad (2.12)$$

$$\varphi_d(W(h))h^{\otimes d} = (D(-L)^{-1})^d(\varphi(W(h))) \quad (2.13)$$

$$\varphi_d(W(h)) = \langle (D(-L)^{-1})^d(\varphi(W(h))), h^{\otimes d} \rangle_{L^2(\mathbb{R}_+^d)}. \quad (2.14)$$

Proof. Using (2.7) and the relation (2.3) between Hermite polynomials and multiple stochastic integrals, we can write

$$\begin{aligned}\varphi(W(h)) &= \sum_{q=d}^{\infty} c_q H_q(W(h)) = \sum_{q=d}^{\infty} c_q I_q(h^{\otimes q}) = \sum_{q=d}^{\infty} c_q \delta^q(h^{\otimes q}) \\ &= \sum_{q=d}^{\infty} c_q \delta^d \left(\delta^{q-d} \left(h^{\otimes q-d} \right) h^{\otimes d} \right) = \delta^d \left(\sum_{q=d}^{\infty} c_q I_{q-d} \left(h^{\otimes q-d} \right) h^{\otimes d} \right) \\ &= \delta^d \left(\sum_{q=d}^{\infty} c_q H_{q-d}(W(h)) h^{\otimes d} \right) = \delta^d(\varphi_d(W(h))h^{\otimes d}),\end{aligned}$$

which is (2.12). On the other hand, we can compute that

$$(D(-L)^{-1})(\varphi(W(h))) = \sum_{q=d}^{\infty} c_q I_{q-1}(h^{\otimes q-1}) h.$$

By iteration, we get

$$(D(-L)^{-1})^d(\varphi(W(h))) = \sum_{q=d}^{\infty} c_q I_{q-d}(h^{\otimes q-d}) h^{\otimes d} = \sum_{q=d}^{\infty} c_q H_{q-d}(W(h)) h^{\otimes d},$$

and the desired conclusions (2.13) and then (2.14) follow. \square

Lemma 2.2. *Suppose that $\varphi \in L^2(\mathbb{R}, \gamma)$ given by (1.1) has Hermite rank $d \geq 1$ and is such that $\mathbb{E}[|\varphi(N)|^p] < \infty$ for some $p > 2$ and $N \sim N(0, 1)$. Then, for any $0 \leq k \leq r \leq d$,*

$$\sup_{\|h\|=1} \mathbb{E}[\|D^k(D(-L)^{-1})^r(\varphi(W(h)))\|_{L^2(\mathbb{R}_+^{r+k})}^p] < \infty,$$

where the supremum runs over the set of all square integrable functions $h \in L^2(\mathbb{R}_+)$ of norm 1.

Proof. The proof is by induction on r . When $r = 0$, one has $k = 0$ and $D^0(D(-L)^{-1})^0$ is the identity operator, so there is nothing to prove.

Suppose now that the conclusion of Lemma 2.2 holds true for some $r - 1 \in \{0, \dots, d - 1\}$, and let us prove that it holds true for $r + 1$ as well.

If $k = 0$, we have $D^0(D(-L)^{-1})^r = (D(-L)^{-1})^r$. But $D(-L)^{-1} = \int_0^\infty DP_t dt$ according to [7, Prop. 2.9.3]. Moreover, according⁶ to [9, Prop. 5.1.5], there exists $c_p > 0$ such that, for any $F \in L^p(\Omega)$,

$$\|DP_t F\|_{L^p(\Omega, L^2(\mathbb{R}_+))} \leq c_p \frac{e^{-t}}{\sqrt{1 - e^{-2t}}} \|F\|_{L^p(\Omega)}. \quad (2.15)$$

It follows from these two facts and the Minkowski inequality that the operator $D(-L)^{-1}$ is bounded from $L^p(\Omega)$ to $L^p(\Omega, L^2(\mathbb{R}_+))$. As a consequence, by iteration one has

$$\sup_{\|h\|=1} \mathbb{E}[\|(D(-L)^{-1})^r(\varphi(W(h)))\|_{L^2(\mathbb{R}_+^r)}^p] \leq c \mathbb{E}[|\varphi(N)|^p] < \infty$$

for any $0 \leq r \leq d$.

Let us finally consider the case $1 \leq k \leq r$. We can write, using among other the left-hand side of (2.9) and then its right-hand side,

$$\begin{aligned} & \sup_{\|h\|=1} \mathbb{E}[\|D^k(D(-L)^{-1})^r(\varphi(W(h)))\|_{L^2(\mathbb{R}_+^{k+r})}^p] \\ &= \sup_{\|h\|=1} \mathbb{E}[\|D^{k+1}(-L)^{-1}(D(-L)^{-1})^{r-1}(\varphi(W(h)))\|_{L^2(\mathbb{R}_+^{k+r})}^p] \\ &\leq \frac{1}{c_1} \sup_{\|h\|=1} \mathbb{E}[\|(-L)^{\frac{k-1}{2}}(D(-L)^{-1})^{r-1}(\varphi(W(h)))\|_{L^2(\mathbb{R}_+^{r-1})}^p] \\ &\leq \frac{c_2}{c_1} \sup_{\|h\|=1} \left(\mathbb{E}[\|D^{k-1}(D(-L)^{-1})^{r-1}\varphi(W(h))\|_{L^2(\mathbb{R}_+^{k+r-2})}^p] \right. \\ &\quad \left. + \mathbb{E}[\|(D(-L)^{-1})^{r-1}\varphi(W(h))\|_{L^2(\mathbb{R}_+^{r-1})}^p] \right), \end{aligned}$$

which is finite by the induction property. \square

3 Proof of Theorem 1.1

Since the point 1 (that is, convergence of the finite-dimensional distributions) follows from the classical Breuer-Major theorem of [3], let us only concentrate on the point 2.

We are thus left to show that the family $(Y_n)_{n \geq 1}$ is tight in the Skorohod space $\mathbf{D}([0, 1])$. Recall from [2, Theorem 15.6] that a sufficient condition for tightness in $\mathbf{D}([0, 1])$ is the existence of $\gamma > 0$ and $c > 0$ such that, for all n ,

$$\mathbb{E}[|Y_n(t) - Y_n(t_1)|^{1+\gamma} |Y_n(t_2) - Y_n(t)|^{1+\gamma}] \leq c(t_2 - t_1)^{1+\gamma}, \quad 0 \leq t_1 \leq t \leq t_2 \leq 1. \quad (3.1)$$

⁶The statement of [9, Prop. 5.1.5] is with $t^{-1/2}$ instead of $\frac{e^{-t}}{\sqrt{1 - e^{-2t}}}$, but the given proof actually provides the estimate stated in (2.15).

We are not going to check (3.1) directly. Instead, we shall use the following lemma, which is not stated in Billingsley book [2] but has nevertheless become part of the folklore. For the sake of completeness, we give its proof.

Lemma 3.1. *Fix $p > 2$ and $c > 0$. If*

$$\|Y_n(t) - Y_n(s)\|_{L^p(\Omega)} \leq C \left(\frac{\lfloor nt \rfloor - \lfloor ns \rfloor}{n} \right)^{1/2}, \quad s, t \in [0, 1] \quad (3.2)$$

for some $p > 2$ and $C > 0$ then (3.1) holds with $\gamma = \frac{p}{2} - 1 > 0$ and $c = 3^{\frac{p}{2}} C^p > 0$.

Proof. Suppose (3.2). Using Cauchy-Schwarz, one has

$$\begin{aligned} & \mathbb{E}[|Y_n(t) - Y_n(t_1)|^{\frac{p}{2}} |Y_n(t_2) - Y_n(t)|^{\frac{p}{2}}] \\ & \leq \|Y_n(t) - Y_n(t_1)\|_{L^p(\Omega)}^{\frac{p}{2}} \|Y_n(t_2) - Y_n(t)\|_{L^p(\Omega)}^{\frac{p}{2}} \\ & \leq c^p \left(\frac{\lfloor nt \rfloor - \lfloor nt_1 \rfloor}{n} \right)^{\frac{p}{4}} \left(\frac{\lfloor nt_2 \rfloor - \lfloor nt \rfloor}{n} \right)^{\frac{p}{4}}. \end{aligned} \quad (3.3)$$

If $\max(n(t - t_1), n(t_2 - t)) < \frac{1}{2}$, then the quantity in (3.3) is zero, and so (3.1) is verified. If $n(t - t_1) \geq \frac{1}{2}$, then

$$\frac{\lfloor nt \rfloor - \lfloor nt_1 \rfloor}{n} \leq \frac{nt - nt_1 + 1}{n} \leq \frac{nt - nt_1 + 2n(t - t_1)}{n} \leq 3(t_2 - t_1),$$

whereas

$$\frac{\lfloor nt_2 \rfloor - \lfloor nt \rfloor}{n} \leq \frac{nt_2 - nt + 1}{n} \leq \frac{nt_2 - nt + 2n(t - t_1)}{n} \leq 3(t_2 - t_1).$$

Similar estimates hold if $n(t_2 - t) \geq \frac{1}{2}$. So, if $\max(n(t - t_1), n(t_2 - t)) \geq \frac{1}{2}$, then the quantity in (3.3) is bounded by $3^{\frac{p}{2}} c^p (t_2 - t_1)^{\frac{p}{2}}$, and the proof of (3.1) is complete. \square

We are now ready to proceed with the proof of point 2 in Theorem 1.1. Combining the previous Lemma 3.1 with [2, Theorem 15.6], we are left to show that (3.2) is satisfied.

We can write

$$\begin{aligned} \|Y_n(t) - Y_n(s)\|_{L^p(\Omega)} &= \frac{1}{\sqrt{n}} \left\| \sum_{i=\lfloor ns \rfloor}^{\lfloor nt \rfloor - 1} \varphi(X_i) \right\|_{L^p(\Omega)} = \frac{1}{\sqrt{n}} \left\| \sum_{i=\lfloor ns \rfloor}^{\lfloor nt \rfloor - 1} \delta^d \left(\varphi_d(X_i) e_i^{\otimes d} \right) \right\|_{L^p(\Omega)} \quad \text{by (2.12)} \\ &\leq c_p \sum_{k=0}^d \frac{1}{\sqrt{n}} \left\| \sum_{i=\lfloor ns \rfloor}^{\lfloor nt \rfloor - 1} D^k \left(\varphi_d(X_i) e_i^{\otimes d} \right) \right\|_{L^p(\Omega; L^2(\mathbb{R}_+^{k+d}))} \quad \text{by (2.8)} \\ &= c_p \sum_{k=0}^d \left\| \frac{1}{n} \sum_{i,j=\lfloor ns \rfloor}^{\lfloor nt \rfloor - 1} D^k(\varphi_d(X_i)) D^k(\varphi_d(X_j)) \langle e_i, e_j \rangle_{L^2(\mathbb{R}_+)}^{d+k} \right\|_{L^{\frac{p}{2}}(\Omega; L^2(\mathbb{R}_+^k))}^{1/2} \\ &=: c_p \sum_{k=0}^d R_k. \end{aligned}$$

On the other hand,

$$\begin{aligned}
& \sup_{i \in \mathbb{Z}} \|D^k(\varphi_d(X_i))\|_{L^p(\Omega; L^2(\mathbb{R}_+^k))} = \sup_{i \in \mathbb{Z}} \mathbb{E} \left[\|D^k(\varphi_d(X_i))\|_{L^2(\mathbb{R}_+^k)}^p \right]^{\frac{1}{p}} \\
&= \sup_{i \in \mathbb{Z}} \mathbb{E} \left[\|D^k(\langle (D(-L)^{-1})^d(\varphi(X_i)), e_i^{\otimes d} \rangle_{L^2(\mathbb{R}_+^d)})\|_{L^2(\mathbb{R}_+^k)}^p \right]^{\frac{1}{p}} \quad \text{by (2.14)} \\
&\leq \sup_{i \in \mathbb{Z}} \mathbb{E} \left[\|D^k(D(-L)^{-1})^d(\varphi(X_i))\|_{L^2(\mathbb{R}_+^{k+d})}^p \right]^{\frac{1}{p}}, \tag{3.4}
\end{aligned}$$

and (3.4) is finite thanks to Lemma 2.2.

Recall from (2.1) that $\langle e_i, e_j \rangle_{L^2(\mathbb{R}_+)} = \rho(i - j)$. Using Minkowski and Hölder inequalities, we can write, for any $0 \leq k \leq d$,

$$\begin{aligned}
R_k &\leq \sup_{i \in \mathbb{Z}} \|D^k(\varphi_d(X_i))\|_{L^p(\Omega; L^2(\mathbb{R}_+^k))} \left(\frac{1}{n} \sum_{i,j=\lfloor ns \rfloor}^{\lfloor nt \rfloor - 1} |\rho(i - j)|^{d+k} \right)^{1/2} \\
&\leq c_k \left(\frac{1}{n} \sum_{i,j=\lfloor ns \rfloor}^{\lfloor nt \rfloor - 1} |\rho(i - j)|^d \right)^{1/2} \quad \text{since } |\rho(k)| \leq 1.
\end{aligned}$$

Finally, the change of indices $(i, j) \rightarrow (i, j + h)$ leads to

$$\frac{1}{n} \sum_{i,j=\lfloor ns \rfloor}^{\lfloor nt \rfloor - 1} |\rho(i - j)|^d \leq C \frac{\lfloor nt \rfloor - \lfloor ns \rfloor}{n} \sum_{h \in \mathbb{Z}} |\rho(h)|^d = C \frac{\lfloor nt \rfloor - \lfloor ns \rfloor}{n},$$

which provides the desired estimate (3.1) and concludes the proof of Theorem 1.1. \square

4 Proof of Theorem 1.2

Since $Z_n(t) = Y_n(t) + \frac{nt - \lfloor nt \rfloor}{\sqrt{n}} \varphi(X_{\lfloor nt \rfloor})$ with $\mathbb{E} \left[\left(\frac{nt - \lfloor nt \rfloor}{\sqrt{n}} \varphi(X_{\lfloor nt \rfloor}) \right)^2 \right] \leq \frac{1}{n} \|\varphi\|_{L^2(\gamma, \mathbb{R})}^2 \rightarrow 0$, point 1 (that is, convergence of the finite-dimensional distribution of Z_n) follows again from the classical Breuer-Major theorem of [3].

Let us now turn to point 2. It remains to show that the family $(Z_n)_{n \geq 1}$ is tight in the space $\mathbf{C}([0, 1])$. Recall from [2, Theorem 12.3] that a sufficient condition for tightness in $\mathbf{C}([0, 1])$ is this time the existence of $\gamma > 0$ and $c > 0$ such that, for all n ,

$$\|Z_n(t) - Z_n(s)\|_{L^p(\Omega)} \leq c|t - s|^{1/2}, \quad s, t \in [0, 1]. \tag{4.1}$$

Using the equivalent representation

$$Z_n(t) = \frac{1}{\sqrt{n}} \int_0^{nt} \varphi(X_{\lfloor u \rfloor}) du,$$

we can write

$$\|Z_n(t) - Z_n(s)\|_{L^p(\Omega)} = \frac{1}{\sqrt{n}} \left\| \int_{ns}^{nt} \varphi(X_{\lfloor u \rfloor}) du \right\|_{L^p(\Omega)}$$

$$\begin{aligned}
&= \frac{1}{\sqrt{n}} \left\| \int_{ns}^{nt} \delta^d \left(\varphi_d(X_{[u]}) e_{[u]}^{\otimes d} \right) du \right\|_{L^p(\Omega)} \quad \text{by (2.12)} \\
&\leq c_p \sum_{k=0}^d \frac{1}{\sqrt{n}} \left\| \int_{ns}^{nt} D^k \left(\varphi_d(X_{[u]}) e_{[u]}^{\otimes d} \right) du \right\|_{L^p(\Omega; L^2(\mathbb{R}_+^{k+d}))} \quad \text{by (2.8)} \\
&= c_p \sum_{k=0}^d \left\| \frac{1}{n} \iint_{[ns, nt]^2} D^k(\varphi_d(X_{[u]})) D^k(\varphi_d(X_{[v]})) \langle e_{[u]}, e_{[v]} \rangle_{L^2(\mathbb{R}_+)}^{d+k} dudv \right\|_{L^{\frac{p}{2}}(\Omega; L^2(\mathbb{R}_+^k))}^{1/2} \\
&=: c_p \sum_{k=0}^d R_k.
\end{aligned}$$

Minkowski and Hölder inequalities yield, for any $0 \leq k \leq d$,

$$\begin{aligned}
R_k &\leq \sup_{u \in \mathbb{R}_+} \|D^k(\varphi_d(X_{[u]}))\|_{L^p(\Omega; L^2(\mathbb{R}_+^k))} \left(\frac{1}{n} \iint_{[ns, nt]^2} |\rho([u] - [v])|^{d+k} dudv \right)^{1/2} \\
&= c_k \left(\frac{1}{n} \iint_{[ns, nt]^2} |\rho([u] - [v])|^{d+k} dudv \right)^{1/2},
\end{aligned}$$

with c_k finite by (3.4).

Finally, since $|\rho(k)| \leq 1$ for all k ,

$$\begin{aligned}
\frac{1}{n} \iint_{[ns, nt]^2} |\rho([u] - [v])|^{d+k} dudv &\leq \frac{1}{n} \int_{ns}^{nt} \left(\int_{ns-v}^{nt-v} |\rho([x+v] - [v])|^d dx \right) dv \\
&\leq \frac{1}{n} \int_{ns}^{nt} \sum_{j \in \mathbb{Z}} |\rho(j)|^d dv = |t-s| \sum_{j \in \mathbb{Z}} |\rho(j)|^d,
\end{aligned}$$

which provides the desired estimate (4.1) and concludes the proof of Theorem 1.2. \square

5 An extension of Theorem 1.1

In this section, our aim is to show that the method we have employed for the proofs of Theorems 1.1 and 1.2 can be easily extended to deal with the case where $\sum_{|j| \leq n} |\rho(j)|^d$ diverges as a slowly varying function when $n \rightarrow \infty$. Instead of stating such a result at a great level of generality, to avoid too much technicalities we prefer to illustrate what happens in a guiding example and only in the setting of Theorem 1.1. The same extension for Theorem 1.2 would follow similar lines; details are left to the interested reader as an exercise.

Consider the fractional Gaussian noise $X_k = B_{k+1} - B_k$ associated with a fractional Brownian motion B of Hurst index $H \in (0, 1)$; in this case, $\rho(k) = \frac{1}{2}(|k+1|^{2H} + |k-1|^{2H} - 2|k|^{2H})$. Also, consider a function $\varphi \in L^2(\mathbb{R}, \gamma)$ with expansion (1.1) and Hermite rank $d \geq 1$. Finally, recall Y_n from (1.3), let $W = \{W_t\}_{t \in [0,1]}$ be a Brownian motion and let σ^2 be defined in (1.4).

Since $\rho(k) \sim c|k|^{2H-2}$ (where c is an explicit constant whose value is useless), in the case where $H \in (0, 1 - \frac{1}{2d})$ one can apply Breuer-Major theorem of [3] to deduce that $Y_n \xrightarrow{\text{f.d.d.}} \sigma W$. If moreover $\varphi \in L^p(\mathbb{R}, \gamma)$ for some $p > 2$, then $Y_n \xrightarrow{\mathbf{D}([0,1])} \sigma W$ thanks to our Theorem 1.1.

In contrast, when $H \in (1 - \frac{1}{2d}, 1)$ Taqqu [10, Theorem 5.6] has shown in the seventies that

$$n^{d(1-H)-\frac{1}{2}} Y_n \xrightarrow{\mathbf{D}([0,1])} Y_\infty,$$

where Y_∞ stands for the Hermite process of index d . Here, note that no additional integrability condition on φ is required for the convergence to hold in $\mathbf{D}([0,1])$; indeed, since the limiting process Y_∞ is α -Hölder continuous with α *strictly* greater than $\frac{1}{2}$, it is enough to bound $\|Y_n(t) - Y_n(s)\|_{L^2(\Omega)}$ (and not $\|Y_n(t) - Y_n(s)\|_{L^p(\Omega)}$ with $p > 2$) to get the tightness, so classical and easy calculations are enough to conclude.

What about the critical case $H = 1 - \frac{1}{2d}$? In this case, $\rho(k) \sim c|k|^{-\frac{1}{d}}$ and so $\sum_{k \in \mathbb{Z}} |\rho(k)| = +\infty$. Nevertheless, since the divergence of the series is slow, the fluctuations are still Brownian after proper normalisation. More precisely, it is shown in [3] that $\frac{Y_n}{\sqrt{\log n}} \xrightarrow{\text{f.d.d.}} \sigma W$, with $\sigma^2 = 2d! \left(\frac{(2d-1)(d-1)}{2d^2} \right)^d$. As far as the convergence in $\mathbf{D}([0,1])$ is concerned, a slight extension of our method leads to the following result.

Theorem 5.1. *Consider a function $\varphi \in L^2(\mathbb{R}, \gamma)$ with expansion (1.1) and Hermite rank $d \geq 1$. Let $X = \{X_n\}_{n \in \mathbb{Z}}$ be the fractional Gaussian noise of index $H = 1 - \frac{1}{2d}$, that is, X is a mean-zero Gaussian stationary sequence with covariance function*

$$\mathbb{E}[X_n X_{n+k}] = \rho(k) = \frac{1}{2} (|k+1|^{2-\frac{1}{d}} + |k-1|^{2-\frac{1}{d}} - 2|k|^{2-\frac{1}{d}}).$$

Finally, recall Y_n from (1.3), let $W = \{W_t\}_{t \in [0,1]}$ be a Brownian motion and let σ be given by $\sigma^2 = 2d! \left(\frac{(2d-1)(d-1)}{2d^2} \right)^d$. Then, as $n \rightarrow \infty$,

1. The finite-dimensional distributions of $\frac{Y_n}{\sqrt{\log n}}$ converge to those of σW ;
2. If $\varphi \in L^p(\mathbb{R}, \gamma)$ for some $p > 2$, then $\frac{Y_n}{\sqrt{\log n}}$ converges in law to σW in $\mathbf{D}([0,1])$ endowed with the Skorohod topology.

Proof. Point 1 follows from Breuer and Major [3]. Combining Lemma 3.1 with [2, Theorem 15.6] (for $\frac{Y_n}{\sqrt{\log n}}$ instead of Y_n), to prove point 2 it is enough to show that (3.2) holds true.

We can write

$$\begin{aligned} & \left\| \frac{Y_n(t)}{\sqrt{\log n}} - \frac{Y_n(s)}{\sqrt{\log n}} \right\|_{L^p(\Omega)} \\ &= \frac{1}{\sqrt{n \log n}} \left\| \sum_{i=\lfloor ns \rfloor}^{\lfloor nt \rfloor-1} \delta^d \left(\varphi_d(X_i) e_i^{\otimes d} \right) \right\|_{L^p(\Omega)} \quad \text{by (2.12)} \\ &\leq c_p \sum_{k=0}^d \frac{1}{\sqrt{n \log n}} \left\| \sum_{i=\lfloor ns \rfloor}^{\lfloor nt \rfloor-1} D^k \left(\varphi_d(X_i) e_i^{\otimes d} \right) \right\|_{L^p(\Omega; L^2(\mathbb{R}_+^{k+d}))} \quad \text{by (2.8)} \end{aligned}$$

$$\begin{aligned}
&= c_p \sum_{k=0}^d \left\| \frac{1}{n \log n} \sum_{i,j=\lfloor ns \rfloor}^{\lfloor nt \rfloor-1} D^k(\varphi_d(X_i)) D^k(\varphi_d(X_j)) \langle e_i, e_j \rangle_{L^2(\mathbb{R}_+)}^{d+k} \right\|_{L^{\frac{p}{2}}(\Omega; L^2(\mathbb{R}_+^k))}^{1/2} \\
&\leq c_p \sum_{k=0}^d \sup_{i \in \mathbb{Z}} \|D^k(\varphi_d(X_i))\|_{L^p(\Omega; L^2(\mathbb{R}_+^k))} \left(\frac{1}{n \log n} \sum_{i,j=\lfloor ns \rfloor}^{\lfloor nt \rfloor-1} |\rho(i-j)|^{d+k} \right)^{1/2} \\
&\leq c \left(\frac{1}{n \log n} \sum_{i,j=\lfloor ns \rfloor}^{\lfloor nt \rfloor-1} |\rho(i-j)|^d \right)^{1/2} \quad \text{since } |\rho(k)| \leq 1 \text{ and using (3.4).}
\end{aligned}$$

Finally, the change of indices $(i, j) \rightarrow (i, j + h)$ and the fact that $|\rho(r)|^d \sim c|r|^{-1}$ as $|r| \rightarrow \infty$ leads to

$$\frac{1}{n \log n} \sum_{i,j=\lfloor ns \rfloor}^{\lfloor nt \rfloor-1} |\rho(i-j)|^d \leq C \frac{\lfloor nt \rfloor - \lfloor ns \rfloor}{n},$$

which provides the desired estimate (3.1) and concludes the proof of Theorem 1.1. \square

References

- [1] S. Ben Hariz (2002): Limit theorems for the non-linear functionals of stationary Gaussian processes. *J. Mult. Anal.* **80**, pp. 191-216.
- [2] P. Billingsley (1968): *Convergence of Probability Measures*. Wiley.
- [3] P. Breuer and P. Major (1983): Central limit theorems for non-linear functionals of Gaussian fields. *J. Mult. Anal.* **13**, pp. 425-441.
- [4] D. Chambers and E. Slud (1989): Central limit theorems for nonlinear functionals of stationary Gaussian processes. *Probab. Th. Rel. Fields* **80**, pp. 323-346.
- [5] A. Jaramillo and D. Nualart (2018): Functional limit theorem for the self-intersection local time of the fractional Brownian motion. *Ann. Inst. H. Poincaré*. To appear.
- [6] P. A. Meyer (1984): Transformations de Riesz pour les lois gaussiennes. *Lecture Notes in Math.* **1059**, pp. 179-193.
- [7] I. Nourdin and G. Peccati (2012): *Normal approximations with Malliavin calculus: from Stein's method to universality*. Cambridge tracts in Mathematics **192**, Cambridge University Press.
- [8] D. Nualart (2006): *The Malliavin calculus and related topics*. 2nd edition. Probability and Its Applications, Springer.
- [9] D. Nualart and E. Nualart (2018): *Introduction to Malliavin Calculus*. Institute of Mathematical Statistics Textbooks, Cambridge University Press.
- [10] M. Taqqu (1979): Convergence of integrated processes of arbitrary Hermite rank. *Z. Wahrsch. verw. Gebiete* **50**, pp. 53-83.