

THE SCHWARZ–VORONOV EMBEDDING OF \mathbb{Z}_2^n -MANIFOLDS

ANDREW JAMES BRUCE, EDUARDO IBARGUENGOYTIA, AND NORBERT PONCIN

ABSTRACT. Informally, \mathbb{Z}_2^n -manifolds are ‘manifolds’ with \mathbb{Z}_2^n -graded coordinates and a sign rule determined by the standard scalar product of their \mathbb{Z}_2^n -degrees. Such manifolds can be understood in a sheaf-theoretic framework, as supermanifolds can, but with significant differences, in particular in integration theory. In this paper, we reformulate the notion of a \mathbb{Z}_2^n -manifold within a categorical framework via the functor of points. We show that it is sufficient to consider \mathbb{Z}_2^n -points, i.e., trivial \mathbb{Z}_2^n -manifolds for which the reduced manifold is just a single point, as ‘probes’ when employing the functor of points. This allows us to construct a fully faithful restricted Yoneda embedding of the category of \mathbb{Z}_2^n -manifolds into a subcategory of contravariant functors from the category of \mathbb{Z}_2^n -points to a category of Fréchet manifolds over algebras. We refer to this embedding as the *Schwarz–Voronov embedding*. We further prove that the category of \mathbb{Z}_2^n -manifolds is equivalent to the full subcategory of locally trivial functors in the preceding subcategory.

Keywords: supergeometry; superalgebra; ringed spaces; higher grading; functor of points

MSC 2010: *Primary:* 58C50; 58D1; 14A22
Secondary: 13F25; 16L30; 17A70

CONTENTS

1. Introduction	1
2. Rudiments of \mathbb{Z}_2^n -graded geometry	3
2.1. The category of \mathbb{Z}_2^n -manifolds	3
2.2. The functor of points	5
3. \mathbb{Z}_2^n -points and the functor of points	7
3.1. The category of \mathbb{Z}_2^n -points	7
3.2. A convenient generating set of \mathbb{Z}_2^n Man	8
3.3. Restricted Yoneda functor and fullness	11
3.4. The manifold structure on the set of Λ -points	19
3.5. The Schwarz–Voronov embedding	22
3.6. Representability and equivalence of categories	25
Appendix A. Generating sets of categories	27
Appendix B. Fréchet spaces, modules and manifolds	28
References	31

1. INTRODUCTION

Various notions of *graded geometry* play an important rôle in mathematical physics and can often provide further insight into classical geometric constructions. For example, supermanifolds, as pioneered by Berezin and collaborators, are essential in describing quasi-classical systems with both bosonic and fermionic degrees of freedom. Very loosely, supermanifolds are ‘manifolds’ for which the structure sheaf is \mathbb{Z}_2 -graded. Such geometries are of fundamental importance in perturbative string theory, supergravity, and the BV-formalism, for example. While the theory of supermanifolds is firmly rooted in theoretical physics, it has since become a respectable area of mathematical research. Indeed, supermanifolds allow for an economical description of Lie algebroids, Courant algebroids as well as various related structures, many of which are of direct interest to physics. We will not elaborate any further and urge the reader to consult the ever-expanding literature.

Interestingly, \mathbb{Z}_2^n -gradings ($\mathbb{Z}_2^n = \mathbb{Z}_2^{\times n}$, $n \geq 2$) can be found in the theory of parastatistics, see for example [21, 25, 26, 61], behind an alternative approach to supersymmetry [55], in relation to the symmetries of the Lévy-Lebony equation [2], and behind the theory of mixed symmetry tensors [12]. Generalizations of the super Schrödinger algebra (see [3]) and the super Poincaré algebra (see [11]) have also appeared in the literature. That said, it is unknown if these ‘higher gradings’ are of the same importance in fundamental physics as \mathbb{Z}_2 -gradings. It must also be remarked that the quaternions and more general Clifford algebras can be understood as \mathbb{Z}_2^n -graded \mathbb{Z}_2^n -commutative (see below) algebras [4, 5]. Thus, one may expect \mathbb{Z}_2^n -gradings to be important in studying Clifford algebras and modules, though the implications for classical and quantum field theory remain

as of yet unexplored. It should be further mentioned that any ‘sign rule’ can be understood in terms of a \mathbb{Z}_2^n -grading (see [16]). A natural question here is to what extent can \mathbb{Z}_2^n -graded geometry be developed.

A locally ringed space approach to \mathbb{Z}_2^n -manifolds has been constructed in a series of papers by Bruce, Covolo, Grabowski, Kwok, Ovsienko & Poncin [20, 16, 17, 18, 19, 40, 12, 14]. It includes the \mathbb{Z}_2^n -differential-calculus, the \mathbb{Z}_2^n -Berezinian, as well as a low dimensional \mathbb{Z}_2^n -integration-theory. Integration on \mathbb{Z}_2^n -manifolds turns out to be fundamentally different from integration on \mathbb{Z}_2^1 -manifolds (i.e., supermanifolds) and is currently being constructed in full generality by authors of the present paper.

Loosely, \mathbb{Z}_2^n -manifolds are ‘manifolds’ for which the structure sheaf has a \mathbb{Z}_2^n -grading and the commutation rule for the local coordinates comes from the standard scalar product of their \mathbb{Z}_2^n -degrees. This is not just a trivial or straightforward generalization of the notion of a supermanifold as one has to deal with formal coordinates that anticommute with other formal coordinates, but are themselves *not* nilpotent. Due to the presence of formal variables that are not nilpotent, formal power series are used rather than polynomials (for standard supermanifolds all functions are polynomial in the Grassmann odd variables). The use of formal power series is unavoidable in order to have a well-defined local theory (see [16]), and a well-defined differential calculus (see [18]). Heuristically, one can view supermanifolds as ‘mild’ noncommutative geometries: the noncommutativity is seen simply as anticommutativity of the odd coordinates. In a similar vein, one can view \mathbb{Z}_2^n -manifolds ($n > 1$) as examples of ‘mild’ nonsupercommutative geometries: the sign rule involved is not determined by the coordinates being even or odd, i.e., by their total degree, but by their \mathbb{Z}_2^n -degree.

The idea of understanding supermanifolds, i.e., \mathbb{Z}_2^1 -manifolds, as ‘Grassmann algebra valued manifolds’ can be traced back to the pioneering work of Berezin [10]. An informal understanding along these lines has continuously been employed in physics, where one chooses a ‘large enough’ Grassmann algebra to capture the aspects to the theory needed. This informal understanding leads to the DeWitt–Rogers approach to supermanifolds which seemed to avoid the theory of locally ringed spaces altogether. However, arbitrariness in the choice of the underlying Grassmann algebra is somewhat displeasing. Furthermore, developing the mathematical consistency of DeWitt–Rogers supermanifolds takes one back to the sheaf-theoretic approach of Berezin & Leites: for a comparison of these approaches, the reader can consult Rogers [43] or Schmitt [47]. From a physics perspective, there seems no compelling reason to think that there is any physical significance to the choice of underlying Grassmann algebra. To quote Schmitt [47]: “However, no one has ever measured a Grassmann number, everyone measures real numbers”. The solution here is, following Schwarz & Voronov [49, 54, 58], not to fix the underlying Grassmann algebra, but rather understand supermanifolds as functors from the category of finite-dimensional Grassmann algebras to, in the first instance, the category of sets. For a given, but arbitrary, Grassmann algebra Λ , one speaks of the set of Λ -points of a supermanifold. It is well known that the set of Λ -points of a given supermanifold comes with the further structure of a Λ_0 -smooth manifold. That is we, in fact, do not only have a set, but also the structure of a finite-dimensional manifold whose tangent spaces are Λ_0 -modules. Moreover, thinking of supermanifolds as functors, not all natural transformations between the Λ -points correspond to genuine supermanifold morphisms, only those that respect the Λ_0 -smooth structure do. The classical roots of these ideas go back to Weil [60] who considered the A -points of a manifold as the set of maps from the algebra of smooth functions on the manifold to a specified finite-dimensional commutative local algebra A . Today one refers to *Weil functors* and these have long been utilised in the theory of jet structures over manifolds, see for example [31].

In this paper, we study *Grothendieck’s functor of points* – from the category opposite to \mathbb{Z}_2^n -manifolds to the category of sets (see [27]) – and restrict it to the category of \mathbb{Z}_2^n -points, i.e., trivial \mathbb{Z}_2^n -manifolds $\mathbb{R}^{0|q}$ that have no degree zero coordinates. More precisely we consider the restricted Yoneda functor from the category of \mathbb{Z}_2^n -manifolds to the category of contravariant functors from \mathbb{Z}_2^n -points to sets. Dual to \mathbb{Z}_2^n -points are what we will call \mathbb{Z}_2^n -Grassmann algebras, which we will also denote as Λ . The aim of this paper is to generalize and carefully prove the main results of Schwarz & Voronov [54, 58] to the ‘higher graded’ setting. In particular, we show that \mathbb{Z}_2^n -points are sufficient to act as ‘probes’ when employing the functor of points. However, not all natural transformations between the sets of Λ -points correspond to morphisms of the underlying \mathbb{Z}_2^n -manifolds. By carefully analyzing the image of the functor of points, we prove that the set of Λ -points of a \mathbb{Z}_2^n -manifold comes with the extra structure of an infinite-dimensional Λ_0 -manifold. By Λ_0 we mean the subalgebra of degree zero elements of the \mathbb{Z}_2^n -Grassmann algebra Λ . Note that we are not trying to define infinite-dimensional \mathbb{Z}_2^n -manifolds, yet infinite-dimensional manifolds, specifically nuclear Fréchet manifolds, are fundamental to the theory. Moreover, we show that natural transformations between sets of Λ -points arise from morphisms of \mathbb{Z}_2^n -manifolds if and only if they respect the Fréchet Λ_0 -manifold structures. Thus, by restricting the natural transformations allowed, we build a full and faithful embedding of the category of \mathbb{Z}_2^n -manifolds into the category of contravariant functors from the category of \mathbb{Z}_2^n -points to the category of nuclear Fréchet Λ_0 -smooth manifolds. This embedding we refer to as the *Schwarz–Voronov embedding*. We finally study representability of such contravariant functors and prove that the category of \mathbb{Z}_2^n -manifolds is equivalent to the full subcategory of locally trivial functors in the just depicted subcategory of contravariant functors from \mathbb{Z}_2^n -points to nuclear Fréchet manifolds.

Methodology: As \mathbb{Z}_2^n -manifolds have well defined local models, we work with \mathbb{Z}_2^n -domains and then ‘globalize’

the results to general \mathbb{Z}_2^n -manifolds. We modify the approach of Schwarz & Voronov [54, 58] and draw on Balduzzi, Carmeli & Fiorese [8, 9] and Konechny & Schwarz [32, 33], making all changes necessary to encompass \mathbb{Z}_2^n -manifolds. The most striking difference between supermanifolds and \mathbb{Z}_2^n -manifolds ($n > 1$) is that we are forced, due to the presence of non-zero degree even coordinates, to work with (infinite-dimensional) Fréchet spaces, algebras and manifolds. Nuclearity is not compulsory.

Arrangement: In Section 2 we review the basic tenets of \mathbb{Z}_2^n -geometry and the theory of \mathbb{Z}_2^n -manifolds. The bulk of this paper is to be found in Section 3. We rely on two appendices: in Appendix A we recall the notion of a generating set of a category, and in Appendix B we review indispensable concepts from the theory of Fréchet spaces, algebras and manifolds.

2. RUDIMENTS OF \mathbb{Z}_2^n -GRADED GEOMETRY

2.1. The category of \mathbb{Z}_2^n -manifolds. The locally ringed space approach to \mathbb{Z}_2^n -manifolds is presented in a series of papers [20, 16, 17, 18, 19, 40] by Covolo, Grabowski, Kwok, Ovsienko, and Poncin. We will draw upon these works heavily and not present proofs of any formal statements.

Definition 2.1. A *locally \mathbb{Z}_2^n -ringed space*, $n \in \mathbb{N}$, is a pair $X := (|X|, \mathcal{O}_X)$, where $|X|$ is a second-countable Hausdorff space, and a \mathcal{O}_X is a sheaf of \mathbb{Z}_2^n -graded \mathbb{Z}_2^n -commutative associative unital \mathbb{R} -algebras, such that the stalks \mathcal{O}_p , $p \in |X|$, are local rings.

In this context, \mathbb{Z}_2^n -commutative means that any two sections $a, b \in \mathcal{O}_X(|U|)$, $|U| \subset |X|$ open, of homogeneous degrees $\deg(a) = \underline{a} \in \mathbb{Z}_2^n$ and $\deg(b) = \underline{b} \in \mathbb{Z}_2^n$ commute according to the sign rule

$$ab = (-1)^{\langle \underline{a}, \underline{b} \rangle} ba,$$

where $\langle \cdot, \cdot \rangle$ is the standard scalar product on \mathbb{Z}_2^n . We will say that a section a is *even* or *odd* if $\langle \underline{a}, \underline{a} \rangle \in \mathbb{Z}_2$ is 0 or 1.

Just as in standard supergeometry, which we recover for $n = 1$, a locally \mathbb{Z}_2^n -ringed space is a \mathbb{Z}_2^n -manifold if it is locally isomorphic to a specific local model. Given the central rôle of (finite dimensional) Grassmann algebras in the theory of supermanifolds, we consider here \mathbb{Z}_2^n -Grassmann algebras.

Remark 2.2. In the following, we order the elements in \mathbb{Z}_2^n *lexicographically*, and refer to this ordering as the *standard ordering*. For example, we thus get

$$\mathbb{Z}_2^2 = \{(0,0), (0,1), (1,0), (1,1)\}.$$

Definition 2.3. A \mathbb{Z}_2^n -Grassmann algebra $\Lambda^{\underline{q}} := \mathbb{R}[[\xi]]$ is the \mathbb{Z}_2^n -graded \mathbb{Z}_2^n -commutative associative unital \mathbb{R} -algebra of all formal power series with coefficients in \mathbb{R} generated by homogeneous parameters ξ^α subject to the commutation relation

$$\xi^\alpha \xi^\beta = (-1)^{\langle \underline{\alpha}, \underline{\beta} \rangle} \xi^\beta \xi^\alpha,$$

where $\underline{\alpha} := \deg(\xi^\alpha) \in \mathbb{Z}_2^n \setminus \underline{0}$, $\underline{0} = (0, \dots, 0)$. The tuple $\underline{q} = (q_1, q_2, \dots, q_N)$, $N = 2^n - 1$, provides the number q_i of generators ξ^α , which have the i -th degree in $\mathbb{Z}_2^n \setminus \underline{0}$ (endowed with its standard order).

A *morphism of \mathbb{Z}_2^n -Grassmann algebras*, $\psi^* : \Lambda^{\underline{q}} \rightarrow \Lambda^{\underline{p}}$, is a map of \mathbb{R} -algebras that preserves the \mathbb{Z}_2^n -grading and the units.

We denote the category of \mathbb{Z}_2^n -Grassmann algebras and corresponding morphisms by $\mathbb{Z}_2^n\text{-GrAlg}$.

Example 2.4. For $n = 0$, we simply get \mathbb{R} considered as an algebra over itself.

Example 2.5. If $n = 1$, we recover the classical concept of Grassmann algebra with the standard supercommutation rule for generators. In this case, all formal power series truncate to polynomials. In particular, the Grassmann algebra generated by a single odd generator is isomorphic to the algebra of dual numbers.

Example 2.6. The \mathbb{Z}_2^2 -Grassmann algebra $\Lambda^{(1,1,1)}$ is described by three generators

$$(\underbrace{\xi}_{(0,1)}, \underbrace{\theta}_{(1,0)}, \underbrace{z}_{(1,1)}),$$

where we have indicated the \mathbb{Z}_2^2 -degree. Note that $\xi\theta = \theta\xi$, while $\xi^2 = 0$ and $\theta^2 = 0$. Moreover, $\xi z = -z\xi$ and $\theta z = -z\theta$, while z is not nilpotent. A general (inhomogeneous) element of $\Lambda^{(1,1,1)}$ is then of the form

$$f(\xi, \theta, z) = f_z(z) + \xi f_\xi(z) + \theta f_\theta(z) + \xi\theta f_{\xi\theta}(z),$$

where $f_z(z)$, $f_\xi(z)$, $f_\theta(z)$ and $f_{\xi\theta}(z)$ are formal power series in z . As a subalgebra we can consider $\Lambda^{(1,1,0)}$, whose generators are ξ and θ . A general element of this subalgebra is a polynomial in these generators.

Within any \mathbb{Z}_2^n -Grassmann algebra $\Lambda := \Lambda^{\underline{q}}$, we have the ideal generated by the generators of Λ , which we will denote as $\mathring{\Lambda}$. In particular we have the decomposition

$$\Lambda = \mathbb{R} \oplus \mathring{\Lambda},$$

which will be used later on. Moreover, the set of degree 0 elements, $\Lambda_0 \subset \Lambda$, is a commutative associative unital \mathbb{R} -algebra.

Very informally, a \mathbb{Z}_2^n -manifold is a smooth manifold whose structure sheaf has been ‘deformed’ to now include the generators of a \mathbb{Z}_2^n -Grassmann algebra.

Definition 2.7. A (smooth) \mathbb{Z}_2^n -manifold of dimension $p|q$ is a locally \mathbb{Z}_2^n -ringed space $M := (|M|, \mathcal{O}_M)$, which is locally isomorphic to the locally \mathbb{Z}_2^n -ringed space $\mathbb{R}^{p|q} := (\mathbb{R}^p, C_{\mathbb{R}^p}^\infty[[\xi]])$. Local sections of \mathcal{O}_M are thus formal power series in the \mathbb{Z}_2^n -graded variables ξ with smooth coefficients,

$$\mathcal{O}_M(|U|) \simeq C_{\mathbb{R}^p}^\infty(|U|)[[\xi]] := \left\{ \sum_{\alpha \in \mathbb{N}^{\sum_i q_i}} \xi^\alpha f_\alpha : f_\alpha \in C_{\mathbb{R}^p}^\infty(|U|) \right\},$$

for ‘small enough’ open subsets $|U| \subset |M|$. A \mathbb{Z}_2^n -morphism, i.e., a morphism between two \mathbb{Z}_2^n -manifolds, say M and N , is a morphism of \mathbb{Z}_2^n -ringed spaces, that is, a pair $\phi = (|\phi|, \phi^*) : (|M|, \mathcal{O}_M) \rightarrow (|N|, \mathcal{O}_N)$ consisting of a continuous map $|\phi| : |M| \rightarrow |N|$ and a sheaf morphism $\phi^* : \mathcal{O}_N \rightarrow |\phi|_* \mathcal{O}_M$, i.e., a family of \mathbb{Z}_2^n -graded unital \mathbb{R} -algebra morphisms $\phi_{|V|}^* : \mathcal{O}_N(|V|) \rightarrow \mathcal{O}_M(|\phi|^{-1}(|V|))$ ($|V| \subset |N|$ open), which commute with restrictions. We will refer to the global sections of the structure sheaf \mathcal{O}_M as *functions* on M and denote them as $C^\infty(M) := \mathcal{O}_M(|M|)$.

Example 2.8 (The local model). The locally \mathbb{Z}_2^n -ringed space $\mathcal{U}^{p|q} := (\mathcal{U}^p, C_{\mathcal{U}^p}^\infty[[\xi]])$, where $\mathcal{U}^p \subset \mathbb{R}^p$ is open, is naturally a \mathbb{Z}_2^n -manifold – we refer to such \mathbb{Z}_2^n -manifolds as \mathbb{Z}_2^n -domains of dimension $p|q$. We can employ (natural) coordinates (x^a, ξ^α) on any \mathbb{Z}_2^n -domain, where the x^a form a coordinate system on \mathcal{U}^p and the ξ^α are formal coordinates.

Canonically associated to any \mathbb{Z}_2^n -graded algebra \mathcal{A} is the homogeneous ideal J of \mathcal{A} generated by all homogeneous elements of \mathcal{A} having nonzero degree. If $f : \mathcal{A} \rightarrow \mathcal{A}'$ is a morphism of \mathbb{Z}_2^n -graded algebras, then $f(J_{\mathcal{A}}) \subset J_{\mathcal{A}'}$. The J -adic topology plays a fundamental rôle in the theory of \mathbb{Z}_2^n -manifolds. In particular, these notions can be ‘sheafified’. That is, for any \mathbb{Z}_2^n -manifold M , there exists an ideal sheaf \mathcal{J}_M , defined by $\mathcal{J}(|U|) = \langle f \in \mathcal{O}_M(|U|) : \deg(f) \neq 0 \rangle$. The \mathcal{J}_M -adic topology on \mathcal{O}_M can then be defined in the obvious way.

Many of the standard results from the theory of supermanifolds pass over to \mathbb{Z}_2^n -manifolds. For example, the topological space $|M|$ comes with the structure of a smooth manifold of dimension p and the continuous base map of any \mathbb{Z}_2^n -morphism is actually smooth. Further, for any \mathbb{Z}_2^n -manifold M , there exists a short exact sequence of sheaves of \mathbb{Z}_2^n -graded \mathbb{Z}_2^n -commutative associative \mathbb{R} -algebras

$$0 \longrightarrow \ker \epsilon \longrightarrow \mathcal{O}_M \xrightarrow{\epsilon} C_{|M|}^\infty \longrightarrow 0,$$

such that $\ker \epsilon = \mathcal{J}_M$.

The immediate problem with \mathbb{Z}_2^n -manifolds is that \mathcal{J}_M is *not* nilpotent – for supermanifolds the ideal sheaf is nilpotent and this is a fundamental property that makes the theory of supermanifolds so well-behaved. However, this loss of nilpotency is compensated by Hausdorff completeness of \mathcal{O}_M with respect to the \mathcal{J}_M -adic topology.

Proposition 2.9. Let M be a \mathbb{Z}_2^n -manifold. Then \mathcal{O}_M is \mathcal{J}_M -adically Hausdorff complete as a sheaf of \mathbb{Z}_2^n -commutative associative unital \mathbb{R} -algebras, i.e., the morphism

$$\mathcal{O}_M \rightarrow \lim_{\leftarrow k} \mathcal{O}_M / \mathcal{J}_M^k,$$

naturally induced by the filtration of \mathcal{O}_M by the powers of \mathcal{J}_M , is an isomorphism.

The presence of formal power series in the coordinate rings of \mathbb{Z}_2^n -manifolds forces one to rely on the Hausdorff-completeness of the \mathcal{J} -adic topology. This completeness replaces the standard fact that supermanifold functions of Grassmann odd variables are always polynomials – a result that is often used in extending results from smooth manifolds to supermanifolds.

What makes \mathbb{Z}_2^n -manifolds a very workable form of noncommutative geometry is the fact that we have well-defined local models. Much like the theory of manifolds, one can construct global geometric concepts via the gluing of local geometric concepts. That is, we can consider a \mathbb{Z}_2^n -manifold as being covered by \mathbb{Z}_2^n -domains together with specified gluing information, i.e., coordinate transformations. Moreover, we have the *chart theorem* ([16, Theorem 7.10]) that says that \mathbb{Z}_2^n -morphisms from a \mathbb{Z}_2^n -manifold $(|M|, \mathcal{O}_M)$ to a \mathbb{Z}_2^n -domain $(\mathcal{U}^p, C_{\mathcal{U}^p}^\infty[[\xi]])$, are completely described by the pullbacks of the coordinates (x^a, ξ^α) . In other words, to define a \mathbb{Z}_2^n -morphism valued in a \mathbb{Z}_2^n -domain, we only need to provide total sections $(s^a, s^\alpha) \in \mathcal{O}_M(|M|)$ of the source structure sheaf, whose degrees coincide with those of the target coordinates (x^a, ξ^α) . Let us stress the condition $(\dots, \epsilon s^a, \dots)(|M|) \subset \mathcal{U}^p$, which is often understood in the literature.

A few words about the atlas definition of a \mathbb{Z}_2^n -manifold are necessary. Let $p|q$ be as above. A $p|q$ -chart (or $p|q$ -coordinate-system) over a (second-countable Hausdorff) smooth manifold $|M|$ is a \mathbb{Z}_2^n -domain

$$\mathcal{U}^{p|q} = (\mathcal{U}^p, C_{\mathcal{U}^p}^\infty[[\xi]]),$$

together with a diffeomorphism $|\psi| : |U| \rightarrow \mathcal{U}^p$, where $|U|$ is an open subset of $|M|$. Given two $p|q$ -charts

$$(2.1) \quad (\mathcal{U}_\alpha^{p|q}, |\psi_\alpha|) \quad \text{and} \quad (\mathcal{U}_\beta^{p|q}, |\psi_\beta|)$$

over $|M|$, we denote by $|\psi_{\beta\alpha}|$ the diffeomorphism

$$(2.2) \quad |\psi_{\beta\alpha}| := |\psi_\beta||\psi_\alpha|^{-1} : V_{\alpha\beta} := |\psi_\alpha|(|U_{\alpha\beta}|) := |\psi_\alpha|(|U_\alpha| \cap |U_\beta|) \rightarrow V_{\beta\alpha} := |\psi_\beta|(|U_{\beta\alpha}|) := |\psi_\beta|(|U_\beta| \cap |U_\alpha|).$$

Whereas in classical differential geometry the coordinate transformations are completely defined by the coordinate systems, in \mathbb{Z}_2^n -geometry, they have to be specified separately. A *coordinate transformation* between two charts, say the ones of (2.1), is an isomorphism of \mathbb{Z}_2^n -manifolds

$$(2.3) \quad \psi_{\beta\alpha} = (|\psi_{\beta\alpha}|, \psi_{\beta\alpha}^*) : \mathcal{U}_\alpha^{p|q}|_{V_{\alpha\beta}} \rightarrow \mathcal{U}_\beta^{p|q}|_{V_{\beta\alpha}},$$

where the source and target are the open \mathbb{Z}_2^n -submanifolds

$$\mathcal{U}_\alpha^{p|q}|_{V_{\alpha\beta}} = (V_{\alpha\beta}, C_{V_{\alpha\beta}}^\infty[[\xi]])$$

(note that the underlying diffeomorphism is (2.2)). A $p|q$ -atlas over $|M|$ is a covering $(\mathcal{U}_\alpha^{p|q}, |\psi_\alpha|)_\alpha$ by charts together with a coordinate transformation (2.3) for each pair of charts, such that the usual cocycle condition $\psi_{\beta\gamma}\psi_{\gamma\alpha} = \psi_{\beta\alpha}$ holds (appropriate restrictions are understood).

Definition 2.10. A (smooth) \mathbb{Z}_2^n -manifold of dimension $p|q$ is a (second-countable Hausdorff) smooth manifold $|M|$ together with a preferred $p|q$ -atlas over it.

As in standard supergeometry, the definitions 2.7 and 2.10 are equivalent [35]. For instance, if $M = (|M|, \mathcal{O}_M)$ is a \mathbb{Z}_2^n -manifold of dimension $p|q$ in the sense of Definition 2.7, there are \mathbb{Z}_2^n -isomorphisms (isomorphisms of \mathbb{Z}_2^n -manifolds)

$$h_\alpha = (|h_\alpha|, h_\alpha^*) : U_\alpha = (|U_\alpha|, \mathcal{O}_M|_{|U_\alpha|}) \rightarrow \mathcal{U}_\alpha^{p|q} = (\mathcal{U}_\alpha^p, C_{\mathbb{R}^p}^\infty|_{\mathcal{U}_\alpha^p}[[\xi]]),$$

such that $(|U_\alpha|)_\alpha$ is an open cover of $|M|$. For any two indices α, β , the restriction $h_\alpha|_{U_{\alpha\beta}}$ of h_α to the open \mathbb{Z}_2^n -submanifold $U_{\alpha\beta} = (|U_{\alpha\beta}|, \mathcal{O}_M|_{|U_{\alpha\beta}|})$, $|U_{\alpha\beta}| = |U_\alpha| \cap |U_\beta|$, is a \mathbb{Z}_2^n -isomorphism between $U_{\alpha\beta}$ and

$$\mathcal{U}_\alpha^{p|q}|_{V_{\alpha\beta}} = (V_{\alpha\beta}, C_{\mathbb{R}^p}^\infty|_{V_{\alpha\beta}}[[\xi]]), \quad V_{\alpha\beta} = |h_\alpha|(|U_{\alpha\beta}|).$$

Therefore, the composite

$$(2.4) \quad \psi_{\beta\alpha} = h_\beta|_{U_{\beta\alpha}} h_\alpha|_{U_{\alpha\beta}}^{-1}$$

is a \mathbb{Z}_2^n -isomorphism

$$\psi_{\beta\alpha} : \mathcal{U}_\alpha^{p|q}|_{V_{\alpha\beta}} \rightarrow \mathcal{U}_\beta^{p|q}|_{V_{\beta\alpha}},$$

such that the cocycle condition is satisfied. As a matter of some formality, \mathbb{Z}_2^n -manifolds and their morphisms form a category. The category of \mathbb{Z}_2^n -manifolds we will denote as $\mathbb{Z}_2^n\text{Man}$. We remark this category is locally small. Moreover, as shown in [14, Theorem 19], the category of \mathbb{Z}_2^n -manifolds admits (finite) products. More precisely, let M_i , $i \in \{1, 2\}$, be \mathbb{Z}_2^n -manifolds. Then there exists a \mathbb{Z}_2^n -manifold $M_1 \times M_2$ and \mathbb{Z}_2^n -morphisms $\pi_i : M_1 \times M_2 \rightarrow M_i$ (with underlying smooth manifold $|M_1 \times M_2| = |M_1| \times |M_2|$ and with underlying smooth morphisms $|\pi_i| : |M_1| \times |M_2| \rightarrow |M_i|$ given by the canonical projections), such that for any \mathbb{Z}_2^n -manifold N and \mathbb{Z}_2^n -morphisms $f_i : N \rightarrow M_i$, there exists a unique morphism $h : N \rightarrow M_1 \times M_2$ making the obvious diagram commute. It follows that, if $\phi \in \text{Hom}_{\mathbb{Z}_2^n\text{Man}}(M, M')$ and $\psi \in \text{Hom}_{\mathbb{Z}_2^n\text{Man}}(N, N')$, there is a unique morphism $\phi \times \psi \in \text{Hom}_{\mathbb{Z}_2^n\text{Man}}(M \times N, M' \times N')$.

Remark 2.11. It is known that an analogue of the Batchelor–Gawędzki theorem holds in the category of (real) \mathbb{Z}_2^n -manifolds, see [17, Theorem 3.2]. That is, any \mathbb{Z}_2^n -manifold is noncanonically isomorphic to a $\mathbb{Z}_2^n \setminus \{0\}$ -graded vector bundle over a smooth manifold. While this result is quite remarkable, we will not exploit it at all in this paper.

2.2. The functor of points. Similar to what happens in classical supergeometry, a \mathbb{Z}_2^n -manifold M is not fully described by its topological points in $|M|$. To remedy this defect, we broaden the notion of ‘point’, as was suggested by Grothendieck in the context of algebraic geometry.

More precisely, set $V = \{z \in \mathbb{C}^n : P(z) = 0\} \in \text{Aff}$, where P denotes a polynomial in n indeterminates with complex coefficients and Aff denotes the category of affine varieties. Grothendieck insisted on solving the equation $P(z) = 0$ not only in \mathbb{C}^n , but in A^n , for any algebra A in the category CA of commutative (associative unital) algebras (over \mathbb{C}). This leads to an arrow

$$\text{Sol}_P : \text{CA} \ni A \mapsto \text{Sol}_P(A) = \{a \in A^n : P(a) = 0\} \in \text{Set},$$

which turns out to be a functor

$$\text{Sol}_P \simeq \text{Hom}_{\text{CA}}(\mathbb{C}[V], -) \in [\text{CA}, \text{Set}],$$

where $\mathbb{C}[V]$ is the algebra of polynomial functions of V . The dual of this functor, whose value $\text{Sol}_P(A)$ is the set of A -points of V , is the functor

$$\underline{\text{Hom}}_{\mathbf{Aff}}(-, V) \in [\mathbf{Aff}^{\text{op}}, \mathbf{Set}] ,$$

whose value $\underline{\text{Hom}}_{\mathbf{Aff}}(W, V)$ is the set of W -points of V .

The latter functor can be considered not only in \mathbf{Aff} , but in any locally small category, in particular in $\mathbb{Z}_2^n\text{Man}$. We thus obtain a covariant functor (functor in \bullet)

$$(2.5) \quad \bullet(-) = \underline{\text{Hom}}(-, \bullet) : \mathbb{Z}_2^n\text{Man} \ni M \mapsto M(-) = \underline{\text{Hom}}_{\mathbb{Z}_2^n\text{Man}}(-, M) \in [\mathbb{Z}_2^n\text{Man}^{\text{op}}, \mathbf{Set}] .$$

As suggested above, the contravariant functor $\underline{\text{Hom}}(-, M)$ (we omit the subscript $\mathbb{Z}_2^n\text{Man}$) (functor in $-$) is referred to as the *functor of points* of M . If $S \in \mathbb{Z}_2^n\text{Man}$, an S -*point* of M is just a morphism $\pi_S \in \underline{\text{Hom}}(S, M)$. One may regard an S -point of M as a ‘family of points of M parameterised by the points of S ’. The functor $\bullet(-)$ is known as the *Yoneda embedding*. For any underlying locally small category \mathbf{C} (here $\mathbf{C} = \mathbb{Z}_2^n\text{Man}$), the functor $\bullet(-)$ is *fully faithful*, what means that, for any $M, N \in \mathbb{Z}_2^n\text{Man}$, the map

$$\bullet_{M, N}(-) : \underline{\text{Hom}}(M, N) \ni \phi \mapsto \underline{\text{Hom}}(-, \phi) \in \text{Nat}(\underline{\text{Hom}}(-, M), \underline{\text{Hom}}(-, N))$$

is bijective (here Nat denotes the set of natural transformations). It can be checked that the correspondence $\bullet_{M, N}(-)$ is natural in M and in N . Moreover, any fully faithful functor is automatically injective up to isomorphism on objects: $M(-) \simeq N(-)$ implies $M \simeq N$. Of course, the functor $\bullet(-)$ is not surjective up to isomorphism on objects, i.e., not every functor $X \in [\mathbb{Z}_2^n\text{Man}^{\text{op}}, \mathbf{Set}]$ is isomorphic to a functor of the type $M(-)$. However, if such M does exist, it is, due to the mentioned injectivity, unique up to isomorphism and it is called ‘*the* representing \mathbb{Z}_2^n -manifold of X . Further, if $X, Y \in [\mathbb{Z}_2^n\text{Man}^{\text{op}}, \mathbf{Set}]$ are two representable functors, represented by M, N respectively, a morphism or natural transformation between them, provides, due to the mentioned bijectivity, a *unique morphism between the representing \mathbb{Z}_2^n -manifolds* M and N . It follows that, instead of studying the category $\mathbb{Z}_2^n\text{Man}$, we can just as well focus on the functor category $[\mathbb{Z}_2^n\text{Man}^{\text{op}}, \mathbf{Set}]$ (which has better properties, in particular it has all limits and colimits). A *generalized \mathbb{Z}_2^n -manifold* is an object in the functor category $[\mathbb{Z}_2^n\text{Man}^{\text{op}}, \mathbf{Set}]$ and morphisms of such objects are natural transformations. The category $[\mathbb{Z}_2^n\text{Man}^{\text{op}}, \mathbf{Set}]$ of generalised \mathbb{Z}_2^n -manifolds has finite products. Indeed, if F, G are two generalized manifolds, we define the functor $F \times G$, given on objects S , by $(F \times G)(S) = F(S) \times G(S)$, and on morphisms $\Psi : S \rightarrow T$, by

$$(F \times G)(\Psi) = F(\Psi) \times G(\Psi) : F(T) \times G(T) \rightarrow F(S) \times G(S) .$$

It is easily seen that $F \times G$ respects compositions and identities. Further, there are canonical natural transformations $\eta_1 : F \times G \rightarrow F$ and $\eta_2 : F \times G \rightarrow G$. If now (H, α_1, α_2) is another functor with natural transformations from it to F and G , respectively, it is straightforwardly checked that there exists a unique natural transformation $\beta : H \rightarrow F \times G$, such that $\alpha_i = \eta_i \circ \beta$.

One passes from the category of \mathbb{Z}_2^n -manifolds to the larger category of generalised \mathbb{Z}_2^n -manifolds in order to understand, for example, the *internal Hom* objects. In particular, there always exists a generalised \mathbb{Z}_2^n -manifold such that the so-called *adjunction formula* holds

$$\underline{\text{Hom}}_{\mathbb{Z}_2^n\text{Man}}(M, N)(-) := \underline{\text{Hom}}_{\mathbb{Z}_2^n\text{Man}}(- \times M, N) .$$

This internal Hom functor is defined on $\phi \in \underline{\text{Hom}}_{\mathbb{Z}_2^n\text{Man}}(P, S)$ by

$$\begin{aligned} \underline{\text{Hom}}_{\mathbb{Z}_2^n\text{Man}}(M, N)(\phi) : \underline{\text{Hom}}_{\mathbb{Z}_2^n\text{Man}}(M, N)(S) &\longrightarrow \underline{\text{Hom}}_{\mathbb{Z}_2^n\text{Man}}(M, N)(P) , \\ \Psi_S &\longmapsto \Psi_P \circ (\phi \times \mathbb{1}_M) . \end{aligned}$$

In general, a mapping \mathbb{Z}_2^n -manifold $\underline{\text{Hom}}_{\mathbb{Z}_2^n\text{Man}}(M, N)$ will not be representable. We will refer to ‘elements’ of a mapping \mathbb{Z}_2^n -manifold as *maps* reserving morphisms for the categorical morphisms of \mathbb{Z}_2^n -manifolds.

Composition of maps between \mathbb{Z}_2^n -manifolds is naturally defined as a natural transformation

$$(2.6) \quad \underline{\circ} : \underline{\text{Hom}}(M, N) \times \underline{\text{Hom}}(N, L) \longrightarrow \underline{\text{Hom}}(M, L) ,$$

defined, for any $S \in \mathbb{Z}_2^n\text{Man}$, by

$$(2.7) \quad \begin{aligned} \underline{\text{Hom}}(S \times M, N) \times \underline{\text{Hom}}(S \times N, L) &\longrightarrow \underline{\text{Hom}}(S \times M, L) \\ (\Psi_S, \Phi_S) &\longmapsto (\Phi \circ \Psi)_S := \Phi_S \circ (\mathbb{1}_S \times \Psi_S) \circ (\Delta \times \mathbb{1}_M) , \end{aligned}$$

where $\Delta : S \rightarrow S \times S$ is the diagonal of S and $\mathbb{1}_S : S \rightarrow S$ is its identity.

Similarly to the cases of smooth manifolds and supermanifolds, morphisms between \mathbb{Z}_2^n -manifolds are completely determined by the corresponding maps between the global functions. We remark that this is not, in general, true for complex (super)manifolds. More carefully, we have the following proposition that was proved in [14, Theorem 9].

Proposition 2.12. *Let $M = (|M|, \mathcal{O}_M)$ and $N = (|N|, \mathcal{O}_N)$ be \mathbb{Z}_2^n -manifolds. Then the natural map*

$$\underline{\text{Hom}}_{\mathbb{Z}_2^n\text{Man}}(M, N) \longrightarrow \underline{\text{Hom}}_{\mathbb{Z}_2^n\text{Alg}}(\mathcal{O}(|N|), \mathcal{O}(|M|)) ,$$

where $\mathbb{Z}_2^n\text{Alg}$ denotes the category of \mathbb{Z}_2^n -graded \mathbb{Z}_2^n -commutative associative unital \mathbb{R} -algebras, is a bijection.

The fact that the functor $\text{Hom}_{\mathbb{Z}_2^n\text{Man}}(S, -)$ respects limits and in particular products directly implies that

$$(2.8) \quad (M \times N)(S) \simeq M(S) \times N(S).$$

The latter result is essential in dealing with \mathbb{Z}_2^n -Lie groups. A (super) Lie group can be defined as a group object in the category of smooth (super)manifolds. This leads us to the following definition.

Definition 2.13. A \mathbb{Z}_2^n -Lie group is a group object in the category of \mathbb{Z}_2^n -manifolds.

A convenient fact here is that, if G is a \mathbb{Z}_2^n -Lie group, then the set $G(S)$ is a group (see (2.8)). In other words, $G(-)$ is a functor from $\mathbb{Z}_2^n\text{Man}^{\text{op}} \rightarrow \text{Grp}$.

Remark 2.14. We leave details and examples of \mathbb{Z}_2^n -Lie groups for future publications. However, we will remark at this point that the idea of “colour supergroup manifolds” has already appeared in the physics literature, albeit without a proper mathematical definition (see [1, 3, 41, 42], for example). Another approach to \mathbb{Z}_2^n -Lie groups is via a generalisation of Harish-Chandra pairs (see [39] for work in this direction).

3. \mathbb{Z}_2^n -POINTS AND THE FUNCTOR OF POINTS

In view of (2.5), we need to ‘probe’ a given \mathbb{Z}_2^n -manifold $M \simeq M(-)$ with all \mathbb{Z}_2^n -manifolds. We will show that this is however not the case, since, much like for the category of supermanifolds, we have a rather convenient generating set that we can employ, namely the set of \mathbb{Z}_2^n -points.

3.1. The category of \mathbb{Z}_2^n -points.

Definition 3.1. A \mathbb{Z}_2^n -point is a \mathbb{Z}_2^n -manifold $\mathbb{R}^{0|m}$ with vanishing ordinary dimension. We denote by $\mathbb{Z}_2^n\text{Pts}$ the full subcategory of $\mathbb{Z}_2^n\text{Man}$, whose collection of objects is the (countable) set of \mathbb{Z}_2^n -points.

Morphisms $\phi : \mathbb{R}^{0|m} \rightarrow \mathbb{R}^{0|n}$ of \mathbb{Z}_2^n -points are exactly morphisms $\phi^* : \Lambda^n \rightarrow \Lambda^m$ of \mathbb{Z}_2^n -Grassmann algebras:

Proposition 3.2. *There is an isomorphism of categories*

$$\mathbb{Z}_2^n\text{Pts} \simeq \mathbb{Z}_2^n\text{GrAlg}^{\text{op}}.$$

We can think of \mathbb{Z}_2^n -points as *formal thickenings* of an ordinary point by the non-zero degree generators. The simplest \mathbb{Z}_2^n -point is the one with trivial formal thickening, $\mathbb{R}^{0|0} := (\mathbb{R}^0, \mathbb{R})$:

Proposition 3.3. *The \mathbb{Z}_2^n -point $\mathbb{R}^{0|0} = \mathbb{R}^0$ is a terminal object in both, $\mathbb{Z}_2^n\text{Man}$ and $\mathbb{Z}_2^n\text{Pts}$.*

Proof. The unique morphism $M \rightarrow \mathbb{R}^{0|0}$ corresponds to the morphism $\mathbb{R} \ni r \cdot 1 \mapsto r \cdot \mathbb{1}_M \in \mathcal{O}_M(|M|)$, where $\mathbb{1}_M$ is the unit function. \square

Proposition 3.4. *The object set $\text{Ob}(\mathbb{Z}_2^n\text{Pts}) \simeq \text{Ob}(\mathbb{Z}_2^n\text{GrAlg})$ is a directed set.*

Proof. Given any $\underline{m} = (m_1, m_1, \dots, m_N)$ and $\underline{n} = (n_1, n_1, \dots, n_N)$, we write $\Lambda^{\underline{m}} \leq \Lambda^{\underline{n}}$ if and only if $m_i \leq n_i$, for all i . This preorder makes the non-empty set of \mathbb{Z}_2^n -Grassmann algebras into a directed set, since, any $\Lambda^{\underline{m}}$ and $\Lambda^{\underline{n}}$ admit $\Lambda^{\underline{p}}$, where $p_i = \sup\{m_i, n_i\}$, as upper bound. \square

We will need the following functional analytic result in later sections of this paper. See Definition B.1 and Definition B.5 for the notion of Fréchet space and Fréchet algebra, respectively.

Proposition 3.5. *The algebra of functions of any \mathbb{Z}_2^n -point is a \mathbb{Z}_2^n -graded \mathbb{Z}_2^n -commutative nuclear Fréchet algebra.*

The proposition is a special case of the fact that the structure sheaf of any \mathbb{Z}_2^n -manifold is a nuclear Fréchet sheaf of \mathbb{Z}_2^n -graded \mathbb{Z}_2^n -commutative algebras [13, Theorem 14].

Moreover, as a direct consequence of [14, Theorem 19, Definition 13], we observe that the category of \mathbb{Z}_2^n -points admits all finite categorical products; in particular: $\mathbb{R}^{0|m} \times \mathbb{R}^{0|n} \simeq \mathbb{R}^{0|m+n}$. By restricting attention to elements of degree 0 $\in \mathbb{Z}_2^n$, we get the following corollary. See Definition B.7 for the concept of Fréchet module.

Corollary 3.1. *The set Λ_0 of degree 0 elements of an arbitrary \mathbb{Z}_2^n -Grassmann algebra Λ is a commutative nuclear Fréchet algebra. Moreover, the algebra Λ can canonically be considered as a Fréchet Λ_0 -module.*

Remark 3.6. Specialising to the $n = 1$ case, we recover the standard and well-known facts about superpoints and their relation with Grassmann algebras.

3.2. A convenient generating set of $\mathbb{Z}_2^n\text{Man}$. It is clear that studying just the underlying topological points of a \mathbb{Z}_2^n -manifold is inadequate to probe the graded structure. Much like the category of supermanifolds, where the set of superpoints forms a generating set, the set of \mathbb{Z}_2^n -points forms a generating set for the category of \mathbb{Z}_2^n -manifolds. For the classical case of standard supermanifolds, see for example [45, Theorem 3.3.3]. For the general notion of a generating set, see Definition A.1.

Theorem 3.7. *The set $\text{Ob}(\mathbb{Z}_2^n\text{Pts})$ constitutes a generating set for $\mathbb{Z}_2^n\text{Man}$.*

Proof. Let $\phi = (|\phi|, \phi^*)$ and $\psi = (|\psi|, \psi^*)$ be two distinct \mathbb{Z}_2^n -morphisms $\phi, \psi : M \rightarrow N$ between two \mathbb{Z}_2^n -manifolds $M = (|M|, \mathcal{O}_M)$ and $N = (|N|, \mathcal{O}_N)$. These morphisms have distinct smooth base maps

$$|\phi|, |\psi| : |M| \rightarrow |N|,$$

or, if $|\phi| = |\psi|$, they have distinct pullback morphisms of sheaves of algebras

$$\phi^*, \psi^* : \mathcal{O}_N \rightarrow |\phi|_* \mathcal{O}_M.$$

If $|\phi| \neq |\psi|$, there is at least one point $m \in |M|$, such that $|\phi|(m) \neq |\psi|(m)$. Let now $s : \mathbb{R}^{0|0} \rightarrow M$ be the \mathbb{Z}_2^n -morphism, which corresponds to the $\mathbb{Z}_2^n\text{Alg}$ morphism $s^* : \mathcal{O}_M(|M|) \ni f \mapsto (\varepsilon f)(m) \in \mathbb{R}$, where ε is the sheaf morphism $\varepsilon : \mathcal{O}_M \rightarrow C_{|M|}^\infty$. It follows from the reconstruction theorem [14, Theorem 9] that the base morphism $|s| : \{\star\} \rightarrow |M|$ maps \star to m . Hence, the \mathbb{Z}_2^n -morphisms $\phi \circ s$ and $\psi \circ s$ have distinct base maps.

Assume now that $|\phi| = |\psi|$, so that there exists $|V| \subset |N|$, such that $\phi^*_{|V|} \neq \psi^*_{|V|}$, i.e., such that $\phi^*_{|V|} f \neq \psi^*_{|V|} f$, for some function $f \in \mathcal{O}_N(|V|)$. A cover of $|V|$ by coordinate patches $(\mathcal{V}_i)_i$, induces a cover $|U_i| := |\phi|^{-1}(\mathcal{V}_i)$ of $|U| := |\phi|^{-1}(|V|)$. It follows that

$$(\phi^*_{|V|} f)|_{|U_i|} \neq (\psi^*_{|V|} f)|_{|U_i|},$$

for some fixed i , i.e., that

$$\phi^*_{\mathcal{V}_i}(f|_{\mathcal{V}_i}) \neq \psi^*_{\mathcal{V}_i}(f|_{\mathcal{V}_i}),$$

so that $\phi^*_{\mathcal{V}_i} \neq \psi^*_{\mathcal{V}_i}$.

recall that, for any open subset $|X| \subset |M|$, there is a \mathbb{Z}_2^n -morphism

$$\iota_X : (|X|, \mathcal{O}_M|_{|X|}) \rightarrow (|M|, \mathcal{O}_M),$$

whose base map $|\iota_X|$ is the inclusion and whose pullback ι_X^* is the obvious restriction. Further, any \mathbb{Z}_2^n -morphism $\phi : M \rightarrow N$, whose base map $|\phi| : |M| \rightarrow |N|$ is valued in an open subset $|Y|$ of $|N|$, induces a \mathbb{Z}_2^n -morphism

$$\phi_Y : (|M|, \mathcal{O}_M) \rightarrow (|Y|, \mathcal{O}_N|_{|Y|}),$$

whose base map $|\phi_Y|$ is the map $|\phi| : |M| \rightarrow |Y|$ and whose pullback ϕ_Y^* is the pullback ϕ^* restricted to $\mathcal{O}_N|_{|Y|}$.

In view of the above, if $(\mathcal{U}_j)_j$ is a cover of $|U_i|$ by coordinate domains, we have

$$(3.1) \quad (\phi^*_{\mathcal{V}_i}(f|_{\mathcal{V}_i}))|_{\mathcal{U}_j} \neq (\psi^*_{\mathcal{V}_i}(f|_{\mathcal{V}_i}))|_{\mathcal{U}_j},$$

for some fixed j . This implies that the \mathbb{Z}_2^n -morphisms $(\phi \circ \iota_{\mathcal{U}_j})_{\mathcal{V}_i}$ and $(\psi \circ \iota_{\mathcal{U}_j})_{\mathcal{V}_i}$ from the \mathbb{Z}_2^n -domain $\mathcal{U}_j = (\mathcal{U}_j, C_{\mathcal{U}_j}^\infty[[\xi]])$ to the \mathbb{Z}_2^n -domain $\mathcal{V}_i = (\mathcal{V}_i, C_{\mathcal{V}_i}^\infty[[\theta]])$ are different. More precisely, they have the same base map $|\phi| = |\psi| : \mathcal{U}_j \rightarrow \mathcal{V}_i$, but their pullbacks are distinct. Indeed, these sheaf morphisms' algebra maps at \mathcal{V}_i are the maps $\iota_{\mathcal{U}_j, |U_i|}^* \circ \phi_{\mathcal{V}_i}^*$ and $\iota_{\mathcal{U}_j, |U_i|}^* \circ \psi_{\mathcal{V}_i}^*$ from $C_{\mathcal{V}_i}^\infty(y)[[\theta]]$ to $C_{\mathcal{U}_j}^\infty(x)[[\xi]]$, where y runs through \mathcal{V}_i and x through \mathcal{U}_j , and the values of these algebra maps at $f|_{\mathcal{V}_i}$ are different (see Equation (3.1)).

In view of Lemma 3.8, there is a \mathbb{Z}_2^n -morphism $s : \mathbb{R}^{0|m} \rightarrow \mathcal{U}_j$, such that

$$(\phi \circ \iota_{\mathcal{U}_j})_{\mathcal{V}_i} \circ s \neq (\psi \circ \iota_{\mathcal{U}_j})_{\mathcal{V}_i} \circ s.$$

However, then the \mathbb{Z}_2^n -morphism $\iota_{\mathcal{U}_j} \circ s : \mathbb{R}^{0|m} \rightarrow M$ separates ϕ and ψ , since the algebra maps at \mathcal{V}_i of the pullbacks $(s^* \circ \iota_{\mathcal{U}_j}^*) \circ \phi^*$ and $(s^* \circ \iota_{\mathcal{U}_j}^*) \circ \psi^*$ differ. Indeed, as the \mathbb{Z}_2^n -morphisms $(\phi \circ \iota_{\mathcal{U}_j})_{\mathcal{V}_i}$ and $(\psi \circ \iota_{\mathcal{U}_j})_{\mathcal{V}_i}$ are fully determined by the pullbacks of the target coordinates, their pullbacks at \mathcal{V}_i differ for at least one coordinate y^b, θ^B . It follows from the proof of Lemma 3.8 that the pullback $s_{\mathcal{U}_j}^* \circ (\iota_{\mathcal{U}_j, |U_i|}^* \circ \phi_{\mathcal{V}_i}^*)$ at \mathcal{V}_i of $(\phi \circ \iota_{\mathcal{U}_j})_{\mathcal{V}_i} \circ s$ and the similar pullback for ψ differ for the same coordinate. However, the pullback at \mathcal{V}_i considered is also the algebra map at \mathcal{V}_i of the pullback $(s^* \circ \iota_{\mathcal{U}_j}^*) \circ \phi^*$, so that the pullbacks $(s^* \circ \iota_{\mathcal{U}_j}^*) \circ \phi^*$ and $(s^* \circ \iota_{\mathcal{U}_j}^*) \circ \psi^*$ are actually distinct. \square

It remains to prove the following

Lemma 3.8. *The statement of Theorem 3.7 holds for any two distinct \mathbb{Z}_2^n -morphisms between \mathbb{Z}_2^n -domains.*

Proof. We consider two \mathbb{Z}_2^n -domains $\mathcal{U}^{p|q}$ and $\mathcal{V}^{r|s}$ together with two distinct \mathbb{Z}_2^n -morphisms

$$\mathcal{U}^{p|q} \xrightarrow[\psi]{\phi} \mathcal{V}^{r|s}.$$

As in the general case above, there are two cases to consider: either $|\phi| \neq |\psi|$, or $|\phi| = |\psi|$ and $\phi^* \neq \psi^*$. In the proof of Theorem 3.7, we showed that in the first case, the maps ϕ and ψ can be separated. In the second case, since a \mathbb{Z}_2^n -morphism valued in a \mathbb{Z}_2^n -domain is fully defined by the pullbacks of the coordinates, these global

\mathbb{Z}_2^n -functions $\phi_{\mathcal{V}^r}^*(Y^i), \psi_{\mathcal{V}^r}^*(Y^i) \in C_{\mathcal{U}^p}^\infty(x)[[\xi]]$ differ for at least one coordinate $Y^i = y^b$ or $Y^i = \theta^B$. Let B be an index, such that

$$\begin{aligned}\phi_{\mathcal{V}^r}^*(\theta^B) &= \sum_{|\alpha|=1}^{\infty} \phi_\alpha^B(x) \xi^\alpha, \\ \psi_{\mathcal{V}^r}^*(\theta^B) &= \sum_{|\alpha|=1}^{\infty} \psi_\alpha^B(x) \xi^\alpha,\end{aligned}$$

where we denoted the coordinates of $\mathcal{U}^{p|q}$ by (x^a, ξ^A) and used the standard multi-index notation, differ. This means that the functions $\phi_\alpha^B(x)$ and $\psi_\alpha^B(x)$ differ for at least one α and at least one $x \in \mathcal{U}^p$, say for $\alpha = \mathfrak{a}$ and $x = \mathfrak{x} \in \mathcal{U}^p \subset \mathbb{R}^p$. From this, we can construct the separating \mathbb{Z}_2^n -morphism

$$\mathbb{R}^{0|q} \xrightarrow{s} \mathcal{U}^{p|q} \xrightarrow[\psi]{\phi} \mathcal{V}^{r|s}.$$

Let us denote the coordinates of $\mathbb{R}^{0|q}$ by χ^A . We then define the \mathbb{Z}_2^n -morphism s by setting

$$\begin{aligned}s_{\mathcal{U}^p}^* x^a &= \mathfrak{x}^a \in \mathbb{R}[[\chi]], \deg(\mathfrak{x}^a) = \deg(x^a), \\ s_{\mathcal{U}^p}^* \xi^A &= \chi^A \in \mathbb{R}[[\chi]], \deg(\chi^A) = \deg(\xi^A).\end{aligned}$$

It is clear that $\phi \circ s \neq \psi \circ s$, since

$$\sum_{|\alpha|=1}^{\infty} \phi_\alpha^B(\mathfrak{x}) \chi^\alpha = s_{\mathcal{U}^p}^*(\phi_{\mathcal{V}^r}^*(\theta^B)) \neq s_{\mathcal{U}^p}^*(\psi_{\mathcal{V}^r}^*(\theta^B)) = \sum_{|\alpha|=1}^{\infty} \psi_\alpha^B(\mathfrak{x}) \chi^\alpha.$$

The case where $\phi_{\mathcal{V}^r}^*(Y^i) \neq \psi_{\mathcal{V}^r}^*(Y^i)$ for $Y^i = y^b$ is almost identical. In particular, we then have

$$\begin{aligned}\phi_{\mathcal{V}^r}^*(y^b) &= |\phi|^b(x) + \sum_{|\alpha|=2}^{\infty} \phi_\alpha^b(x) \xi^\alpha, \\ \psi_{\mathcal{V}^r}^*(y^b) &= |\psi|^b(x) + \sum_{|\alpha|=2}^{\infty} \psi_\alpha^b(x) \xi^\alpha.\end{aligned}$$

Since we know that $|\phi| = |\psi|$, we can proceed as for $Y^i = \theta^B$. □

In view of Proposition A.3, we get the

Corollary 3.2. *The restricted Yoneda functor*

$$\mathcal{Y}_{\mathbb{Z}_2^n \text{Pts}} : \mathbb{Z}_2^n \text{Man} \ni M \mapsto \text{Hom}_{\mathbb{Z}_2^n \text{Man}}(-, M) \in [\mathbb{Z}_2^n \text{Pts}^{\text{op}}, \text{Set}],$$

is faithful.

Above, we wrote $M(-) \in [\mathbb{Z}_2^n \text{Man}^{\text{op}}, \text{Set}]$ for the image of $M \in \mathbb{Z}_2^n \text{Man}$ by the non-restricted Yoneda functor. If no confusion arises, we will use the same notation $M(-)$ for the image $\mathcal{Y}_{\mathbb{Z}_2^n \text{Pts}}(M) \in [\mathbb{Z}_2^n \text{Pts}^{\text{op}}, \text{Set}]$ of M by the restricted Yoneda functor.

Definition 3.9. Let M be an object of $\mathbb{Z}_2^n \text{Man}$ and $\Lambda \simeq \mathbb{R}^{0|m}$ an object of $\mathbb{Z}_2^n \text{GrAlg} \simeq \mathbb{Z}_2^n \text{Pts}^{\text{op}}$. We refer to the set

$$(3.2) \quad M(\Lambda) := \text{Hom}_{\mathbb{Z}_2^n \text{Man}}(\mathbb{R}^{0|m}, M) \simeq \text{Hom}_{\mathbb{Z}_2^n \text{Alg}}(\mathcal{O}(|M|), \Lambda)$$

as the set of Λ -points of M .

Proposition 3.10. *Let*

$$m^* \in \text{Hom}_{\mathbb{Z}_2^n \text{Alg}}(\mathcal{O}(|M|), \Lambda)$$

be a Λ -point of M and let $s \in \mathcal{O}(|M|)$. The Λ -point m^* can equivalently be viewed as a \mathbb{Z}_2^n -morphism

$$m = (|m|, m^*) \in \text{Hom}_{\mathbb{Z}_2^n \text{Man}}(\mathbb{R}^{0|m}, M)$$

and therefore it defines a unique topological point $x := |m|(\star) \in |M|$. If $|U| \subset |M|$ is an open neighbourhood of x , such that $s|_{|U|} = 0$, then $m^*(s) = 0$.

Proof. We have

$$m^*(s) = m^*(s)|_{\{\star\}} = m_{\{\star\}}^*(s)|_{|m|^{-1}(|U|)} = m_{|U|}^*(s|_{|U|}) = 0.$$

□

Lemma 3.11. *There is a 1:1 correspondence*

$$M(\Lambda) \simeq \bigcup_{x \in |M|} \text{Hom}_{\mathbb{Z}_2^n \text{Alg}}(\mathcal{O}_{M,x}, \Lambda)$$

between the set of Λ -points of M and the set of morphisms from the stalks of \mathcal{O}_M to Λ . The set

$$M_x(\Lambda) := \text{Hom}_{\mathbb{Z}_2^n \text{Alg}}(\mathcal{O}_{M,x}, \Lambda)$$

is referred to as the set of Λ -points near x .

Proof. Any Λ -point m^* or $m = (|m|, m^*)$ defines a topological point $x = |m|(\star) \in |M|$, as well as a $\mathbb{Z}_2^n \text{Alg}$ -morphism $\phi_x \in \text{Hom}_{\mathbb{Z}_2^n \text{Alg}}(\mathcal{O}_{M,x}, \Lambda)$ between stalks. This morphism is given, for any $t_U \in \mathcal{O}(|U|)$ defined in some neighborhood $|U|$ of x in $|M|$, by

$$\phi_x[t_U]_x = m^*_x[t_U]_x = [m^*_{|U|}t_U]_\star = m^*_{|U|}t_U.$$

Conversely, any morphism $\psi_y \in \text{Hom}_{\mathbb{Z}_2^n \text{Alg}}(\mathcal{O}_{M,y}, \Lambda)$ ($y \in |M|$) between stalks defines a Λ -point $\mu^* \in \text{Hom}_{\mathbb{Z}_2^n \text{Alg}}(\mathcal{O}(|M|), \Lambda)$. It suffices to set

$$\mu^*t = \psi_y[t]_y \in \Lambda,$$

for all $t \in \mathcal{O}(|M|)$.

It remains to check that the composites $m^* \mapsto \phi_x \mapsto \mu^*$ and $\psi_y \mapsto \mu^* \mapsto \phi_x$ are identities. In the first case, for any $t \in \mathcal{O}(|M|)$, we get $\mu^*t = \phi_x[t]_x = m^*t$, so that $\mu^* = m^*$. In the second case, we need the following reconstruction results. Let $|U| \subset |M|$ be an open subset and set

$$S_U = \{s \in \mathcal{O}^0(|M|) : (\varepsilon s)|_{|U|} \text{ is invertible in } C^\infty(|U|)\}.$$

Then the localization map $\lambda_U : \mathcal{O}(|M|) \cdot S_U^{-1} \rightarrow \mathcal{O}(|U|)$ is an isomorphism in $\mathbb{Z}_2^n \text{Alg}$. More precisely, for any $t_U \in \mathcal{O}(|U|)$, there is a unique $Fs^{-1} \in \mathcal{O}(|M|) \cdot S_U^{-1}$, such that $t_U = F|_{|U|}s|_{|U|}^{-1}$ (if $s \in S_U$, then $s|_{|U|}$ is invertible in $\mathcal{O}(|U|)$), and we identify Fs^{-1} with t_U . For the proof of these statements or more details on them, see [14]. It is further clear from the results of [14] that $x = |\mu|(\star)$ is the topological point y .

We now compute the second composite above. For any t_U defined in a neighborhood $|U|$ of x , we get

$$\begin{aligned} \phi_x[t_U]_x &= \mu^*_{|U|}(Fs^{-1}) = \mu^*(F)\mu^*(s)^{-1} = \\ \psi_x[F]_x (\psi_x[s]_x)^{-1} &= \psi_x[F]_x \psi_x([s]_x^{-1}) = \psi_x([F]_{|U|}x[s]_{|U|}^{-1}) = \psi_x[t_U]_x, \end{aligned}$$

where the second equality is part of the reconstruction theorem of \mathbb{Z}_2^n -morphisms [14]. \square

Let us consider an open cover $(|U_I|)_{I \in \mathcal{A}}$ of the smooth manifold $|M|$, as well as the open \mathbb{Z}_2^n -submanifolds $U_I := (|U_I|, \mathcal{O}_M|_{|U_I|})$ of the \mathbb{Z}_2^n -manifold M (which need *not* be coordinate charts).

Proposition 3.12. *For any \mathbb{Z}_2^n -Grassmann algebra Λ and \mathbb{Z}_2^n -manifold $M = (|M|, \mathcal{O}_M)$, we have a natural 1:1 correspondence*

$$M(\Lambda) \simeq \bigcup_{I \in \mathcal{A}} U_I(\Lambda),$$

so that the family of sets $(U_I(\Lambda))_{I \in \mathcal{A}}$ is a cover of the set $M(\Lambda)$.

Proof. Since it is clear from the definition of a stalk that $\mathcal{O}_{U_I,x} = \mathcal{O}_{M,x}$, for any $x \in |U_I|$, it follows from Lemma 3.11 that

$$\bigcup_{I \in \mathcal{A}} U_I(\Lambda) \simeq \bigcup_{I \in \mathcal{A}} \bigcup_{x \in |U_I|} \text{Hom}_{\mathbb{Z}_2^n \text{Alg}}(\mathcal{O}_{M,x}, \Lambda) = \bigcup_{x \in |M|} \text{Hom}_{\mathbb{Z}_2^n \text{Alg}}(\mathcal{O}_{M,x}, \Lambda) \simeq M(\Lambda).$$

\square

Recall that

$$\text{Hom}_{\mathbb{Z}_2^n \text{Man}}(-, -) \in [\mathbb{Z}_2^n \text{Man}, [\mathbb{Z}_2^n \text{Pts}^{\text{op}}, \text{Set}]],$$

so that,

(i) any \mathbb{Z}_2^n -morphism $\phi = (|\phi|, \phi^*) : M \rightarrow N$ is mapped (injectively) to a natural transformation

$$\phi \simeq \text{Hom}_{\mathbb{Z}_2^n \text{Man}}(-, \phi) : \text{Hom}_{\mathbb{Z}_2^n \text{Man}}(-, M) \rightarrow \text{Hom}_{\mathbb{Z}_2^n \text{Man}}(-, N),$$

whose Λ -component ($\Lambda \simeq \mathbb{R}^{0|m}$) is the **Set**-map given by

$$(3.3) \quad \phi_\Lambda := \text{Hom}_{\mathbb{Z}_2^n \text{Man}}(\Lambda, \phi) : M(\Lambda) = \text{Hom}_{\mathbb{Z}_2^n \text{Man}}(\mathbb{R}^{0|m}, M) \simeq \text{Hom}_{\mathbb{Z}_2^n \text{Alg}}(\mathcal{O}(|M|), \Lambda) \ni m^* \mapsto$$

$$m^* \circ \phi^* \in \text{Hom}_{\mathbb{Z}_2^n \text{Alg}}(\mathcal{O}(|N|), \Lambda) \simeq \text{Hom}_{\mathbb{Z}_2^n \text{Man}}(\mathbb{R}^{0|m}, N) = N(\Lambda), \text{ and ,}$$

(ii) for any fixed $M \in \mathbb{Z}_2^n\text{Man}$, given a morphism $\psi = (|\psi|, \psi^*) : \mathbb{R}^{0|m'} \rightarrow \mathbb{R}^{0|m}$ of \mathbb{Z}_2^n -points, or, equivalently, a morphism $\psi^* : \Lambda \rightarrow \Lambda'$ of \mathbb{Z}_2^n -Grassmann algebras, we get the induced **Set**-map

$$(3.4) \quad M(\psi^*) := \text{Hom}_{\mathbb{Z}_2^n\text{Man}}(\psi, M) : M(\Lambda) = \text{Hom}_{\mathbb{Z}_2^n\text{Man}}(\mathbb{R}^{0|m}, M) \simeq \text{Hom}_{\mathbb{Z}_2^n\text{Alg}}(\mathcal{O}(|M|), \Lambda) \ni m^* \mapsto \psi^* \circ m^* \in \text{Hom}_{\mathbb{Z}_2^n\text{Alg}}(\mathcal{O}(|M|), \Lambda') \simeq \text{Hom}_{\mathbb{Z}_2^n\text{Man}}(\mathbb{R}^{0|m'}, M) = M(\Lambda') .$$

When reading the maps ϕ_Λ and $M(\psi^*)$ through the 1:1 correspondence

$$M(\Lambda) \ni m^* \mapsto (x, m_\star^*) \in \bigcup_{y \in |M|} \text{Hom}_{\mathbb{Z}_2^n\text{Alg}}(\mathcal{O}_{M,y}, \Lambda) ,$$

where $x = |m|(\star)$, we obtain

$$(3.5) \quad \begin{aligned} \phi_\Lambda : M(\Lambda) &\longrightarrow N(\Lambda) \\ (x, m_\star^*) &\mapsto (|\phi|(x), m_\star^* \circ \phi_x^*) , \text{ and} , \end{aligned}$$

$$(3.6) \quad \begin{aligned} M(\psi^*) : M(\Lambda) &\longrightarrow M(\Lambda') \\ (x, m_\star^*) &\mapsto (x, \psi^* \circ m_\star^*) . \end{aligned}$$

3.3. Restricted Yoneda functor and fullness. The Yoneda functor from any locally small category \mathbf{C} into the category of **Set**-valued contravariant functors on \mathbf{C} , is fully faithful. This holds in particular for $\mathbf{C} = \mathbb{Z}_2^n\text{Man}$. When we restrict the contravariant functors to the generating set $\mathbb{Z}_2^n\text{Pts}$, the resulting restricted Yoneda functor is automatically faithful. In the following, we show that it is not full, i.e., that not all natural transformations are induced by a \mathbb{Z}_2^n -morphism.

Naturality of any transformation $\phi : M(-) \rightarrow N(-)$ between **Set**-valued contravariant (resp., covariant) functors on $\mathbb{Z}_2^n\text{Pts}$ (resp., $\mathbb{Z}_2^n\text{GrAlg}$), means that the diagram

$$(3.7) \quad \begin{array}{ccc} M(\Lambda) & \xrightarrow{\phi_\Lambda} & N(\Lambda) \\ M(\psi^*) \downarrow & & \downarrow N(\psi^*) \\ M(\Lambda') & \xrightarrow{\phi_{\Lambda'}} & N(\Lambda') \end{array}$$

commutes, for any morphism $\psi^* : \Lambda \rightarrow \Lambda'$ of \mathbb{Z}_2^n -Grassmann algebras.

A Λ -point of a \mathbb{Z}_2^n -manifold M is denoted by m^* or $m = (|m|, m^*)$. If the manifold is a \mathbb{Z}_2^n -domain $\mathcal{U}^{p|q}$, we use the notation x^* or $x = (|x|, x^*)$. If (x^a, ξ^A) are the coordinates of $\mathcal{U}^{p|q}$, a Λ -point x^* in $\mathcal{U}^{p|q}$ is completely determined by the degree-respecting pullbacks

$$(x_\Lambda^a, \xi_\Lambda^A) := (x^*(x^a), x^*(\xi^A)) .$$

Since $x_\Lambda^a \in \Lambda_0 = \mathbb{R} \oplus \mathring{\Lambda}_0$, we write $x_\Lambda^a = (x_{||}^a, \dot{x}_\Lambda^a)$. Hence, any Λ -point x^* in $\mathcal{U}^{p|q}$ can be identified with

$$(3.8) \quad x^* \simeq (x_\Lambda^a, \xi_\Lambda^A) = (x_{||}^a, \dot{x}_\Lambda^a, \xi_\Lambda^A) \in \mathbb{R}^p \times \mathring{\Lambda}_0^p \times \mathring{\Lambda}_{\gamma_1}^{q_1} \times \cdots \times \mathring{\Lambda}_{\gamma_N}^{q_N} ,$$

where

$$x_{||} = (x_{||}^a) = (..., x_{||}^a, ...) \in \mathcal{U}^p ,$$

and where $\gamma_1, \dots, \gamma_N$ denote the non-zero \mathbb{Z}_2^n -degrees in standard order. Here the \dot{x}_Λ^a (resp., the ξ_Λ^A) are formal power series containing at least 2 (resp., at least 1) of the generators (θ^C) of the \mathbb{Z}_2^n -Grassmann algebra Λ .

As mentioned above, any \mathbb{Z}_2^n -morphism, in particular any morphism $\phi : \mathcal{U}^{p|q} \rightarrow \mathcal{V}^{r|s}$ between \mathbb{Z}_2^n -domains, naturally induces a natural transformation, with Λ -component

$$\phi_\Lambda : \mathcal{U}^{p|q}(\Lambda) \ni x^* \mapsto x^* \circ \phi^* \in \mathcal{V}^{r|s}(\Lambda) .$$

If (y^b, η^B) are the coordinates of $\mathcal{V}^{r|s}$, the morphism ϕ reads

$$(3.9a) \quad \phi^*(y^b) = \sum_{|\alpha| \geq 0} \phi_\alpha^b(x) \xi^\alpha ,$$

$$(3.9b) \quad \phi^*(\eta^B) = \sum_{|\alpha| > 0} \phi_\alpha^B(x) \xi^\alpha ,$$

where the right-hand sides have the appropriate degrees and where $\phi_0(\mathcal{U}^p) \subset \mathcal{V}^r$. Further, the image Λ -point $x^* \circ \phi^*$ in $\mathcal{V}^{r|\underline{s}}$ by ϕ_Λ of the Λ -point $x^* \simeq (x^*(x^a); x^*(\xi^A)) = (x_{||}^a, \dot{x}_\Lambda^a; \xi_\Lambda^A)$ in $\mathcal{U}^{p|\underline{q}}$, is given by

$$(3.10a) \quad y_\Lambda^b = \sum_{|\alpha| \geq 0} \sum_{|\beta| \geq 0} \frac{1}{\beta!} (\partial_x^\beta \phi_\alpha^b)(x_{||}) \dot{x}_\Lambda^\beta \xi_\Lambda^\alpha,$$

$$(3.10b) \quad \eta_\Lambda^B = \sum_{|\alpha| > 0} \sum_{|\beta| \geq 0} \frac{1}{\beta!} (\partial_x^\beta \phi_\alpha^B)(x_{||}) \dot{x}_\Lambda^\beta \xi_\Lambda^\alpha.$$

Let us recall that there is no convergence issue with terms in $x_{||}$ [16]. Thus the components of a natural transformation implemented by a \mathbb{Z}_2^n -morphism between \mathbb{Z}_2^n -domains, are very particular formal power series in the formal variables \dot{x}_Λ^a and ξ_Λ^A , which are themselves formal power series in the generators (θ^C) of Λ .

We are now able to prove that not all natural transformations between the restricted functors $M(-), N(-) \in [\mathbb{Z}_2^n \text{Pts}, \text{Set}]$ associated with $M, N \in \mathbb{Z}_2^n \text{Man}$, arise from a \mathbb{Z}_2^n -morphism $M \rightarrow N$. Since it suffices to give one counter-example, we choose $M = N = \mathbb{R}^{p|\underline{0}} = \mathbb{R}^p$.

Example 3.13. Consider an arbitrary diffeomorphism $\phi : \mathbb{R}^p \rightarrow \mathbb{R}^p$. The Λ -component of the associated natural transformation is

$$\begin{aligned} \phi_\Lambda : \mathbb{R}^{p|\underline{0}}(\Lambda) &\longrightarrow \mathbb{R}^{p|\underline{0}}(\Lambda) \\ (x_\Lambda^b, 0) &\mapsto \left(\phi^b(x_{||}) + \sum_{|\beta| > 0} \frac{1}{\beta!} (\partial_x^\beta \phi^b)(x_{||}) \dot{x}_\Lambda^\beta, 0 \right). \end{aligned}$$

From this data we obtain another natural transformation

$$\begin{aligned} \alpha_\Lambda : \mathbb{R}^{p|\underline{0}}(\Lambda) &\longrightarrow \mathbb{R}^{p|\underline{0}}(\Lambda) \\ (x_\Lambda^b, 0) &\mapsto (\phi^b(x_{||}), 0). \end{aligned}$$

The natural transformation α is not implemented by a morphism $\psi : \mathbb{R}^p \rightarrow \mathbb{R}^p$. Indeed, otherwise $\alpha_\Lambda = \psi_\Lambda$, for all Λ . This means that

$$(\phi^b(x_{||}), 0) = \left(\psi^b(x_{||}) + \sum_{|\beta| > 0} \frac{1}{\beta!} (\partial_x^\beta \psi^b)(x_{||}) \dot{x}_\Lambda^\beta, 0 \right),$$

for all Λ and all Λ -points. Since $\phi^b(x) \equiv \psi^b(x)$, we have $\partial_x^\beta \phi^b \equiv \partial_x^\beta \psi^b$. Take now any $\beta : |\beta| = 1$, so that $\beta_a = 1$, for some fixed $a \in \{1, \dots, p\}$. As we can choose Λ and x_Λ^b , for all $b \in \{1, \dots, p\}$, arbitrarily, we can choose $\dot{x}_\Lambda^b = 0$, for all $b \neq a$, and $\dot{x}_\Lambda^a = \theta^D \theta^E$, where θ^D and θ^E are two different generators of Λ that have the same degree. The coefficient of $\theta^D \theta^E$ in the sum over all β is then $(\partial_{x^a} \psi^b)(x_{||})$, hence $\partial_{x^a} \phi^b \equiv \partial_{x^a} \psi^b \equiv 0$. The latter observation is a contradiction, since the Jacobian determinant of ϕ does not vanish anywhere in \mathbb{R}^p .

We now generalise a technical result [58, Theorem 1] to \mathbb{Z}_2^n -domains $\mathcal{U}^{p|\underline{q}}$. Let

$$\mathcal{B}_{p|\underline{q}}(\mathcal{U}^p) := \mathcal{F}(\mathcal{U}^p, \mathbb{R})[[X, \Xi]],$$

be the \mathbb{Z}_2^n -graded \mathbb{Z}_2^n -commutative associative unital \mathbb{R} -algebra of formal power series in p parameters X^a of \mathbb{Z}_2^n -degree 0 and q_1, \dots, q_N parameters Ξ^A of non-zero \mathbb{Z}_2^n -degree $\gamma_1, \dots, \gamma_N$, and with coefficients in arbitrary \mathbb{R} -valued functions on \mathcal{U}^p , i.e., we do *not* ask that these functions be continuous let alone smooth. Following [49, 54, 58], we will refer to this algebra as a \mathbb{Z}_2^n -Berezin algebra. Any element of this algebra is of the form

$$(3.11) \quad F = \sum_{|\alpha| \geq 0} \sum_{|\beta| \geq 0} F_{\alpha\beta}(x) X^\beta \Xi^\alpha,$$

where the x^a are coordinates in \mathcal{U}^p .

Theorem 3.14. *For any \mathbb{Z}_2^n -domains $\mathcal{U}^{p|\underline{q}}$ and $\mathcal{V}^{r|\underline{s}}$, there is a 1:1 correspondence*

$$\text{Nat}(\mathcal{U}^{p|\underline{q}}, \mathcal{V}^{r|\underline{s}}) \rightarrow (\mathcal{B}_{p|\underline{q}}(\mathcal{U}^p))^{r|\underline{s}}$$

between

- the set of natural transformations in $[\mathbb{Z}_2^n \text{Pts}^{\text{op}}, \text{Set}]$ between $\mathcal{U}^{p|\underline{q}}(-)$ and $\mathcal{V}^{r|\underline{s}}(-)$, and
- the set of ‘vectors’ \mathbf{F} with r (resp., with s_1, \dots, s_N) components F^b of degree 0 (resp., components F^B of degrees $\gamma_1, \dots, \gamma_N$) of the type (3.11), such that the r -tuple (F_{00}^b) made of the coefficients $F_{00}^b(x)$ of the r series F^b satisfies

$$(F_{00}^b)(\mathcal{U}^p) \subset \mathcal{V}^r.$$

Proof. Let \mathbf{F} be such a ‘vector’. For any Λ , we define the map

$$\beta_\Lambda : \mathcal{U}^{p|q}(\Lambda) \ni (x_{||}^a, \dot{x}_\Lambda^a, \xi_\Lambda^A) \mapsto (y_\Lambda^b, \eta_\Lambda^B) \in \mathcal{V}^{r|s}(\Lambda) ,$$

where

$$(3.12) \quad y_\Lambda^b := \sum_{|\alpha| \geq 0} \sum_{|\beta| \geq 0} F_{\alpha\beta}^b(x_{||}) \dot{x}_\Lambda^\beta \xi_\Lambda^\alpha \quad \text{and} \quad \eta_\Lambda^B := \sum_{|\alpha| \geq 0} \sum_{|\beta| \geq 0} F_{\alpha\beta}^B(x_{||}) \dot{x}_\Lambda^\beta \xi_\Lambda^\alpha .$$

Since $\dot{x}_\Lambda^a, \xi_\Lambda^A$ have the same degrees as X^a, Ξ^A , the right-hand sides of (3.12) have the same degrees as F^b, F^B , hence, $y_\Lambda^b, \eta_\Lambda^B$ have the degrees required to be a Λ -point in $\mathcal{V}^{r|s}$. Moreover, we have

$$y_{||}^b = F_{00}^b(x_{||}) ,$$

so that $y_{||} \in \mathcal{V}^r$. The target of the map β_Λ is thus actually $\mathcal{V}^{r|s}(\Lambda)$. The naturalness of β under morphisms of \mathbb{Z}_2^n -Grassmann algebras is obvious: β is a natural transformation in $[\mathbb{Z}_2^n \text{Pts}^{\text{op}}, \text{Set}]$ between $\mathcal{U}^{p|q}(-)$ and $\mathcal{V}^{r|s}(-)$. Finally, we defined a map

$$\mathcal{I} : (\mathcal{B}_{p|q}(\mathcal{U}^p))^{r|s} \rightarrow \text{Nat}(\mathcal{U}^{p|q}, \mathcal{V}^{r|s}) .$$

We will explain now that any natural transformation $\beta : \mathcal{U}^{p|q}(-) \rightarrow \mathcal{V}^{r|s}(-)$ is the image by \mathcal{I} of a unique ‘vector’ \mathbf{F} . We first show that, for any $\Lambda \simeq \mathbb{R}^{0|m}$, the image $\beta_\Lambda(x^*) \in \mathcal{V}^{r|s}(\Lambda)$ of any Λ -point

$$x^* \simeq (x_{||}^a, \dot{x}_\Lambda^a, \xi_\Lambda^A) \in \mathcal{U}^p \times \mathring{\Lambda}_0^p \times \mathring{\Lambda}_{\gamma_1}^{q_1} \times \cdots \times \mathring{\Lambda}_{\gamma_N}^{q_N}$$

in $\mathcal{U}^{p|q}$, has components y_Λ^b and η_Λ^B of the type (3.12).

Step 1. We prove that any Λ -point in $\mathcal{U}^{p|q}$ is the image by a \mathbb{Z}_2^n -Grassmann algebra map $\varphi^* : \Lambda' \rightarrow \Lambda$ of a Λ' -point in $\mathcal{U}^{p|q}$, some of whose defining series are series in formal pairings.

Let (θ^C) be the generators of Λ . The Λ -point x^* then reads

$$x^* \simeq (x_{||}^a, \theta^\lambda \theta^\kappa K_{\kappa\lambda}^a, \xi_\Lambda^A) ,$$

where the degree of $K_{\kappa\lambda}^a \in \Lambda$ is the sum of the degrees of θ^λ and θ^κ . Recall that a (resp., A) runs through $\{1, \dots, p\}$ (resp., through $\{1, \dots, |\underline{q}|\}$), and that λ, κ run through $\{1, \dots, |\underline{m}|\}$. Consider now the set S of generators

$$\theta' = (\eta^{a\lambda}, \zeta_\lambda^b, \psi^A) ,$$

where b has the same range as a , and define their (non-zero) \mathbb{Z}_2^n -degrees by

$$\deg(\eta^{a\lambda}) = \deg(\theta^\lambda), \quad \deg(\zeta_\lambda^b) = \deg(\theta^\kappa), \quad \deg(\psi^A) = \deg(\xi_\Lambda^A) = \deg(\xi^A) .$$

Let Λ' be the \mathbb{Z}_2^n -Grassmann algebra defined by S , and set

$$(3.13) \quad x'^* \simeq (x_{||}^a, \eta^{a\lambda} \zeta_\lambda^a, \psi^A) \in \mathcal{U}^p \times \mathring{\Lambda}_0'^p \times \mathring{\Lambda}_{\gamma_1}'^{q_1} \times \cdots \times \mathring{\Lambda}_{\gamma_N}'^{q_N}$$

(no sum over a in the formal pairings $\eta^{a\lambda} \zeta_\lambda^a$). The degree-respecting equalities

$$\varphi^*(\eta^{a\lambda}) = \theta^\lambda, \quad \varphi^*(\zeta_\lambda^b) = \theta^\lambda K_{\lambda\kappa}^b, \quad \varphi^*(\psi^A) = \xi_\Lambda^A$$

define a morphism of \mathbb{Z}_2^n -Grassmann algebras $\varphi^* : \Lambda' \rightarrow \Lambda$. It suffices to set

$$\varphi^* \left(\sum_\varepsilon r_\varepsilon \theta^\varepsilon \right) := \sum_\varepsilon r_\varepsilon (\varphi^* \theta')^\varepsilon .$$

Indeed, any term of the right-hand side is a series in θ whose terms contain at least $|\varepsilon|$ generators. Hence, for any ε , only the terms $|\varepsilon| \leq |\varepsilon|$ can contribute to θ^ε , and therefore there is no convergence issue with the coefficient of θ^ε . Since the Λ -point $\varphi^* \circ x'^*$ in $\mathcal{U}^{p|q}$ reads

$$\varphi^* \circ x'^* \simeq \varphi^*(x_{||}^a, \eta^{a\lambda} \zeta_\lambda^a, \psi^A) = (x_{||}^a, \theta^\lambda \theta^\kappa K_{\kappa\lambda}^a, \xi_\Lambda^A) \simeq x^* ,$$

naturality of the transformation $\beta : \mathcal{U}^{p|q}(-) \rightarrow \mathcal{V}^{r|s}(-)$ implies that

$$(3.14) \quad (y_\Lambda^b, \eta_\Lambda^B) := \beta_\Lambda(x^*) = \beta_\Lambda(\varphi^* \circ x'^*) = \beta_\Lambda(\mathcal{U}^{p|q}(\varphi^*)(x'^*)) = \mathcal{V}^{r|s}(\varphi^*)(\beta_{\Lambda'}(x'^*)) = \varphi^* \circ (\beta_{\Lambda'}(x'^*)) \simeq \varphi^*(y_{\Lambda'}^b, \eta_{\Lambda'}^B) ,$$

where $y_{\Lambda'}^b$ and $\eta_{\Lambda'}^B$ are series in the generators of Λ' .

Step 2. We define formal rotations under which the formal pairings are invariant. Moreover, we show that any formal series that is invariant under the formal rotations is a series in the formal pairings.

The formal part of each degree 0 component of x'^* can be viewed as a formal pairing $\eta^a \cdot \zeta^a = \eta^{a\lambda} \zeta_\lambda^a$, which is stable under formal rotations R^* . More precisely, we set

$$R^*(\eta^{a\lambda}) = \eta^{a\kappa} (O^a)_\kappa^\lambda, \quad R^*(\zeta_\lambda^b) = (O^{b\kappa})_\kappa^\lambda \zeta_\lambda^b, \quad R^*(\psi^A) = \psi^A ,$$

where O^a and $O^{b\mathbf{t}}$ are any $(m_1 + \dots + m_N) \times (m_1 + \dots + m_N)$ block-diagonal matrices with entries in \mathbb{R} that satisfy

$$(3.15) \quad (O^a)_\rho^\lambda (O^{a\mathbf{t}})_\lambda^\omega = \delta_\rho^\omega.$$

Since, for any fixed a (resp., b), the components $\eta^{a\lambda}$ (resp., ζ_κ^b) are ordered such that the m_1 first components have degree γ_1 , the next m_2 degree γ_2 , and so on, these equalities are degree-preserving. Hence, they define a \mathbb{Z}_2^n -Grassmann algebra morphism $R^* : \Lambda' \rightarrow \Lambda'$ via

$$R^* \left(\sum_\epsilon r_\epsilon \theta'^\epsilon \right) = R^* \left(\sum_{\alpha\beta\gamma} r_{\alpha\beta\gamma} \eta^\alpha \zeta^\beta \psi^\gamma \right) := \sum_{\alpha\beta\gamma} r_{\alpha\beta\gamma} (R^* \eta)^\alpha (R^* \zeta)^\beta \psi^\gamma.$$

Since the images $R^*(\eta^{a\lambda})$ (resp., $R^*(\zeta_\kappa^b)$) are linear in the $\eta^{a\kappa}$ (resp., ζ_λ^b) (of the same degree), the term indexed by $\alpha\beta\gamma$ is a homogeneous polynomial of order $|\alpha| + |\beta| + |\gamma|$ in the generators θ' . Hence, for any ϵ , only the terms $|\alpha| + |\beta| + |\gamma| = |\epsilon|$ can contribute to θ'^ϵ , so that no convergence problems arise. In view of (3.15), it is clear that, as mentioned above, the formal pairing $\eta^a \cdot \zeta^a = \eta^{a\lambda} \zeta_\lambda^a$ is invariant under R^* . As any \mathbb{Z}_2^n -Grassmann algebra morphism, the formal rotation R^* induces maps $\mathcal{U}^{p|q}(R^*)$ and $\mathcal{V}^{r|s}(R^*)$, and due to naturality of β , we find

$$\mathcal{V}^{r|s}(R^*)(\beta_{\Lambda'} \mathbf{x}'^*) = \beta_{\Lambda'} (\mathcal{U}^{p|q}(R^*)(\mathbf{x}'^*)) = \beta_{\Lambda'} (R^* \circ \mathbf{x}'^*) \simeq \beta_{\Lambda'} (R^*(x_{||}^a, \eta^{a\lambda} \zeta_\lambda^a, \psi^A)) \simeq \beta_{\Lambda'} \mathbf{x}'^*,$$

so that $\beta_{\Lambda'} \mathbf{x}'^*$ is invariant under rotations.

We are now prepared to continue the computation (3.14). Since

$$(3.16) \quad \beta_{\Lambda'}(\mathbf{x}'^*) \simeq (y_{\Lambda'}^b, \eta_{\Lambda'}^B) = (y_{||}^b, \dot{y}_{\Lambda'}^b, \eta_{\Lambda'}^B)$$

is invariant under the rotations R^* , the series $\dot{y}_{\Lambda'}^b, \eta_{\Lambda'}^B$ in the generators θ' are invariant. More explicitly, for each series, we have an equality of the type

$$\sum_\gamma \left(\sum_{k,\ell} \sum_{|\alpha|=k, |\beta|=\ell} F_{\alpha\beta\gamma} \eta^\alpha \zeta^\beta \right) \psi^\gamma = \sum_\gamma \left(\sum_{k,\ell} \sum_{|\alpha|=k, |\beta|=\ell} F_{\alpha\beta\gamma} (R^* \eta)^\alpha (R^* \zeta)^\beta \right) \psi^\gamma,$$

which is equivalent to

$$\begin{aligned} \sum_{|\alpha|=k, |\beta|=\ell} F_{\alpha\beta\gamma} \dots \eta^{a\lambda} \eta^{b\mu} \zeta_\nu^c \dots &= \sum_{|\alpha|=k, |\beta|=\ell} F_{\alpha\beta\gamma} \eta^\alpha \zeta^\beta = \\ \sum_{|\alpha|=k, |\beta|=\ell} F_{\alpha\beta\gamma} (R^* \eta)^\alpha (R^* \zeta)^\beta &= \sum_{|\alpha|=k, |\beta|=\ell} F_{\alpha\beta\gamma} \dots \eta^{a\delta} (O^a)_\delta^\lambda \eta^{b\delta'} (O^b)_{\delta'}^\mu (O^{c\mathbf{t}})_\nu^{\delta''} \zeta_\delta^c \dots, \end{aligned}$$

and holds for all (!) formal rotations. This is only possible, if the power series considered, i.e., the series $\dot{y}_{\Lambda'}^b$ and $\eta_{\Lambda'}^B$, are series in pairings $\eta^a \cdot \zeta^a = \eta^{a\lambda} \zeta_\lambda^a$ [58]. In view of (3.14), we thus get

$$(y_{||}^b, \dot{y}_{\Lambda'}^b, \eta_{\Lambda'}^B) = \beta_{\Lambda'}(\mathbf{x}'^*) = \beta_{\Lambda'}(x_{||}^a, \dot{x}_\Lambda^a, \xi_\Lambda^A) = (y_{||}^b, \varphi^*(\dot{y}_{\Lambda'}^b), \varphi^*(\eta_{\Lambda'}^B)),$$

where any image by φ^* is of the type

$$\sum_{(\alpha,\beta) \neq (0,0)} F_{\alpha\beta} \varphi^*((\eta \cdot \zeta)^\beta) \varphi^*(\psi^\alpha) = \sum_{(\alpha,\beta) \neq (0,0)} F_{\alpha\beta} \dot{x}_\Lambda^\beta \xi_\Lambda^\alpha.$$

It is clear from (3.16) and (3.13) that the coefficients

$$F_{\alpha\beta}^b, F_{\alpha\beta}^B \ ((\alpha, \beta) \neq (0, 0)), \text{ and } F_{00}^b := y_{||}^b$$

depend (only) on $x_{||} \in \mathcal{U}^p$. Hence, the image

$$(y_{\Lambda'}^b, \eta_{\Lambda'}^B) = \beta_{\Lambda'}(\mathbf{x}'^*) = (F^b(x_{||}, \dot{x}_\Lambda, \xi_\Lambda), F^B(x_{||}, \dot{x}_\Lambda, \xi_\Lambda))$$

is actually of the type (3.12). Since $\beta_{\Lambda'}(\mathbf{x}'^*)$ is a Λ -point in $\mathcal{V}^{r|s}$, the r series $F^b(x_{||}, \dot{x}_\Lambda, \xi_\Lambda)$ and the s_i series $F^B(x_{||}, \dot{x}_\Lambda, \xi_\Lambda)$ are of degree 0 and degree γ_i , respectively, i.e., the r series $F^b(x, X, \Xi)$ and the s_i series $F^B(x, X, \Xi)$ are of degree 0 and degree γ_i , respectively. For the same reason, we have $F_{00}(x_{||}) \in \mathcal{V}^r$, for all $x_{||} \in \mathcal{U}^p$, so that we constructed a ‘vector’ $\mathbf{F} \in (\mathcal{B}_{p|q}(\mathcal{U}^p))^{r|s}$, whose image by \mathcal{I} is obviously β .

Step 3. We show that \mathbf{F} is unique (which concludes the proof). If there is another ‘vector’ \mathbf{F}' , such that $\mathcal{I}(\mathbf{F}') = \beta$, we have

$$(3.17) \quad \sum_{|\alpha| \geq 0, |\beta| \geq 0} F_{\alpha\beta}^b(x_{||}) \dot{x}_\Lambda^\beta \xi_\Lambda^\alpha = \sum_{|\alpha| \geq 0, |\beta| \geq 0} F'_{\alpha\beta}^b(x_{||}) \dot{x}_\Lambda^\beta \xi_\Lambda^\alpha,$$

for all $b \in \{b, B\}$, all Λ , and all \mathbf{x}'^* . Notice first that any \dot{x}_Λ^a (resp., any ξ_Λ^A) is a series of degree 0 (resp., of degree $\deg(\xi_\Lambda^A) = \gamma_A$) in the θ -s that contains at least two parameters $\theta^C \theta^{C'}$ (resp., at least one parameter $\theta^{C''}$). Hence, both sides are series in θ , and the left-hand side and right-hand side coefficients of any monomial θ^ϵ coincide. A term $(\alpha, \beta) \neq (0, 0)$ cannot contribute to the independent term θ^0 . Hence $F_{00}^b(x_{||}) = F'_{00}^b(x_{||})$. We now show that $F_{\alpha\beta}^b(x_{||}) = F'_{\alpha\beta}^b(x_{||})$, for an arbitrarily fixed $(\alpha, \beta) \neq (0, 0)$. Since Λ is arbitrary, we can choose as many different generators θ in each non-zero degree as necessary, and, since \mathbf{x}'^* is arbitrary, we can

choose $x_{||}$ arbitrarily in \mathcal{U}^p and we can choose the coefficients of the series \dot{x}_Λ^a and ξ_Λ^A arbitrarily (except that we have to observe that the coefficient of a monomial θ^ε , which does not have the required degree, *must be zero*). Let now $\alpha_1, \dots, \alpha_\mu$ and $\beta_1, \dots, \beta_\nu$ be the non-zero components in the fixed α and β . For each factor $\xi_\Lambda^{A_i}$ of

$$\xi_\Lambda^\alpha = (\xi_\Lambda^{A_1})^{\alpha_1} \dots (\xi_\Lambda^{A_\mu})^{\alpha_\mu},$$

we choose a monomial in one generator θ^{C_i} of degree γ_{A_i} , set its coefficient r_{C_i} to 1, and all the other coefficients in the series $\xi_\Lambda^{A_i}$ to zero. Further, for different $\xi_\Lambda^{A_i}$, we choose different generators θ^{C_i} . Similarly, for each factor $\dot{x}_\Lambda^{a_j}$ of

$$\dot{x}_\Lambda^\beta = (\dot{x}_\Lambda^{a_1})^{\beta_1} \dots (\dot{x}_\Lambda^{a_\nu})^{\beta_\nu},$$

we choose monomials $\theta^{D_{jk}} \theta^{E_{jk}}$ ($k \in \{1, \dots, \beta_j\}$) in two generators of the same odd degree (for all $n \geq 1$, there is at least one odd degree), set their coefficient $r_{D_{jk} E_{jk}}$ to 1, and all the other coefficients in the series $\dot{x}_\Lambda^{a_j}$ to zero. Further, we choose the generators so that all generators θ^{C_i} , $\theta^{D_{jk}}$, and $\theta^{E_{jk}}$ are different. When setting

$$\theta^\omega = \prod_{j=1}^{\nu} \theta^{D_{j1}} \theta^{E_{j1}} \dots \theta^{D_{j\beta_j}} \theta^{E_{j\beta_j}} \prod_{i=1}^{\mu} (\theta^{C_i})^{\alpha_i} \neq 0,$$

the terms indexed by (the fixed) (α, β) in both sides of (3.17), read

$$\beta! F_{\alpha\beta}^b(x_{||}) \theta^\omega \text{ and } \beta! F_{\alpha\beta}'^b(x_{||}) \theta^\omega.$$

For any term $(\alpha', \beta') \neq (\alpha, \beta)$, we either get a new series ξ_Λ^A or \dot{x}_Λ^a , or we get an old series a different number of times. In the second case, the term (α', β') does not contribute to the coefficient of θ^ω ; in the first, we set all the coefficients of the new series to 0, so that the term (α', β') vanishes. Finally, we obtain $F_{\alpha\beta}^b(x_{||}) = F_{\alpha\beta}'^b(x_{||})$, for any $x_{||} \in \mathcal{U}^p$. \square

We now show that $\mathbb{R}^{p|q}(\Lambda)$ is a Fréchet space and that $\mathcal{U}^{p|q}(\Lambda)$ is an open subset of $\mathbb{R}^{p|q}(\Lambda)$. This means that we have a notion of directional derivative, as well as a notion of smoothness of continuous maps between the Λ -points of \mathbb{Z}_2^n -domains. For more details on Fréchet objects, we refer the reader to Appendix B.

Proposition 3.15. *For any $\Lambda \in \mathbb{Z}_2^n \text{GrAlg}$, the set $\mathbb{R}^{p|q}(\Lambda)$ is a nuclear Fréchet space and a Fréchet Λ_0 -module. Moreover, the set $\mathcal{U}^{p|q}(\Lambda)$ is an open subset of $\mathbb{R}^{p|q}(\Lambda)$.*

Proof. Let $\Lambda \in \mathbb{Z}_2^n \text{GrAlg}$. As explained above, there is a 1:1 correspondence between the Λ -points x^* of $\mathbb{R}^{p|q}$ (resp., of $\mathcal{U}^{p|q}$) and the $(p + |q|)$ -tuples

$$x^* \simeq (x_\Lambda^a, \xi_\Lambda^A) \in \Lambda_0^p \times \Lambda_{\gamma_1}^{q_1} \times \dots \times \Lambda_{\gamma_N}^{q_N}$$

(resp., the same $(p + |q|)$ -tuples, but with the additional requirement that the p -tuple $(x_{||}^a)$ made of the independent terms of (x_Λ^a) be a point in $\mathcal{U}^p \subset \mathbb{R}^p$). Note now that Λ is the \mathbb{Z}_2^n -graded \mathbb{Z}_2^n -commutative nuclear Fréchet \mathbb{R} -algebra of global \mathbb{Z}_2^n -functions of some $\mathbb{R}^{0|m}$. Hence, all its homogeneous subspaces Λ_{γ_i} ($i \in 0, \dots, N$, $\gamma_0 = 0$) are nuclear Fréchet vector spaces. Since any product (resp., any countable product) of nuclear (resp., Fréchet) spaces is nuclear (resp., Fréchet), the set $\mathbb{R}^{p|q}(\Lambda)$ of Λ -points of $\mathbb{R}^{p|q}$ is nuclear Fréchet. The latter statements can be found in [13].

As for the second claim in Proposition 3.15, recall that Λ_0 is a (commutative) Fréchet algebra, see Corollary 3.1. The Fréchet Λ_0 -module structure on $\mathbb{R}^{p|q}(\Lambda)$ is then defined by

$$(3.18) \quad m : \Lambda_0 \times \mathbb{R}^{p|q}(\Lambda) \ni (a, x^*) \mapsto (a \cdot x_\Lambda^a, a \cdot \xi_\Lambda^A) \in \mathbb{R}^{p|q}(\Lambda).$$

Since this action is defined using the continuous associative multiplication $\cdot : \Lambda_{\gamma_i} \times \Lambda_{\gamma_j} \rightarrow \Lambda_{\gamma_i + \gamma_j}$ of the Fréchet algebra Λ , it is (jointly) continuous.

As any closed subspace of a Fréchet space is itself a Fréchet space, the space

$$\mathring{\Lambda}_0 \simeq \{0\} \times \mathring{\Lambda}_0 \subset \mathbb{R} \times \mathring{\Lambda}_0 = \Lambda_0$$

is Fréchet. We thus see that

$$(3.19) \quad \mathcal{U}^{p|q}(\Lambda) \simeq \mathcal{U}^p \times \mathring{\Lambda}_0^p \times \prod_{i=1}^N \Lambda_{\gamma_i}^{q_i} \subset \mathbb{R}^p \times \mathring{\Lambda}_0^p \times \prod_{i=1}^N \Lambda_{\gamma_i}^{q_i} \simeq \mathbb{R}^{p|q}(\Lambda)$$

is open. \square

Remark 3.16. In the following, we will use the isomorphisms (3.19) (and similar ones) without further reference.

The just described Λ_0 -module structure is vital in understanding the structure of the Λ -points of any \mathbb{Z}_2^n -manifold. In particular, morphisms between \mathbb{Z}_2^n -domains induce natural transformations between the associated functors that respect this module structure. The converse is also true, that is, any natural transformation between the associated functors that respects the Λ_0 -module structure comes from a morphism between the underlying \mathbb{Z}_2^n -domains. More carefully, we have the following proposition.

Theorem 3.17. *Let $\mathcal{U}^{p|q}$ and $\mathcal{V}^{r|s}$ be \mathbb{Z}_2^n -domains. A natural transformation $\beta : \mathcal{U}^{p|q}(-) \rightarrow \mathcal{V}^{r|s}(-)$ comes from a \mathbb{Z}_2^n -manifold morphism $\mathcal{U}^{p|q} \rightarrow \mathcal{V}^{r|s}$ if and only if $\beta_\Lambda : \mathcal{U}^{p|q}(\Lambda) \rightarrow \mathcal{V}^{r|s}(\Lambda)$ is Λ_0 -smooth, for all $\Lambda \in \mathbb{Z}_2^n \text{GrAlg}$. That is, for all Λ , the map β_Λ must be a smooth map (from the open subset $\mathcal{U}^{p|q}(\Lambda)$ of the Fréchet space $\mathbb{R}^{p|q}(\Lambda)$ to the Fréchet space $\mathbb{R}^{r|s}(\Lambda)$, see Appendix B) and its Gâteaux derivative (see Appendix B) must be Λ_0 -linear, i.e.,*

$$d_{x^*} \beta_\Lambda(a \cdot v) = a \cdot d_{x^*} \beta_\Lambda(v) ,$$

for all $x^* \in \mathcal{U}^{p|q}(\Lambda)$, $a \in \Lambda_0$, and $v \in \mathbb{R}^{p|q}(\Lambda)$.

Proof. Part I. Let $\beta : \mathcal{U}^{p|q}(-) \rightarrow \mathcal{V}^{r|s}(-)$ be a natural transformation with Λ_0 -smooth components β_Λ , $\Lambda \in \mathbb{Z}_2^n \text{GrAlg}$. From Theorem 3.14, we know that β_Λ is completely specified by the systems

$$(3.20) \quad y_\Lambda^b = \sum_{|\alpha| \geq 0, |\beta| \geq 0} F_{\alpha\beta}^b(x_{||}) \dot{x}_\Lambda^\beta \xi_\Lambda^\alpha \quad \text{and} \quad \eta_\Lambda^B = \sum_{|\alpha| > 0, |\beta| \geq 0} F_{\alpha\beta}^B(x_{||}) \dot{x}_\Lambda^\beta \xi_\Lambda^\alpha ,$$

where the coefficients $F_{\alpha\beta}^b$ ($b \in \{b, B\}$) are set-theoretical maps from \mathcal{U}^p to \mathbb{R} .

Part Ia. Smoothness of β_Λ implies that these coefficients are smooth. Indeed, we will show that $F_{\alpha\beta}^b \in C^0(\mathcal{U}^p)$ and that, if $F_{\alpha\beta}^b \in C^k(\mathcal{U}^p)$ ($k \geq 0$), then $F_{\alpha\beta}^b \in C^{k+1}(\mathcal{U}^p)$.

Step 1. Since

$$\beta_\Lambda : \mathcal{U}^{p|q}(\Lambda) \rightarrow \Lambda_0^r \times \prod_{i=1}^N \Lambda_{\gamma_i}^{s_i}$$

is continuous, any of its components

$$y_\Lambda^b : \mathcal{U}^{p|q}(\Lambda) \rightarrow \Lambda_{\gamma_i(b)} = \mathbb{R}[[\theta]]_{\gamma_i(b)} \simeq \prod_{\gamma_i(b)} \mathbb{R}$$

is continuous. For simplicity, we wrote y_Λ^B instead of η_Λ^B , and we will continue doing so. Moreover, the target space are the formal power series in θ with coefficients in \mathbb{R} , all whose terms have the degree $\gamma_i(b)$ of y^b , and this space is identified with the corresponding space of families of reals. For any ω such that θ^ω has the degree $\gamma_i(b)$, the corresponding real coefficient gives rise to a continuous map

$$y_\Lambda^{b,\omega} : \mathcal{U}^{p|q}(\Lambda) \rightarrow \mathbb{R} .$$

Since this joint continuity implies separate continuity with respect to $x_{||} \in \mathcal{U}^p$, for any fixed $(\dot{x}_\Lambda, \xi_\Lambda)$ and any Λ , we can proceed as at the end of the proof of Theorem 3.14. More precisely, select any (α, β) and select (for an appropriate Λ) the pair $(\dot{x}_\Lambda, \xi_\Lambda)$ such that $\dot{x}_\Lambda^\beta \xi_\Lambda^\alpha = \beta! \theta^\omega$, where θ^ω is now the degree $\gamma_i(b)$ monomial defined in the proof just mentioned. The real coefficient of this monomial is $\beta! F_{\alpha\beta}^b(x_{||})$, which, as said, is an \mathbb{R} -valued continuous map on \mathcal{U}^p , so that $F_{\alpha\beta}^b \in C^0(\mathcal{U}^p)$, for all b and all (α, β) .

Step 2. Since

$$\mathcal{U}^{p|q}(\Lambda) \subset \mathbb{R} \times (\mathbb{R}^{p-1} \times \Lambda_0^p \times \prod_{i=1}^N \Lambda_{\gamma_i}^{q_i})$$

is an open subset of a product of two Fréchet spaces, smoothness of β_Λ implies (via an iterated application of Proposition B.4) that, for any $b \in \{b, B\}$, any $\ell \in \mathbb{N}$ and any $\gamma \in \mathbb{N}^p$ ($|\gamma| = \ell$), the partial derivative

$$d_{x_{||}}^\gamma y_\Lambda^b : \mathcal{U}^{p|q}(\Lambda) \times \mathbb{R}^{\times \ell} \rightarrow \prod_{\gamma_i(b)} \mathbb{R}$$

is continuous.

Assume now that $F_{\alpha\beta}^b \in C^k(\mathcal{U}^p)$ ($k \geq 0$), for any b and any (α, β) , as well as that, for any $\gamma \in \mathbb{N}^p$ ($|\gamma| = k$) and any b , the continuous partial Gâteaux derivative

$$d_{x_{||}}^\gamma y_\Lambda^b(1, \dots, 1) : \mathcal{U}^{p|q}(\Lambda) \rightarrow \prod_{\gamma_i(b)} \mathbb{R}$$

is given by

$$(3.21) \quad d_{x_{||}, x^*}^\gamma y_\Lambda^b(1, \dots, 1) = \sum_{\alpha\beta} (\partial_x^\gamma F_{\alpha\beta}^b)(x_{||}) \dot{x}_\Lambda^\beta \xi_\Lambda^\alpha .$$

Observe that for $k = 0$, this condition is automatically satisfied. We will now show that, under these assumptions, the same statements hold at order $k + 1$. In view of (3.21), any order $k + 1$ continuous partial Gâteaux derivative

$$d_{x_{||}}^a d_{x_{||}}^\gamma y_\Lambda^b(1, \dots, 1) : \mathcal{U}^{p|q}(\Lambda) \rightarrow \prod_{\gamma_i(b)} \mathbb{R}$$

$(a \in \{1, \dots, p\}, |\gamma| = k)$ is given, at any $\mathbf{x}^* \simeq (x_{||}, \dot{x}_\Lambda, \xi_\Lambda) \in \mathcal{U}^{p|q}(\Lambda)$, by

$$(3.22) \quad \sum_{\alpha\beta} \lim_{t \rightarrow 0} \frac{1}{t} \left((\partial_x^\gamma F_{\alpha\beta}^b)(x_{||}^1, \dots, x_{||}^a + t, \dots, x_{||}^p) - (\partial_x^\gamma F_{\alpha\beta}^b)(x_{||}^1, \dots, x_{||}^a, \dots, x_{||}^p) \right) \dot{x}_\Lambda^\beta \xi_\Lambda^\alpha .$$

When proceeding as in Step 1, we get that the limit is an \mathbb{R} -valued continuous function in \mathcal{U}^p . In other words, the partial derivative $\partial_{x^a} \partial_x^\gamma F_{\alpha\beta}^b$ exists and is continuous in \mathcal{U}^p , i.e., $F_{\alpha\beta}^b \in C^{k+1}(\mathcal{U}^p)$. Moreover, Formula (3.21) pertaining to order k derivatives, extends to the order $k+1$ derivatives, see (3.22).

Part Ib. We examine the further consequences of Λ_0 -smoothness, in particular those of Λ_0 -linearity. Since y_Λ^b is of class C^1 , its components $y_\Lambda^b : \mathcal{U}^{p|q}(\Lambda) \rightarrow \prod_{\gamma_i(b)} \mathbb{R}$ are of class C^1 . Further, as

$$\mathcal{U}^{p|q}(\Lambda) \subset (\mathbb{R} \times \mathring{\Lambda}_0) \times (\mathbb{R}^{p-1} \times \mathring{\Lambda}_0^{p-1} \times \prod_{i=1}^N \Lambda_{\gamma_i}^{q_i})$$

is an open subset of a product of two Fréchet spaces, the partial Gâteaux derivative

$$d_{(x_{||}^a, \dot{x}_\Lambda^a), \mathbf{x}^*} y_\Lambda^b : \mathcal{U}^{p|q}(\Lambda) \times (\mathbb{R} \times \mathring{\Lambda}_0) \rightarrow \prod_{\gamma_i(b)} \mathbb{R}$$

is continuous. It is given by

$$d_{(x_{||}^a, \dot{x}_\Lambda^a), \mathbf{x}^*} y_\Lambda^b(v_{||}, \dot{v}_\Lambda) = d_{x_{||}^a, \mathbf{x}^*} y_\Lambda^b(v_{||}) + d_{\dot{x}_\Lambda^a, \mathbf{x}^*} y_\Lambda^b(\dot{v}_\Lambda) =$$

$$v_{||} \sum_{\alpha\beta} (\partial_{x^a} F_{\alpha\beta}^b)(x_{||}) \dot{x}_\Lambda^\beta \xi_\Lambda^\alpha + \sum_{\alpha\beta} F_{\alpha\beta}^b(x_{||}) \lim_{t \rightarrow 0} \frac{1}{t} ((\dot{x}_\Lambda^a + t \dot{v}_\Lambda)^{\beta_a} - (\dot{x}_\Lambda^a)^{\beta_a}) \prod_{b \neq a} (\dot{x}_\Lambda^b)^{\beta_b} \xi_\Lambda^\alpha =: v_{||} T_1 + \mathfrak{T}_2 .$$

As $\mathring{\Lambda}_0$ is a commutative algebra, it follows from the binomial formula that

$$\mathfrak{T}_2 = \dot{v}_\Lambda \sum_{\alpha\beta} \beta_a F_{\alpha\beta}^b(x_{||}) \dot{x}_\Lambda^{\beta - e_a} \xi_\Lambda^\alpha =: \dot{v}_\Lambda T_2 ,$$

where $(e_a)_a$ is the canonical basis of \mathbb{R}^p . Observe now that, in view of (3.18), the Λ_0 -linearity of the total Gâteaux derivative of y_Λ^b with respect to \mathbf{x}^* is equivalent to the Λ_0 -linearity of all its partial Gâteaux derivatives with respect to the $x_\Lambda^a = (x_{||}^a, \dot{x}_\Lambda^a)$ and the ξ_Λ^A . For $a = 0 + \dot{v}_\Lambda \in \Lambda_0$ and $\mathbf{v} = 1 + 0 \in \mathbb{R} + \mathring{\Lambda}_0 = \Lambda_0$, this implies that

$$\dot{v}_\Lambda T_2 = d_{(x_{||}^a, \dot{x}_\Lambda^a), \mathbf{x}^*} y_\Lambda^b(\dot{v}_\Lambda \cdot 1) = \dot{v}_\Lambda \cdot d_{(x_{||}^a, \dot{x}_\Lambda^a), \mathbf{x}^*} y_\Lambda^b(1) = \dot{v}_\Lambda T_1 ,$$

i.e., that

$$\dot{v}_\Lambda \sum_{\alpha\beta} (\beta_a + 1) F_{\alpha, \beta + e_a}^b(x_{||}) \dot{x}_\Lambda^\beta \xi_\Lambda^\alpha = \dot{v}_\Lambda \sum_{\alpha, \gamma: \gamma_a \neq 0} \gamma_a F_{\alpha\gamma}^b(x_{||}) \dot{x}_\Lambda^{\gamma - e_a} \xi_\Lambda^\alpha = \dot{v}_\Lambda \sum_{\alpha\beta} (\partial_{x^a} F_{\alpha\beta}^b)(x_{||}) \dot{x}_\Lambda^\beta \xi_\Lambda^\alpha .$$

Since $\Lambda \in \mathbb{Z}_2^n \mathbf{GrAlg}$, $\dot{v}_\Lambda \in \mathring{\Lambda}_0$, and $\mathbf{x}^* \in \mathcal{U}^{p|q}(\Lambda)$ are arbitrary, we can repeat the θ^ω -argument used above. More precisely, we select (α, β) , select $(\dot{x}_\Lambda, \xi_\Lambda)$ such that $\dot{x}_\Lambda^\beta \xi_\Lambda^\alpha = \beta! \theta^\omega$, and select $\dot{v}_\Lambda = \theta^D \theta^E \in \mathring{\Lambda}_0$ such that $\theta^D \theta^E \theta^\omega \neq 0$. The coefficients of the latter monomial in the left and right hand sides do coincide, which means that

$$(3.23) \quad (\beta_a + 1) F_{\alpha, \beta + e_a}^b(x_{||}) = (\partial_{x^a} F_{\alpha\beta}^b)(x_{||}), \text{ or, equivalently, } F_{\alpha\gamma}^b(x_{||}) = \frac{1}{\gamma_a!} (\partial_{x^a} F_{\alpha, \gamma - e_a}^b)(x_{||}) ,$$

for all \mathbf{b}, α, a , all $\gamma : \gamma_a \neq 0$, and all $x_{||} \in \mathcal{U}^p$. For any \mathbf{b} , α , and $x_{||}$, we now set

$$\phi_\alpha^b(x_{||}) := F_{\alpha 0}^b(x_{||}) \in C^\infty(\mathcal{U}^p) .$$

An iterated application of (3.23) shows that

$$F_{\alpha\gamma}^b(x_{||}) = \frac{1}{\gamma!} (\partial_x^\gamma \phi_\alpha^b)(x_{||}) .$$

Hence, the y_Λ^b have the form (3.10a) and (3.10b). This means that the natural transformation β is implemented by the ϕ_α^b , which define actually a \mathbb{Z}_2^n -morphism from $\mathcal{U}^{p|q}$ to $\mathcal{V}^{r|s}$. Indeed, the property $(\phi_0^b)(\mathcal{U}^p) \subset \mathcal{V}^r$ follows from the similar property of (F_{00}^b) . On the other hand, the pullback

$$\phi^*(y^b) := \sum_\alpha \phi_\alpha^b(x) \xi_\Lambda^\alpha$$

must have the same degree as y^b . However, if $\deg(\xi_\Lambda^\alpha) \neq \deg(y^b)$, then $\deg(\xi_\Lambda^\alpha) \neq \deg(y_\Lambda^b)$, whatever ξ_Λ . It follows therefore from (3.20) that $\phi_\alpha^b = F_{\alpha 0}^b = 0$.

Part II. The proof of the converse implication is less demanding. Let $\beta : \mathcal{U}^{p|q}(-) \rightarrow \mathcal{V}^{r|s}(-)$ be a natural transformation that is induced by a \mathbb{Z}_2^n -morphism $\phi : \mathcal{U}^{p|q} \rightarrow \mathcal{V}^{r|s}$, i.e., that is of the form (3.10a) and (3.10b). For any $\Lambda \in \mathbb{Z}_2^n \mathbf{GrAlg}$, the map β_Λ is smooth and its derivative is Λ_0 -linear if and only if its components y_Λ^b

have these properties. The total derivative of y_Λ^b with respect to x^* exists, is continuous, and is Λ_0 -linear if and only if its partial derivatives with respect to the x_Λ^a and the ξ_Λ^A exist, are continuous, and are Λ_0 -linear. When computing the derivative y_Λ^b with respect to $\xi_\Lambda^{A_i} \in \Lambda_{\gamma_i}$ at $x^* \in \mathcal{U}^{p|q}(\Lambda)$ in the direction of $w_\Lambda \in \Lambda_{\gamma_i}$, we get

$$\sum_{\alpha\beta} \frac{1}{\beta!} (\partial_x^\beta \phi_\alpha^b)(x_{||}) \dot{x}_\Lambda^\beta (\xi_\Lambda^{A_1})^{\alpha_1} \dots \lim_{t \rightarrow 0} \frac{1}{t} ((\xi_\Lambda^{A_i} + tw_\Lambda)^{\alpha_i} - (\xi_\Lambda^{A_i})^{\alpha_i}) \dots (\xi_\Lambda^{A_{|\underline{q}|}})^{\alpha_{|\underline{q}|}}.$$

If γ_i is odd, the exponent α_i is 0 or 1. In the first (resp., the second) case, the limit vanishes (resp., is w_Λ). If γ_i is even, the multiplication of vectors in Λ_{γ_i} is commutative and the binomial formula shows that the limit is $w_\Lambda \alpha_i (\xi_\Lambda^{A_i})^{\alpha_i-1}$. The derivative thus exists, is continuous, and is Λ_0 -linear. Similarly, the derivative of y_Λ^b with respect to x_Λ^a exists if and only if its derivatives with respect to $x_{||}^a$ and with respect to \dot{x}_Λ^a exist. The (standard) computation of the derivative with respect to x_Λ^a at x^* in the direction of

$$v_\Lambda = (v_{||}, \dot{v}_\Lambda) \in \mathbb{R} \times \mathring{\Lambda}_0$$

thus leads to the sum of the terms

$$v_{||} \sum_{\alpha\beta} \frac{1}{\beta!} (\partial_x^{\beta+e_a} \phi_\alpha^b)(x_{||}) \dot{x}_\Lambda^\beta \xi_\Lambda^\alpha$$

and

$$\dot{v}_\Lambda \sum_{\alpha, \gamma: \gamma_a \neq 0} \frac{1}{\gamma!} (\partial_x^\gamma \phi_\alpha^b)(x_{||}) \gamma_a \dot{x}_\Lambda^{\gamma-e_a} \xi_\Lambda^\alpha = \dot{v}_\Lambda \sum_{\alpha\beta} \frac{1}{\beta!} (\partial_x^{\beta+e_a} \phi_\alpha^b)(x_{||}) \dot{x}_\Lambda^\beta \xi_\Lambda^\alpha.$$

The derivative considered does therefore exist, is continuous, and is Λ_0 -linear (note that it is essential that the derivative is the series over $\alpha\beta$ multiplied by v_Λ – as $a \in \Lambda_0$ does not act on $v_{||}$). \square

Remark 3.18. The Λ_0 -linearity is a strong constraint that takes us from the category of generalized \mathbb{Z}_2^n -manifolds to the one of \mathbb{Z}_2^n -manifolds. A similar phenomenon exists in complex analysis. Indeed, for any real differentiable function $f = u + i v : \Omega \subset \mathbb{C} \simeq \mathbb{R}^2 \rightarrow \mathbb{C} \simeq \mathbb{R}^2$, the Jacobian is an \mathbb{R} -linear map $J_f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$. However, if we further insist that the Jacobian be \mathbb{C} -linear, then we see that f must be holomorphic, that is, it must satisfy the Cauchy–Riemann equations on Ω . Imposing \mathbb{C} -linearity thus greatly restricts class of functions and takes us from real analysis to complex analysis.

It will also be important to understand what happens to the Λ -points of a given \mathbb{Z}_2^n -domain under morphisms of \mathbb{Z}_2^n -Grassmann algebras.

Proposition 3.19. *Let $\mathcal{U}^{p|q}$ be a \mathbb{Z}_2^n -domain and let $\psi^* : \Lambda \rightarrow \Lambda'$ be a morphism of \mathbb{Z}_2^n -Grassmann algebras. The induced map (see (3.4))*

$$\Psi := \mathcal{U}^{p|q}(\psi^*) : \mathcal{U}^{p|q}(\Lambda) \ni x^* \simeq (x_\Lambda, \xi_\Lambda) \mapsto \psi^* \circ x^* \simeq \psi^*(x_\Lambda, \xi_\Lambda) \in \mathcal{U}^{p|q}(\Lambda')$$

is a smooth map from the open subset $\mathcal{U}^{p|q}(\Lambda)$ of the Fréchet space and Fréchet Λ_0 -module $\mathbb{R}^{p|q}(\Lambda)$ to the open subset $\mathcal{U}^{p|q}(\Lambda')$ of the Fréchet space and Fréchet Λ'_0 -module $\mathbb{R}^{p|q}(\Lambda')$, such that

$$d_{x^*} \Psi(a \cdot v) = \psi^*(a) \cdot d_{x^*} \Psi(v),$$

for all $x^ \in \mathcal{U}^{p|q}(\Lambda)$, $v \in \mathbb{R}^{p|q}(\Lambda)$ and $a \in \Lambda_0$.*

Proof. Since $\Lambda = \mathcal{O}_{\mathbb{R}^{0|m}}(\{\star\})$, so that

$$\psi^* \in \text{Hom}_{\mathbb{Z}_2^n \text{Alg}}(\mathcal{O}_{\mathbb{R}^{0|m}}(\{\star\}), \mathcal{O}_{\mathbb{R}^{0|m'}}(\{\star\})),$$

there is a unique morphism

$$\Phi = (|\phi|, \phi^*) \in \text{Hom}_{\mathbb{Z}_2^n \text{Man}}(\mathbb{R}^{0|m'}, \mathbb{R}^{0|m}),$$

such that $\psi^* = \phi^*_{\{\star\}}$. Hence, the morphism ψ^* is continuous from $\Lambda = \mathbb{R}[[\theta]]$ to $\Lambda' = \mathbb{R}[[\theta']]$ endowed with their standard locally convex topologies [13], and so are its restrictions $\psi^*|_{\Lambda_{\gamma_i}}$ from Λ_{γ_i} to Λ'_{γ_i} . We thus see that the induced map

$$\Psi = (\psi^*|_{\Lambda_0})^{\times p} \times \prod_{i=1}^N (\psi^*|_{\Lambda_{\gamma_i}})^{\times q_i}$$

is continuous.

At $x^* \simeq (x_\Lambda, \xi_\Lambda) =: u_\Lambda \in \mathcal{U}^{p|q}(\Lambda)$ and $v \simeq v_\Lambda \in \mathbb{R}^{p|q}(\Lambda)$, the derivative

$$d\Psi : \mathcal{U}^{p|q}(\Lambda) \times \mathbb{R}^{p|q}(\Lambda) \longrightarrow \mathbb{R}^{p|q}(\Lambda')$$

is defined as

$$\begin{aligned} d_{x^*}\Psi(v) &= \lim_{t \rightarrow 0} \frac{\Psi(x^* + tv) - \Psi(x^*)}{t} \\ &= \lim_{t \rightarrow 0} \left(\dots, \frac{\psi^*(u_\Lambda^a + tv_\Lambda^a) - \psi^*(u_\Lambda^a)}{t}, \dots \right) \\ &= \left(\dots, \psi^*(v_\Lambda^a), \dots \right) \\ &=: (\psi^*(v_\Lambda^a)), \end{aligned}$$

where a is the label $a \in \{1, \dots, p\}$ or $A \in \{1, \dots, |q|\}$ of any coordinate in $\mathbb{R}^{p|q}(\Lambda)$, and where we used the \mathbb{R} -linearity of the \mathbb{Z}_2^n -algebra morphism $\psi^* : \Lambda \rightarrow \Lambda'$. Hence, for any $a \in \Lambda_0$, we get

$$d_{x^*}\psi(a \cdot v) = (\psi^*(a \cdot v_\lambda^a)) = (\psi^*(a) \cdot \psi^*(v_\lambda^a)) = \psi^*(a) \cdot d_{x^*}\psi(v).$$

Since the higher order derivatives of Ψ vanish, all its derivatives exist and are continuous, hence, the map Ψ is actually smooth. \square

3.4. The manifold structure on the set of Λ -points. The next theorem generalizes Propositions 3.15 and 3.19. For information about Fréchet manifolds, we refer to Appendix B. We recall that the Λ -points $M(\Lambda)$ of a \mathbb{Z}_2^n -manifold M can be equivalently viewed as the maps $m = (|m|, m^*) \in \text{Hom}_{\mathbb{Z}_2^n \text{Man}}(\mathbb{R}^{0|m}, M)$, as the global pullbacks $m^* = m_{|M|}^* \in \text{Hom}_{\mathbb{Z}_2^n \text{Alg}}(\mathcal{O}_M(|M|), \Lambda)$, or as the induced morphisms

$$m_\star^* \in \text{Hom}_{\mathbb{Z}_2^n \text{Alg}}(\mathcal{O}_{M,x}, \Lambda),$$

where $x = |m|(\star) \in |M|$. If $M = \mathcal{U}^{p|q}$ is a \mathbb{Z}_2^n -domain, we often write x instead of m and we can identify $x \simeq x^* \simeq x_\star^*$ with the pullbacks

$$(u_{||}, \dot{u}_\Lambda, \rho_\Lambda) \in \mathcal{U}^p \times \dot{\Lambda}_0^p \times \prod_i \Lambda_{\gamma_i}^{q_i}$$

by x^* of the coordinate functions (u, ρ) in $\mathcal{U}^{p|q}$. Recall as well that \mathbb{Z}_2^n -morphisms $\phi : M \rightarrow N$ are mapped injectively to natural transformations $\phi : M(-) \rightarrow N(-)$ with Λ -component

$$(3.24) \quad \phi_\Lambda : M(\Lambda) \ni (x, m_\star^*) \mapsto (|\phi|(x), m_\star^* \circ \phi_x^*) \in N(\Lambda),$$

and that, for any fixed M , a \mathbb{Z}_2^n -Grassmann algebra morphism $\psi^* : \Lambda \rightarrow \Lambda'$ induces a map

$$M(\psi^*) : M(\Lambda) \ni (x, m_\star^*) \mapsto (x, \psi^* \circ m_\star^*) \in M(\Lambda').$$

Theorem 3.20. *Let M be a \mathbb{Z}_2^n -manifold, and let Λ and Λ' be \mathbb{Z}_2^n -Grassmann algebras. Then*

- (i) *$M(\Lambda)$ has the structure of a nuclear Fréchet Λ_0 -manifold, and,*
- (ii) *given a morphism of \mathbb{Z}_2^n -Grassmann algebras $\psi^* : \Lambda \rightarrow \Lambda'$, the induced mapping $M(\psi^*)$ is ψ^* -smooth.*

Proof.

- (i) Let $p|q$ be the dimension of the \mathbb{Z}_2^n -manifold M . The local \mathbb{Z}_2^n -isomorphisms

$$h_\alpha = (|h_\alpha|, h_\alpha^*) : U_\alpha = (|U_\alpha|, \mathcal{O}_M|_{|U_\alpha|}) \rightarrow \mathcal{U}_\alpha^{p|q} = (\mathcal{U}_\alpha^p, C_{\mathbb{R}^p}^\infty|_{\mathcal{U}_\alpha^p}[[\rho]]),$$

where α varies in some \mathcal{A} and where $|U_\alpha| \subset |M|$ is open, provide an atlas on M (see paragraph below Definition 2.10). As recalled above, the \mathbb{Z}_2^n -isomorphisms

$$h_\alpha : U_\alpha \rightarrow \mathcal{U}_\alpha^{p|q}$$

implement natural isomorphisms h_α with Λ -components

$$(3.25) \quad h_{\alpha, \Lambda} : U_\alpha(\Lambda) \ni (x, m_\star^*) \mapsto (|h_\alpha|(x), m_\star^* \circ (h_\alpha)_x^*) \in \mathcal{U}_\alpha^{p|q}(\Lambda),$$

whose inverses are the similar maps defined using

$$|h_\alpha^{-1}| = |h_\alpha|^{-1} \quad \text{and} \quad (h_\alpha^{-1})_y^* = ((h_\alpha)_y^*|_{h_\alpha|^{-1}(y)})^{-1} \quad (y \in \mathcal{U}_\alpha^p).$$

The family $(U_\alpha(\Lambda), h_{\alpha, \Lambda})$ ($\alpha \in \mathcal{A}$) is an atlas that endows $M(\Lambda)$ with a nuclear Fréchet Λ_0 -manifold structure. Indeed:

- (a) Any $h_{\alpha, \Lambda} : U_\alpha(\Lambda) \rightarrow \mathcal{U}_\alpha^{p|q}(\Lambda)$ is a bijection valued in the open subset $\mathcal{U}_\alpha^{p|q}(\Lambda)$ of the nuclear Fréchet vector space $\mathbb{R}^{p|q}(\Lambda)$, which is also a Fréchet module over the nuclear Fréchet algebra Λ_0 . Moreover, as the $|U_\alpha|$ are an open cover of $|M|$, we have

$$M(\Lambda) = \bigcup_{\alpha \in \mathcal{A}} U_\alpha(\Lambda),$$

in view of Proposition 3.12.

(b) The image $h_{\alpha,\Lambda}(U_\alpha(\Lambda) \cap U_\beta(\Lambda))$ is open in $\mathbb{R}^{p|q}(\Lambda)$. To see this, set $|U_{\alpha\beta}| = |U_\alpha| \cap |U_\beta| \subset |U_\alpha|$ and consider the open \mathbb{Z}_2^n -submanifold $U_{\alpha\beta} = (|U_{\alpha\beta}|, \mathcal{O}_M|_{|U_{\alpha\beta}|})$ of U_α . The \mathbb{Z}_2^n -isomorphism h_α restricts to a \mathbb{Z}_2^n -isomorphism

$$h_\alpha : U_{\alpha\beta} \rightarrow \mathcal{U}_{\alpha\beta}^{p|q},$$

where the target is the open \mathbb{Z}_2^n -subdomain $\mathcal{U}_{\alpha\beta}^{p|q}$ of $\mathcal{U}_\alpha^{p|q}$ defined over the open subset

$$\mathcal{U}_{\alpha\beta}^p := |h_\alpha|(|U_{\alpha\beta}|) \subset \mathcal{U}_\alpha^p,$$

obtained as the image of the open subset $|U_{\alpha\beta}| \subset |U_\alpha|$ by the diffeomorphism $|h_\alpha|$. The restricted \mathbb{Z}_2^n -isomorphism h_α induces a natural isomorphism h_α , whose Λ -component is a bijection

$$h_{\alpha,\Lambda} : U_{\alpha\beta}(\Lambda) \rightarrow \mathcal{U}_{\alpha\beta}^{p|q}(\Lambda).$$

Further, we have

$$\begin{aligned} U_{\alpha\beta}(\Lambda) &= \bigcup_{x \in |U_{\alpha\beta}|} \text{Hom}_{\mathbb{Z}_2^n \text{Alg}}(\mathcal{O}_{M,x}, \Lambda) = \\ &\bigcup_{x \in |U_\alpha|} \text{Hom}_{\mathbb{Z}_2^n \text{Alg}}(\mathcal{O}_{M,x}, \Lambda) \cap \bigcup_{x \in |U_\beta|} \text{Hom}_{\mathbb{Z}_2^n \text{Alg}}(\mathcal{O}_{M,x}, \Lambda) = U_\alpha(\Lambda) \cap U_\beta(\Lambda). \end{aligned}$$

Hence, the image $h_{\alpha,\Lambda}(U_\alpha(\Lambda) \cap U_\beta(\Lambda)) = \mathcal{U}_{\alpha\beta}^{p|q}(\Lambda) \subset \mathbb{R}^{p|q}(\Lambda)$ is open.

(c) We have still to prove that the transition bijections

$$h_{\beta,\Lambda}(h_{\alpha,\Lambda})^{-1} : \mathcal{U}_{\alpha\beta}^{p|q}(\Lambda) \rightarrow \mathcal{U}_{\beta\alpha}^{p|q}(\Lambda)$$

are Λ_0 -smooth. In view of Theorem 3.17, the \mathbb{Z}_2^n -isomorphism

$$h_\beta h_\alpha^{-1} : \mathcal{U}_{\alpha\beta}^{p|q} \rightarrow \mathcal{U}_{\beta\alpha}^{p|q}$$

induces a natural isomorphism $h_\beta h_\alpha^{-1}$ with a Λ_0 -smooth Λ -component

$$(h_\beta h_\alpha^{-1})_\Lambda : \mathcal{U}_{\alpha\beta}^{p|q}(\Lambda) \rightarrow \mathcal{U}_{\beta\alpha}^{p|q}(\Lambda).$$

In view of Equations (3.24) and (3.25), we get

$$\begin{aligned} (h_\beta h_\alpha^{-1})_\Lambda(u, x_\star^*) &= (|h_\beta h_\alpha^{-1}|(u), x_\star^* \circ (h_\beta \circ h_\alpha^{-1})_u^*) = \\ (|h_\beta|(|h_\alpha|^{-1}(u)), x_\star^* \circ ((h_\alpha)_{|h_\alpha|^{-1}(u)}^*)^{-1} \circ (h_\beta)_{|h_\alpha|^{-1}(u)}^*) &= h_{\beta,\Lambda}((h_{\alpha,\Lambda})^{-1}(u, x_\star^*)), \end{aligned}$$

for any $(u, x_\star^*) \in \mathcal{U}_{\alpha\beta}^{p|q}(\Lambda)$. It follows that $h_{\beta,\Lambda}(h_{\alpha,\Lambda})^{-1} = (h_\beta h_\alpha^{-1})_\Lambda$ is Λ_0 -smooth.

(ii) The statement of part (ii) is purely local, see Appendix B. Let $(x, m_\star^*) \in M(\Lambda)$, let $(U_\alpha(\Lambda), h_{\alpha,\Lambda})$ be a chart of $M(\Lambda)$ around (x, m_\star^*) , and let $(U_\beta(\Lambda'), h_{\beta,\Lambda'})$ be a chart of $M(\Lambda')$, such that $M(\psi^*)(U_\alpha(\Lambda)) \subset U_\beta(\Lambda')$. We must show that the local form

$$h_{\beta,\Lambda'} \circ M(\psi^*) \circ (h_{\alpha,\Lambda})^{-1}$$

of $M(\psi^*)$ is ψ^* -smooth. Actually, we can choose $(U_\alpha(\Lambda'), h_{\alpha,\Lambda'})$ as second chart, since the image by $M(\psi^*)$ of a point (y, n_\star^*) in $U_\alpha(\Lambda)$, i.e., a point

$$(y, n_\star^*) \in \text{Hom}_{\mathbb{Z}_2^n \text{Alg}}(\mathcal{O}_{M,y}, \Lambda)$$

with $y \in |U_\alpha|$, is the point

$$(y, \psi^* \circ n_\star^*) \in \text{Hom}_{\mathbb{Z}_2^n \text{Alg}}(\mathcal{O}_{M,y}, \Lambda'),$$

i.e., in $U_\alpha(\Lambda')$. From here, we omit subscript α . Since $h : U(-) \rightarrow \mathcal{U}^{p|q}(-)$ is a natural transformation, the diagram

$$\begin{array}{ccc} U(\Lambda) & \xrightarrow{M(\psi^*)} & U(\Lambda') \\ h_\Lambda \downarrow & & \downarrow h_{\Lambda'} \\ \mathcal{U}^{p|q}(\Lambda) & \xrightarrow{\quad} & \mathcal{U}^{p|q}(\Lambda') \\ & & \mathcal{U}^{p|q}(\psi^*) \end{array}$$

commutes. Since h is in fact a natural isomorphism, we get that

$$h_{\Lambda'} \circ M(\psi^*) \circ (h_\Lambda)^{-1} = \mathcal{U}^{p|q}(\psi^*).$$

From Proposition 3.19 we conclude that this local form is indeed ψ^* -smooth. □

In view of (3.8), the local model $\mathbb{R}^{p|q}(\Lambda)$ of $M(\Lambda)$ is infinite-dimensional, due to the non-zero degree even coordinates in Λ . Further, we have the

Corollary 3.3. *For any \mathbb{Z}_2^n -manifold M , the associated functor*

$$M(-) \in [\mathbb{Z}_2^n \mathbf{Pts}^{\text{op}}, \mathbf{Set}]$$

can be considered as a functor

$$M(-) \in [\mathbb{Z}_2^n \mathbf{Pts}^{\text{op}}, \mathbf{A(N)FMan}] ,$$

where the target category is either the category **AFMan** of Fréchet manifolds over a Fréchet algebra or the category **ANFMan** of nuclear Fréchet manifolds over a nuclear Fréchet algebra, see Appendix B. Therefore, the faithful restricted Yoneda functor $\mathcal{Y}_{\mathbb{Z}_2^n \mathbf{Pts}}$, see Corollary 3.2, can be viewed as a faithful functor

$$\mathcal{Y}_{\mathbb{Z}_2^n \mathbf{Pts}} : \mathbb{Z}_2^n \mathbf{Man} \rightarrow [\mathbb{Z}_2^n \mathbf{Pts}^{\text{op}}, \mathbf{A(N)FMan}] .$$

The latter statement requires that the natural transformation $\phi : M(-) \rightarrow N(-)$ induced by a \mathbb{Z}_2^n -morphism $\phi : M \rightarrow N$ have components $\phi_\Lambda : M(\Lambda) \rightarrow N(\Lambda)$ that are morphisms in **A(N)FMan** between the Fréchet Λ_0 -manifolds $M(\Lambda)$ and $N(\Lambda)$, i.e., that the ϕ_Λ be ρ -smooth for some morphism $\rho : \Lambda_0 \rightarrow \Lambda_0$ of Fréchet algebras. We will show in the next subsection that this condition is satisfied for $\rho = \text{id}_{\Lambda_0}$, i.e., we will show that:

Proposition 3.21. *Any natural transformation $\phi : M(-) \rightarrow N(-)$ that is implemented by a \mathbb{Z}_2^n -morphism $\phi : M \rightarrow N$ has Λ_0 -smooth components $\phi_\Lambda : M(\Lambda) \rightarrow N(\Lambda)$.*

Theorem 3.22. *Let $M \in \mathbb{Z}_2^n \mathbf{Man}$ be of dimension $p|q$ and let $\Lambda \in \mathbb{Z}_2^n \mathbf{GrAlg}$.*

(i) *The nuclear Fréchet Λ_0 -manifold $M(\Lambda)$ is a fiber bundle in the category **ANFMan**. Its base is the nuclear Fréchet \mathbb{R} -manifold $M(\mathbb{R})$, i.e., the smooth manifold $|M|$, and its typical fiber is the nuclear Fréchet Λ_0 -manifold*

$$(3.26) \quad \Lambda^{p|q} := \mathring{\Lambda}_0^p \times \prod_{i=1}^N \Lambda_{\gamma_i}^{q_i} .$$

(ii) *The topology of $M(\Lambda)$, which is defined, as in the case of smooth manifolds, by the atlas providing the nuclear Fréchet Λ_0 -structure, is a Hausdorff topology, so that $M(\Lambda)$ is a genuine Fréchet manifold.*

Proof. (i) We think of fiber bundles in **ANFMan** exactly as of fiber bundles in the category of smooth manifolds. Of course, in such a fiber bundle, all objects and arrows are **ANFMan**-objects and **ANFMan**-morphisms.

Let $p^* : \Lambda \rightarrow \mathbb{R}$ be, as above, the canonical $\mathbb{Z}_2^n \mathbf{GrAlg}$ -morphism. The induced map

$$\pi := M(p^*) : M(\Lambda) \ni (x, m_\star^*) \mapsto (x, p^* \circ m_\star^*) \simeq x \in M(\mathbb{R}) \simeq |M|$$

is p^* -smooth, i.e., is a morphism in the category **ANFMan**.

We will show that π is surjective and that the local triviality condition is satisfied.

Let $z \in |M|$. There is a \mathbb{Z}_2^n -chart (U, h) of M , such that $|U| \subset |M|$ is a neighborhood of z . The \mathbb{Z}_2^n -isomorphism $h : U \rightarrow \mathcal{U}^{p|q}$ induces a natural isomorphism h , whose Λ -components are Λ_0 -diffeomorphisms, i.e., Λ_0 -smooth maps that have a Λ_0 -smooth inverse. We have the following commutative diagram:

$$\begin{array}{ccc} & h_\Lambda & \\ U(\Lambda) & \xleftarrow{\quad} & \mathcal{U}^{p|q}(\Lambda) \simeq \mathcal{U}^p \times \Lambda^{p|q} \\ U(p^*) = \pi|_{U(\Lambda)} \downarrow & & \downarrow \mathcal{U}^{p|q}(p^*) \simeq \text{prj}_1 \\ U(\mathbb{R}) \simeq |U| & \xleftarrow[h_{\mathbb{R}} = |h|]{\quad} & \mathcal{U}^{p|q}(\mathbb{R}) \simeq \mathcal{U}^p \end{array} ,$$

where prj_1 is the canonical projection. Let us explain that $\mathcal{U}^{p|q}(p^*) \simeq \text{prj}_1$, when read through $\flat : \mathcal{U}^p \times \Lambda^{p|q} \leftrightarrow \mathcal{U}^{p|q}(\Lambda)$. We need a more explicit description of the equivalent views on Λ -points of a \mathbb{Z}_2^n -domain, see beginning of Subsection 3.4. As elsewhere in this text, we denote a \mathbb{Z}_2^n -morphism $\mathbb{R}^{0|m} \rightarrow \mathcal{U}^{p|q}$ by $x = (|x|, x^*)$ and we denote the morphism it induces between the stalks $\mathcal{O}_{\mathcal{U}^{p|q}, |x|(\star)} \rightarrow \Lambda$ by x_\star^* . The morphism \flat is the succession of identifications

$$(3.27) \quad \mathcal{U}^p \times \Lambda^{p|q} \ni (x_{||}, \dot{x}_\Lambda, \xi_\Lambda) \simeq x = (|x|, x^*) \simeq (|x|(\star), x_\star^*) \in \mathcal{U}^{p|q}(\Lambda) ,$$

where the components of the base morphism $|x|$ are obtained (see [16]) by applying the base projection $\varepsilon_\star : \Lambda \rightarrow \mathbb{R}$ of $\mathbb{R}^{0|m}$, i.e., the canonical morphism p^* , to the components $x_\Lambda^a = (x_{||}^a, \dot{x}_\Lambda^a) \in \Lambda_0$. Hence, we get

$$(3.28) \quad |x|(\star) = |x| = (\dots, p^*(x_\Lambda^a), \dots) = x_{||} .$$

Therefore, we actually obtain that

$$\mathcal{U}^{p|q}(p^*)(\flat(x_{||}, \dot{x}_\Lambda, \xi_\Lambda)) = (|x|(\star), p^* \circ x_\star^*) \simeq |x|(\star) = x_{||} = \text{prj}_1(x_{||}, \dot{x}_\Lambda, \xi_\Lambda) .$$

Since $\pi|_{U(\Lambda)} = |h|^{-1} \circ \text{prj}_1 \circ h_\Lambda$, the local projection $\pi|_{U(\Lambda)}$ is surjective, so that z is in the image of π , which is thus surjective as well.

As just mentioned, we started from $z \in |M|$ and found a neighborhood $|U|$ of z and a Λ_0 -diffeomorphism h_Λ . When identifying $|U|$ with \mathcal{U}^p via $|h|$ (which then becomes id), we get the Λ_0 -diffeomorphism

$$(3.29) \quad h_\Lambda : \pi^{-1}(|U|) \simeq U(\Lambda) \ni (y, m_\star^*) \mapsto (y, m_\star^* \circ h_y^*) \in |U| \times \Lambda^{p|q}.$$

Observe that in Equation (3.29) we used \flat^{-1} defined in Equations (3.27) and (3.28), thus identifying

$$(y, m_\star^* \circ h_y^*) \in \text{Hom}_{\mathbb{Z}_2^n \text{Man}}(\mathcal{O}_{\mathcal{U}^{p|q}, y}, \Lambda) \subset \mathcal{U}^{p|q}(\Lambda)$$

with $h \circ m \in \text{Hom}_{\mathbb{Z}_2^n \text{Man}}(\mathbb{R}^{0|m}, \mathcal{U}^{p|q})$, and then with

$$(y, \text{pr}_2(m^*(h^*(x))), m^*(h^*(\xi))) \in |U| \times \Lambda^{p|q},$$

where we denoted the projection of Λ_0 onto $\mathring{\Lambda}_0$ by pr_2 . Notice also that the conclusion that $\Lambda^{p|q}$ is a nuclear Fréchet Λ_0 -manifold comes from the facts that any subspace (resp., any closed subspace) of a nuclear (resp., a Fréchet) space is a nuclear (resp., a Fréchet) space.

Hence, the trivialization condition is satisfied as well, and $M(\Lambda)$ is a fiber bundle in ANFMan , as announced.

(ii) Now consider two different Λ -points $m^* = (x, m_\star^*)$ and $n^* = (y, n_\star^*)$ in $M(\Lambda)$. If $x \neq y$, then, as $|M|$ is Hausdorff, there exist open neighborhoods $|U|$ of x and $|V|$ of y , such that $|U| \cap |V| = \emptyset$. When denoting the corresponding open \mathbb{Z}_2^n -submanifolds by U and V , respectively, we get open neighborhoods $U(\Lambda)$ and $V(\Lambda)$ of m^* and n^* , such that $U(\Lambda) \cap V(\Lambda) = \emptyset$. We have of course to check that, for any \mathbb{Z}_2^n -chart (U_α, h_α) , the image

$$h_{\alpha, \Lambda}(U_\alpha(\Lambda) \cap U(\Lambda))$$

is open in $\mathbb{R}^{p|q}(\Lambda)$, and similarly for $V(\Lambda)$. To see this, it suffices to proceed as in the proof of Theorem 3.20.

Next, consider the situation where $x = y =: z \in |M|$, use the trivialization constructed in (i), and denote the canonical projection from $\mathcal{U}^p \times \Lambda^{p|q}$ onto $\Lambda^{p|q}$ by prj_2 . As $m^* \neq n^*$, we have $h_\Lambda(m^*) \neq h_\Lambda(n^*)$, i.e.,

$$(|h|(z), \text{prj}_2(h_\Lambda(m^*))) \neq (|h|(z), \text{prj}_2(h_\Lambda(n^*))).$$

Since $\text{prj}_2(h_\Lambda(m^*)) \neq \text{prj}_2(h_\Lambda(n^*))$ are points in the Hausdorff space $\Lambda^{p|q}$, there are open neighborhoods V_{m^*} and V_{n^*} of these projections that do not intersect. The preimages U_{m^*} and U_{n^*} of V_{m^*} and V_{n^*} by the continuous map

$$\text{prj}_2 \circ h_\Lambda : U(\Lambda) \rightarrow \Lambda^{p|q}$$

are then open neighborhoods of m^* and n^* that do not intersect.

Finally, the space $M(\Lambda)$ is indeed a Hausdorff topological space. \square

3.5. The Schwarz–Voronov embedding. In order to get a fully faithful functor, hence, to embed the category $\mathbb{Z}_2^n \text{Man}$ as full subcategory into a functor category, we need to replace the target category $[\mathbb{Z}_2^n \text{Pts}^{\text{op}}, \text{A(N)FMan}]$ by a subcategory that we denote by $[\mathbb{Z}_2^n \text{Pts}^{\text{op}}, \text{A(N)FMan}]$ and that we define as follows:

Definition 3.23. The category $[\mathbb{Z}_2^n \text{Pts}^{\text{op}}, \text{A(N)FMan}]$ is the subcategory of the category $[\mathbb{Z}_2^n \text{Pts}^{\text{op}}, \text{A(N)FMan}]$,

- (i) whose objects are the functors \mathcal{F} , such that, for any $\Lambda \in \mathbb{Z}_2^n \text{Pts}^{\text{op}}$, the value $\mathcal{F}(\Lambda)$ is a (nuclear) Fréchet Λ_0 -manifold, and
- (ii) whose morphisms are natural transformations $\eta : \mathcal{F} \rightarrow \mathcal{G}$, such that, for any Λ , the component $\eta_\Lambda : \mathcal{F}(\Lambda) \rightarrow \mathcal{G}(\Lambda)$ is Λ_0 -smooth.

Proposition 3.24. The restricted Yoneda functor $\mathcal{Y}_{\mathbb{Z}_2^n \text{Pts}}$ can be considered as a faithful functor

$$\mathcal{S} : \mathbb{Z}_2^n \text{Man} \rightarrow [\mathbb{Z}_2^n \text{Pts}^{\text{op}}, \text{A(N)FMan}].$$

Proof. The image $\mathcal{Y}_{\mathbb{Z}_2^n \text{Pts}}(M)$ of an object $M \in \mathbb{Z}_2^n \text{Man}$ is a functor $M(-) \in [\mathbb{Z}_2^n \text{Pts}^{\text{op}}, \text{A(N)FMan}]$, such that, for any Λ , the value $M(\Lambda)$ is a (nuclear) Fréchet Λ_0 -manifold. Further, the image $\mathcal{Y}_{\mathbb{Z}_2^n \text{Pts}}(\phi)$ of a \mathbb{Z}_2^n -morphism $\phi : M \rightarrow N$ is a natural transformation $\phi : M(-) \rightarrow N(-)$, such that, for any Λ , the component $\phi_\Lambda : M(\Lambda) \rightarrow N(\Lambda)$ is Λ_0 -smooth.

The proof of Λ_0 -smoothness uses the following construction, which we will also need later on. Let $M, N \in \mathbb{Z}_2^n \text{Man}$ be manifolds of dimension $p|q$ and $r|s$, respectively, let $|\phi| \in C^\infty(|M|, |N|)$, and let $(|V_\beta|)_\beta$ be an open cover of $|N|$ by \mathbb{Z}_2^n -charts

$$g_\beta : V_\beta \rightarrow \mathcal{V}_\beta^{r|s}, \text{ where } V_\beta = (|V_\beta|, \mathcal{O}_N|_{|V_\beta|}).$$

The open subsets $|U_\beta| := |\phi|^{-1}(|V_\beta|) \subset |M|$ cover $|M|$, and each $|U_\beta|$ can be covered by \mathbb{Z}_2^n -charts

$$h_{\beta\alpha} : U_{\beta\alpha} \rightarrow \mathcal{U}_{\beta\alpha}^{p|q}, \text{ where } U_{\beta\alpha} = (|U_{\beta\alpha}|, \mathcal{O}_M|_{|U_{\beta\alpha}|}).$$

The \mathbb{Z}_2^n -morphism $\phi : M \rightarrow N$ restricts to a \mathbb{Z}_2^n -morphism $\phi|_{U_{\beta\alpha}} : U_{\beta\alpha} \rightarrow V_\beta$. In particular, the composite

$$g_\beta \circ \phi|_{U_{\beta\alpha}} \circ (h_{\beta\alpha})^{-1} : \mathcal{U}_{\beta\alpha}^{p|q} \rightarrow \mathcal{V}_\beta^{r|s}$$

is a \mathbb{Z}_2^n -morphism.

We now show that ϕ_Λ is Λ_0 -smooth. Therefore, let $(x, m_\star^*) \in M(\Lambda)$. There is a \mathbb{Z}_2^n -chart (V_β, g_β) of N such that $|\phi|(x) \in |V_\beta|$, and there is a \mathbb{Z}_2^n -chart $(U_{\beta\alpha}, h_{\beta\alpha})$ of M such that $x \in |U_{\beta\alpha}|$. These charts (we omit in the following the subscripts β and α) induce charts $(U(\Lambda), h_\Lambda)$ of $M(\Lambda)$ around (x, m_\star^*) , and $(V(\Lambda), g_\Lambda)$ of $N(\Lambda)$ such that $\phi_\Lambda(U(\Lambda)) \subset V(\Lambda)$. It suffices to show (see Appendix B) that the local form

$$g_\Lambda \circ \phi_\Lambda \circ (h_\Lambda)^{-1} = (g \circ \phi|_U \circ h^{-1})_\Lambda$$

is Λ_0 -smooth. This is the case in view of Theorem 3.17. This completes the proof. \square

We will prove that the functor \mathcal{S} is fully faithful, hence, injective (up to isomorphism) on objects. Therefore, it embeds the category $\mathbb{Z}_2^n\text{Man}$ of \mathbb{Z}_2^n -manifolds as full subcategory into the larger functor category $[\mathbb{Z}_2^n\text{Pts}^{\text{op}}, \mathbf{A}(\mathbb{N})\text{FMan}]$.

Definition 3.25. We refer to the faithful functor

$$\mathcal{S} : \mathbb{Z}_2^n\text{Man} \longrightarrow [\mathbb{Z}_2^n\text{Pts}^{\text{op}}, \mathbf{A}(\mathbb{N})\text{FMan}]$$

as the *Schwarz–Voronov embedding*.

Theorem 3.26. *The Schwarz–Voronov embedding \mathcal{S} is a fully faithful functor. That is, given two \mathbb{Z}_2^n -manifolds M and N , the injective map*

$$\mathcal{S}_{M,N} : \text{Hom}_{\mathbb{Z}_2^n\text{Man}}(M, N) \rightarrow \text{Hom}_{[\mathbb{Z}_2^n\text{Pts}^{\text{op}}, \mathbf{A}(\mathbb{N})\text{FMan}]}(M(-), N(-))$$

is bijective.

Proof. Notice first that it follows from the results of [14] and Lemma 3.11 that there is a 1:1 correspondence

$$|M| \simeq \text{Hom}_{\mathbb{Z}_2^n\text{Alg}}(\mathcal{O}_M(|M|), \mathbb{R}) \simeq \bigcup_{x \in |M|} \text{Hom}_{\mathbb{Z}_2^n\text{Alg}}(\mathcal{O}_{M,x}, \mathbb{R}) = M(\mathbb{R}),$$

which is given by

$$x \mapsto \varepsilon_x \mapsto (x, \varepsilon_x),$$

where ε_x is the evaluation map $\varepsilon_x(f) = (\varepsilon f)(x)$ ($f \in \mathcal{O}_M(|M|)$) and where ε is the base map $\varepsilon : \mathcal{O}_M \rightarrow C_{|M|}^\infty$. Hence, any $(x, m_\star^*) \in M(\mathbb{R})$ is equal to (x, ε_x) and can be identified with x . In view of (3.25), this 1:1-correspondence identifies the nuclear Fréchet \mathbb{R} -manifold structure on $M(\mathbb{R})$ with the smooth manifold structure on $|M|$.

Let now

$$\eta : M(-) \rightarrow N(-)$$

be a natural transformation in the target set of $\mathcal{S}_{M,N}$, i.e., a natural transformation such that, for any Λ , the Λ -component η_Λ is Λ_0 -smooth. In particular, the map

$$|\phi| := \eta_{\mathbb{R}} : |M| \rightarrow |N|,$$

is a smooth map between the reduced manifolds.

As in the proof of Proposition 3.24, let $(V_\beta, g_\beta)_\beta$ be an open cover of $|N|$ by \mathbb{Z}_2^n -charts, and, for any β , let $(U_{\beta\alpha}, h_{\beta\alpha})_\alpha$ be an open cover of $|U_\beta| := |\phi|^{-1}(|V_\beta|)$ by \mathbb{Z}_2^n -charts. When denoting the canonical \mathbb{Z}_2^n -Grassmann algebra morphism $\Lambda \rightarrow \mathbb{R}$ by p^* , we get the commutative diagram

$$(3.30) \quad \begin{array}{ccc} \bigcup_{\beta\alpha} U_{\beta\alpha}(\Lambda) & \xrightarrow{\eta_\Lambda} & \bigcup_\beta V_\beta(\Lambda) \\ M(p^*) \downarrow & & \downarrow N(p^*) \\ \bigcup_{\beta\alpha} |U_{\beta\alpha}| & \xrightarrow{|\phi|} & \bigcup_\beta |V_\beta| \end{array},$$

which shows that, for any β, α , we get the Λ_0 -smooth map

$$(\eta_\Lambda)|_{U_{\beta\alpha}(\Lambda)} : U_{\beta\alpha}(\Lambda) \rightarrow V_\beta(\Lambda).$$

Indeed, if, for $(x, m_\star^*) \in U_{\beta\alpha}(\Lambda)$, we set $\eta_\Lambda(x, m_\star^*) = (y, n_\star^*)$, the commutativity of the diagram implies that

$$y \simeq (y, p^* \circ n_\star^*) = (N(p^*) \circ \eta_\Lambda)(x, m_\star^*) = (\eta_{\mathbb{R}} \circ M(p^*))(x, m_\star^*) = \eta_{\mathbb{R}}(x, p^* \circ m_\star^*) \simeq |\phi|(x) \in |V_\beta|.$$

Therefore, the restriction

$$\eta|_{U_{\beta\alpha}(-)} : U_{\beta\alpha}(-) \rightarrow V_\beta(-)$$

is a natural transformation with Λ_0 -smooth components.

Note that

$$h_{\beta\alpha} : U_{\beta\alpha} \rightarrow \mathcal{U}_{\beta\alpha}^{p|q} \quad \text{and} \quad g_\beta : V_\beta \rightarrow \mathcal{V}_\beta^{r|s}$$

are \mathbb{Z}_2^n -isomorphisms and induce natural isomorphisms, also denoted by $h_{\beta\alpha}$ and g_β , whose components are chart diffeomorphisms

$$h_{\beta\alpha,\Lambda} : U_{\beta\alpha}(\Lambda) \rightarrow \mathcal{U}_{\beta\alpha}^{p|q}(\Lambda) \quad \text{and} \quad g_{\beta,\Lambda} : V_\beta(\Lambda) \rightarrow \mathcal{V}_\beta^{r|s}(\Lambda)$$

of nuclear Fréchet Λ_0 -manifolds. The local form

$$g_{\beta,\Lambda} \circ (\eta_\Lambda)|_{U_{\beta\alpha}(\Lambda)} \circ (h_{\beta\alpha,\Lambda})^{-1} : \mathcal{U}_{\beta\alpha}^{p|q}(\Lambda) \rightarrow \mathcal{V}_\beta^{r|s}(\Lambda)$$

of η_Λ is thus Λ_0 -smooth. In other words, any Λ -component of the natural transformation

$$(3.31) \quad \varphi_{\beta\alpha} := g_\beta \circ \eta|_{U_{\beta\alpha}(-)} \circ h_{\beta\alpha}^{-1} : \mathcal{U}_{\beta\alpha}^{p|q}(-) \rightarrow \mathcal{V}_\beta^{r|s}(-)$$

between functors associated to \mathbb{Z}_2^n -domains, is Λ_0 -smooth. It therefore follows from 3.17 that $\varphi_{\beta\alpha}$ is implemented by a \mathbb{Z}_2^n -morphism

$$\varphi_{\beta\alpha} : \mathcal{U}_{\beta\alpha}^{p|q} \rightarrow \mathcal{V}_\beta^{r|s},$$

so that the composite

$$(3.32) \quad \phi_{\beta\alpha} := g_\beta^{-1} \circ \varphi_{\beta\alpha} \circ h_{\beta\alpha} : U_{\beta\alpha} \rightarrow N$$

is a \mathbb{Z}_2^n -morphism that is defined on an open \mathbb{Z}_2^n -submanifold of M . The question is whether we can patch together these locally defined \mathbb{Z}_2^n -morphisms, which are inherited from η , and get a globally defined \mathbb{Z}_2^n -morphism $\phi : M \rightarrow N$ that induces η .

Let $\phi_{\beta\alpha}|_{U_{\beta\alpha,\nu\mu}}$ and $\phi_{\nu\mu}|_{U_{\beta\alpha,\nu\mu}}$ be the \mathbb{Z}_2^n -morphisms obtained by restriction to the open \mathbb{Z}_2^n -submanifold $U_{\beta\alpha,\nu\mu}$ with base manifold $|U_{\beta\alpha,\nu\mu}| := |U_{\beta\alpha}| \cap |U_{\nu\mu}|$. They coincide as \mathbb{Z}_2^n -morphisms, if they do as associated natural transformations, i.e., if all Λ -components of those transformations coincide. This is the case since both Λ -components are equal to $\eta_\Lambda|_{U_{\beta\alpha,\nu\mu}(\Lambda)}$. It follows that the \mathbb{Z}_2^n -algebra morphisms

$$\phi_{\beta\alpha}|_{U_{\beta\alpha,\nu\mu}}^*, \phi_{\nu\mu}|_{U_{\beta\alpha,\nu\mu}}^* : \mathcal{O}_N(|N|) \rightarrow \mathcal{O}_M(|U_{\beta\alpha,\nu\mu}|)$$

coincide. This implies that we can glue the \mathbb{Z}_2^n -algebra morphisms $\phi_{\beta\alpha}^* : \mathcal{O}_N(|N|) \rightarrow \mathcal{O}_M(|U_{\beta\alpha}|)$ and get a \mathbb{Z}_2^n -algebra morphism

$$\phi^* : \mathcal{O}_N(|N|) \rightarrow \mathcal{O}_M(|M|).$$

Indeed, for any $f \in \mathcal{O}_N(|N|)$, the $\phi_{\beta\alpha}^*(f) \in \mathcal{O}_M(|U_{\beta\alpha}|)$ are a family of \mathbb{Z}_2^n -functions on an open cover of $|M|$, which do coincide on the intersections. To see this, note that

$$(\phi_{\beta\alpha}^*(f))|_{U_{\beta\alpha,\nu\mu}} = \phi_{\beta\alpha}|_{U_{\beta\alpha,\nu\mu}}^*(f) = \phi_{\nu\mu}|_{U_{\beta\alpha,\nu\mu}}^*(f) = (\phi_{\nu\mu}^*(f))|_{U_{\beta\alpha,\nu\mu}}.$$

Hence, there is a unique global section $F \in \mathcal{O}_M(|M|)$ of the sheaf \mathcal{O}_M , such that $F|_{U_{\beta\alpha}} = \phi_{\beta\alpha}^*(f)$. The **Set**-morphism, which is defined by

$$\phi_{|N|}^* : \mathcal{O}_N(|N|) \ni f \mapsto F \in \mathcal{O}_M(|M|),$$

is actually a morphism of \mathbb{Z}_2^n -algebras. Indeed, note that

$$\rho_{|U_{\beta\alpha}|}^{|M|} \circ \phi_{|N|}^* = \phi_{\beta\alpha}^*$$

(ρ is the restriction) and observe that, for any element $|U_{\beta\alpha}|$ of the open cover of $|M|$ considered, we have

$$(\phi_{|N|}^*(f \cdot g))|_{U_{\beta\alpha}} = \phi_{\beta\alpha}^*(f) \cdot \phi_{\beta\alpha}^*(g) = (\phi_{|N|}^*(f) \cdot \phi_{|N|}^*(g))|_{U_{\beta\alpha}}.$$

The \mathbb{Z}_2^n -algebra morphism $\phi_{|N|}^*$ fully characterizes a \mathbb{Z}_2^n -morphism $\phi = (||\phi||, \phi^*) : M \rightarrow N$. We will show that ϕ induces the natural transformation η , which then completes the proof.

Since ϕ is glued from the \mathbb{Z}_2^n -morphisms $\phi_{\beta\alpha}$, we get, in view of Equations (3.31) and (3.32), in particular that

$$(3.33) \quad ||\phi|||_{U_{\beta\alpha}} = |\phi_{\beta\alpha}| = \eta_{\mathbb{R}}|_{U_{\beta\alpha}(\mathbb{R})} = |\phi||_{U_{\beta\alpha}},$$

so that $||\phi|| = |\phi|$. Further, for any $|V_\beta|$,

$$(3.34) \quad \rho_{|U_{\beta\alpha}|}^{|U_\beta|} \circ \phi_{|V_\beta|}^* = \phi_{\beta\alpha,|V_\beta|}^* : \mathcal{O}_N(|V_\beta|) \rightarrow \mathcal{O}_M(|U_{\beta\alpha}|).$$

Let now Λ be any \mathbb{Z}_2^n -Grassmann algebra and let $(x, m_\star^*) \in U_{\beta\alpha}(\Lambda)$. As $x \in |U_{\beta\alpha}|$ and $|\phi|(x) \in |V_\beta|$, it follows from Equations (3.33), (3.34), (3.31), and (3.32), that the image of (x, m_\star^*) by the Λ -component of the natural transformation induced by ϕ is

$$\phi_\Lambda(x, m_\star^*) = (|\phi|(x), m_\star^* \circ \phi_x^*) = (|\phi_{\beta\alpha}|(x), m_\star^* \circ \phi_{\beta\alpha,x}^*) = (\phi_{\beta\alpha})_\Lambda(x, m_\star^*) = \eta_\Lambda(x, m_\star^*).$$

□

The following theorem is of importance in the study of \mathbb{Z}_2^n -Lie groups.

Theorem 3.27. *The Schwarz–Voronov embedding \mathcal{S} sends \mathbb{Z}_2^n -Lie groups G to functors $\mathcal{S}(G) = G(-)$ from the category $\mathbb{Z}_2^n\text{Pts}^{\text{op}}$ of \mathbb{Z}_2^n -Grassmann algebras to the category ANFLg of nuclear Fréchet Lie groups over nuclear Fréchet algebras.*

Proof. The proof is not entirely straightforward and will be given in a paper on \mathbb{Z}_2^n -Lie groups, which is currently being written down. \square

3.6. Representability and equivalence of categories. As the Schwarz–Voronov embedding is fully faithful, the category $\mathbb{Z}_2^n\text{Man}$ can be viewed as a full subcategory of the category $[\mathbb{Z}_2^n\text{Pts}^{\text{op}}, \mathbf{A}(\mathbb{N})\text{FMan}]$. Functor categories are known to be well-suited for geometric constructions. Hence, when trying to build a \mathbb{Z}_2^n -manifold M (possibly from other \mathbb{Z}_2^n -manifolds M_ι), it is often easier to build a functor \mathcal{F} in $[\mathbb{Z}_2^n\text{Pts}^{\text{op}}, \mathbf{A}(\mathbb{N})\text{FMan}]$ (from the given \mathbb{Z}_2^n -manifolds interpreted as functors $M_\iota(-)$). However, one has then to show that \mathcal{F} can be represented by a \mathbb{Z}_2^n -manifold M , i.e., that there is a \mathbb{Z}_2^n -manifold M , such that $M(-) \simeq \mathcal{F}$.

Definition 3.28. A functor

$$\mathcal{F} \in [\mathbb{Z}_2^n\text{Pts}^{\text{op}}, \mathbf{A}(\mathbb{N})\text{FMan}]$$

is said to be *representable*, if there exists a \mathbb{Z}_2^n -manifold $M \in \mathbb{Z}_2^n\text{Man}$ (which is then unique up to unique isomorphism), such that

$$M(-) \simeq \mathcal{F} \quad \text{in } [\mathbb{Z}_2^n\text{Pts}^{\text{op}}, \mathbf{A}(\mathbb{N})\text{FMan}].$$

We define the restriction $\mathcal{F}|_{|U|}$ of a functor $\mathcal{F} \in [\mathbb{Z}_2^n\text{Pts}^{\text{op}}, \mathbf{A}(\mathbb{N})\text{FMan}]$ to an open subset $|U| \subset \mathcal{F}(\mathbb{R}) \in (\mathbb{N})\text{FMan}$.

For any $\Lambda \in \mathbb{Z}_2^n\text{GrAlg}$, let

$$p_\Lambda^* : \Lambda \longrightarrow \mathbb{R}$$

be the canonical projection, let

$$\mathcal{F}(p_\Lambda^*) : \mathcal{F}(\Lambda) \longrightarrow \mathcal{F}(\mathbb{R})$$

be the corresponding smooth map. The preimage

$$(3.35) \quad \mathcal{F}|_{|U|}(\Lambda) := (\mathcal{F}(p_\Lambda^*))^{-1}(|U|)$$

is an open (nuclear) Fréchet Λ_0 -submanifold of $\mathcal{F}(\Lambda)$.

Consider now a morphism $\varphi^* : \Lambda \longrightarrow \Lambda'$ in $\mathbb{Z}_2^n\text{GrAlg}$. As $p_{\Lambda'}^* \circ \varphi^* = p_\Lambda^*$, we get the restriction

$$(3.36) \quad \mathcal{F}|_{|U|}(\varphi^*) := \mathcal{F}(\varphi^*)|_{\mathcal{F}|_{|U|}(\Lambda)} : \mathcal{F}|_{|U|}(\Lambda) \longrightarrow \mathcal{F}|_{|U|}(\Lambda'),$$

which is a morphism in $\mathbf{A}(\mathbb{N})\text{FMan}$.

Definition 3.29. For any functor

$$\mathcal{F} \in [\mathbb{Z}_2^n\text{Pts}^{\text{op}}, \mathbf{A}(\mathbb{N})\text{FMan}]$$

and any open subset $|U| \subset \mathcal{F}(\mathbb{R})$, the *restriction of \mathcal{F} to $|U|$* is the functor

$$\mathcal{F}|_{|U|} \in [\mathbb{Z}_2^n\text{Pts}^{\text{op}}, \mathbf{A}(\mathbb{N})\text{FMan}]$$

that is defined by Equations (3.35) and (3.36).

Example 3.30. Let $M \in \mathbb{Z}_2^n\text{Man}$, let $M(-)$ be the corresponding functor, and let $|U| \subset |M| \simeq M(\mathbb{R})$ be an open subset. The restriction $M(-)|_{|U|}$ is given:

(i) on objects Λ , by

$$(3.37) \quad M(-)|_{|U|}(\Lambda) := \{(x, m_\star^*) \in M(\Lambda) : (x, p_\Lambda^* \circ m_\star^*) \simeq x \in |U|\} = U(\Lambda),$$

where $U = (|U|, \mathcal{O}_M|_{|U|})$ is the open \mathbb{Z}_2^n -submanifold of M over $|U|$, and

(ii) on morphisms $\varphi^* : \Lambda \rightarrow \Lambda'$, by

$$(3.38) \quad M(-)|_{|U|}(\varphi^*) := M(\varphi^*)|_{U(\Lambda)} = U(\varphi^*),$$

since both maps are given by

$$U(\Lambda) \ni (x, m_\star^*) \mapsto (x, \varphi^* \circ m_\star^*) \in U(\Lambda').$$

Let \mathcal{F} be representable, let M be ‘its’ representing \mathbb{Z}_2^n -manifold, and let

$$(3.39) \quad \eta : \mathcal{F} \rightarrow M(-)$$

be the corresponding natural isomorphism in $[\mathbb{Z}_2^n\text{Pts}^{\text{op}}, \mathbf{A}(\mathbb{N})\text{FMan}]$. The maps η_Λ and η_Λ^{-1} are then Λ_0 -smooth, i.e., η_Λ is a Λ_0 -diffeomorphism, for any Λ . In particular, the map $\eta_{\mathbb{R}} : \mathcal{F}(\mathbb{R}) \rightarrow M(\mathbb{R})$ is a diffeomorphism of (nuclear) Fréchet manifolds. This means that the (nuclear) Fréchet structures on $\mathcal{F}(\mathbb{R}) \simeq M(\mathbb{R})$ coincide. Further, if one identifies $\mathcal{F}(\mathbb{R}) \simeq M(\mathbb{R})$ with $|M|$, the (nuclear) Fréchet structure on $\mathcal{F}(\mathbb{R}) \simeq M(\mathbb{R})$ coincides with the smooth structure on $|M|$. We can therefore view $\mathcal{F}(\mathbb{R})$ as being the smooth manifold $|M|$. Consider now a \mathbb{Z}_2^n -atlas $(U_\alpha, h_\alpha)_\alpha$ of M . If we denote the dimension of M by $p|q$, the \mathbb{Z}_2^n -chart map h_α is a \mathbb{Z}_2^n -isomorphism

$$h_\alpha : U_\alpha \rightarrow \mathcal{U}_\alpha^{p|q}$$

valued in a \mathbb{Z}_2^n -domain of dimension $p|q$, which implies that

$$(3.40) \quad h_\alpha : U_\alpha(-) \rightarrow \mathcal{U}_\alpha^{p|q}(-)$$

is a natural isomorphism in $[\mathbb{Z}_2^n \text{Pts}^{\text{op}}, \mathbf{A}(\mathbb{N})\text{FMan}]$. In view Equations (3.39), (3.37), (3.38), and (3.40), the family $(|U_\alpha|)_\alpha$ is an open cover of $|M| \simeq \mathcal{F}(\mathbb{R})$, such that, for any α , we have

$$\mathcal{F}|_{|U_\alpha|} \simeq M(-)|_{|U_\alpha|} = U_\alpha(-) \simeq \mathcal{U}_\alpha^{p|q}(-)$$

in $[\mathbb{Z}_2^n \text{Pts}^{\text{op}}, \mathbf{A}(\mathbb{N})\text{FMan}]$.

Theorem 3.31. *A functor $\mathcal{F} \in [\mathbb{Z}_2^n \text{Pts}^{\text{op}}, \mathbf{A}(\mathbb{N})\text{FMan}]$ is representable if and only if there exists an open cover $(|U_\alpha|)_\alpha$ of $\mathcal{F}(\mathbb{R})$, such that, for each α , we have*

$$(3.41) \quad \mathcal{F}|_{|U_\alpha|} \simeq \mathcal{U}_\alpha^{p|q}(-)$$

in $[\mathbb{Z}_2^n \text{Pts}^{\text{op}}, \mathbf{A}(\mathbb{N})\text{FMan}]$, where $\mathcal{U}_\alpha^{p|q}$ is a \mathbb{Z}_2^n -domain in a fixed $\mathbb{R}^{p|q}$.

Proof. We showed already that the condition is necessary. Assume now that Condition (3.41) is satisfied, i.e., that we have natural isomorphisms

$$k_\alpha : \mathcal{F}|_{|U_\alpha|} \rightarrow \mathcal{U}_\alpha^{p|q}(-)$$

in $[\mathbb{Z}_2^n \text{Pts}^{\text{op}}, \mathbf{A}(\mathbb{N})\text{FMan}]$. This means that the Λ -components

$$k_{\alpha, \Lambda} : \mathcal{F}|_{|U_\alpha|}(\Lambda) \rightarrow \mathcal{U}_\alpha^{p|q}(\Lambda)$$

are Λ_0 -diffeomorphisms.

In particular, we have a diffeomorphism

$$|h_\alpha| := k_{\alpha, \mathbb{R}} : \mathcal{F}|_{|U_\alpha|}(\mathbb{R}) = (\mathcal{F}(p_{\mathbb{R}}^*))^{-1}(|U_\alpha|) = |U_\alpha| \rightarrow \mathcal{U}_\alpha^{p|q}(\mathbb{R}) \simeq \mathcal{U}_\alpha^p.$$

Notice that $(|U_\alpha|, |h_\alpha|)_\alpha$ can be interpreted as a smooth atlas on $|M| := \mathcal{F}(\mathbb{R})$. The direct image of the structure sheaf $\mathcal{O}_{\mathcal{U}_\alpha^{p|q}}$ over \mathcal{U}_α^p by the continuous map $|h_\alpha|^{-1} : \mathcal{U}_\alpha^p \rightarrow |U_\alpha|$ is a sheaf over $|U_\alpha|$, which we denote by \mathcal{O}_{U_α} :

$$\mathcal{O}_{U_\alpha} := (|h_\alpha|^{-1})_* \mathcal{O}_{\mathcal{U}_\alpha^{p|q}}.$$

The \mathbb{Z}_2^n -ringed space

$$U_\alpha := (|U_\alpha|, \mathcal{O}_{U_\alpha})$$

is isomorphic to the \mathbb{Z}_2^n -domain $\mathcal{U}_\alpha^{p|q}$. The isomorphism is $h_\alpha := (|h_\alpha|, h_\alpha^*)$, where h_α^* is the identity map (a composite of direct images is the direct image by the composite). In other words, we have an isomorphism of \mathbb{Z}_2^n -manifolds

$$h_\alpha : U_\alpha \rightarrow \mathcal{U}_\alpha^{p|q}.$$

Consider now an overlap $|U_{\alpha\beta}| := |U_\alpha| \cap |U_\beta| \neq \emptyset$. Omitting restrictions, we get that $k_\beta k_\alpha^{-1}$ is a natural isomorphism (in $[\mathbb{Z}_2^n \text{Pts}^{\text{op}}, \mathbf{A}(\mathbb{N})\text{FMan}]$)

$$k_{\beta\alpha} := k_\beta k_\alpha^{-1} : \mathcal{U}_{\alpha\beta}^{p|q}(-) \rightarrow \mathcal{U}_{\beta\alpha}^{p|q}(-)$$

between functors corresponding to \mathbb{Z}_2^n -domains (defined as usual). In view of Theorem 3.17, the natural isomorphism $k_{\beta\alpha}$ is implemented by a \mathbb{Z}_2^n -isomorphism

$$k_{\beta\alpha} : \mathcal{U}_{\alpha\beta}^{p|q} \rightarrow \mathcal{U}_{\beta\alpha}^{p|q}.$$

It follows that

$$\psi_{\beta\alpha} := h_\beta^{-1} k_{\beta\alpha} h_\alpha : U_{\alpha\beta} \rightarrow U_{\beta\alpha}$$

is an isomorphism of \mathbb{Z}_2^n -manifolds, where $U_{\alpha\beta} := (|U_{\alpha\beta}|, \mathcal{O}_{U_{\alpha\beta}}|_{|U_{\alpha\beta}|})$. The \mathbb{Z}_2^n -manifolds U_α can thus be glued and provide then a \mathbb{Z}_2^n -manifold M over $|M| = \mathcal{F}(\mathbb{R})$, such that there are \mathbb{Z}_2^n -isomorphisms $(|U_\alpha|, \mathcal{O}_M|_{|U_\alpha|}) \rightarrow U_\alpha$, if the $\psi_{\beta\alpha}$ satisfy the cocycle condition.

Since the Schwarz–Voronov embedding is fully faithful, we have that $\psi_{\gamma\beta}\psi_{\beta\alpha} = \psi_{\gamma\alpha}$ as \mathbb{Z}_2^n -morphisms if and only if the induced natural transformations coincide. However, for any Λ , we get

$$(\psi_{\gamma\beta}\psi_{\beta\alpha})_\Lambda = (h_{\gamma, \Lambda})^{-1} k_{\gamma, \Lambda} (k_{\beta, \Lambda})^{-1} h_{\beta, \Lambda} (h_{\beta, \Lambda})^{-1} k_{\beta, \Lambda} (k_{\alpha, \Lambda})^{-1} h_{\alpha, \Lambda} = \psi_{\gamma\alpha, \Lambda}.$$

It remains to show that M actually represents \mathcal{F} , i.e., that we can find a natural isomorphism $\eta : M(-) \rightarrow \mathcal{F}$ in the category $[\mathbb{Z}_2^n \text{Pts}^{\text{op}}, \mathbf{A}(\mathbb{N})\text{FMan}]$, i.e., that, for any $\Lambda \in \mathbb{Z}_2^n \text{GrAlg}$, there is a Λ_0 -diffeomorphism $\eta_\Lambda : M(\Lambda) \rightarrow \mathcal{F}(\Lambda)$ that is natural in Λ . As $(|U_\alpha|)_\alpha$ is an open cover of $|M|$, the source decomposes as

$$M(\Lambda) = \bigcup_\alpha U_\alpha(\Lambda),$$

the $U_\alpha(\Lambda)$ being open (nuclear) Fréchet Λ_0 -submanifolds. On any $U_\alpha(\Lambda)$, we define η_Λ by setting

$$\eta_\Lambda|_{U_\alpha(\Lambda)} := (k_{\alpha, \Lambda})^{-1} h_{\alpha, \Lambda} : U_\alpha(\Lambda) \rightarrow \mathcal{F}|_{|U_\alpha|}(\Lambda) \subset \mathcal{F}(\Lambda).$$

These restrictions provide a well-defined map

$$\eta_\Lambda : M(\Lambda) \rightarrow \mathcal{F}(\Lambda).$$

Indeed, if $(x, m_\star^*) \in U_\alpha(\Lambda) \cap U_\beta(\Lambda)$, we have

$$(k_{\alpha, \Lambda})^{-1}(h_{\alpha, \Lambda}(x, m_\star^*)) = (k_{\beta, \Lambda})^{-1}(h_{\beta, \Lambda}(x, m_\star^*)) \quad \text{if and only if} \quad \psi_{\beta, \alpha}(x, m_\star^*) = (x, m_\star^*).$$

However, since we glued M from the U_α , the gluing \mathbb{Z}_2^n -isomorphisms $\psi_{\beta, \alpha}$ became identities and so did the induced natural isomorphisms. The definition of η_Λ^{-1} is similar. The source $\mathcal{F}(\Lambda)$ decomposes as

$$\mathcal{F}(\Lambda) = \bigcup_\alpha \mathcal{F}|_{|U_\alpha|}(\Lambda),$$

the $\mathcal{F}|_{|U_\alpha|}(\Lambda)$ being open (nuclear) Fréchet Λ_0 -submanifolds. On any $\mathcal{F}|_{|U_\alpha|}(\Lambda)$, we define η_Λ^{-1} by setting

$$\eta_\Lambda^{-1}|_{\mathcal{F}|_{|U_\alpha|}(\Lambda)} := (h_{\alpha, \Lambda})^{-1}k_{\alpha, \Lambda} : \mathcal{F}|_{|U_\alpha|}(\Lambda) \rightarrow U_\alpha(\Lambda) \subset M(\Lambda).$$

The condition for these restrictions to give a well-defined map

$$\eta_\Lambda^{-1} : \mathcal{F}(\Lambda) \rightarrow M(\Lambda)$$

is equivalent to the condition for η_Λ . Clearly, the maps η_Λ and η_Λ^{-1} are inverses. Naturality and Λ_0 -smoothness are local questions and are therefore consequences of the naturality and the Λ_0 -smoothness of $(k_{\alpha, \Lambda})^{-1}h_{\alpha, \Lambda}$ and of $(h_{\alpha, \Lambda})^{-1}k_{\alpha, \Lambda}$. \square

We are now prepared to show that the category $\mathbb{Z}_2^n\text{Man}$ is equivalent to a functor category.

Theorem 3.32. *The category $\mathbb{Z}_2^n\text{Man}$ of \mathbb{Z}_2^n -manifolds (defined as \mathbb{Z}_2^n -ringed spaces that are locally isomorphic to \mathbb{Z}_2^n -domains) and \mathbb{Z}_2^n -morphisms (defined as morphisms of \mathbb{Z}_2^n -ringed spaces) is equivalent to the full subcategory $[\mathbb{Z}_2^n\text{Pts}^{\text{op}}, \mathbf{A}(\mathbf{N})\text{FMan}]_{\text{rep}}$ of representable functors in $[\mathbb{Z}_2^n\text{Pts}^{\text{op}}, \mathbf{A}(\mathbf{N})\text{FMan}]$.*

In other words, the category $\mathbb{Z}_2^n\text{Man}$ is equivalent to the category of locally trivial functors in the subcategory of the functor category $[\mathbb{Z}_2^n\text{Pts}^{\text{op}}, \mathbf{A}(\mathbf{N})\text{FMan}]$, whose objects \mathcal{F} have values $\mathcal{F}(\Lambda)$ in (nuclear) Fréchet Λ_0 -manifolds and whose morphisms are the natural transformations with Λ_0 -smooth components.

Remark 3.33. This result is reminiscent of the identification of schemes with those contravariant functors from affine schemes to sets that are sheaves (for the Zariski topology on affine schemes) and have a cover by open immersions of affine schemes.

Proof. The Schwarz–Voronov embedding viewed as functor valued in $[\mathbb{Z}_2^n\text{Pts}^{\text{op}}, \mathbf{A}(\mathbf{N})\text{FMan}]_{\text{rep}}$ is obviously fully faithful and essentially surjective. It thus induces an equivalence of categories. \square

APPENDIX A. GENERATING SETS OF CATEGORIES

We will freely use Mac Lane’s book [36] as our source of categorical notions and proofs of general statements. For completeness, we recall the concept of generating set of a category.

Definition A.1 ([36], page 127). Let \mathbf{C} be a category. A set $S = \{S_i \in \text{Ob}(\mathbf{C}) : i \in I\}$, where I is any index set, is said to be a *generating set* of \mathbf{C} , if, for any pair of distinct \mathbf{C} -morphisms

$$\phi, \psi : A \longrightarrow B,$$

i.e., $\phi \neq \psi$, there exists some $i \in I$ and a \mathbf{C} -morphism

$$s : S_i \longrightarrow A,$$

such that the compositions

$$S_i \xrightarrow{s} A \rightrightarrows \begin{matrix} \phi \\ \psi \end{matrix} B,$$

are distinct, i.e., $\phi \circ s \neq \psi \circ s$. In this case, we say that the object S_i *separates* the morphisms ϕ and ψ , and that the set S *generates* the category \mathbf{C} .

Example A.2. The set $\{\mathbb{R}\}$ is a generating set of the category of finite dimensional real vector spaces. This is easily seen, as, if we have two distinct linear maps $\phi, \psi : V \rightarrow W$, then there exists a vector $v \in V$ ($v \neq 0$), such that $\phi(v) \neq \psi(v)$. Thus, the two linear maps differ on the one dimensional subspace generated by v . Now let z be a basis of \mathbb{R} . Then, the linear map $s : \mathbb{R} \rightarrow V$ given by $s(z) = v$, keeps ϕ and ψ separate.

Proposition A.3. *For any locally small category \mathbf{C} , a set $S \subset \text{Ob}(\mathbf{C})$ generates \mathbf{C} if and only if the restricted Yoneda embedding*

$$\mathcal{Y}_S : \mathbf{C} \rightarrow [S^{\text{op}}, \mathbf{Set}],$$

where S is viewed as full subcategory of \mathbf{C} , is faithful.

Proof. The restricted embedding is defined on objects by

$$\mathcal{Y}_S(A) = \mathbf{Hom}_{\mathbf{C}}(-, A) \in [S^{\text{op}}, \mathbf{Set}]$$

and on morphisms by

$$\mathcal{Y}_S(\phi) = \mathbf{Hom}_{\mathbf{C}}(-, \phi) : \mathcal{Y}_S(A) \rightarrow \mathcal{Y}_S(B) ,$$

where

$$(\mathcal{Y}_S(\phi))_{S_i} : \mathbf{Hom}_{\mathbf{C}}(S_i, A) \ni s \mapsto \phi \circ s \in \mathbf{Hom}_{\mathbf{C}}(S_i, B) .$$

The embedding \mathcal{Y}_S is faithful if and only if, for any different $\phi, \psi : A \rightarrow B$, the corresponding natural transformations are distinct, i.e., there is at least one $i \in I$ and one $s \in \mathbf{Hom}_{\mathbf{C}}(S_i, A)$, such that $\phi \circ s \neq \psi \circ s$. \square

APPENDIX B. FRÉCHET SPACES, MODULES AND MANIFOLDS

Manifolds over algebras A , also known as A -manifolds, are manifolds for which the tangent spaces are endowed with a module structure over a given finite-dimensional commutative algebra. For details, the reader may consult Shurygin [51, 52, 53], and for a discussion of the specific case of (the even part of) Grassmann algebras one may consult Azarmi [7]. A comprehensive introduction to the subject can be found in the book by Vishnevskii, Shirokov, and Shurygin [57] (in Russian). The concept needed in this paper is a infinite-dimensional generalisation of an A -manifold to the category of Fréchet algebras and Fréchet manifolds. For an introduction to locally convex spaces, including Fréchet vector spaces, we refer the reader to Conway [15, Chapter IV], Trèves [56, Part I], or Rudin [44, Chapter 1]. A brief introduction to Fréchet algebras can be found in Waelbroeck [59, Chapter VII]. For Fréchet manifolds, the reader can consult Saunders [46, Chapter 7] and Hamilton [29, Part I.4].

Definition B.1. A *Fréchet (vector) space* is a complete Hausdorff metrizable locally convex topological vectorspace.

There exist a few other, equivalent, definitions of Fréchet spaces. The topology on a locally convex space is metrizable if and only if it can be derived from a countable family of semi-norms $\| - \|_k$, $k \in \mathbb{N}$. The topology is Hausdorff if and only if the family of semi-norms is separating, i.e., if $\|x\|_k = 0$, for all k , implies $x = 0$. Given such a family of semi-norms, one defines a translationally invariant metric that induces the topology by setting

$$d(x, y) = \sum_{k=0}^{\infty} 2^{-k} \frac{\|x - y\|_k}{1 + \|x - y\|_k} ,$$

for all x and y .

Example B.2. Let $M = (|M|, \mathcal{O})$ be a \mathbb{Z}_2^n -manifold. For any open subset $U \subset |M|$, the space $\mathcal{O}(U)$ of \mathbb{Z}_2^n -functions f on U is a Fréchet space. An inducing family of semi-norms is given by

$$\|f\|_{C,D} = \sup_{x \in C} |\varepsilon(D(f))(x)| ,$$

where ε is the projection $\varepsilon : \mathcal{O}(U) \rightarrow C^\infty(U)$ of \mathbb{Z}_2^n -functions to base functions, where C is any compact subset of U , and where D is any \mathbb{Z}_2^n -differential operator over U [13].

Given two Fréchet spaces $(F, (\| - \|_k^F)_{k \in \mathbb{N}})$ and $(G, (\| - \|_k^G)_{k \in \mathbb{N}})$, a linear map

$$\phi : F \longrightarrow G ,$$

is continuous if and only if, for every semi-norm $\| - \|_k^G$, there exists a semi-norm $\| - \|_l^F$ and a positive real number $C > 0$, such that

$$\|\phi(x)\|_k^G \leq C \|x\|_l^F ,$$

for every $x \in F$. A similar result holds for continuous bilinear maps

$$\phi : F \times G \rightarrow H .$$

The *morphisms of Fréchet spaces* are the continuous linear maps, so that the category of Fréchet spaces is a full subcategory of the category of topological vector spaces.

What makes Fréchet spaces interesting, is the fact that they have just enough structure to define a derivative of a mapping between such spaces. This leads to a meaningful notion of a smooth map between Fréchet spaces, and so much of finite dimensional differential geometry can be transferred to the infinite dimensional setting, using Fréchet spaces as local models. The well known *Gâteaux (directional) derivative* is defined as follows.

Definition B.3. Let F and G be Fréchet spaces and $U \subset F$ be open, and let $\phi : U \rightarrow G$ be a (nonlinear) continuous map. Then the *derivative of ϕ in the direction of $v \in F$ at $x \in U$* is defined as

$$d_x \phi(v) := \lim_{t \rightarrow 0} \frac{\phi(x + tv) - \phi(x)}{t}$$

provided the limit exists. We say that ϕ is *continuously differentiable*, if the limit exists for all $x \in U$ and $v \in F$, and if the mapping

$$d\phi : U \times F \longrightarrow G$$

is (jointly) continuous.

Higher order derivatives are defined inductively, i.e.,

$$d_x^{k+1} \phi(v_1, v_2, \dots, v_{k+1}) := \lim_{t \rightarrow 0} \frac{d_{x+tv_{k+1}}^k \phi(v_1, v_2, \dots, v_k) - d_x^k \phi(v_1, v_2, \dots, v_k)}{t}.$$

A continuous map $\phi : U \rightarrow G$ is then said to be k times continuously differentiable or to be of class C^k , if

$$d^k \phi : U \times F^{\times k} \longrightarrow G$$

is continuous (or, more explicitly, if all its derivatives of order $\leq k$ exist everywhere and are continuous). If ϕ is of class C^k , its derivative $d_x^k \phi(v_1, v_2, \dots, v_k)$ is multilinear and symmetric in $F^{\times k}$ [50]. Furthermore, we say that ϕ is *smooth*, if it is of class C^k , for all k .

Proposition B.4. Let F_1, F_2 be Fréchet spaces and let $U \subset F_1 \times F_2$ be an open subset. A continuous map $\phi : U \rightarrow G$ valued in a Fréchet space G is of class C^1 if and only if its (total) derivative

$$d\phi : U \times (F_1 \times F_2) \ni ((f_1, f_2), (v_1, v_2)) \mapsto d_{(f_1, f_2)} \phi(v_1, v_2) \in G$$

is continuous, which is the case if and only if the naturally defined partial derivatives

$d_{f_1} \phi : U \times F_1 \ni ((f_1, f_2), v_1) \mapsto d_{f_1, (f_1, f_2)} \phi(v_1) \in G$ and $d_{f_2} \phi : U \times F_2 \ni ((f_1, f_2), v_2) \mapsto d_{f_2, (f_1, f_2)} \phi(v_2) \in G$ are continuous. In this case, we have

$$d_{(f_1, f_2)} \phi(v_1, v_2) = d_{f_1, (f_1, f_2)} \phi(v_1) + d_{f_2, (f_1, f_2)} \phi(v_2).$$

The Gâteaux or Fréchet–Gâteaux derivative gives a rather weak notion of differentiation, however, most of the standard results from calculus in the finite dimensional setting remain true. Specifically, the fundamental theorem of calculus and the chain rule still hold. However, the inverse function theorem is in general lost. For a special class of Fréchet spaces, known as ‘tame’ Fréchet spaces, there is an analogue of the inverse function theorem known as the Nash–Moser inverse function theorem, see Hamilton [29] for details.

A *nuclear space* is a locally convex topological vector space F , such that, for any locally convex topological vector space G , the natural map $F \widehat{\otimes}_\pi G \longrightarrow F \widehat{\otimes}_\iota G$ from the projective to the injective tensor product of F and G is an isomorphism of locally convex topological vector spaces. In particular, a *nuclear Fréchet space* is a locally convex topological vector space that is a nuclear space and a Fréchet space. Loosely, if a space F is nuclear, then, for any locally convex space G , the complete topological vector space $F \widehat{\otimes} G$ is independent of the locally convex topology considered on $F \otimes G$. Because of this, and their nice dual properties, nuclear spaces provide a reasonable setting for infinite dimensional analysis. All the Fréchet spaces we encounter in this paper are in fact nuclear.

The following definition is standard.

Definition B.5. A *Fréchet algebra* is a Fréchet vector space A , which is equipped with an associative bilinear and (jointly) continuous multiplication $\cdot : A \times A \rightarrow A$. If $(p_i)_{i \in I}$ is a family of semi-norms that induces the topology on A , (joint) continuity is equivalent to the existence, for any $i \in I$, of $j \in I$, $k \in I$, and $C > 0$, such that

$$p_i(x \cdot y) \leq C p_j(x) p_k(y), \quad \forall x, y \in A.$$

We can always consider an equivalent increasing countable family of semi-norms $(\|\cdot\|_n)_{n \in \mathbb{N}}$. The preceding condition then requires that, for any $n \in \mathbb{N}$, there is $r \in \mathbb{N}$ ($r \geq n$) and $C > 0$, such that

$$\|x \cdot y\|_n \leq C \|x\|_r \|y\|_r \quad \forall x, y \in A.$$

In particular, the topology can be induced by a countable family of submultiplicative semi-norms, i.e., by a family $(q_n)_{n \in \mathbb{N}}$ that satisfies

$$q_n(x \cdot y) \leq q_n(x) q_n(y), \quad \forall n \in \mathbb{N}, \forall x, y \in A.$$

Note that many authors define a Fréchet algebra simply as a Fréchet vector space, which carries an associative bilinear multiplication, and whose topology can be induced by a countable family of submultiplicative semi-norms. This latter definition is equivalent to the former.

In general, a Fréchet algebra need not be unital, and, if it is, one does not require $p_i(1_A) = 1$, in contrast to what is usually required for Banach algebras.

Example B.6 (Formal power series). Consider the space

$$\mathbb{R}[[z_1, z_2, \dots, z_q]]$$

of formal power series in q parameters and coefficients in reals. We set $j := (j_1, j_2, \dots, j_q) \in \mathbb{N}^{\times q}$ and $|j| := j_1 + j_2 + \dots + j_q$. A general series x now reads

$$x = \sum_j z^j a_j = \sum_j z_1^{j_1} z_2^{j_2} \dots z_q^{j_q} a_{j_q \dots j_2 j_1},$$

with no question on the convergence. The algebra structure is the standard multiplication of formal power series. The topology of coordinate-wise convergence is metrizable and given by the family of semi-norms

$$\|x\|_k := \sum_{|j| \leq k} |a_j|, \quad \forall k \in \mathbb{N}.$$

This algebra is unital with the obvious unit, and it is submultiplicative.

Let us denote the category of *Fréchet algebras* (resp., *commutative Fréchet algebras*) as \mathbf{FAlg} (resp., \mathbf{CAlg}). Morphisms in this category are defined to be continuous algebra morphisms. If we restrict attention to nuclear Fréchet algebras (resp., commutative nuclear Fréchet algebras), then we work in the full subcategory \mathbf{NAlg} (resp., \mathbf{CNAlg}).

Definition B.7. Fix $A \in \mathbf{FAlg}$. A *Fréchet A -module* is a Fréchet vector space F , together with a continuous action

$$\begin{aligned} A \times F &\xrightarrow{\mu} F \\ (a, v) &\mapsto \mu(a, v), \end{aligned}$$

which we will write as $\mu(a, v) := a \cdot v$ (and which is of course compatible with the multiplication in A).

We give a short survey on Fréchet manifolds.

Definition B.8. Let \mathcal{M} be a set. An F -chart of \mathcal{M} is a bijective map $\phi : U \rightarrow \phi(U) \subset F$, where $U \subset \mathcal{M}$ and $\phi(U)$ is an open subset of a Fréchet space F .

A Fréchet atlas can be defined using charts valued in various Fréchet spaces. For our purposes, it is sufficient to consider a fixed Fréchet model.

Definition B.9. Let \mathcal{M} be a set. A *smooth F -atlas* on \mathcal{M} is a collection of F -charts $((U_\alpha, \phi_\alpha))_{\alpha \in \mathcal{A}}$, such that

- (i) the subsets U_α cover the set \mathcal{M} ,
- (ii) the subsets $\phi_\alpha(U_\alpha \cap U_\beta)$ are open in F ,
- (iii) the transition maps

$$\phi_{\beta\alpha} := \phi_\beta \circ \phi_\alpha^{-1} : \phi_\alpha(U_\alpha \cap U_\beta) \subset F \longrightarrow \phi_\beta(U_\beta \cap U_\alpha) \subset F$$

are smooth.

A new F -chart (U, ϕ) on \mathcal{M} is *compatible* with a given smooth F -atlas, if and only if their union is again a smooth F -atlas, i.e., the subsets $\phi(U \cap U_\alpha) \subset F$ and $\phi_\alpha(U_\alpha \cap U) \subset F$ are open, and the transition maps

$$\phi_\alpha \circ \phi^{-1} : \phi(U \cap U_\alpha) \longrightarrow \phi_\alpha(U_\alpha \cap U) \quad \text{and} \quad \phi \circ \phi_\alpha^{-1} : \phi_\alpha(U_\alpha \cap U) \longrightarrow \phi(U \cap U_\alpha)$$

are smooth (for every $\alpha \in \mathcal{A}$). Similarly, two smooth F -atlases are compatible provided their union is also a smooth F -atlas. Compatibility is an equivalence relation on all possible smooth F -atlases on \mathcal{M} .

Definition B.10. A *smooth F -structure* on a set \mathcal{M} is a choice of an equivalence class of smooth F -atlases on \mathcal{M} . We say that \mathcal{M} is a *Fréchet manifold* modelled on the Fréchet space F , if \mathcal{M} comes equipped with a smooth F -structure. If the model vector space F is nuclear, we speak of a *nuclear Fréchet manifold*.

A smooth F -atlas on a Fréchet manifold \mathcal{M} allows us to define in the obvious way a topology on \mathcal{M} , which is independent of the atlas considered in the chosen equivalence class. The domain U of an F -chart (U, ϕ) is open in this topology and the bijective map $\phi : U \subset \mathcal{M} \rightarrow \phi(U) \subset F$ is a homeomorphism for the induced topologies. Most authors confine themselves to Fréchet manifolds, whose topology is Hausdorff.

Morphisms between two Fréchet manifolds are the *smooth maps* between them, where smoothness is defined, just as in the finite dimensional case of smooth manifolds, in terms of charts and smoothness of local representatives of the maps. We denote the category of Fréchet manifolds and the morphisms between them by \mathbf{FMan} .

Further, the *tangent space* $\mathbf{T}_f \mathcal{M}$ to a Fréchet manifold \mathcal{M} at a point $f \in \mathcal{M}$ can be defined as usual, using the tangency equivalence relation for the smooth curves of \mathcal{M} that pass through f at time 0. One can easily see that $\mathbf{T}_f \mathcal{M}$ is a Fréchet space. The concept of *Fréchet vector bundle* is the natural generalization of the notion

of smooth vector bundle to the category of Fréchet manifolds. The *tangent bundle* $\mathbf{T}\mathcal{M}$ of a Fréchet manifold \mathcal{M} is an example of a Fréchet vector bundle.

In general, we must make a distinction between the (*kinematic*) *tangent bundle* as defined here and the *operational tangent bundle* defined in terms of derivations of the algebra of functions of a Fréchet manifold. Indeed, the two notions do not, in general, coincide, there are derivations that do not correspond to tangent vectors. However, it is known that for nuclear Fréchet manifolds the two concepts do coincide.

Let $\mathfrak{F} : \mathcal{M} \rightarrow \mathcal{N}$ be a smooth map between Fréchet manifolds modelled on Fréchet spaces F and G , respectively. There is a *tangent map* $\mathbf{T}\mathfrak{F}$ of \mathfrak{F} , which is a smooth map

$$\mathbf{T}\mathfrak{F} : \mathbf{T}\mathcal{M} \rightarrow \mathbf{T}\mathcal{N},$$

and restricts, for any $f \in \mathcal{M}$, to a linear map

$$\mathbf{T}_f\mathfrak{F} : \mathbf{T}_f\mathcal{M} \rightarrow \mathbf{T}_{\mathfrak{F}(f)}\mathcal{N}.$$

As in the finite dimensional case, the local representative of $\mathbf{T}_f\mathfrak{F}$ is the derivative $d_{\phi(f)}(\psi\mathfrak{F}\phi^{-1})$ of the corresponding local representative

$$\psi\mathfrak{F}\phi^{-1} : \phi(U) \subset F \rightarrow G$$

of \mathfrak{F} at the point $\phi(f)$.

Fundamental to the work in this paper are Fréchet manifolds with a further module structure on their tangent bundle.

Definition B.11. Let \mathcal{M} be a Fréchet manifold, whose model Fréchet space F is a module over a Fréchet algebra A . We say that \mathcal{M} is a *Fréchet A-manifold*, if and only if all transition maps are A -linear, i.e.,

$$d_{\phi_\alpha(f)}\phi_{\beta\alpha}(a \cdot v) = a \cdot d_{\phi_\alpha(f)}\phi_{\beta\alpha}(v),$$

for all $f \in U_\alpha \cap U_\beta$, $a \in A$, and $v \in F$.

Morphisms between Fréchet A-manifolds \mathcal{M} and \mathcal{N} are the A -smooth maps between them, i.e., are the smooth maps $\mathfrak{F} : \mathcal{M} \rightarrow \mathcal{N}$ that are A -linear at every point. This means that, for any point $f \in \mathcal{M}$, there is an \mathcal{M} -chart (U, ϕ) around f and an \mathcal{N} -chart (V, ψ) around $\mathfrak{F}(f)$ that contains $\mathfrak{F}(U)$, such that the local representative

$$d_{\phi(f)}(\psi\mathfrak{F}\phi^{-1})$$

of the derivative $\mathbf{T}_f\mathfrak{F}$ is an A -linear endomorphism of the A -module F . The requirement actually means that the derivative $\mathbf{T}_f\mathfrak{F}$ must be A -linear at any point $f \in \mathcal{M}$. In this way, we obtain the category of Fréchet A -manifolds, which we denote as **AFMan**.

In this paper, we will use the category **AFMan**, whose objects are the Fréchet A -manifolds, where A is not a fixed Fréchet algebra, but any Fréchet algebra. The definition of **AFMan**-morphisms generalizes the definition of **AFMan**-morphisms. Suppose that \mathcal{M} is a Fréchet A -manifold modelled on an A -module F and \mathcal{N} is a Fréchet B -manifold modelled on a B -module G . The **AFMan**-morphisms from \mathcal{M} to \mathcal{N} are the A -smooth maps between them, i.e., those smooth maps $\mathfrak{F} : \mathcal{M} \rightarrow \mathcal{N}$ that are at any point compatible with the module structures of F and G . This means that there is a Fréchet algebra morphism $\rho : A \rightarrow B$, and, for any $f \in \mathcal{M}$, there exist charts (U, ϕ) and (V, ψ) as above, such that

$$d_{\phi(f)}(\psi\mathfrak{F}\phi^{-1})(a \cdot v) = \rho(a) \cdot d_{\phi(f)}(\psi\mathfrak{F}\phi^{-1})(v),$$

for any $a \in A$ and $v \in F$. This requirement actually means that, for any f , the derivative $\mathbf{T}_f\mathfrak{F}$ is compatible with the induced actions on the tangent spaces. We will refer to an A -smooth map with associated Fréchet algebra morphism ρ , as a ρ -smooth map. If we restrict our attention to nuclear objects, i.e, the model Fréchet vector space and the Fréchet algebra are both nuclear, then we denote the corresponding category as **ANFMan**.

REFERENCES

- [1] N. Aizawa, P.S. Isaac & J. Segar, $\mathbb{Z}_2 \times \mathbb{Z}_2$ generalizations of $\mathcal{N} = 1$ superconformal Galilei algebras and their representations, preprint, 1808.09112 [[math-ph](#)] (2018).
- [2] N. Aizawa, Z. Kuznetsova, H. Tanaka & F. Toppan, $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded Lie symmetries of the Lévy-Leblond equations, *Prog. Theo. Exp. Phys.* **12** (2016), 123A01.
- [3] N. Aizawa & J. Segar, $\mathbb{Z}_2 \times \mathbb{Z}_2$ generalizations of $\mathcal{N} = 2$ super Schrödinger algebras and their representations, *J. Math. Phys.* **58** (2017), no. 11, 113501, 14 pp.
- [4] H. Albuquerque & S. Majid, Quasialgebra structure of the octonions, *J. Alg.* **220** (1999), 188–224.
- [5] H. Albuquerque & S. Majid, Clifford algebras obtained by twisting of group algebras, *J. Pure Appl. Alg.* **171** (2002), 133–148.
- [6] A. Alldridge & M. Laubinger, Infinite-dimensional supermanifolds over arbitrary base fields, *Forum Math.* **24** (2012), no. 3, 565–608.
- [7] S. Azarmi, Foliations associated with the structure of a manifold over a Grassmann algebra of exterior forms of even degree, *Russian Math.* **56** (2012), no. 1, 76–78.
- [8] L. Balduzzi, C. Carmeli & R. Fioresi, The local functors of points of supermanifolds, *Expo. Math.* **28** (2010), no. 3, 201–217.

- [9] L. Balduzzi, C. Carmeli & R. Fioresi, A comparison of the functors of points of supermanifolds, *J. Alg. Appl.* **12** (2013), no. 3, 1250152, 41 pp.
- [10] F.A. Berezin, “Introduction to superanalysis”, vol. 9 of Math. Phys. and Appl. Math., *D. Reidel Publishing Co.*, Dordrecht, 1987.
- [11] A.J. Bruce, On a \mathbb{Z}_2^n -graded version of supersymmetry, *Symmetry* **11**(1) (2019), 116.
- [12] A.J. Bruce & E. Ibarguengoytia, The Graded Differential Geometry of Mixed Symmetry Tensors, to appear in *Archivum Mathematicum* (2019).
- [13] A.J. Bruce & N. Poncin, Functional analytic issues in \mathbb{Z}_2^n -Geometry, to appear in *Revista de la Unión Matemática Argentina*.
- [14] A.J. Bruce & N. Poncin, Products in the category of \mathbb{Z}_2^n -manifolds, *J. Nonlin. Math. Phys.* **26** (2019), no. 3.
- [15] J.B. Conway, *A course in functional analysis*, second edition, *Graduate Texts in Mathematics*, vol. 96, Springer-Verlag, New York, 1990. xvi+399 pp.
- [16] T. Covolo, J. Grabowski & N. Poncin, The category of \mathbb{Z}_2^n -supermanifolds, *J. Math. Phys.* **57** (2016), no. 7, 073503, 16 pp.
- [17] T. Covolo, J. Grabowski & N. Poncin, Splitting theorem for \mathbb{Z}_2^n -supermanifolds, *J. Geo. and Phys.* **110** (2016), 393–401.
- [18] T. Covolo, S. Kwok & N. Poncin, Differential calculus on \mathbb{Z}_2^n -supermanifolds, preprint, [arXiv:1608.00949 \[math.DG\]](https://arxiv.org/abs/1608.00949) (2016).
- [19] T. Covolo, S. Kwok & N. Poncin, The Frobenius theorem for \mathbb{Z}_2^n -supermanifolds, preprint, [arXiv:1608.00961 \[math.DG\]](https://arxiv.org/abs/1608.00961) (2016).
- [20] T. Covolo, V. Ovsienko & N. Poncin, Higher trace and Berezinian of matrices over a Clifford algebra, *J. Geo. and Phys.* **62** (2012), no. 11, 2294–2319.
- [21] K. Drühl, R. Haag & J.E. Roberts, On parastatistics, *Comm. Math. Phys.* **18** (1970), 204–226.
- [22] M. Fairon, Introduction to graded geometry, *Eur. J. Math.* **3** (2017), no. 2, 208–222.
- [23] R. Fioresi & F. Zanchetta, Representability in supergeometry, *Expo. Math.* **35** (2017), no. 3, 315–325.
- [24] R. Goodman & N.R. Wallach, *Symmetry, representations, and invariants*, vol. 255 of *Graduate Texts in Mathematics*, Springer, Dordrecht, 2009.
- [25] H.S. Green, A generalized method of field quantization, *Phys. Rev.* **90** (1953), 270.
- [26] O.W. Greenberg & A.M.L. Messiah, Selection rules for parafields and the absence of paraparticles in nature, *Phys. Rev. (2)* **138** (1965).
- [27] A. Grothendieck, *Introduction to functorial algebraic geometry, part 1: affine algebraic geometry*, summer school in Buffalo, 1973, lecture notes by Federico Gaeta.
- [28] A. Grothendieck, *Topological vector spaces*, Notes on Mathematics and its Applications, Gordon and Breach Science Publishers, New York-London-Paris, 1973.
- [29] R.S. Hamilton, The inverse function theorem of Nash and Moser, *Bull. Amer. Math. Soc. (N.S.)* **7** (1982), no. 1, 65–222.
- [30] F. Hanisch, A Supermanifold Structure on Spaces of Morphisms between Supermanifolds, [1406.7484 \[math.DG\]](https://arxiv.org/abs/1406.7484) (2014).
- [31] I. Kolář, P.W. Michor & J. Slovák, *Natural operations in differential geometry*, Springer-Verlag, Berlin, 1993. vi+434 pp.
- [32] A. Konechny & A. Schwarz, On $(k \oplus l|q)$ -dimensional supermanifolds, in *Supersymmetry and quantum field theory* (Kharkov, 1997), 201–206, *Lecture Notes in Phys.*, **509**, Springer, Berlin, 1998.
- [33] A. Konechny & A. Schwarz, Theory of $(k \oplus l|q)$ -dimensional supermanifolds, *Selecta Math. (N.S.)* **6** (2000), no. 4, 471–486.
- [34] A. Kriegl & P.W. Michor, *The convenient setting of global analysis*, Mathematical Surveys and Monographs, 53, American Mathematical Society, Providence, RI, 1997. x+618 pp.
- [35] D. Leites (ed.), Seminar on supersymmetry (v. 1. Algebra and calculus: Main chapters) (J. Bernstein, D. Leites, V. Molotkov, V. Shander), MCCME, Moscow, 2011, 410 pp. (in Russian; an English version is in preparation but available for perusal).
- [36] S. Mac Lane, *Categories for the working mathematician*, vol. 5 of *Graduate Texts in Mathematics*, Springer-Verlag, New York, second edition, 1998.
- [37] S. Mac Lane, & I. Moerdijk, *Sheaves in geometry and logic. A first introduction to topos theory*. Corrected reprint of the 1992 edition, *Universitext*, Springer-Verlag, New York, 1994.
- [38] K.-H. Neeb, Towards a Lie theory of locally convex groups, *Jpn. J. Math.* **1** (2006), no. 2, 291–468.
- [39] M. Mohammadi & H. Salmasian, The Gelfand-Naimark-Segal construction for unitary representations of \mathbb{Z}_2^n -graded Lie supergroups, in: *50th Seminar “Sophus Lie”*, 263–274, Banach Center Publ., 113, *Polish Acad. Sci., Inst. Math., Warsaw*, 2017.
- [40] N. Poncin, Towards integration on colored supermanifolds, *Banach Center Publ.* **110** (2016), 201–217.
- [41] V. Rittenberg & D. Wyler, Generalized superalgebras, *Nuclear Phys. B.* **139** (1978), no. 3, 189–202.
- [42] V. Rittenberg & D. Wyler, Sequences of $\mathbb{Z}_2 \oplus \mathbb{Z}_2$ graded Lie algebras and superalgebras, *J. Math. Phys.* **19** (1978), no. 10, 2193–2200.
- [43] A. Rogers, *Supermanifolds. Theory and applications*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2007. xii+251 pp.
- [44] W. Rudin, *Functional analysis*, second edition, *International Series in Pure and Applied Mathematics*, McGraw-Hill, Inc., New York, 1991. xviii+424 pp.
- [45] C. Sachse, Global Analytic Approach to Super Teichmüller Spaces, PhD thesis, Universität Leipzig, 2007.

- [46] D.J. Saunders, *The geometry of jet bundles*, London Mathematical Society, Lecture Note Series, 142, Cambridge University Press, Cambridge, 1989. viii+293 pp.
- [47] T. Schmitt, Supergeometry and quantum field theory, or: what is a classical configuration? *Rev. Math. Phys.* **9** (1997), no. 8, 993–1052.
- [48] Jakob Schütt, Infinite-Dimensional Supermanifolds via Multilinear Bundles, [arXiv:1810.05549 \[math.DG\]](https://arxiv.org/abs/1810.05549) (2018).
- [49] A.S. Schwarz, Supergravity, complex geometry and G-structures, *Comm. Math. Phys.* **87** (1982/83), no. 1, 37–63.
- [50] V. V. Sharko, Functions on Manifolds: Algebraic and Topological Aspects, Amer. Math. Soc. (1993).
- [51] V.V. Shurygin, On the cohomology of manifolds over local algebras, *Russian Math.* **40** (1996), no. 9, 67–81.
- [52] V.V. Shurygin, The structure of smooth mappings over Weil algebras and the category of manifolds over algebras, *Lobachevskii J. Math.* **5** (1999), 29–55.
- [53] V.V. Shurygin, Smooth manifolds over local algebras and Weil bundles, *J. Math. Sci.* **108** (2002), no. 2, 249–294.
- [54] A.S. Shvarts, On the definition of superspace, *Teoret. Mat. Fiz.* **60** (1984), no. 1, 37–42.
- [55] V.N. Tolstoy, Super-de Sitter and alternative super-Poincaré symmetries, in: Lie theory and its applications in physics, Selected papers based on the presentations at the 10th international workshop, LT 10, Varna, Bulgaria, June 17–23, 2013.
- [56] F. Trèves, *Topological vector spaces, distributions and kernels*, Academic Press, New York-London, 1967.
- [57] V.V. Vishnevskii, A.P. Shirokov & V.V. Shurygin, *Spaces over algebras* (Russian), Kazanskii Gosudarstvennyi Universitet, Kazan, 1985. 264 pp.
- [58] A. Voronov, Maps of supermanifolds, *Teoret. Mat. Fiz.* **60** (1984), no. 1, 43–48.
- [59] L. Waelbroeck, *Topological vector spaces and algebras*, Lecture Notes in Mathematics, vol. 230, Springer-Verlag, Berlin-New York, 1971. vii+158 pp.
- [60] A. Weil, Théorie des points proches sur les variétés différentiables, in: *Géométrie différentielle*, Colloques Internationaux du Centre National de la Recherche Scientifique, Strasbourg, 1953, pp. 111–117.
- [61] W. Yang & S. Jing, A new kind of graded Lie algebra and parastatistical supersymmetry, *Sci. China Ser. A* **44** (2001), no. 9, 1167–1173.

MATHEMATICS RESEARCH UNIT, UNIVERSITY OF LUXEMBOURG, MAISON DU NOMBRE, 6 AVENUE DE LA FONTE, L-4364 ESCH-SUR-ALZETTE, LUXEMBURG

Email address: andrewjamesbruce@googlemail.com, eduardo.ibarguengoytia@uni.lu, norbert.poncin@uni.lu