

Adult disinhibited social engagement in adoptees exposed to extreme institutional deprivation: examination of its clinical status and functional impact

Mark Kennedy, Jana Kreppner, Nicky Knights, Robert Kumsta, Barbara Maughan, Dennis Golm, Jonathan Hill, Michael Rutter, Wolff Schlotz and Edmund Sonuga-Barke

Background

Early-life institutional deprivation produces disinhibited social engagement (DSE). Portrayed as a childhood condition, little is known about the persistence of DSE-type behaviours into, presentation during, and their impact on, functioning in adulthood.

Aims

We examine these issues in the young adult follow-up of the English and Romanian Adoptees study.

Method

A total of 122 of the original 165 Romanian adoptees who had spent up to 43 months as children in Ceaușescu's Romanian orphanages and 42 UK adoptees were assessed for DSE behaviours, neurodevelopmental and mental health problems, and impairment between ages 2 and 25 years.

Results

Young adult DSE behaviour was strongly associated with early childhood deprivation, with a sixfold increase for those who spent more than 6 months in institutions. However, although DSE overlapped with autism spectrum disorder and attention-deficit hyperactivity disorder symptoms it was not, in itself, related to broader patterns of mental health

problems or impairments in daily functioning in young adulthood.

Conclusions

DSE behaviour remained a prominent, but largely clinically benign, young adult feature of some adoptees who experienced early deprivation.

Declaration of interest

Over the past 3 years E.S.-B. has received speaker fees, consultancy, research funding and conference support from Shire Pharma and speaker fees from Janssen Cilag. He has received consultancy fees from Neurotech solutions, Aarhus University, Copenhagen University and Berhanderling, Skolerne, Copenhagen, KU Leuven. J.K. has received honoraria for invited presentations at professional associations from Association of Child and Adolescent Mental Health and the Association of Young People's Health.

Copyright and usage

© The Royal College of Psychiatrists 2017. This is an open access article distributed under the terms of the Creative Commons Attribution (CC BY) licence.

Adversity experienced early in childhood can create deep psychological and neurobiological vulnerabilities.¹ In this regard, institutional care settings pose rather particular risks and have distinctive effects on development.² In such settings there may be limited personalised care, restricting the chance for children to develop close, stable and loving relationships with individual carers.³ There may also be a lack of linguistic and intellectual stimulation.⁴ In more extreme cases, nutrition and hygiene are compromised.³ The effects of these institutional exposures can be seen in subsequent elevated rates of neurodevelopmental disorders and mental health problems – with more severe institutional deprivation associated with worse outcomes persisting into adulthood.⁵ Some of the difficulties observed in individuals who have been previously institutionalised represent deprivation-related variants of more general, common childhood problems, such as attention-deficit hyperactivity disorder (ADHD),⁶ autism spectrum disorder (ASD)⁷ and behavioural and emotional problems.⁸ In contrast, disinhibited social engagement (DSE, as defined in DSM-5⁹) is a diagnosis restricted to individuals who have experienced institutional deprivation or other forms of gross early social neglect. It presents as an inappropriate over-familiarity with, and lack of wariness of, strangers, and a failure to observe appropriate physical and verbal boundaries during social interaction. For example, it can involve an inappropriate seeking out of physical proximity and/or asking overly intrusive and overly

familiar questions.^{10,11} Within the social ecology of poor-quality care in institutions, DSE behaviours may be adaptive – they may fulfil a need for social intimacy and increase access to care from staff.¹² However, post-institutionally they represent a potential source of vulnerability and impairment.¹³ Individuals with DSE may be perceived by peers as tiresome and/or hostile, and the indiscriminate nature of their social behaviours may undermine friendship development.¹⁴ Their overly trusting nature may increase risk of exploitation within dysfunctional relationships.¹¹

DSE is considered a condition of early childhood and although recent evidence suggests that DSE can persist into later childhood and adolescence,^{15–18} little is known about its long-term persistence into adulthood.¹⁹ To address this point, we have recently reported childhood-to-adult developmental trajectories for DSE behaviours in the English and Romanian Adoptees (ERA) study cohort²⁰ of individuals who experienced extreme early deprivation in the institutions of the latter years of the Ceaușescu regime, before adoption into UK families.⁵ Perhaps against expectation, DSE behaviours showed strong persistence into adulthood alongside more expected persistence of ASD and inattention/overactivity (representing a subset of symptoms of ADHD) symptoms – 35% of young adult adoptees who experienced deprivation for longer than the first 6 months of life displayed evidence of DSE behaviours compared with 6% in a comparison group of UK adoptees. In the current paper, we

extend this initial analysis of the persistence of DSE behaviours. Our aim is to address the following questions. What are the characteristic features of the young adult DSE presentation in individuals who have experienced early deprivation? Can DSE behaviours still be directly observed? To what extent are affected adults aware of their own DSE behaviours? Do DSE behaviours overlap with continuing neurodevelopmental disorders? Does young adult DSE have an impact on social functioning, mental health, service use and quality of life (QoL)? Given prior findings we predicted that DSE behaviours would persist into adulthood but be less observable than at younger ages and affected individuals would show some awareness of their own tendency towards DSE. We also predicted that DSE behaviours would be associated with impaired social functioning and reduced QoL.

We were also interested in exploring the relationship between DSE behaviours and parent-child relationships – especially their link with attachment security in adulthood. This is of interest because the DSE pattern was initially conceptualised in terms of attachment. However, against this view, DSE has since been shown to persist even after the formation of secure attachment relationships.¹¹ Our own earlier data suggest that the core of the DSE pattern is linked to unmodulated and indiscriminate social behaviours with relative strangers rather than a lack of selectivity of attachment behaviours with regard to carers. At age 6, for instance, there was only a weak negative association between DSE and attachment security; 53% of those displaying DSE were coded as secure compared with 79% of those without the DSE pattern.¹¹ As a result of this Rutter *et al*²¹ concluded that the DSE pattern should not be conceptualised as an attachment disorder – a view consistent with the work of Zeanah and colleagues,¹⁰ and the eventual designation of DSE as a separate disorder in DSM-5.⁹

Method

Participants

In total, 165 Romanian (91 females) and 52 UK adoptees (18 females – no deprivation history) and their adoptive families entered the study in the mid-1990s. In the UK group, data were available for 47 (90%) and 39 (75%) individuals in adolescence and in young adulthood respectively. In the Romanian group these figures were 148 (90%) and 107 (65%) individuals respectively. The average age at young adult assessment was 23.2 (UK) and 23.6 years (Romanian). A comparison of age 15 characteristics of those that dropped out in young adulthood and those that remained in the study provided no evidence of selective attrition. The proportion of UK and Romanian individuals was similar at the two ages and there was no difference between those remaining in the study at the young adult follow-up and those dropping out in terms of gender of child, age, IQ or the proportion with deprivation-related problems at age 15 years (data available from authors on request). A total of 18.4% of the retained sample met the criteria for DSE whereas 21.6% of those who dropped out did. This difference was non-significant ($P=0.648$).

Measures

DSE

The assessment of DSE behaviours was based on interviewers' ratings of parents' answers based on three codes used at previous assessment waves.^{7,11,18} These were: (a) too friendly with strangers or too eager to approach strangers; (b) makes very personal comments or asked intrusive questions of others they have just met; and (c) unaware of social boundaries, or the closeness of interaction with unfamiliar people. Two additional questions

were included in the young adult assessment: (d) excessive self-disclosure and (e) not stranger-aware. A rating between 0 and 2 was made – with 0 representing 'no evidence of disinhibition' and 2 'definite evidence of disinhibition'. Interrater agreement in adulthood was high, with a mean intraclass correlation of 0.91 (range 0.81–0.97). Interviewers masked to prior DSE status also directly observed the adoptees' behaviour during their assessment visit for evidence of the following: (a) social disinhibition, (b) inappropriate physical contact, (c) inappropriate/intrusive comments, (d) violation of physical social boundaries and (e) violation of verbal social boundaries. Ratings were made using the same coding scheme as for the parent-based ratings.

Self-awareness of their DSE behaviours by young adults was assessed using specific DSE items from the Social Emotional Functioning Interview.²² These items were: general social disinhibition with others; a lack of the concept of friendship; an inability to differentiate between friends and non-friends; a lack of awareness of any problems with peers/others; and a lack of the concept of loneliness. Items were rated on a three-point scale (0–2) with the higher rating reflecting definite difficulty. Difficulties were deemed present when any item was positively endorsed at the level of 1 or 2 (i.e. some or definite). Given the non-specific nature of social difficulties, this scale had reasonable internal consistency ($\alpha=0.58$). This scale also had high interrater reliability between two raters on a randomly selected 20% subsample, with an average kappa of 0.890 (range 0.71–1). Further information on individual items used available on request from the authors.

Co-occurring young adult developmental and mental health problems

ADHD, conduct and mood problems. ADHD (18 items), generalised anxiety (13 items), major depression (15 items) and conduct disorder (14 items) symptoms were assessed using the Conners Behaviour Rating Scales (CBRS) using standardised T-scores.²³ ADHD assessment was based on parent report whereas the other outcomes were based on self-report, in line with established guidelines.²⁴ Callous-unemotional traits were measured using the parent report Inventory of Callous-Unemotional traits (ICU),²⁵ which measures the affective personality features of psychopathy. It contains 24 items assessed on a 0- to 3-point rating scale with higher scores reflecting increased levels of callous-unemotional traits. The ICU had good internal consistency ($\alpha=0.90$).

ASD. The Social Communication Questionnaire (SCQ)²⁶ is a parent-completed and clinically validated 35-item screen for ASD symptoms that maps onto DSM diagnostic criteria. To ensure its developmental appropriateness in young adulthood our analysis was based on a 15-item version with five items from each scale (social reciprocal interaction; communication and repetitive and stereotyped behaviours).⁵ Items were rated as either 0 for 'No' or 1 for 'Yes'. This revised 15-item scale demonstrated high reliability ($\alpha=0.88$).

IQ. A shortened version of the Wechsler Abbreviated Scale of Intelligence (WASI, two-subscale (vocabulary and block design) version)²⁷ was administered in early adulthood.

Clinical service use

Information about lifetime history of clinical service use was gathered from parents and young people during interviews at ages 6, 11 and 15 years and during the young adult follow-up.

Young adult-parent attachment security

Young adults' attachment was measured using the Parental Attachment Questionnaire (PAQ)²⁸ and the Inventory of Parent and Peer Attachment (IPPA)²⁹ – validated in this age group.³⁰ PAQ has three subscales: affective quality (27 items), facilitation of independence (14 items) and support (13 items), with items rated on a scale of 1 to 5. In the present sample, scale reliability was high for each subscale ($\alpha=0.96$, 0.90 and 0.85 respectively). IPPA is a 25-item measure of mother- and father-child attachment. Items are rated on a scale of 1 to 5, with higher scores indicating greater attachment security. Both mother and father attachment subscales showed high scale reliability in this sample ($\alpha=0.97$ and 0.96 respectively).

QoL

Participants self-rated QoL using the five-item Satisfaction with Life Scale³¹ (1, strongly disagree to 5, strongly agree). They reported whether they (a) see their current life as ideal; (b) are satisfied with life; (c) would live life again in the same way; (d) find life excellent and; (d) think they have secured the important things in life. This scale had high internal consistency ($\alpha=0.90$).

Functioning within different social domains

Social functioning was assessed with the work, romantic relationships, friendships, non-specific social contacts and coping subscales of the Revised Adult Personality Functioning Assessment (RAPFA),³² each on a scale of 1–9 – where a score of 3 or more represents the presence of dysfunction.³³ Interrater reliability was high, with intraclass correlations ranging from 0.78 to 0.96, based on a randomly selected 20% of the whole sample. Participants were grouped as 0 for 'no significant dysfunction' or 1 for 'evidence of significant dysfunction'.

Education and employment status

Key indicators of young adult functioning were derived from young adult and parent reports. These were (a) currently being unemployed and (b) having lower educational achievement (i.e. GCSEs or less). These were coded in a binary form (0, does not apply; 1, applies).

Adoptive family demographics

Adoptive family socioeconomic status (SES) was based on data on parents' occupation at the age 15 follow-up.²⁰ Families were divided into high and low SES groups based on the Registrar General's classification.³⁴ Manual and unskilled occupations were classified as low SES and skilled, managerial/technical and professional occupations as high SES. Parental marital status was derived from parental reports made at young adult follow-up to create a binary code (marriage intact, 1; original adoptive parents divorced, separated or widowed, 0).

Statistical analysis

First, we used correlational analysis to examine the structure of parent-rated young adult DSE. Second, we used χ^2 and Fisher's exact tests to examine the relationship between deprivation and DSE. As in previous papers we divided the Romanian adoptees into two groups – those who had experienced less than 6 months of deprivation (Rom<6 group) and those who experienced between 6 and 43 months deprivation (Rom>6 group). This was based on evidence for a step increase in risk between 6 and 12 months of deprivation exposure and the finding of equally

low risk of problems in the UK and the Rom<6 groups.³⁵ These groups were compared in terms of the proportion of individuals with a score of 1 (probable DSE) for each of the five parent-rated items. Third, we estimated the proportion of individuals meeting criteria for DSE based on interviewer coding of parent interviews using thresholds equivalent to those used in previous studies.¹¹ More specifically, DSE was deemed present if an individual had two or more positive endorsements (a score of either 1 or 2) from the five items. These proportions were then compared across the three groups. Fourth, we split the Rom>6 group into those with and without DSE (DSE+ v. DSE–) and then validated this DSE categorisation against (a) interviewer observations and (b) self-ratings. For these measures, separate binary outcomes were created so that those with no self-reported or observed disinhibition could be compared with those with at least one positive item. Fifth, we examined patterns of continuity between adolescence and young adulthood using McNemar's statistical test, which compared the proportion of individuals meeting threshold at our two time points. Sixth, using the same group comparison we examined patterns of co-occurring neurodevelopmental, mental health problems and clinical service use as well as QoL, functioning within different social domains, employment and educational status. Seventh, where the above analysis revealed significant differences between DSE+ and DSE– groups we ran a series of multiple logistic regression analyses with other deprivation-specific problems (ADHD, ASD), callous-unemotional traits and IQ entered as covariates to assess the extent to which poor outcomes were specific to DSE.

Results

The five parent-reported adult DSE items were significantly inter-correlated (Spearman's $rs=0.39$ to 0.83) creating a single scale with high internal consistency ($\alpha=0.82$). Figure 1 presents the proportion of positive DSE-item endorsements for young adults within the UK, Rom<6 and Rom>6 groups. Adult DSE behaviours were strongly related to levels of prior deprivation. There were equally low rates in the UK and Rom<6 groups ($\chi^2(1)=0.07$, $P=0.79$), but they were significantly greater in the Rom>6 group compared with the Rom<6 group ($\chi^2(1)=11.8$, $P=0.001$). Individuals in the Rom>6 group were seven and six times more likely to meet our thresholds than those in either the UK ($\chi^2(1)=11.8$, $P=0.001$) or the Rom<6 group respectively ($\chi^2(1)=12.1$, $P=0.001$). Furthermore, there was significant persistence of DSE behaviours, with 64% of those in the Rom>6 group meeting criteria for DSE at 15 also meeting criteria in young adulthood (McNemar, $P=0.79$).

Given the similarity in terms of low parent-derived DSE levels in the UK and Rom<6 groups we then tested whether these groups differed in terms of self-rated and observed DSE behaviours: there were no significant differences ($P=0.054$ and $P=0.34$ respectively). The UK and Rom<6 groups were therefore combined for subsequent analyses into a low deprivation group (LoDep group). This group was then compared with the Rom>6 group divided into DSE+ v. DSE– groups. Figure 2 compares these three groups in terms of observed and self-rated DSE behaviours. The Rom>6 group DSE+ had significantly higher levels of any self-rated (46.7% v. 4.9%; Fisher's Exact, $P<0.001$) and observed DSE (41.2% v. 1.2%, Fisher's Exact, $P<0.001$).

Table 1 presents group differences in family demographics and levels of young adult co-occurring neurodevelopmental problems, mental health symptoms, clinical service use, educational attainment, employment status, functioning within different social domains and QoL. There was no difference between the




Fig. 1 The proportion of individuals in each group scoring positively for each disinhibited social engagement (DSE) trait (on left) and the proportion of individuals who met criteria for the presence of DSE overall as coded by the interviewer based on parental descriptions (on right).

UK, UK adoptees; <6, Romanian adoptees with less than 6 month of deprivation; >6, Romanian adoptees with more than 6 months of deprivation.

DSE- and DSE+ groups in terms of gender, SES or the intactness of the adoptive family. Levels of ASD and ADHD symptoms and callous-unemotional traits and clinical service use were significantly greater in the Rom>6 DSE+ group than either the LoDep or the Rom>6 DSE- group. The same pattern was seen for clinical service use, lower educational attainment and unemployment. DSE+ and DSE- did not differ in terms of anxiety, depression or IQ. The groups did not differ in terms of QoL or RAPFA measures of functioning within any social domains.

Multiple logistic regression analyses, with service use, educational attainment and unemployment as dependent variables showed that associations of DSE with service use ($\text{Exp}(B) = 1.52$, $P = 0.64$), low educational attainment ($\text{Exp}(B) = 2.55$, $P = 0.31$) or unemployment ($\text{Exp}(B) = 1.03$, $P = 0.97$) all dropped to non-significant levels when ASD and ADHD symptoms, callous-unemotional traits and IQ were

added as predictors alongside DSE. Interestingly, the effects relating to service use were driven by the specific association with co-occurring ADHD symptoms ($\text{Exp}(B) = 1.14$, $P = 0.011$). For employment, IQ drove the association ($\text{Exp}(B) = 0.91$, $P = 0.019$). ADHD symptoms and IQ accounted jointly for educational performance; (ADHD: $\text{Exp}(B) = 1.10$, $P = 0.035$; IQ: $\text{Exp}(B) = 0.92$, $P = 0.044$). Finally, DSE was not related to adult attachment security, whether measured via the IPPA or the PAQ ($P > 0.05$; see Table 1).

Discussion

Main findings

The term DSE has been used to describe an inappropriate, overfamiliar and socially intrusive pattern of behaviour displayed by children who have experienced early social adversity – especially neglect.³⁶ Past research has shown DSE to be at the core of a pattern of deprivation-specific problems observed in children and adolescents who have experienced institutional deprivation including those in the ERA study.⁷ In such individuals we recently reported the persistence of DSE behaviours into adulthood alongside ADHD and ASD symptoms.⁵ Here we present the first detailed characterisation of deprivation-related DSE behaviours in adulthood. There were a number of findings of note.

First, each of the five parent-reported young adult DSE descriptions were similarly elevated in individuals who experienced more than 6 months of deprivation – including the new items ‘lack of stranger wariness’ and ‘excessive self-disclosure’ – items that extend the DSE construct in potentially important ways. In particular, an overtrusting nature and willingness to divulge private and confidential information could increase the risk of abuse and exploitation.

Second, high DSE status, established by pooling across interviewer-codings of parent-derived information, was corroborated by investigator-based observations and self-ratings by the young adults themselves. This provided external validation of the young adult DSE construct. This is important because it is possible that parental descriptions could be conditioned by the child’s history of DSE behaviours rather than reflecting the current situation.




Fig. 2 The proportion of individuals with and without disinhibited social engagement (DSE) who met criteria for self-reported and observed social disinhibition.

LoDep, combined UK adoptees with Romanian adoptees with less than 6 months of deprivation. DSE-, Romanian adoptees with more than 6 months of deprivation not meeting DSE criteria; DSE+, Romanian adoptees with more than 6 months of deprivation meeting DSE criteria

Table 1 Demographic characteristics and clinical outcomes

	Rom > 6 group			Group contrasts		
	LoDep (n = 92)	DSE- (n = 51)	DSE+ (n = 21)	LoDep v. DSE-	LoDep v. DSE+	DSE- v. DSE+
Female, n (%)	43 (46.7)	24 (47.1)	15 (71.4)	$\chi^2 (1) = 0.00$ $P = 0.971$	$\chi^2 (1) = 4.17$ $P = 0.041$	$\chi^2 (1) = 3.56$ $P = 0.059$
Low socioeconomic status, ^a n (%)	11 (12.2)	5 (10.4)	4 (21.1)	$\chi^2 (1) = 0.10$ $P = 0.752$	$\chi^2 (1) = 1.03$ $P = 0.310$	$\chi^2 (1) = 1.32$ $P = 0.250$
Intact families, n (%)	71 (78.0)	30 (75.0)	13 (61.9)	$\chi^2 (1) = 0.14$ $P = 0.705$	$\chi^2 (1) = 2.36$ $P = 0.124$	$\chi^2 (1) = 1.14$ $P = 0.287$
Neurodevelopmental problems, mean (s.d.)						
ADHD symptoms ^b	50.47 (10.57)	56.24 (14.39)	73.55 (12.18)	$t(57.96) = -2.24$ $P = 0.029$	$t(102) = -8.67$ $P < 0.001$	$t(58) = -4.68$ $P < 0.001$
ASD symptoms ^c	0.93 (2.12)	1.73 (2.76)	3.98 (3.36)	$t(120) = -1.77$ $P = 0.79$	$t(22.77) = -3.87$ $P < 0.001$	$t(57) = -2.74$ $P = 0.008$
IQ	102.68 (16.10)	96.36 (11.03)	91.00 (14.21)	$t(95.72) = 2.43$ $P = 0.79$	$t(90) = 2.62$ $P = 0.017$	$t(49) = 1.45$ $P = 0.153$
Callous-unemotional traits ^d	19.59 (10.43)	21.41 (11.99)	32.84 (13.22)	$t(118) = -0.85$ $P = 0.79$	$t(100) = -4.83$ $P < 0.001$	$t(56) = -3.33$ $P = 0.002$
Mental health in early adulthood, mean (s.d.)						
Depression ^b	54.29 (13.96)	58.22 (12.03)	63.31 (18.22)	$t(106) = -1.44$ $P = 0.152$	$t(19.10) = -1.86$ $P = 0.078$	$t(21.03) = -1.02$ $P = 0.318$
Anxiety ^b	54.15 (13.63)	58.50 (11.44)	62.69 (16.67)	$t(107) = -1.55$ $P = 0.102$	$t(87) = -2.18$ $P = 0.032$	$t(21.53) = -0.92$ $P = 0.371$
Conduct disorder ^b	46.38 (10.78)	47.25 (9.33)	53.31 (16.54)	$t(106) = -0.42$ $P = 0.679$	$t(17.93) = -1.60$ $P = 0.126$	$t(19.38) = -1.37$ $P = 0.186$
Clinical service use, young adulthood: n (%)	15 (16.7)	12 (30.8)	13 (68.4)	$\chi^2 (1) = 3.27$ $P = 0.071$	$\chi^2 (1) = 22.01$ $P < 0.011$	$\chi^2 (1) = 7.39$ $P = 0.007$
Young adult functioning, quality of life, ^e mean (s.d.)	16.92 (5.38)	16.47 (6.21)	16.87 (3.96)	$t(102) = -0.37$ $P = 0.710$	$t(85) = 0.03$ $P = 0.973$	$t(40.52) = -0.27$ $P = 0.792$
RAPFA – social domain dysfunction, n (%)						
Work	23 (28.4)	24 (55.8)	11 (68.8)	$\chi^2 (1) = 8.97$ $P = 0.003$	$\chi^2 (1) = 9.56$ $P = 0.002$	$\chi^2 (1) = 0.81$ $P = 0.369$
Romantic relationships	37 (46.3)	27 (64.3)	12 (80.0)	$\chi^2 (1) = 3.59$ $P = 0.058$	$\chi^2 (1) = 5.76$ $P = 0.116$	$\chi^2 (1) = 1.26$ $P = 0.261$
Friendships	20 (24.7)	24 (55.8)	10 (62.5)	$\chi^2 (1) = 11.89$ $P = 0.001$	$\chi^2 (1) = 8.94$ $P = 0.003$	$\chi^2 (1) = 0.21$ $P = 0.644$
Non-specific social contacts	11 (13.6)	10 (23.3)	7 (43.8)	$\chi^2 (1) = 1.87$ $P = 0.172$	$\chi^2 (1) = 8.05$ $P = 0.005$	$\chi^2 (1) = 2.39$ $P = 0.122$
Coping	18 (23.1)	17 (40.5)	8 (53.3)	$\chi^2 (1) = 4.00$ $P = 0.045$	$\chi^2 (1) = 5.72$ $P = 0.017$	$\chi^2 (1) = 0.74$ $P = 0.389$
Unemployed	11 (12.0)	12 (23.5)	14 (66.7)	$\chi^2 (1) = 2.26$ $P = 0.071$	$\chi^2 (1) = 29.70$ $P < 0.001$	$\chi^2 (1) = 12.00$ $P = 0.001$
Low education (GCSE or less)	24 (26.4)	15 (29.4)	14 (66.7)	$\chi^2 (1) = 0.15$ $P = 0.700$	$\chi^2 (1) = 12.36$ $P < 0.001$	$\chi^2 (1) = 8.58$ $P = 0.003$
Attachment security, mean (s.d.)						
IPPA mother	98.26 (19.06)	95.98 (27.29)	93.11 (23.48)	$t(43.36) = 0.43$ $P = 0.672$	$t(83) = 0.91$ $P = 0.366$	$t(45) = 0.35$ $P = 0.727$
IPPA father	96.71 (20.32)	99.96 (21.79)	94.62 (18.36)	$t(96) = -0.72$ $P = 0.473$	$t(77) = 0.33$ $P = 0.741$	$t(41) = 0.75$ $P = 0.457$
PAQ affective quality	108.38 (19.72)	103.85 (26.67)	101.15 (26.72)	$t(50.89) = 0.88$ $P = 0.382$	$t(84) = 1.21$ $P = 0.230$	$t(47) = 0.33$ $P = 0.746$
PAQ facilitators of independence	51.52 (11.08)	51.84 (14.19)	48.22 (12.53)	$t(96) = -0.13$ $P = 0.899$	$t(84) = 1.03$ $P = 0.308$	$t(47) = 0.85$ $P = 0.398$
PAQ source of support	45.32 (9.49)	44.07 (11.31)	43.67 (9.51)	$t(103) = 0.59$ $P = 0.557$	$t(84) = 0.61$ $P = 0.543$	$t(47) = 0.12$ $P = 0.904$

LoDep, combined UK adoptees with Romanian adoptees with less than 6 months of deprivation. DSE-, Romanian adoptees with >6 months of deprivation individuals not meeting DSE criteria; DSE+, Romanian adoptees with more than 6 months of deprivation meeting DSE criteria; ADHD, attention-deficit hyperactivity disorder; ASD, autism spectrum disorder; RAPFA, Revised Adult Personality Functioning Assessment; IPPA, Inventory of Parent and Peer Attachment; PAQ, Parental Attachment Questionnaire.

a. Based on family occupational status at age 15.

b. Self-rated Conners Behaviour Rating Scales T-score.

c. Parent-rated Social Communication Questionnaire.

d. Parent-rated.

e. Self-rated Satisfaction with Life Scale score.

Note: n's may not sum due to missing data. Where degrees of freedom have decimal places, corrected values have been used to account for unequal variances.

The fact that specific DSE behaviours could be observed during a short interview session is quite striking given the expectation for social conformity in such situations. Noteworthy here is that none of the interviewers of young adults had visited participants in the past phases of the project. The evidence that individuals with DSE had insight into their own gregarious nature and open social

interpersonal style is also potentially important – because self-knowledge can represent a necessary, although not sufficient, condition for the development of coping and defensive strategies that can reduce social risks.

Third, DSE behaviours still strongly overlapped with the two other core deprivation-specific problems features (ASD and

ADHD symptoms), providing further evidence for the idea of a loosely configured deprivation syndrome as previously proposed by Rutter *et al.*¹⁹ The fourth element of the original deprivation-specific problems cluster, cognitive impairment, was more weakly related to DSE than in the previous follow-up assessments – largely because in general individuals in the Rom>6 group displayed cognitive recovery by young adulthood with IQs in the normal range even for the most deprived.⁵ We also provide the first evidence that adult DSE behaviours overlap to a degree with callous–unemotional traits. Callous–unemotional traits were examined in the sample for the first time at the adolescent follow-up. We have recently reported that callous–unemotional traits are also linked to ADHD in young adults who were exposed to early deprivation.⁶ These data together raise the possibility that callous–unemotional traits might be a core feature of a deprivation syndrome alongside ADHD, ASD and DSE – at least in adulthood.

Fourth, and perhaps most strikingly, adult DSE behaviours appeared to be largely clinically benign. They were unrelated to both the development of anxiety and depression, despite the increase in these aspects of mental health problems observed between adolescence and early adulthood more generally⁵ and appeared not to have a significant impact on either functioning within different social domains or QoL – at least as perceived by the young adults themselves. Where there were effects (on educational and employment status) it appeared that these were due to co-occurring neurodevelopmental problems (particularly ADHD symptoms and low IQ), rather than to DSE itself. This lack of an association with impairment casts doubt on whether, at least in adulthood, DSE on its own should be considered a disorder as characterised in the DSM-5 revisions. It is even possible that certain features of disinhibited social behaviours may be beneficial in certain settings and occupations. Indeed, in discussion with the young adults with DSE it became clear that some of them had found a successful niche in jobs in for instance sales that exploited their ‘natural’ openness and friendliness. Despite these specific examples, there was no suggestion in our current analysis that DSE had an overall positively protective quality in the face of co-occurring problems. There is one important caveat to the conclusion that DSE is benign – in our previous work, we have proposed that it is in combination with other deprivation-specific problems, especially ASD, that DSE becomes especially impairing. This question will be explored in future papers.

Finally, DSE was unrelated to both young adults’ own and their adoptive parents’ perceptions of attachment security. This is significant because initially DSE was cast clinically as an attachment disorder with its origins in the absence of appropriate attachment figures in institutions leading to later disrupted attachment behaviours.³⁷ However, our data in the past suggested that the association between attachment security/insecurity and the DSE pattern was not so clear-cut.^{18,21,38} Specifically, over time, it became apparent that individuals with DSE were often also securely attached – although a somewhat lower proportion of children with DSE, compared with those without DSE, were rated as secure.¹¹ To date, it is unclear whether persistence in DSE into young adulthood would have a negative impact on parent–child attachment relationships. Our data suggest that there were no differences between those with DSE and those without it in the way the young adults, or their parents, perceived their attachment relationship.

Strengths and limitations

This paper is the first to present a systematic characterisation of DSE behaviours in adulthood. It has a number of strengths

including its prospective longitudinal nature, natural experimental adoption design and the stratification of deprivation-related risk (by duration) to increase statistical power. However, there are also a number of limitations that need to be considered when interpreting the findings. First, DSE was assessed by a limited number of items at each age. However, these appeared to be psychometrically robust and to capture important deprivation-related features that were validated by direct observation and self-reports. Second, attrition at the young adult assessment was higher than the very low rates seen at previous ERA follow-up assessments. This, although not surprising given the age range of the young adults, limited the statistical power of the analysis especially in the comparison of DSE+ and DSE– groups. However, attrition appeared to be non-selective in nature. Third, we had limited information on pre-institutional risks such as maternal smoking or stress exposures during pregnancy and these could have been elevated in the current sample. However, it seems unlikely that Rom>6 children would have been exposed to sufficiently greater risk prenatally than their Rom<6 group counterparts, in part at least because the timing of adoption was largely determined by the fall of the Ceausescu regime. Fourth, because duration and timing of deprivation overlapped, we were unable to address the key issue of sensitive developmental windows in any detail.

In summary, DSE behaviours continue to be core to the deprivation-related pattern in young adulthood in the ERA study – it is strongly related to the duration of deprivation and other deprivation-specific patterns such as ADHD and ASD. However, unlike these other patterns, DSE in and of itself appears to be relatively benign – apparently not, in and of itself, affecting QoL, functioning within different social domains, or educational and employment status.

Mark Kennedy, PhD, Developmental Brain-Behaviour Laboratory, Department of Psychology, University of Southampton, Southampton and Department of Child & Adolescent Psychiatry, Institute of Psychiatry, Psychology and Neuroscience, King’s College London, London, UK; **Jana Kreppner**, PhD, Developmental Brain-Behaviour Laboratory, Department of Psychology, University of Southampton, Southampton, UK; **Nicky Knights**, PhD, The Amy Winehouse Foundation, London, UK; **Robert Kumsta**, PhD, Department of Genetic Psychology, Faculty of Psychology, Ruhr-University Bochum, Germany; **Barbara Maughan**, PhD, MRC Social, Genetic & Developmental Psychiatry Centre, Institute of Psychiatry, Psychology and Neuroscience, King’s College London, London, UK; **Dennis Golm**, PhD, Department of Child & Adolescent Psychiatry, Institute of Psychiatry, Psychology and Neuroscience, King’s College London, London, UK; **Jonathan Hill**, PhD, School of Psychology and Clinical Language Sciences, University of Reading, Reading, UK; **Michael Rutter**, MD, MRC Social, Genetic & Developmental Psychiatry Centre, Institute of Psychiatry, Psychology and Neuroscience, King’s College London, London, UK; **Wolff Schlotz**, PhD, Max-Planck-Institute for Empirical Aesthetics, Frankfurt am Main, Germany; **Edmund Sonuga-Barke**, PhD, Department of Child & Adolescent Psychiatry, Institute of Psychiatry, Psychology and Neuroscience, King’s College London, London, UK

Correspondence: Edmund J. S. Sonuga-Barke, Department of Child and Adolescent Psychiatry, PO85, Institute of Psychiatry, Psychology and Neuroscience, 16 De Crespigny Park, London SE5 8AF, UK.
Email: edmund.sonuga-barke@kcl.ac.uk

First received 27 Feb 2017, final revision 3 May 2017, accepted 8 May 2017

Funding

This work was funded by a project grant from the UK Economic Social Research Council (ESRC; RES-062-23-3300).

Acknowledgements

We would like to express our sincere gratitude to all the families and young people who have participated in this study over the many years it has been running. Without them this work would not be possible. We are also grateful for the comments of Dr Graeme Fairchild during initial planning meetings. Thanks go to all the previous researchers on the project especially Dr Celia Beckett, Jenny Castle, Dr Suzanne Stevens, Dr Emma Colvert, Christine Groothues and Amanda Hawkins who were involved in the collection of data during the mid-adolescent phase of the project. We thank our advisory board for all their useful

suggestions especially Professor Megan Gunnar, Professor Trevor Robbins and Dr John Simmonds.

References

- 1 Teicher MH, Samson JA. Annual research review: enduring neurobiological effects of childhood abuse and neglect. *J Child Psychol Psychiatry* 2016; **57**: 241–66.
- 2 Kumsta R, Kreppner J, Kennedy M, Knights N, Rutter M, Sonuga-Barke E. Psychological consequences of early global deprivation. *Eur Psychol* 2015; **20**: 138–51.
- 3 Rutter M. Developmental catch-up, and deficit, following adoption after severe global early privation. *J Child Psychol Psychiatry* 1998; **39**: 465–76.
- 4 Castle J, Groothues C, Bredenkamp D, Beckett C, O'Connor T, Rutter M. Effects of qualities of early institutional care on cognitive attainment. *Am J Orthopsychiatry* 1999; **69**: 424–37.
- 5 Sonuga-Barke E, Kennedy M, Kreppner J, Knights N, Kumsta R, Maughan B, et al. Child-to-adult neurodevelopmental and mental health trajectories after early life deprivation: the young adult follow-up of the longitudinal English and Romanian Adoptees study. *Lancet* 2017; **389**: 1539–48.
- 6 Kennedy M, Kreppner J, Knights N, Kumsta R, Maughan B, Golm D, et al. Early severe institutional deprivation is associated with a persistent variant of adult attention-deficit/hyperactivity disorder: clinical presentation, developmental continuities and life circumstances in the English and Romanian Adoptees study. *J Child Psychol Psychiatry* 2016; **57**: 1113–25.
- 7 Kumsta R, Kreppner J, Rutter M, Beckett C, Castle J, Stevens S, et al. Deprivation-specific psychological patterns. *Monogr Soc Res Child Dev* 2010; **75**: 48–78.
- 8 Sonuga-Barke EJ, Schlotz W, Kreppner JV. Differentiating developmental trajectories for conduct, emotion, and peer problems following early deprivation. *Monogr Soc Res Child Dev* 2010; **75**: 102–24.
- 9 American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (DSM-5)*. APA, 2013.
- 10 Zeanah CH, Gleason MM. *Reactive Attachment Disorder: A Review for DSM-V. Report Presented to the American Psychiatric Association*. APA, 2010.
- 11 Rutter M, Colvert E, Kreppner J, Beckett C, Castle J, Groothues C, et al. Early adolescent outcomes for institutionally-deprived and non-deprived adoptees. I: disinhibited attachment. *J Child Psychol Psychiatry* 2007; **48**: 17–30.
- 12 Minnis H, Fleming G, Cooper SA. Reactive attachment disorder symptoms in adults with intellectual disabilities. *J Appl Res Intellect Disabil* 2010; **23**: 398–403.
- 13 Zeanah CH, Gleason MM. Annual research review: attachment disorders in early childhood—clinical presentation, causes, correlates, and treatment. *J Child Psychol Psychiatry* 2015; **56**: 207–22.
- 14 Lyons-Ruth K, Bureau JF, Riley CD, Atlas-Corbett AF. Socially indiscriminate attachment behavior in the Strange Situation: convergent and discriminant validity in relation to caregiving risk, later behavior problems, and attachment insecurity. *Dev Psychopathol* 2009; **21**: 355–72.
- 15 Humphreys KL, Nelson CA, Fox NA, Zeanah CH. Signs of reactive attachment disorder and disinhibited social engagement disorder at age 12 years: effects of institutional care history and high-quality foster care. *Dev Psychopathol* 2017; **29**: 675–84.
- 16 Lawler JM, Koss KJ, Doyle CM, Gunnar MR. The course of early disinhibited social engagement among post-institutionalized adopted children. *J Child Psychol Psychiatry* 2016; **57**: 1126–34.
- 17 Smyke AT, Zeanah CH, Gleason MM, Drury SS, Fox NA, Nelson CA, et al. A randomized controlled trial comparing foster care and institutional care for children with signs of reactive attachment disorder. *Am J Psychiatry* 2016; **169**: 508–14.
- 18 Kreppner J, Kumsta R, Rutter M, Beckett C, Castle J, Stevens S, et al. IV. Developmental course of deprivation-specific psychological patterns: early manifestations, persistence to age 15, and clinical features. *Monogr Soc Res Child Dev* 2010; **75**: 79–101.
- 19 Rutter M, Sonuga-Barke EJ, Beckett C, Castle J, Kreppner J, Kumsta R, et al. III. Deprivation-specific psychological patterns: effects of institutional deprivation. *Monogr Soc Res Child Dev* 2010; **75**: 1–253.
- 20 Rutter M, Sonuga-Barke EJ. X. Conclusions: overview of findings from the era study, inferences, and research implications. *Monogr Soc Res Child Dev* 2010; **75**: 212–29.
- 21 Rutter M, Kreppner J, Sonuga-Barke E. Emanuel Miller Lecture: attachment insecurity, disinhibited attachment, and attachment disorders: where do research findings leave the concepts? *J Child Psychol Psychiatry* 2009; **50**: 529–43.
- 22 Rutter M, LeCouteur A, Lord C, Macdonald H, Rios P, Folstein S. Diagnosis and subclassification of autism. In *Diagnosis and Assessment in Autism* (eds E Schopler, GB Mesibov): 239–59. Springer, 1988.
- 23 Conners CK, Pitkänen J, Rzepa SR. *Conners Comprehensive Behavior Rating Scale*. Springer, 2011.
- 24 American Psychiatric Association. *American Psychiatric Association Practice Guidelines for the Treatment of Psychiatric Disorders: Compendium*. APA, 2006.
- 25 Essau CA, Sasagawa S, Frick PJ. Callous-unemotional traits in a community sample of adolescents. *Assessment* 2006; **13**: 454–69.
- 26 Rutter M, Bailey A, Lord C. *The Social Communication Questionnaire: Manual*. Western Psychological Services, 2003.
- 27 Wechsler D. *Manual for the Wechsler Abbreviated Intelligence Scale (WASI)*. The Psychological Corporation, 1999.
- 28 Kenny ME. Homepage of Maureen E. Kenny, PhD. Boston College, 1990. Available from: <http://www2.bc.edu/Maureen-Kenny> (accessed 29 August 2017).
- 29 Armsden GC, Greenberg MT. The inventory of parent and peer attachment: individual differences and their relationship to psychological well-being in adolescence. *J Youth Adolesc* 1987; **16**: 427–54.
- 30 Kenny ME, Sirin SR. Parental attachment, self-worth, and depressive symptoms among emerging adults. *J Counsel Develop* 2006; **84**: 61–71.
- 31 Diener ED, Emmons RA, Larsen RJ, Griffin S. The Satisfaction with Life Scale. *J Personal Assess* 1985; **49**: 71–5.
- 32 Hill J, Pilkonis P, Morse J, Feske U, Reynolds S, Hope H, et al. Social domain dysfunction and disorganization in borderline personality disorder. *Psychol Med* 2008; **38**: 135–46.
- 33 Mackie E, Hill J, Kondry H, McNally R. Adult psychosocial outcomes in long-term survivors of acute lymphoblastic leukaemia and Wilms' tumour: a controlled study. *Lancet* 2000; **355**: 1310–4.
- 34 General R. *Registrar General's Decennial Supplement, England and Wales, 1961. Occupational Mortality Tables*. HMSO, 1971.
- 35 Kreppner JM, Rutter M, Beckett C, Castle J, Colvert E, Groothues C, et al. Normality and impairment following profound early institutional deprivation: a longitudinal follow-up into early adolescence. *Dev Psychol* 2007; **43**: 931–46.
- 36 Kay C, Green J, Sharma K. Disinhibited attachment disorder in UK adopted children during middle childhood: prevalence, validity and possible developmental origin. *J Abnorm Child Psychol* 2016; **44**: 1375–86.
- 37 Lyons-Ruth K. Commentary: should we move away from an attachment framework for understanding disinhibited social engagement disorder (DSED)? A commentary on Zeanah and Gleason (2015). *J Child Psychol Psychiatry* 2015; **56**: 223–7.
- 38 Kreppner J, Rutter M, Marvin R, O'Connor TG, Sonuga-Barke E. Assessing the concept of the insecure-other category in the Marvin scheme: changes between 4 and 6 years in the English and Romanian adoptee study. *Soc Dev* 2011; **20**: 1–16.

