

The Tor Spectral Sequence and Flat Morphisms in Homotopical \mathcal{D} -Geometry

Alisa Govzmann, Damjan Pištalo, and Norbert Poncin

December 23, 2024

Abstract

Homotopical algebraic \mathcal{D} -geometry combines aspects of homotopical algebraic geometry [32, 33] and \mathcal{D} -geometry [3]. It was introduced in [10] as a suitable framework for a coordinate-free study of the Batalin-Vilkovisky complex and more generally for the study of non-linear partial differential equations and their symmetries [5, 25]. In order to consolidate the foundation of the theory, we have to prove that the standard methods of linear and commutative algebra are available in the context of homotopical algebraic \mathcal{D} -geometry, and we must show that in this context the étale topology is a kind of homotopical Grothendieck topology and that the notion of smooth morphism is, roughly speaking, local for the étale topology. The first half of this work was done in [10]. The remaining part covers the study of étale and flat morphisms in the category of differential graded \mathcal{D} -algebras and is based on the Tor spectral sequence which connects the graded Tor functors in homology with the homology of the derived tensor product of two differential graded \mathcal{D} -modules over a differential graded \mathcal{D} -algebra.

MSC 2020: 18G15, 18E35, 18N40, 14A30

Keywords: model category, homotopy theory, derived functor, \mathcal{D} -module, \mathcal{D} -geometry, homotopical geometry, Tor functor, spectral sequence

Contents

1	Introduction	2
2	Notation and preliminary results	6

3	Derived tensor product in modules over a DG \mathcal{D}-algebra	12
4	Graded Tor functors in modules over a graded \mathcal{D}-algebra	15
5	The \mathcal{D}-generalization of Quillen's Tor spectral sequence	21
6	Exactness of the derived tensor product	31
7	A characterization of flat modules	37
8	Outlook	48

1 Introduction

Although deep insights into theoretical physics are often gained from calculations in coordinates, since the discovery of the general theory of relativity, geometers have been striving again for coordinate-free concepts and results. As many mathematical models reduce real-world problems to solving systems of partial differential equations (PDE-s), a coordinate-independent theory of PDE-s and related aspects is essential. In [35], A. Vinogradov and his coworkers propose a cohomological analysis of PDE-s over smooth manifolds, A. Beilinson and V. Drinfeld advocate in [3] working in the algebraic \mathcal{D} -geometric setting over smooth schemes, while Kashiwara [20] and his co-workers work with \mathcal{D} -modules over complex manifolds. The solutions of a system of non-linear PDE-s are related to diffieties in the first approach and to \mathcal{D}_X -schemes in the second. In the third approach, solutions to systems of linear PDE-s are related to left \mathcal{D} -modules. To treat the moduli space of solutions modulo symmetries, we have to replace diffieties or \mathcal{D}_X -schemes with appropriate objects from derived geometry. In the algebraic approach, this suggests combining the \mathcal{D} -geometry of Beilinson and Drinfeld with the homotopical algebraic geometry in the sense of B. Toën and G. Vezzosi [32], [33]. In a series of papers [9], [10], [27] and [14], the authors present homotopical algebraic \mathcal{D} -geometry as a suitable framework for the study of PDE-s and their symmetries. In fact, this new geometry provides in particular a convenient method for encoding total derivatives and leads to a covariant description of the classical Batalin-Vilkovisky complex, which appears as a special case of general constructions. Further evidence for this point of view can be found in [25], [26]. Similar projects could be pursued in the smooth setting combining recent advances in derived smooth geometry ([31], [19], [2], ...) with the work of Vinogradov and his followers, or in the derived complex geometry ([29], [23], ...). Further generalization, which is outside

of the scope of the present work, would be to extend the theory by allowing singularities. For example, in the complex setting differential operators are generalized to singular spaces via Kashiwara's theorem, and in the algebraic setting via crystals ([13]).

In order to implement the homotopical algebraic \mathcal{D} -geometry, we have to prove that the tuple $(\mathbf{DGDM}, \mathbf{DGDM}, \mathbf{DGDA}, \tau, \mathbf{P})$ is a homotopical algebraic geometric context (HAG context) in the sense of [33]. Here, \mathcal{D} denotes the non-commutative ring of total sections of the sheaf of differential operators of a smooth affine algebraic variety [10], \mathbf{DGDM} is the symmetric monoidal model category of differential non-negatively graded \mathcal{D} -modules (for \mathcal{D} -modules, see for instance [20]), \mathbf{DGDA} is its subcategory of commutative monoids, τ is a suitable model pre-topology on the opposite category $\mathbf{DGDA}^{\text{op}}$ (thought of as the category of derived affine \mathcal{D} -schemes), and \mathbf{P} is a class of morphisms of $\mathbf{DGDA}^{\text{op}}$ that is compatible with τ (we can think of τ as the étale topology and of \mathbf{P} as the class of smooth morphisms). Apart from enabling us to by-pass sheaf-theoretical aspects of the theory, the affines of the underlying smooth variety is at this point necessary. Namely, the axioms of a HAG context require in particular the existence of a projective model structure on \mathbf{DGDA} transferred from \mathbf{DGDM} via the free-forgetful adjunction – which in turn requires the category \mathbf{DGDM} to have enough projectives. Later requirement is simply not satisfied in the analogous sheaf-theoretical category (section 2.1) over a not necessarily affine scheme. However, the assumption that X is affine could still be relaxed in the later stages of the project. Indeed, the homotopical algebraic geometry context, as introduced by [33], is intended to describe the affine picture, global aspects being addressed only once it is established.

In [10] the first part of the HAG context conditions was proved, i.e., we showed that the triplet $(\mathbf{DGDM}, \mathbf{DGDM}, \mathbf{DGDA})$ is a homotopical algebraic context (HA context) [33], which guarantees that essential tools from linear and commutative algebra are still available. Although in an HA context there exist abstract notions of étale and smooth maps, they are not suitable for showing that the abstractized étale topology satisfies the required axioms, or that it is compatible with the class \mathbf{P} of "smooth" morphisms. Similarly to the HA context of derived algebraic geometry, in the sense of [33, Chapter 2], we expect a concrete characterization of those properties to hold. The conjecture is that a morphism $f : A \rightarrow B$ of \mathbf{DGDA} is flat/étale/smooth if and only if $H_0(f)$ is such (in the appropriate sense, as a map of commutative \mathcal{D} -algebras), and a certain strongness property (Definition 5) is satisfied. Extension of the Quillen's Tor spectral sequence [30, II 6.8] to \mathcal{D} -algebraic framework is the essential tool when proving such conjectures. In the paper's final section, the conjecture is proven true for flat morphisms.

The generalization of the Tor spectral sequence connects the homology of the derived tensor product of two modules $M, N \in \mathbf{DGDM}$ over an algebra $\mathcal{A} \in \mathbf{DGDA}$ with the Tor of the corresponding homologies, which appears as the second sheet of a first quadrant spectral sequence in \mathbf{DM} . If M is flat in the sense that the derived tensor product with M preserves homotopy pullbacks, the Tor spectral sequence collapses at its second sheet, giving rise to the isomorphism of $H\mathcal{A}$ -modules

$$HM \otimes_{H\mathcal{A}} HN \cong H(M \otimes_{\mathcal{A}}^{\mathbb{L}} N).$$

Quillen's original construction of the Tor spectral sequence for modules over a simplicial algebra has since been adopted to other contexts, such as modules over a differential graded or even A_∞ -algebra ([21]), \mathbb{E}_1 -ring spectrum ([23, Proposition 7.2.1.19.], [11, Theorem 4.1]), and its $K(1)$ -local analogue ([34]). Nevertheless, a generalization to differential graded algebras over differential operators (or a more general abstract context) is new to the best of our knowledge. Although the general philosophy is similar in all these versions of the Tor spectral sequence, the technical reasoning is specific to each setting.

The paper is organized as follows.

Section 2 establishes the conventions used throughout this text. First subsection deals with the notation, also providing definitions of categories of (differential) graded \mathcal{D} modules and algebras, and their properties essential for the present paper. The second subsection reviews model categories, their homotopy categories, derived functors, as well as the replacements used to compute them. Theorem 1 makes it easier to navigate the thicket of similar but different concepts, definitions, and results – see for instance [12, 17, 18, 22]. Indeed, although the facts presented in this theorem are by no means new, to our knowledge it is difficult to find a similar result in the literature.

In Section 3 we prove that in our category $\mathbf{Mod}(\mathcal{A})$ of modules M, N in \mathbf{DGDM} over $\mathcal{A} \in \mathbf{DGDA}$ (i.e., of modules in the category of differential non-negatively graded \mathcal{D} -modules over a commutative monoid in this symmetric monoidal category), in order to compute the derived tensor product $M \otimes_{\mathcal{A}}^{\mathbb{L}} N$, we can choose a cofibrant replacement of M and N , of M only, or N only, just as in classical homological algebra.

The family of Tor functors is usually defined in the setting of modules over a ring. Considering what we said above, this family appears here in the context of (homology) modules in \mathbf{GDM} over $\mathfrak{A} \in \mathbf{GDA}$ (see the section 2.1 for the notation). Hence the value on objects of each one of the Tor functors is a graded \mathcal{D} -module over the graded \mathcal{D} -algebra \mathfrak{A} . The modules

of this category $\text{Mod}(\mathfrak{A})$ are *not* graded modules over the single ring of differential operators with coefficients in \mathfrak{A} . Instead, we are dealing in addition to the \mathfrak{A} -action with a compatible action by the non-commutative ring \mathcal{D} of differential operators. Further, the \mathcal{D} -context is a rather special one, among other things because the tensor product $M \otimes_{\mathfrak{A}} N$ (for notations, see preceding paragraph) is a quotient of the tensor product $M \otimes N$ of differential graded \mathcal{D} -modules, which however is not the tensor product over \mathcal{D} , but that over the commutative ring \mathcal{O} of functions. In Section 4 we show that the just mentioned graded Tor functors do exist, which includes checking that the categories $\text{Mod}(\mathcal{A})$ and $\text{Mod}(\mathfrak{A})$ are abelian, that the latter has enough projectives, and that the partial tensor product functor over it is right exact.

In Section 5 we give the \mathcal{D} -geometric generalization of Quillen's Tor spectral sequence – a particular Künneth spectral sequence used by Quillen within the framework of simplicial modules over a simplicial ring. We explain why Quillen's original proof can be generalized to our setting. In particular, we use Sullivan differential graded \mathcal{D} -modules over differential graded \mathcal{D} -algebras [9, 10] and graded \mathcal{D} -modules over graded \mathcal{D} -algebras which are free over a graded set. In addition, we prove a strong Künneth theorem in this context.

In Section 6 we give all the details needed for a rigorous homotopy invariant definition of flatness (see above). More precisely, on the one hand, the homotopy pullback in $\text{Mod}(\mathcal{A})$ is a functor from the homotopy category $\text{Ho}(\text{CoSpan}(\text{Mod}(\mathcal{A})))$ of cospan diagrams in \mathcal{A} -modules to the category $\text{Ho}(\text{Mod}(\mathcal{A}))$. On the other hand, the derived tensor product with $M \in \text{Mod}(\mathcal{A})$ is an endofunctor of $\text{Ho}(\text{Mod}(\mathcal{A}))$ and induces an endofunctor of $\text{CoSpan}(\text{Ho}(\text{Mod}(\mathcal{A})))$, but not an endofunctor of the non-equivalent homotopy category $\text{Ho}(\text{CoSpan}(\text{Mod}(\mathcal{A})))$. Therefore, it is not clear at first what it means for the derived tensor product to preserve homotopy pullbacks. Also, similar to usual pullbacks, for a functor \mathcal{F} to preserve homotopy pullbacks it is not sufficient that the \mathcal{F} -image of the homotopy pullback of a diagram is isomorphic to the homotopy pullback of the \mathcal{F} -image, but this ‘isomorphism’ must be realized by a specific universal weak equivalence. Section 6 solves these difficulties and leads to a completely rigorous concept of flatness (Definition 3). It is further shown (Corollary 1) that for an \mathcal{A} -module M to be flat it is sufficient that the derived tensor product with M preserves homotopy fiber sequences in the sense of [14]. As with the preservation of homotopy pullbacks, a few subtleties have to be considered in order to give the preservation of homotopy fiber sequences a rigorous meaning (Definition 4).

In Section 7 we prove that an \mathcal{A} -module in the category of differential graded \mathcal{D} -modules is flat if and only if it is strongly flat (for a similar result in the case of a commutative base ring, see [33, Lemma 2.2.2.]).

We conclude with a brief outlook in Section 8.

Acknowledgments

Authors are grateful to the anonymous referee for his constructive criticism.

2 Notation and preliminary results

2.1 Differential graded \mathcal{D} -modules and algebras.

This subsection fixes the notation, summarizing along the way the results from [9] and [10] relevant for the present work.

Throughout the paper, we work over a field k of characteristic zero, using exclusively homological convention. All the rings considered in the paper are unital.

We fix a smooth affine algebraic variety X over k . In particular, this allows us to replace quasi-coherent sheaves of \mathcal{O}_X -modules by their global sections. Denoting the sheaf of differential operators over X by \mathcal{D}_X , and the algebra of its global sections by $\mathcal{D} := \Gamma(X, \mathcal{D}_X)$, the category of differential non-negatively graded \mathcal{D} -modules (denoted by $\mathbf{D}\mathcal{G}\mathcal{D}\mathcal{M}$) is equivalent to the category of sheaves of differential non-negatively graded \mathcal{O}_X -quasi-coherent \mathcal{D}_X -modules. Both categories are closed symmetric monoidal: tensor products are respectively over \mathcal{O}_X , and $\mathcal{O} := \Gamma(X, \mathcal{O}_X)$, with the action of differential operators given by the (non-graded) Leibniz rule. The categorical equivalence descends to the respective subcategories of symmetric monoids. Category of symmetric monoids in $\mathbf{D}\mathcal{G}\mathcal{D}\mathcal{M}$ is referred to as the category of dg \mathcal{D} -algebras, and denoted by $\mathbf{D}\mathcal{G}\mathcal{D}\mathcal{A}$. Its objects are differential graded commutative \mathcal{O} -algebras equipped with the action of differential operators which respects the (non-graded) Leibniz rule. As the category of non-negatively graded chain complexes over a unital ring, $\mathbf{D}\mathcal{G}\mathcal{D}\mathcal{M}$ carries a cofibrantly generated projective model structure. With the above tensor product it is a monoidal model category. The model structure transfers via the free-forgetful adjunction to $\mathbf{D}\mathcal{G}\mathcal{D}\mathcal{A}$.

Given $\mathcal{A} \in \mathbf{D}\mathcal{G}\mathcal{D}\mathcal{A}$, denote by $\mathbf{Mod}(\mathcal{A})$ the category of \mathcal{A} -actions in $\mathbf{D}\mathcal{G}\mathcal{D}\mathcal{M}$ in the sense of [24, VII 4.]. Throughout the paper, we refer to it as the category of \mathcal{A} -modules. It is again a cofibrantly generated monoidal model category whose weak equivalences are quasi-isomorphisms, and whose fibrations are surjections in strictly positive degrees. Cofibrant objects in $\mathbf{Mod}(\mathcal{A})$ are exactly the retracts of the Sullivan \mathcal{A} -modules, i.e., of the \mathcal{A} -modules

$\mathcal{A} \otimes_{\mathcal{O}} (\bigoplus_{i \in I} \mathcal{D} g_i[n_i])$, where I is a well-ordered set (equivalently, an ordinal) and the $g_i[n_i]$ are symbols of degree $n_i \in \mathbb{N}$, and whose differential is lowering in the sense of Definition 2.3.6. of [10]. This explicit description of cofibrations is repeatedly used throughout the paper. We denote the symmetric monoidal structure by $(\text{Mod}(\mathcal{A}), \otimes_{\mathcal{A}}, \mathcal{A})$.

Category of \mathbb{N} -graded \mathcal{D} -modules is denoted by \mathbf{GDM} . Category of \mathbb{N} -graded-commutative \mathcal{D} -algebras (the subcategory of symmetric monoids) is denoted by \mathbf{GDA} , and its objects are denoted by the fraktur font. Given $\mathfrak{A} \in \mathbf{GDA}$ the category of \mathfrak{A} -actions in \mathbf{GDM} is denoted by $\text{Mod}(\mathfrak{A})$. There are no noteworthy differences to the similar category $\text{Mod}(\mathcal{A})$ ($\mathcal{A} \in \mathbf{DGDA}$) introduced above. In particular the category $\text{Mod}(\mathfrak{A})$ is symmetric monoidal with tensor product $-\otimes_{\mathfrak{A}}-$.

\mathcal{DM} denotes the category of \mathcal{D} -modules, and \mathcal{DA} denotes the category of commutative \mathcal{D} -algebras. Finally, a reoccurring property of the ring \mathcal{D} is that it is projective as an \mathcal{O} -module [10, Proposition 1.0.11.]. This allows the use of Künneth formula for complexes, which is repeatedly invoked throughout the paper.

2.2 Zooming in on derivatives in model categories

This section establishes notations and conventions and ensures that the present text is self-contained. Theorem 1 is a model categorical generalization of the well-known homological algebraic fact that the values of the left derived functors of a covariant right exact functor from an abelian category with enough projectives to an abelian category, are independent, up to canonical isomorphism, of the projective resolution used.

We assume that the reader is familiar with model categories. We adopt the definition of a model category that is used in [17]. More precisely, a model category is a category \mathbf{M} that is equipped with three classes of morphisms called weak equivalences, fibrations and cofibrations. The category \mathbf{M} has all *small* limits and colimits and the 2-out-of-3 axiom, the retract axiom and the lifting axiom are satisfied. Moreover \mathbf{M} admits a *functorial* cofibration - trivial fibration factorization system (Cof - TrivFib factorization) and a functorial trivial cofibration - fibration factorization system (TrivCof - Fib factorization).

Let now (α, β) be any functorial Cof - TrivFib factorization. For every $X \in \mathbf{M}$, it factors the map $i_X : 0 \rightarrow X$ out of the initial object of \mathbf{M} into a cofibration $\alpha(i_X)$ followed by a trivial fibration $q_X := \beta(i_X)$:

$$i_X : 0 \rightarrowtail QX \xrightarrow{\sim} X .$$

Regardless of the factorization

$$i_X : 0 \rightarrowtail CX \xrightarrow{\sim} X$$

of i_X into a cofibration followed by a weak equivalence c_X considered, we refer to CX as a *cofibrant replacement* of X . The object QX we call a *cofibrant F-replacement* of X (or just a cofibrant replacement if we do not want to stress that q_X is a fibration). From the fact that the factorization (α, β) is functorial it follows that Q is an endofunctor of \mathbf{M} . Moreover $q_X : QX \rightarrow X$ is functorial in X : q is a natural transformation $q : Q \Rightarrow \text{id}_{\mathbf{M}}$ from the *cofibrant replacement functor* Q to the identity functor $\text{id}_{\mathbf{M}}$ [18]. Instead of the cofibrant F-replacement functor Q that is **globally** defined by the functorial factorization (α, β) , we will also use **local** / object-wise cofibrant replacements CX or *local cofibrant F-replacements* $\tilde{C}X$ such that the map c_X in the factorization

$$i_X : 0 \rightarrowtail \tilde{C}X \xrightarrow{\sim} X$$

is id_X if X is already cofibrant [30]. It is important to remember that if for every X we choose such a local cofibrant F-replacement and if $f : X \rightarrow Y$, there exists a lifting $\tilde{C}f : \tilde{C}X \rightarrow \tilde{C}Y$:

$$\begin{array}{ccccc} 0 & \xrightarrow{\quad} & \tilde{C}Y & \xleftarrow{\quad} & \\ \downarrow & \nearrow \tilde{C}f & \downarrow \sim c_Y & & \\ \tilde{C}X & \xrightarrow{\sim c_X} & X & \xrightarrow{f} & Y \end{array} \tag{1}$$

By homotopy category of a model category \mathbf{M} in this work we mean the Quillen homotopy category $\mathbf{Ho}(\mathbf{M})$, which is the strong localization $\mathbf{M}[[W^{-1}]]$ of \mathbf{M} at its class W of weak equivalences [18]. Like any localization, the category $\mathbf{Ho}(\mathbf{M})$ comes along with a localization functor $L_{\mathbf{M}} : \mathbf{M} \rightarrow \mathbf{Ho}(\mathbf{M})$ that sends weak equivalences to isomorphisms. By strong localization we mean that if \mathbf{C} is another category and $F : \mathbf{M} \rightarrow \mathbf{C}$ another functor that sends weak equivalences to isomorphisms, then there exists a *unique* functor $\mathbf{Ho}(F) : \mathbf{Ho}(\mathbf{M}) \rightarrow \mathbf{C}$ such that the resulting triangle commutes *on the nose*, i.e., $F = \mathbf{Ho}(F) \circ L_{\mathbf{M}}$. The objects of $\mathbf{Ho}(\mathbf{M})$ are the objects of \mathbf{M} and the morphisms of $\mathbf{Ho}(\mathbf{M})$ are the alternating finite strings

$$[S] = [\rightarrow \rightsquigarrow \rightarrow \rightsquigarrow \dots] \quad \text{and} \quad [S] = [\rightsquigarrow \rightarrow \rightsquigarrow \rightarrow \dots]$$

of morphisms $f : X \rightarrow Y$ of \mathbf{M} and formal reversals $w^{-1} : Y \rightsquigarrow Z$ of weak equivalences $w : Z \rightarrow Y$ of \mathbf{M} . Here the class $[\cdot]$ refers to the identifications of the empty string 1_X at X , the concatenation string $f, g : X \rightarrow Y \rightarrow Z$ of composable \mathbf{M} -morphisms, the concatenation

strings $w, w^{-1} : Z \rightarrow Y \rightsquigarrow Z$ and $w^{-1}, w : Y \rightsquigarrow Z \rightarrow Y$ with the \mathbf{M} -morphisms id_X , $g \circ f$, id_Z and id_Y , respectively. The unique factorization $\text{Ho}(F)$ of F through $\text{Ho}(\mathbf{M})$ is defined on objects $X \in \text{Ho}(\mathbf{M})$ by $\text{Ho}(F)(X) = F(X)$ and on $\text{Ho}(\mathbf{M})$ -morphisms $[S]$ by

$$\text{Ho}(F)[f : X \rightarrow Y] = F(f : X \rightarrow Y) \quad \text{and} \quad \text{Ho}(F)[w^{-1} : Y \rightsquigarrow Z] = (F(w : Z \rightarrow Y))^{-1}.$$

From what has been said in the previous paragraph it follows that a functor $F : \mathbf{M} \rightarrow \mathbf{N}$ between model categories \mathbf{M} and \mathbf{N} , which preserves *all* weak equivalences can be viewed canonically as a functor between the homotopy categories. Indeed, the composite $L_{\mathbf{N}} \circ F : \mathbf{M} \rightarrow \text{Ho}(\mathbf{N})$ sends weak equivalences to isomorphisms and therefore factors uniquely through the homotopy category of \mathbf{M} , i.e., there is a unique functor

$$\text{Ho}(L_{\mathbf{N}} \circ F) : \text{Ho}(\mathbf{M}) \rightarrow \text{Ho}(\mathbf{N}) \tag{2}$$

such that

$$\text{Ho}(L_{\mathbf{N}} \circ F) \circ L_{\mathbf{M}} = L_{\mathbf{N}} \circ F.$$

However, requiring that functors between model categories respect the *entire* model structure is far too strong a requirement. Usually, functors F between model categories are left (or right) Quillen functors, i.e., they preserve all cofibrations and all trivial cofibrations (resp., all fibrations and all trivial fibrations). Since a left Quillen functor F thus sends trivial cofibrations between cofibrant objects to weak equivalences, it follows from Brown's lemma that it sends all *weak equivalences between cofibrant objects* to weak equivalences. Hence, if $Q_{\mathbf{M}}$ denotes a cofibrant replacement functor of \mathbf{M} , since due to the 2-out-of-3 axiom $Q_{\mathbf{M}}$ preserves weak equivalences, the functor $L_{\mathbf{N}} \circ F \circ Q_{\mathbf{M}} : \mathbf{M} \rightarrow \text{Ho}(\mathbf{N})$ sends weak equivalences to isomorphisms and therefore factors uniquely through the homotopy category of \mathbf{M} , i.e., there is a unique functor

$$\text{Ho}(L_{\mathbf{N}} \circ F \circ Q_{\mathbf{M}}) : \text{Ho}(\mathbf{M}) \rightarrow \text{Ho}(\mathbf{N}) \tag{3}$$

such that

$$\text{Ho}(L_{\mathbf{N}} \circ F \circ Q_{\mathbf{M}}) \circ L_{\mathbf{M}} = L_{\mathbf{N}} \circ F \circ Q_{\mathbf{M}}.$$

We refer to this functor as ‘the’ left derived functor of F . Since its definition is based on the universal property of the strong localization $\text{Ho}(\mathbf{M})$ and it is computed using the cofibrant replacement functor $Q_{\mathbf{M}}$, we also speak of the **strongly universal left derived functor** of F and denote this functor $\mathbb{L}_Q^S F$:

$$\mathbb{L}_Q^S F := \text{Ho}(L_{\mathbf{N}} \circ F \circ Q_{\mathbf{M}}). \tag{4}$$

Alternatively one can define the left derived functor of F as the right Kan extension of $L_N \circ F$ along L_M . We refer to this derived functor as the **Kan extension left derived functor** of F and denote it by $\mathbb{L}^K F$. The right derived functors $\mathbb{R}_R^S G$ ($R = R_M$: fibrant replacement functor) and $\mathbb{R}^K G$ of a right Quillen functor G are defined dually.

We first clarify the relationship between (2) and (3):

Proposition 1. *If $F \in \text{Fun}(M, N)$ is a functor between model categories that preserves all weak equivalences, the whiskering $\theta := L_N \star (F \star q_M)$ is a canonical natural isomorphism*

$$\theta : \text{Ho}(L_N \circ F \circ Q_M) \xrightarrow{\cong} \text{Ho}(L_N \circ F) .$$

We refer to the latter by simply writing

$$\mathbb{L}_Q^S F := \text{Ho}(L_N \circ F \circ Q_M) \doteq \text{Ho}(L_N \circ F) .$$

Proof. As $q_M : Q_M \xrightarrow{\cong} \text{id}_M$ is a natural weak equivalence, i.e., a natural transformation that is objectwise a weak equivalence, the whiskering $\theta := L_N \star (F \star q_M)$ is a natural isomorphism

$$\theta : \text{Ho}(L_N \circ F \circ Q_M) \circ L_M \xrightarrow{\cong} \text{Ho}(L_N \circ F) \circ L_M ,$$

because for every $X \in M$ the morphism $\theta_X := L_N(F(q_{M,X}))$ is an isomorphism. That θ is a natural isomorphism

$$\theta : \text{Ho}(L_N \circ F \circ Q_M) \xrightarrow{\cong} \text{Ho}(L_N \circ F)$$

follows from [15, Lemma 1] which is a rather straightforward consequence of the above description of $\text{Ho}(M)$. \square

The next theorem addresses the question of stability of a derived functor with respect to a change of definition (Kan extension versus strongly universal) and with respect to a change of the type of cofibrant replacement used to compute it (local versus global):

Theorem 1 ([16]). *If $F \in \text{Fun}(M, N)$ is a functor between model categories that sends weak equivalences between cofibrant objects to weak equivalences, its Kan extension left derived functor*

$$\mathbb{L}^K F \in \text{Fun}(\text{Ho}(M), \text{Ho}(N))$$

and its strongly universal left derived functor

$$\mathbb{L}^S F \in \text{Fun}(\text{Ho}(M), \text{Ho}(N))$$

([15]) exist and we have

$$\mathbb{L}^K F \doteq \text{Ho}(L_N \circ F \circ \tilde{C}_M) \doteq \mathbb{L}_Q^S F := \text{Ho}(L_N \circ F \circ Q_M) \xrightarrow{\cong} \mathbb{L}^S F, \quad (5)$$

where \tilde{C}_M is a local cofibrant F -replacement, Q_M is a cofibrant F -replacement functor and Ho the unique on-the-nose factorization through $\text{Ho}(M)$. This implies that

$$\mathbb{L}^K F \circ L_M \doteq L_N \circ F \circ \tilde{C}_M \doteq \mathbb{L}_Q^S F \circ L_M = L_N \circ F \circ Q_M \xrightarrow{\cong} \mathbb{L}^S F \circ L_M, \quad (6)$$

where \doteq denotes a canonical natural isomorphism and $\xrightarrow{\cong}$ a not necessarily canonical natural isomorphism.

Hence, for every $X \in M$, the value of the derived functor at $L_M X = X \in \text{Ho}(M)$ is

$$\mathbb{L}^K F(X) \doteq F(\tilde{C}_M X) \doteq \mathbb{L}_Q^S F(X) = F(Q_M X) \cong \mathbb{L}^S F(X), \quad (7)$$

where \doteq is a canonical isomorphism in $\text{Ho}(N)$ and \cong a not necessarily canonical isomorphism. Moreover, ([16, Proposition 1]) if $C_M X$ is any cofibrant replacement of X , there is a canonical $\text{Ho}(N)$ -isomorphism

$$F(\tilde{C}_M X) \doteq F(C_M X). \quad (8)$$

In view of (7) and (8) the value of a derived functor at an object is well-defined only up to isomorphism of the target homotopy category. The isomorphism class is independent of the type of derived functor considered, Kan extension or strongly universal, as well as independent of the type of cofibrant F -replacement considered, local or global. Also the choice of another local or another global replacement does not change the isomorphism class. If we compute the value of the derived functor using a local cofibrant replacement that is not necessarily an F -replacement, we get again the same class. Finally, the three representatives considered of the value of the derived functor, namely $L_N(F(\tilde{C}_M X))$, $L_N(F(Q_M X))$ and $L_N(F(C_M X))$, are related by canonical isomorphisms when viewed as objects of $\text{Ho}(N)$ and by zigzags of weak equivalences when viewed as objects of N . All of this, of course, is consistent with the idea underlying homotopy theory that instead of requiring two objects to be equal, we should simply ask that they are related by a weak equivalence.

Remark 1. Let us stress that the symbols \sim (resp., \approx , \simeq , \cong , \doteq) denote in this text a weak equivalence (resp., a zigzag of weak equivalences, a homotopy, an isomorphism, a canonical isomorphism).

3 Derived tensor product in modules over a DG \mathcal{D} -algebra

Given a commutative ring R , the derived tensor product of R -modules can be computed by taking the projective resolution – cofibrant replacements in the category of dg R -modules – in any of its two arguments. In contrast, computing the derived tensor product in a general monoidal model category requires cofibrant replacements in both arguments. The present section shows that computing derived tensor product in $\mathbf{DGD}\mathbf{M}$ requires cofibrant replacement in a single argument, as over a commutative ring. Argumentation relies on earlier joint work of the second and third author with G. di Brino.

It is well-known that if we fix one argument in a bifunctor, i.e., in a functor $\Pi : \mathbf{C} \times \mathbf{D} \rightarrow \mathbf{E}$ out of a product category, we get a functor in the other argument. More precisely, for every $c \in \mathbf{C}$ and $d \in \mathbf{D}$, the partial functors $L_c := \Pi(c, -) : \mathbf{D} \rightarrow \mathbf{E}$ and $R_d := \Pi(-, d) : \mathbf{C} \rightarrow \mathbf{E}$ satisfy

$$L_c(d) = \Pi(c, d) = R_d(c) , \quad (9)$$

and for every morphisms $f : c' \rightarrow c''$ and $g : d' \rightarrow d''$, we have

$$L_{c''}(g) \circ R_{d'}(f) = \Pi(\text{id}_{c''}, g) \circ \Pi(f, \text{id}_{d'}) = \Pi(f, g) = \Pi(f, \text{id}_{d''}) \circ \Pi(\text{id}_{c'}, g) = R_{d''}(f) \circ L_{c'}(g) . \quad (10)$$

Conversely, if for every object (c, d) of $\mathbf{C} \times \mathbf{D}$ there are functors L_c and R_d that satisfy (9) and if for every morphism (f, g) of $\mathbf{C} \times \mathbf{D}$ the condition (10) is satisfied, then there exists a bifunctor Π whose partial functors are the L_c and R_d [24]. We use this connection between a bifunctor and its partial functors below.

In view of the HA properties [10], the universal arrow $f \square g$ in the next pushout-diagram is a cofibration if $f : M \rightarrow N$ and $g : P \rightarrow S$ are and it is a trivial cofibration if one of f and g is a cofibration and the other a trivial cofibration:

$$\begin{array}{ccccc}
 M \otimes_{\mathcal{A}} P & \xrightarrow{f \otimes \text{id}} & N \otimes_{\mathcal{A}} P & & \\
 \text{id} \otimes g \downarrow & & \downarrow & & \\
 M \otimes_{\mathcal{A}} S & \xrightarrow{\quad} & (N \otimes_{\mathcal{A}} P) \coprod_{M \otimes_{\mathcal{A}} P} (M \otimes_{\mathcal{A}} S) & \xrightarrow{\text{id} \otimes g} & \\
 & \searrow & \text{---} & \nearrow & \\
 & & f \square g & & \\
 & \searrow & \text{---} & \nearrow & \\
 & & f \otimes \text{id} & & \\
 & & \searrow & & \\
 & & N \otimes_{\mathcal{A}} S & &
 \end{array}$$

If $U \in \mathbf{Mod}(\mathcal{A})$ is cofibrant, the unique arrow $f : \{0\} \rightarrow U$ from the initial \mathcal{A} -module to U is a cofibration, so the universal arrow

$$f \square g = \text{id}_U \otimes_{\mathcal{A}} g : U \otimes_{\mathcal{A}} P \rightarrow U \otimes_{\mathcal{A}} S$$

is a cofibration (resp., a trivial cofibration), if $g : P \rightarrow S$ is a cofibration (resp., a trivial cofibration). Hence, the functor $U \otimes_{\mathcal{A}} -$ respects cofibrations and trivial cofibrations and is therefore a left Quillen endofunctor of the model category $\mathbf{Mod}(\mathcal{A})$. Given what we said above, the functor

$$U \otimes_{\mathcal{A}} - : \mathbf{Mod}(\mathcal{A}) \rightarrow \mathbf{Mod}(\mathcal{A}) \quad (11)$$

sends weak equivalences between cofibrant objects to weak equivalences and of course so does the functor

$$- \otimes_{\mathcal{A}} U : \mathbf{Mod}(\mathcal{A}) \rightarrow \mathbf{Mod}(\mathcal{A}) . \quad (12)$$

Now, if $(h, k) : (U, U') \xrightarrow{\sim} (V, V')$ is a weak equivalence between cofibrant objects of the product model category $\mathbf{Mod}(\mathcal{A}) \times \mathbf{Mod}(\mathcal{A})$, i.e., if (h, k) is a pair of weak equivalences between cofibrant objects of $\mathbf{Mod}(\mathcal{A})$, then in view of (10), (11) and (12), the $\mathbf{Mod}(\mathcal{A})$ -morphism

$$h \otimes_{\mathcal{A}} k = (V \otimes_{\mathcal{A}} k) \circ (h \otimes_{\mathcal{A}} U')$$

is a weak equivalence. Hence the functor

$$- \otimes_{\mathcal{A}} - : \mathbf{Mod}(\mathcal{A}) \times \mathbf{Mod}(\mathcal{A}) \rightarrow \mathbf{Mod}(\mathcal{A})$$

meets the requirements of Theorem 1 and all reasonable definitions of its left derived functor lead to isomorphic results.

Remark 2. Let us fix a functorial *Cof - TrivFib* factorization system of $\mathbf{Mod}(\mathcal{A})$, so that the cofibrant F-replacement functor Q and the natural weak equivalence $q : Q \xrightarrow{\sim} \text{id}$ are fixed as well. In the following, we write

$$- \otimes_{\mathcal{A}}^{\mathbb{L}} - := \mathbb{L}_{Q \times Q}^S(- \otimes_{\mathcal{A}} -) .$$

However, the left Quillen endofunctor $U \otimes_{\mathcal{A}} -$ (U : cofibrant) not only sends weak equivalences between cofibrant objects to weak equivalences, but it preserves *all* weak equivalences ([10, Proposition 3.4.1.]). Even better:

Proposition 2. *Let $M, P, S \in \mathbf{Mod}(\mathcal{A})$ and let $g : P \rightarrow S$ be a weak equivalence.*

1. *If M is cofibrant, the map $\text{id}_M \otimes g : M \otimes_{\mathcal{A}} P \rightarrow M \otimes_{\mathcal{A}} S$ is a weak equivalence.*

2. If P and S are cofibrant, the map $\text{id}_M \otimes g : M \otimes_{\mathcal{A}} P \rightarrow M \otimes_{\mathcal{A}} S$ is a weak equivalence.

Proof. We must still prove Item 2. In view of Item 1 and Equation (10), we have the commutative diagram

$$\begin{array}{ccccc} QM \otimes_{\mathcal{A}} P & \xrightarrow{\sim} & QM \otimes_{\mathcal{A}} S & \xrightarrow{\sim} & M \otimes_{\mathcal{A}} S \\ \downarrow \sim & & & \nearrow & \\ M \otimes_{\mathcal{A}} P & & & & \end{array}$$

From the 2-out-of-3 property it follows that $M \otimes_{\mathcal{A}} P \rightarrow M \otimes_{\mathcal{A}} S$ is a weak equivalence as well. \square

The previous proposition implies that in our model categorical context it suffices, just like in classical homological algebra, to resolve one of the two arguments:

Theorem 2. *The composites $L \star \otimes_{\mathcal{A}} \star (q \times \text{id}_Q)$ and $L \star \otimes_{\mathcal{A}} \star (\text{id}_Q \times q)$, where \star denotes whiskering and id_Q is the natural automorphism of Q , are canonical natural isomorphisms*

$$\text{Ho}(L \circ (\text{id}_{\text{Mod}(\mathcal{A})} - \otimes_{\mathcal{A}} Q-)) \doteq - \otimes_{\mathcal{A}}^L - := \text{Ho}(L \circ (Q - \otimes_{\mathcal{A}} Q-)) \doteq \text{Ho}(L \circ (Q - \otimes_{\mathcal{A}} \text{id}_{\text{Mod}(\mathcal{A})} -)) . \quad (13)$$

Hence, for every $M, N \in \text{Mod}(\mathcal{A})$, we have

$$M \otimes_{\mathcal{A}}^L N := QM \otimes_{\mathcal{A}} QN \approx QM \otimes_{\mathcal{A}} N \approx M \otimes_{\mathcal{A}} QN \quad (14)$$

in $\text{Mod}(\mathcal{A})$. Further:

$$M \otimes_{\mathcal{A}}^L N := QM \otimes_{\mathcal{A}} QN \approx \tilde{C}M \otimes_{\mathcal{A}} \tilde{C}N \approx CM \otimes_{\mathcal{A}} CN \approx QM \otimes_{\mathcal{A}} CN , \quad (15)$$

where \tilde{C} (resp., C) denotes any local cofibrant F -replacement (resp., any local cofibrant replacement).

Proof. Of course, it suffices to prove one of the two statements of (13), for example the second. If $f : M \xrightarrow{\sim} N$ and $g : P \xrightarrow{\sim} S$ are two weak equivalences in $\text{Mod}(\mathcal{A})$, it follows from

$$Qf \otimes_{\mathcal{A}} g = (Qf \otimes_{\mathcal{A}} S) \circ (QM \otimes_{\mathcal{A}} g)$$

and Proposition 2 that the functor $L \circ (Q - \otimes_{\mathcal{A}} \text{id}_{\text{Mod}(\mathcal{A})} -)$ sends every weak equivalence of $\text{Mod}(\mathcal{A})^{\times 2}$ to an isomorphism of $\text{Ho}(\text{Mod}(\mathcal{A}))$, so that its factorization $\text{Ho}(L \circ (Q - \otimes_{\mathcal{A}} \text{id}_{\text{Mod}(\mathcal{A})} -))$ is well-defined. Due to [15, Lemma 1] we only need to build a canonical natural isomorphism

$$L \circ (Q - \otimes_{\mathcal{A}} Q-) \doteq L \circ (Q - \otimes_{\mathcal{A}} \text{id}_{\text{Mod}(\mathcal{A})} -) .$$

Since $\text{id}_Q : Q \Rightarrow Q$ is a natural automorphism and $q : Q \Rightarrow \text{id}_{\text{Mod}(\mathcal{A})}$ is a natural weak equivalence, the horizontal composite $\theta := L \star \otimes_{\mathcal{A}} \star (\text{id}_Q \times q)$ is a canonical natural transformation

$$\theta : L \circ (Q - \otimes_{\mathcal{A}} Q -) \Rightarrow L \circ (Q - \otimes_{\mathcal{A}} \text{id}_{\text{Mod}(\mathcal{A})} -)$$

whose components $\theta_{M,N} = L(QM \otimes_{\mathcal{A}} q_N)$ are isomorphisms. Equation (14) is a direct consequence of [16, Proposition 2] and Equation (15) follows from Theorem 1 and Equation (8). \square

4 Graded Tor functors in modules over a graded \mathcal{D} -algebra

To make sense of the graded Tor functors in $\text{Mod}(\mathfrak{A})$ one has to show that the category in hands is abelian with enough projectives, and that the graded tensor product is right exact. In short, the first claim is proven using general results on the category of modules over a monoid in an abelian symmetric monoidal category, while right exactness essentially follows from the fact that the tensor product in question is merely the tensor product over the underlying graded commutative ring \mathfrak{A} , additionally endowed with the action of differential operators via the Liebniz rule.

Remark 3. It is well-known [3, 2.3.5] that the category $\text{Mod}(\mathbf{A})$ ($\mathbf{A} \in \mathcal{D}\mathbf{A}$) coincides with the category $\mathbf{A}[\mathcal{D}]M$ of modules over the ring $\mathbf{A}[\mathcal{D}] := \mathbf{A} \otimes_{\mathcal{O}} \mathcal{D}$ (tensor product over functions) of linear differential operators with coefficients in \mathbf{A} , whose multiplication is defined for instance in the section 5 of [27]. Moreover, the category $\text{Mod}(\mathbf{A})$ coincides with the category $\mathbf{DGA}[\mathcal{D}]M$ of (non-negatively) graded chain complexes in $\mathbf{A}[\mathcal{D}]M$. However, the fact that $\mathbf{A} \in \mathcal{D}\mathbf{A}$ is not graded is crucial in the proof of these results. Therefore we will not use notations like $\mathbf{G}\mathfrak{A}[\mathcal{D}]M$ and $\mathbf{DGA}[\mathcal{D}]M$ for the categories $\text{Mod}(\mathfrak{A})$ and $\text{Mod}(\mathcal{A})$, respectively.

The category $\text{Mod}(\mathbf{A})$ is a category of modules over a ring and is therefore abelian and has enough projectives (at least if we assume the axiom of choice, what we do systematically). Further:

Proposition 3. *The categories $\text{Mod}(\mathfrak{A})$ and $\text{Mod}(\mathcal{A})$ are abelian and $\text{Mod}(\mathfrak{A})$ has enough projectives.*

Proof. The categories \mathbf{GDM} and $\mathbf{DGD}\mathbf{M}$ are abelian symmetric monoidal categories (see The Stacks project, Section 12.16.2 and see [9]). It therefore follows from [1, Theorem 3.6] that

the categories $\text{Mod}(\mathfrak{A})$ and $\text{Mod}(\mathcal{A})$ are also abelian if the functors

$$\mathfrak{A} \otimes_{\mathcal{O}} - : \mathbf{GDM} \rightarrow \mathbf{GDM} \quad \text{and} \quad \mathcal{A} \otimes_{\mathcal{O}} - : \mathbf{DGDM} \rightarrow \mathbf{DGDM}$$

are additive and preserve cokernels. As the \mathcal{O} -module isomorphisms

$$M \otimes_{\mathcal{O}} \{0\} \cong \{0\} \quad \text{and} \quad M \otimes_{\mathcal{O}} (N \oplus P) \cong M \otimes_{\mathcal{O}} N \oplus M \otimes_{\mathcal{O}} P$$

respect the gradings and the actions of vector fields, as well as the differentials if they are present, both left tensoring functors are additive. As for cokernels, notice that both tensoring functors are right exact in \mathcal{O} -modules, hence commute with all finite colimits, in particular with cokernels in \mathcal{O} -modules. However, cokernels in (differential) graded \mathcal{D} -modules are computed degree-wise in \mathcal{D} -modules and cokernels in \mathcal{D} -modules coincide with cokernels in \mathcal{O} -modules and are just the quotients by the set-theoretical images. From here it follows that the functors $\mathfrak{A} \otimes_{\mathcal{O}} -$ and $\mathcal{A} \otimes_{\mathcal{O}} -$ commute with cokernels in \mathbf{GDM} and \mathbf{DGDM} , respectively, and that the categories $\text{Mod}(\mathfrak{A})$ and $\text{Mod}(\mathcal{A})$ are abelian.

Before we go any further, we recall some general definitions.

Definition 1. 1. An object C of a category \mathbf{C} is *projective* if it has the LLP with respect to epimorphisms, or, equivalently, if its covariant Hom-functor $\text{Hom}(C, -)$ respects epimorphisms.

2. A category \mathbf{C} has *enough projectives* if for any $C \in \mathbf{C}$ there is an epimorphism $P \rightarrow C$ from a projective object P .
3. A *projective resolution* of an object A of an abelian category \mathbf{A} is a chain complex $P_{\bullet} \in \mathbf{Ch}_+(\mathbf{A})$ made of projective objects P_i together with a quasi-isomorphism $P_{\bullet} \rightarrow A$, or, equivalently, it is an exact sequence

$$\cdots \rightarrow P_1 \rightarrow P_0 \rightarrow A \rightarrow 0$$

with projective nodes P_i .

Remark 4. The projective resolutions of $A \in \mathbf{A}$ are exactly the cofibrant replacements of $A \in \mathbf{Ch}_+(\mathbf{A})$ in the projective model structure of $\mathbf{Ch}_+(\mathbf{A})$. If \mathbf{A} has enough projectives any of its objects has a projective resolution.

We still have to prove that $\text{Mod}(\mathfrak{A})$ has enough projectives. It is clear that the projective objects in \mathbf{GDM} are exactly the direct sums over \mathbb{N} of projective objects in \mathbf{DM} (projective

objects in \mathbf{GDM} are more complicated) and that the category $\mathcal{D}\mathbf{M}$ of modules over \mathcal{D} has enough projectives. If $M = \bigoplus_i M_i \in \mathbf{GDM}$, there exists for each $M_i \in \mathcal{D}\mathbf{M}$ a surjective \mathcal{D} -linear map $P_i \rightarrow M_i$ out of a projective $P_i \in \mathcal{D}\mathbf{M}$, hence, there exists a surjective degree-respecting \mathcal{D} -linear map $h : P \rightarrow M$ out of the projective $P := \bigoplus_i P_i \in \mathbf{GDM}$, so that also \mathbf{GDM} has enough projectives. To see that the same holds for $\mathbf{Mod}(\mathfrak{A})$, recall the free-forgetful adjunction

$$\mathfrak{A} \otimes_{\mathcal{O}} - : \mathbf{GDM} \rightleftarrows \mathbf{Mod}(\mathfrak{A}) : \text{For} ,$$

i.e., the functorial bijections

$$\Phi_{NM} : \text{Hom}_{\mathbf{Mod}(\mathfrak{A})}(\mathfrak{A} \otimes_{\mathcal{O}} N, M) \cong \text{Hom}_{\mathbf{GDM}}(N, \text{For } M) ,$$

which are rooted in the observation that \mathfrak{A} -linear maps

$$k(a \otimes n) = a \triangleleft k(1_{\mathfrak{A}} \otimes n)$$

are fully defined by their values on the elements $1_{\mathfrak{A}} \otimes n \cong n \in N$. From here it follows that for $M \in \mathbf{Mod}(\mathfrak{A})$ the above surjective \mathbf{GDM} -map $h : P \rightarrow M$ is the image by Φ of a unique $\mathbf{Mod}(\mathfrak{A})$ -map $k : \mathfrak{A} \otimes_{\mathcal{O}} P \rightarrow M$. As k is obviously surjective, we can conclude that $\mathbf{Mod}(\mathfrak{A})$ has enough projectives as soon as we have shown that $\mathfrak{A} \otimes_{\mathcal{O}} P \in \mathbf{Mod}(\mathfrak{A})$ is projective since $P \in \mathbf{GDM}$ is. However, this is obvious. Indeed, if we choose in $\mathbf{Mod}(\mathfrak{A})$ a surjective morphism $f : R \rightarrow S$ and a morphism $g_{\mathfrak{A}} : \mathfrak{A} \otimes_{\mathcal{O}} P \rightarrow S$, its image $g := \Phi(g_{\mathfrak{A}}) : P \rightarrow S$ lifts to R , i.e., there is a \mathbf{GDM} -morphism $\ell : P \rightarrow R$ such that $f \circ \ell = g$. The image $\ell_{\mathfrak{A}} := \Phi^{-1}(\ell) : \mathfrak{A} \otimes_{\mathcal{O}} P \rightarrow R$ then lifts $g_{\mathfrak{A}}$, i.e., $f \circ \ell_{\mathfrak{A}} = g_{\mathfrak{A}}$. \square

For $M \in \mathbf{Mod}(\mathfrak{A})$, the tensor product $M \otimes_{\mathfrak{A}} - : \mathbf{Mod}(\mathfrak{A}) \rightarrow \mathbf{Mod}(\mathfrak{A})$ is a covariant functor between abelian categories, whose source has enough projectives. We can therefore consider its classical left derived functors $L_p(M \otimes_{\mathfrak{A}} -) : \mathbf{Mod}(\mathfrak{A}) \rightarrow \mathbf{Mod}(\mathfrak{A})$ ($p \in \mathbb{N}$), if the tensor product is right exact.

Remark 5. In Section 3, we defined the left derived functor $M \otimes_{\mathcal{A}}^{\mathbb{L}} -$ (in the model categorical sense) of the tensor product in $\mathbf{Mod}(\mathcal{A})$, for $\mathcal{A} \in \mathbf{DGA}$. In the present section, we are considering the left derived functors $L_p(M \otimes_{\mathfrak{A}} -)$ (in the classical homological algebraic sense) of the tensor product in $\mathbf{Mod}(\mathfrak{A})$, for $\mathfrak{A} \in \mathbf{GDA}$. We will define the graded Tor functor using the functors $L_p(M \otimes_{\mathfrak{A}} -)$, and the Tor spectral sequence will connect the graded Tor functor with the derived functor $M \otimes_{\mathcal{A}}^{\mathbb{L}} -$.

Remark 6. Let us also stress the so far implicit fact that in this text we consider left module structures, left \mathcal{D} -modules and left modules over the graded-commutative algebras \mathcal{A} and \mathfrak{A}

– although a left action implements a right action via $m \triangleright a = (-1)^{am}a \triangleleft m$ (the numbers a and m in the exponent of -1 are the degrees of the vectors a and m) and vice versa.

Proposition 4. *The tensor product functor*

$$M \otimes_{\mathfrak{A}} - : \mathbf{Mod}(\mathfrak{A}) \rightarrow \mathbf{Mod}(\mathfrak{A})$$

is right exact for every $M \in \mathbf{Mod}(\mathfrak{A})$. The same is true for $- \otimes_{\mathfrak{A}} M$.

Proof. Notice first that the cokernels, kernels and images of the morphisms $\ell : M \rightarrow N$ of $\mathbf{Mod}(\mathfrak{A})$ are those of the underlying morphisms of \mathbf{GDM} (i.e., the direct sums of the cokernels, kernels and images of the component morphisms of \mathcal{D} -modules). Indeed, the set-theoretical image $\text{im } \ell \in \mathbf{GDM}$ is an \mathfrak{A} -submodule of N , so that the cokernel $\text{coker } \ell = N / \text{im } \ell \in \mathbf{GDM}$ is also an \mathfrak{A} -module. It is easily seen that $\text{coker } \ell \in \mathbf{Mod}(\mathfrak{A})$ is the cokernel of ℓ in $\mathbf{Mod}(\mathfrak{A})$. Similarly, kernels in $\mathbf{Mod}(\mathfrak{A})$ are given by the corresponding set-theoretical kernels in \mathbf{GDM} equipped with their induced \mathfrak{A} -module structure. The same statement is therefore also valid for images.

We are now ready to prove that $M \otimes_{\mathfrak{A}} -$ is a right exact endofunctor of $\mathbf{Mod}(\mathfrak{A})$. Let therefore

$$P \xrightarrow{f} R \xrightarrow{g} S \rightarrow 0 \tag{16}$$

be an exact sequence in the abelian category $\mathbf{Mod}(\mathfrak{A})$, i.e., a sequence in $\mathbf{Mod}(\mathfrak{A})$ such that at each spot the image of the incoming map coincides with the kernel of the outgoing one, i.e., in view of what we just said, a sequence such that at each spot the set-theoretical image coincides with the set-theoretical kernel.

Observe now that $\mathfrak{A} \in \mathbf{GDA}$ is a graded-commutative ring $\tilde{\mathfrak{A}}$ when equipped with its additive group structure and its multiplication. Any object $P \in \mathbf{Mod}(\mathfrak{A})$ is in particular an additive group that comes equipped with a degree zero, \mathcal{D} -linear $\nu : \mathfrak{A} \otimes_0 P \rightarrow P$ that satisfies the usual associativity and unitality requirements. Since ν is 0 -linear on \otimes_0 , it is 0 -bilinear on \times , so in particular biadditive. Hence ν is an action of the ring $\tilde{\mathfrak{A}}$ on P , which is therefore an $\tilde{\mathfrak{A}}$ -module $\tilde{P} : \tilde{P} \in \mathbf{Mod}(\tilde{\mathfrak{A}})$. Finally, a $\mathbf{Mod}(\mathfrak{A})$ -morphism $f : P \rightarrow R$ is a degree zero, \mathcal{D} - and \mathfrak{A} -linear map, so it is in particular a $\mathbf{Mod}(\tilde{\mathfrak{A}})$ -morphism $\tilde{f} : \tilde{P} \rightarrow \tilde{R}$. From what we just said follows that the exact $\mathbf{Mod}(\mathfrak{A})$ -sequence (16) can be interpreted as an exact $\mathbf{Mod}(\tilde{\mathfrak{A}})$ -sequence

$$\tilde{P} \xrightarrow{\tilde{f}} \tilde{R} \xrightarrow{\tilde{g}} \tilde{S} \rightarrow 0. \tag{17}$$

Actually, the $\mathbf{Mod}(\mathfrak{A})$ -sequence (16) is exact if and only if the corresponding $\mathbf{Mod}(\tilde{\mathfrak{A}})$ -sequence (17) is.

We have to show that $M \otimes_{\mathfrak{A}} -$ respects exactness of (16), i.e., that the $\text{Mod}(\mathfrak{A})$ -sequence

$$M \otimes_{\mathfrak{A}} P \xrightarrow{M \otimes f} M \otimes_{\mathfrak{A}} R \xrightarrow{M \otimes g} M \otimes_{\mathfrak{A}} S \longrightarrow 0 \quad (18)$$

is exact, or, equivalently, that the corresponding $\text{Mod}(\tilde{\mathfrak{A}})$ -sequence

$$(M \otimes_{\mathfrak{A}} P) \xrightarrow{\sim} (M \otimes_{\mathfrak{A}} R) \xrightarrow{\sim} (M \otimes_{\mathfrak{A}} S) \longrightarrow 0 \quad (19)$$

is, i.e., that at each of its spots the set-theoretical image coincides with the set-theoretical kernel. However, the tensor product $- \otimes_{\tilde{\mathfrak{A}}} -$ in $\text{Mod}(\tilde{\mathfrak{A}})$ over the graded-commutative ring $\tilde{\mathfrak{A}}$ being the standard tensor product over the non-commutative ring $\tilde{\mathfrak{A}}$, the functor $\tilde{M} \otimes_{\tilde{\mathfrak{A}}} -$ is right exact, so that the sequence

$$\tilde{M} \otimes_{\tilde{\mathfrak{A}}} \tilde{P} \xrightarrow{\tilde{M} \otimes \tilde{f}} \tilde{M} \otimes_{\tilde{\mathfrak{A}}} \tilde{R} \xrightarrow{\tilde{M} \otimes \tilde{g}} \tilde{M} \otimes_{\tilde{\mathfrak{A}}} \tilde{S} \rightarrow 0 \quad (20)$$

is exact in $\text{Mod}(\tilde{\mathfrak{A}})$, i.e., at each spot the set-theoretical image coincides with the set-theoretical kernel. It is now sufficient to explain why Sequence (19) and Sequence (20) are made of the same sets and the same set-theoretical maps. The maps have obviously both the same values as $M \otimes f$ and $M \otimes g$, respectively. It is therefore enough to prove that $(M \otimes_{\mathfrak{A}} P) \xrightarrow{\sim} \tilde{M} \otimes_{\tilde{\mathfrak{A}}} \tilde{P}$. By definition the latter set is made of the finite sums of decomposed tensors $m \otimes p$, where \otimes is weakly \mathfrak{A} -bilinear, i.e., is biadditive and such that

$$(a \triangleleft m) \otimes p = (-1)^{am} m \otimes (a \triangleleft p), \quad (21)$$

where we denoted the vectors by the same symbols as their degree. The first tensor product $M \otimes_{\mathfrak{A}} P$ is defined as the cokernel in \mathbf{GDM} of the map

$$\alpha : M \otimes_{\mathfrak{O}} \mathfrak{A} \otimes_{\mathfrak{O}} P \ni (m, a, p) \mapsto (a \triangleleft m) \otimes p - (-1)^{am} m \otimes (a \triangleleft p) \in M \otimes_{\mathfrak{O}} P.$$

As said above $\text{coker } \alpha \in \mathbf{GDM}$ is the quotient $(M \otimes_{\mathfrak{O}} P) / \text{im } \alpha$, so it is given by the finite sums of decomposed tensors $m \otimes p$, where \otimes is \mathfrak{O} -bilinear and weakly \mathfrak{A} -bilinear. This graded \mathcal{D} -module inherits a compatible \mathfrak{A} -action

$$a \triangleleft (m \otimes p) = (a \triangleleft m) \otimes p = (-1)^{am} m \otimes (a \triangleleft p).$$

When passing to the $\tilde{\mathfrak{A}}$ -module $(M \otimes_{\mathfrak{A}} P) \xrightarrow{\sim} \tilde{M} \otimes_{\tilde{\mathfrak{A}}} \tilde{P}$, we forget in particular the \mathfrak{O} -action, so this set is made of the finite sums of decomposed tensors $m \otimes p$, where \otimes is weakly \mathfrak{A} -bilinear: $(M \otimes_{\mathfrak{A}} P) \xrightarrow{\sim} \tilde{M} \otimes_{\tilde{\mathfrak{A}}} \tilde{P}$. \square

Definition 2. Let $\mathfrak{A} \in \mathbf{GDA}$ and $M \in \mathbf{Mod}(\mathfrak{A})$. We refer to the p -th left derived functor of the covariant right exact endofunctor $M \otimes_{\mathfrak{A}} -$ as the p -th graded Tor endofunctor $\mathrm{Tor}_p^{\mathfrak{A}}(M, -)$ of $\mathbf{Mod}(\mathfrak{A})$. For any $N \in \mathbf{Mod}(\mathfrak{A})$, we have $\mathrm{Tor}_p^{\mathfrak{A}}(M, N) \in \mathbf{Mod}(\mathfrak{A}) \subset \mathbf{GDM}$ and we denote the q -th homogeneous component by $\mathrm{Tor}_p^{\mathfrak{A}}(M, N)_q \in \mathbf{DM}$.

Let us recall that in order to compute $\mathrm{Tor}_p^{\mathfrak{A}}(M, N)$, we choose a projective resolution $P_{\bullet} \rightarrow N$ of N in the abelian category $\mathbf{Mod}(\mathfrak{A})$, i.e., we choose a cofibrant replacement $\mathcal{Q}N$ of N in the projective model structure of $\mathbf{Ch}_+(\mathbf{Mod}(\mathfrak{A}))$ (in particular the cofibrant replacement $\mathcal{Q}N$ that is provided by a functorial ‘cofibration – trivial fibration’ factorization), and we compute the p -th homology \mathfrak{A} -module of $M \otimes_{\mathfrak{A}} \mathcal{Q}N \in \mathbf{Ch}_+(\mathbf{Mod}(\mathfrak{A}))$. In other words, we have

$$\mathrm{Tor}_p^{\mathfrak{A}}(M, -) : \mathbf{Mod}(\mathfrak{A}) \xrightarrow{i} \mathbf{Ch}_+(\mathbf{Mod}(\mathfrak{A})) \xrightarrow{M \otimes \mathcal{Q}-} \mathbf{Ch}_+(\mathbf{Mod}(\mathfrak{A})) \xrightarrow{H_p} \mathbf{Mod}(\mathfrak{A}).$$

Here the first arrow i is the functor that sends a module (resp., a module morphism) to the corresponding chain complex concentrated in degree zero (resp., the corresponding chain map concentrated in degree zero), the second arrow $M \otimes \mathcal{Q}-$ is the composite of a cofibrant replacement functor in the projective model structure of chain complexes and the tensor product functor, whereas the last arrow H_p is the p -th homology functor. Finally, as in the classical situation, if we resolve M instead of N , we get the same result.

In view of Theorem 2, we have in $\mathbf{Mod}(\mathcal{A})$ ($\mathcal{A} \in \mathbf{DGDA}$) that

$$M \otimes_{\mathcal{A}}^{\mathbb{L}} N := QM \otimes_{\mathcal{A}} QN \approx QM \otimes_{\mathcal{A}} N \approx M \otimes_{\mathcal{A}} QN \in \mathbf{Mod}(\mathcal{A}) \subset \mathbf{DGDM} = \mathbf{Ch}_+(\mathbf{DM}) \quad (22)$$

(Q : cofibrant replacement functor in $\mathbf{Mod}(\mathcal{A})$), whereas in $\mathbf{Mod}(\mathfrak{A})$ ($\mathfrak{A} \in \mathbf{GDA}$) we have that

$$\mathrm{Tor}_p^{\mathfrak{A}}(M, N) = H_p(M \otimes_{\mathfrak{A}} \mathcal{Q}N) \in \mathbf{Mod}(\mathfrak{A}) \subset \mathbf{GDM} \quad (23)$$

(\mathcal{Q} : cofibrant replacement functor in $\mathbf{Ch}_+(\mathbf{Mod}(\mathfrak{A}))$). Notice further that, since $M \in \mathbf{Mod}(\mathcal{A}) \subset \mathbf{DGDM} = \mathbf{Ch}_+(\mathbf{DM})$ is not a chain complex of $\mathcal{A}[\mathcal{D}]$ -modules (see Remark 3), the homology module $H(M)$ is a priori just a graded \mathcal{D} -module. However, the homology module $H(\mathcal{A}) \in \mathbf{GDM}$ inherits the obvious multiplication, which makes it a graded \mathcal{D} -algebra \mathfrak{A} . It is also easily seen that the homology module $H(M) \in \mathbf{GDM}$ can be endowed with the canonical $H(\mathcal{A})$ -action, which makes it a module

$$H(M) \in \mathbf{Mod}(H(\mathcal{A})) = \mathbf{Mod}_{\mathbf{GDM}}(H(\mathcal{A}));$$

in fact, the functor $H : \mathbf{DGDM} \rightarrow \mathbf{GDM}$ restricts to a functor

$$H : \mathbf{Mod}(\mathcal{A}) \rightarrow \mathbf{Mod}(H(\mathcal{A})). \quad (24)$$

A possible relationship between $\otimes_{\mathcal{A}}^{\mathbb{L}}$ in (22) and $\text{Tor}^{\mathfrak{A}}$ in (23) should therefore involve

$$\text{Tor}_p^{H(\mathcal{A})}(H(M), H(N))_q \in \mathcal{DM} \quad \text{and} \quad H_{p+q}(M \otimes_{\mathcal{A}}^{\mathbb{L}} N) \in \mathcal{DM},$$

where $\mathcal{A} \in \mathsf{DGDA}$, $M, N \in \mathsf{Mod}(\mathcal{A})$ and $p, q \in \mathbb{N}$. The \mathcal{D} -generalization of Quillen's Tor spectral sequence specifies this relationship.

5 The \mathcal{D} -generalization of Quillen's Tor spectral sequence

Theorem 3. *For every $\mathcal{A} \in \mathsf{DGDA}$ and every $M, N \in \mathsf{Mod}(\mathcal{A}) := \mathsf{Mod}_{\mathsf{DGDM}}(\mathcal{A})$, there is a first quadrant spectral sequence E_{pq}^{\bullet} in the abelian category \mathcal{DM} , whose second sheet is $E_{pq}^2 = \text{Tor}_p^{H(\mathcal{A})}(H(M), H(N))_q$ and which converges to $H_{p+q}(M \otimes_{\mathcal{A}}^{\mathbb{L}} N)$:*

$$E_{pq}^2 = \text{Tor}_p^{H(\mathcal{A})}(H(M), H(N))_q \Rightarrow H_{p+q}(M \otimes_{\mathcal{A}}^{\mathbb{L}} N). \quad (25)$$

What we call Quillen's Tor spectral sequence is a similar spectral sequence in the category Ab of abelian groups, for a simplicial ring \mathcal{A} and a simplicial right (resp., left) \mathcal{A} -module M (resp., N) (for simplicial modules, see for instance 'The Stacks project, 21.40'; see also [4], where another type of modules over a varying ring appears). Roughly speaking the result connects the derived tensor product of the homology and the homology of the derived tensor product.

The proof of Theorem 3 relies on two propositions.

Proposition 5. *For every $\mathcal{A} \in \mathsf{DGDA}$, every $M \in \mathsf{DGDM}$ and every $N \in \mathsf{Mod}(\mathcal{A})$, there is a natural isomorphism*

$$(M \otimes_{\mathcal{O}} \mathcal{A}) \otimes_{\mathcal{A}} N \cong M \otimes_{\mathcal{O}} N$$

in $\mathsf{Mod}(\mathcal{A})$, where the \mathcal{A} -module structure of $M \otimes_{\mathcal{O}} N$ is canonically induced by the \mathcal{A} -module structure of N :

$$(\text{id}_M \otimes \nu_N) \circ (\text{com} \otimes \text{id}_N) : \mathcal{A} \otimes_{\mathcal{O}} M \otimes_{\mathcal{O}} N \ni a \otimes m \otimes n \mapsto (-1)^{am} m \otimes (a \triangleleft n) \in M \otimes_{\mathcal{O}} N,$$

where ν_N is the \mathcal{A} -module structure \triangleleft of N and com the braiding of DGDM . The same result is valid for $\mathfrak{A} \in \mathsf{GDA}$, $M \in \mathsf{GDM}$ and $N \in \mathsf{Mod}(\mathfrak{A})$.

Similar results exist in other settings. The main difference here is that $\mathcal{A} \in \mathbf{DGA}$ and $N \in \mathbf{DGM}$, but the \mathbf{DGM} -map $\nu_N : \mathcal{A} \otimes_{\mathcal{O}} N \rightarrow N$ is not \mathcal{D} -bilinear as the tensor product is over \mathcal{O} .

For the proof of Proposition 5, we refer the reader to Lemma 3.1.1. in [10].

For the next proposition we need some preparation. Let \mathbf{GSet} be the category whose objects are the (non-negatively) graded sets, i.e., the disjoint unions $I = \sqcup_{k \in \mathbb{N}} I_k$ of sets I_k indexed by the non-negative integers $k \in \mathbb{N}$, and whose morphisms are the grading preserving set-theoretical mappings. We systematically and implicitly consider such sets as well-ordered. Namely, we well-order all the sets I_k and then well-order I by declaring that the elements of I_k are smaller than the elements of I_ℓ , if $k < \ell$. Every $i \in I$ belongs to a unique I_k , so that we get the degree assigning map $n : I \ni i \mapsto n_i = k \in \mathbb{N}$. The functors

$$\mathfrak{F} : \mathbf{GSet} \rightleftarrows \mathbf{GDM} : \text{For}_{\mathcal{D}} \quad (26)$$

form a free-forgetful adjunction. More precisely, the free graded \mathcal{D} -module over a graded set I is the free \mathcal{D} -module

$$\mathfrak{F}(I) = \bigoplus_{i \in I} \mathcal{D} i = \bigoplus_{k \in \mathbb{N}} \bigoplus_{i_k \in I_k} \mathcal{D} i_k$$

over the set I , but with i_k in degree $n_{i_k} = k$, or, equivalently, is the direct sum over all degrees $k \in \mathbb{N}$ of the free \mathcal{D} -modules over the sets I_k . In other words, we set

$$\mathfrak{F}(I) = \bigoplus_{i \in I} \mathcal{D} \mathbf{1}_i[n_i] = \bigoplus_{i \in I} S_i^{n_i} \in \mathbf{GDM} ,$$

where $\mathbf{1}[n_i]$ is the generator of the n_i -sphere S^{n_i} (the n_i -sphere is concentrated on \mathcal{D} in degree $n_i \in \mathbb{N}$). When composing, for $\mathfrak{A} \in \mathbf{GDA}$, the adjunction (26) with the adjunction

$$\mathfrak{A} \otimes_{\mathcal{O}} - : \mathbf{GDM} \rightleftarrows \mathbf{Mod}(\mathfrak{A}) : \text{For}_{\mathfrak{A}} ,$$

we see that the free \mathfrak{A} -module in \mathbf{GDM} over a graded set I with degree assigning map n is

$$\mathfrak{A} \otimes_{\mathcal{O}} (\bigoplus_{i \in I} S_i^{n_i}) = \mathfrak{A} \triangleleft (1_{\mathfrak{A}} \otimes_{\mathcal{O}} (\bigoplus_{i \in I} S_i^{n_i})) \in \mathbf{Mod}(\mathfrak{A}) .$$

Hence:

Lemma 1. *If $P \in \mathbf{Mod}(\mathfrak{A})$, a $\mathbf{Mod}(\mathfrak{A})$ -morphism*

$$q : \mathfrak{A} \otimes_{\mathcal{O}} (\bigoplus_{i \in I} S_i^{n_i}) \rightarrow P \quad (27)$$

is uniquely defined by its values

$$q(1_{\mathfrak{A}} \otimes \mathbf{1}_i[n_i]) \in P_{n_i} \quad (i \in I) .$$

A similar result holds in the differential graded setting. More precisely, from item (i) of Lemma 2.3.1. in [10] it follows that, if $\mathcal{A} \in \mathbf{DGDA}$, the graded \mathcal{D} -module $\mathcal{A} \otimes_{\mathcal{O}} (\bigoplus_{i \in I} S_i^{n_i})$ equipped with its natural differential $d_{\mathcal{A}} \otimes \text{id}$ and its natural \mathcal{A} -action is a module in $\mathbf{Mod}(\mathcal{A})$. From item (ii) of the same lemma we get:

Lemma 2. *If $P \in \mathbf{Mod}(\mathcal{A})$, a $\mathbf{Mod}(\mathcal{A})$ -morphism*

$$q : \mathcal{A} \otimes_{\mathcal{O}} (\bigoplus_{i \in I} S_i^{n_i}) \rightarrow P \quad (28)$$

is uniquely defined by its values

$$q(1_{\mathcal{A}} \otimes \mathbf{1}_i[n_i]) \in P_{n_i} \cap \ker d_P \quad (i \in I)$$

(provided the source module is equipped with its natural differential $\text{id}_{\mathcal{A}} \otimes \text{id}$).

Indeed, it is then enough to extend q as a \mathcal{D} -linear map to the direct sum and as an \mathcal{A} -linear map to the tensor product.

Proposition 6. *For $\mathcal{A} \in \mathbf{DGDA}$ and $P, N \in \mathbf{Mod}(\mathcal{A}) := \mathbf{Mod}_{\mathbf{GDM}}(\mathcal{A})$, with P a cofibrant object whose homology $H(P)$ is a free $H(\mathcal{A})$ -module in \mathbf{GDM} , there exists an isomorphism*

$$H(P) \otimes_{H(\mathcal{A})} H(N) \cong H(P \otimes_{\mathcal{A}} N)$$

in the category $\mathbf{Mod}(H(\mathcal{A})) := \mathbf{Mod}_{\mathbf{GDM}}(H(\mathcal{A}))$.

Proof. It is natural to interpret the graded \mathcal{D} -module

$$S := \bigoplus_{i \in I} S_i^{n_i}$$

as a differential graded \mathcal{D} -module with zero differential (since each term is such a differential graded \mathcal{D} -module – we have used this fact already above). Hence S is a chain complex of \mathcal{O} -modules that is in each degree k a direct sum $S(k)$ of copies of \mathcal{D} . Since \mathcal{D} is \mathcal{O} -flat ([10, Proposition 1.0.11.]), this implies that $S(k)$, $d(S(k)) = 0$ and $H_k(S) = S(k)$ are \mathcal{O} -flat, so that for every $M \in \mathbf{GDM}$, we have $H(S \otimes_{\mathcal{O}} M) \cong S \otimes_{\mathcal{O}} H(M)$, in view of Künneth's theorem (the homology does of course not depend on whether we interpret a complex as complex of \mathcal{D} -modules or as complex of \mathcal{O} -modules). In particular:

Lemma 3. *If $\mathcal{A} \in \mathbf{DGDA}$ and $N \in \mathbf{Mod}(\mathcal{A})$, we have*

$$S \otimes_{\mathcal{O}} H(N) \cong H(S \otimes_{\mathcal{O}} N) \quad \text{and} \quad H(\mathcal{A}) \otimes_{\mathcal{O}} S \cong H(\mathcal{A} \otimes_{\mathcal{O}} S) . \quad (29)$$

The first \mathbf{GOM} -isomorphism $s \otimes [n] \mapsto [s \otimes n]$ is also a $\mathbf{Mod}(H(\mathcal{A}))$ -isomorphism. A similar statement holds for the second isomorphism.

By assumption P is a cofibrant \mathcal{A} -module whose homology is

$$H(P) = H(\mathcal{A}) \otimes_{\mathcal{O}} S \cong H(\mathcal{A} \otimes_{\mathcal{O}} (\bigoplus_{i \in I} S_i^{n_i})) = H(\mathcal{A} \otimes_{\mathcal{O}} (\bigoplus_{i \in I} \mathcal{D} \mathbf{1}_i[n_i])) .$$

As mentioned above, we view I as a well-ordered set; moreover, we have $d(\mathbf{1}_i[n_i]) = 0$, so that the differential is obviously lowering. Hence, the \mathcal{A} -module $\mathcal{A} \otimes_{\mathcal{O}} (\bigoplus_{i \in I} \mathcal{D} \mathbf{1}_i[n_i])$ is a Sullivan module and therefore a cofibrant module.

In the following, we denote homology classes as usual with $[-]$. Confusion with the degree shift $[n_i]$ is excluded, since the meaning of $[-]$ always comes from the context. Since $[1_{\mathcal{A}}] \otimes \mathbf{1}_i[n_i] \in H_{n_i}(P)$ for every i , we have $[1_{\mathcal{A}}] \otimes \mathbf{1}_i[n_i] = [p_i]$, with $p_i \in P_{n_i} \cap \ker d_P$. Choosing a representative p_i for each $i \in I$, we get values

$$q(1_{\mathcal{A}} \otimes \mathbf{1}_i[n_i]) = p_i \in P_{n_i} \cap \ker d_P \quad (i \in I) ,$$

which (see Lemma (2)) define a $\text{Mod}(\mathcal{A})$ -morphism

$$q : \mathcal{A} \otimes_{\mathcal{O}} (\bigoplus_{i \in I} S_i^{n_i}) \rightarrow P .$$

If

$$H(q) : H(\mathcal{A}) \otimes_{\mathcal{O}} (\bigoplus_{i \in I} S_i^{n_i}) \rightarrow H(P)$$

denotes the induced $\text{Mod}(H(\mathcal{A}))$ -morphism in homology (see Equation (24)), we have

$$H(q)([1_{\mathcal{A}}] \otimes \mathbf{1}_i[n_i]) \cong H(q)[1_{\mathcal{A}} \otimes \mathbf{1}_i[n_i]] = [q(1_{\mathcal{A}} \otimes \mathbf{1}_i[n_i])] = [p_i] = [1_{\mathcal{A}}] \otimes \mathbf{1}_i[n_i] \in H_{n_i}(P)$$

($i \in I$). We know (see Lemma 1) with $\mathfrak{A} = H(\mathcal{A})$) that these values define a unique $\text{Mod}(H(\mathcal{A}))$ -morphism $h : H(\mathcal{A}) \otimes_{\mathcal{O}} S \rightarrow H(P)$. In the situation under consideration this morphism h is identity. Hence $H(q) = \text{id}$, so that q is a weak equivalence between cofibrant \mathcal{A} -modules. Recall that a weak equivalence $f : K \rightarrow L$ in $\text{Mod}(\mathcal{A})$ is a $\text{Mod}(\mathcal{A})$ -morphism whose underlying $\text{DGD}\mathcal{M}$ -morphism is a quasi-isomorphism. It is easy to check though that $H(f) : H(K) \rightarrow H(L)$ is not only an isomorphism in $\text{G}\mathcal{D}\mathcal{M}$, but also an isomorphism in $\text{Mod}(H(\mathcal{A}))$. From Proposition 2 (with $g = q$) we now get a $\text{Mod}(H(\mathcal{A}))$ -isomorphism

$$H((\mathcal{A} \otimes_{\mathcal{O}} S) \otimes_{\mathcal{A}} N) = H((\mathcal{A} \otimes_{\mathcal{O}} (\bigoplus_{i \in I} S_i^{n_i})) \otimes_{\mathcal{A}} N) \cong H(P \otimes_{\mathcal{A}} N) . \quad (30)$$

Using Proposition 5 and Equation 29, we get $\text{Mod}(H(\mathcal{A}))$ -isomorphisms

$$H((\mathcal{A} \otimes_{\mathcal{O}} S) \otimes_{\mathcal{A}} N) \cong H(S \otimes_{\mathcal{O}} N) \cong S \otimes_{\mathcal{O}} H(N) , \quad (31)$$

and

$$H(P) \otimes_{H(\mathcal{A})} H(N) = (H(\mathcal{A}) \otimes_{\mathcal{O}} S) \otimes_{H(\mathcal{A})} H(N) \cong S \otimes_{\mathcal{O}} H(N) . \quad (32)$$

Equations 30, 31 and 32 finally give the desired result. \square

We are now prepared to prove Theorem 3.

Let $\mathcal{A} \in \mathbf{DGDA}$ and $M, N \in \mathbf{Mod}(\mathcal{A})$. Our goal is to prove that there is a first quadrant spectral sequence in the abelian category \mathbf{DM} such that

$$E_{pq}^2 = \mathrm{Tor}_p^{H(\mathcal{A})}(H(M), H(N))_q \Rightarrow H_{p+q}(M \otimes_{\mathcal{A}}^{\mathbb{L}} N), \quad (33)$$

for each $p, q \in \mathbb{N}$.

Recall first that to any first quadrant double complex $(C_{\bullet\bullet}, d^h, d^v)$ in an abelian category, one associates its total complex $(\mathrm{Tot}(C)_{\bullet}, d)$, whose grading is $\mathrm{Tot}(C)_n = \bigoplus_{p+q=n} C_{pq}$ and whose differential is $d = d^h + (-1)^p d^v$. The total complex admits two filtrations, the horizontal filtration $F_p^h(\mathrm{Tot}(C)) = \bigoplus_{r \leq p} C_{r\bullet}$ and the vertical filtration $F_p^v(\mathrm{Tot}(C)) = \bigoplus_{s \leq p} C_{\bullet s}$. The spectral sequence of the resulting horizontally filtered chain complex $(\mathrm{Tot}(C)_{\bullet}, d, F_{\bullet}^h)$ (resp., the vertically filtered chain complex $(\mathrm{Tot}(C)_{\bullet}, d, F_{\bullet}^v)$) is the horizontal (resp., vertical) spectral sequence ${}^h E_{pq}^{\bullet}$ (resp., ${}^v E_{pq}^{\bullet}$) of the double complex considered. The second sheet of the horizontal (resp., vertical) spectral sequence is

$${}^h E_{pq}^2 = H_p^h(H_q^v(C_{\bullet\bullet})) \quad (\text{resp., } {}^v E_{pq}^2 = H_p^v(H_q^h(C_{\bullet\bullet}))).$$

Moreover, this spectral sequence converges to the homology of the total complex:

$${}^h E_{pq}^2 = H_p^h(H_q^v(C_{\bullet\bullet})) \Rightarrow H_{p+q}(\mathrm{Tot}(C)) \quad (\text{resp., } {}^v E_{pq}^2 = H_p^v(H_q^h(C_{\bullet\bullet})) \Rightarrow H_{p+q}(\mathrm{Tot}(C))).$$

This means that for any $p, q \in \mathbb{N}$, the horizontal spectral sequence ${}^h E_{pq}^r$ ($r \in \mathbb{N}$) stabilizes at some $r(p, q) \in \mathbb{N}$ and

$${}^h E_{pq}^{r(p,q)} \cong {}^h G_p(H_{p+q}(\mathrm{Tot}(C))), \quad (34)$$

where the RHS is the p -th term of the graded space that is associated to the filtered space ${}^h F_{\bullet}(H_{p+q}(\mathrm{Tot}(C)))$, whose filtration is induced by the filtration F_{\bullet}^h of $\mathrm{Tot}(C)_{\bullet}$. The dual result holds for the vertical spectral sequence and the vertical filtration. If exactly one row or column of the grid ${}^h E_{pq}^2$ ($p, q \in \mathbb{N}$) does not vanish, the horizontal spectral sequence collapses at its second sheet and $H_n(\mathrm{Tot}(C))$ ($n \in \mathbb{N}$) is the unique non-zero ${}^h E_{pq}^2$ such that $p + q = n$ [36, Definition 5.2.7]. Again, the dual result holds for the vertical spectral sequence.

Also remember (see Section 4) that in order to compute the LHS of (33), we compute the p -th homology space of the q -th term of the tensor product of $H(N)$ with a projective resolution of $H(M)$ in $\mathbf{Mod}(H(\mathcal{A}))$. On the other hand, in order to compute the derived tensor product in the RHS of (33) (up to a zigzag of weak equivalences in $\mathbf{Mod}(\mathcal{A})$), we can

(see Theorem 2) tensor over \mathcal{A} the cofibrant replacement $X := QM$ of M in $\text{Mod}(\mathcal{A})$ with a cofibrant replacement of N .

Since $\text{Mod}(\mathcal{A})$ is a cofibrantly generated model category [10, Theorem 2.2.3.] (we denote its set of generating cofibrations by I), the small object argument gives a functorial factorization of $\text{Mod}(\mathcal{A})$ -morphisms $f : R \rightarrow S$ into an I -cell $i : R \rightarrow \mathfrak{Q}$ and a trivial fibration $p : \mathfrak{Q} \xrightarrow{\sim} S$. As the specific cofibrations we call I -cells are known to be relative Sullivan \mathcal{A} -modules [10, Definition 2.3.6.], we get in the case of $f : 0 \rightarrow N$ a cofibrant replacement \mathfrak{Q} of N that is a Sullivan \mathcal{A} -module

$$\mathfrak{Q} = \mathcal{A} \otimes_{\mathcal{O}} V := \mathcal{A} \otimes_{\mathcal{O}} \bigoplus_{k \in K} \mathcal{D} g_k$$

(where the set K is well-ordered, the generators g_k have homogeneous degrees $\deg(g_k) := n_k \in \mathbb{N}$ and the differential on \mathfrak{Q} is lowering).

In the following, we use the cofibrant replacement X of M and the cofibrant replacement \mathfrak{Q} of N . Further, we will construct a projective resolution of $H(M) \cong H(X)$ in $\text{Mod}(H(\mathcal{A}))$ as the sequence induced in homology by a resolution P_{\bullet} of X in $\text{Mod}(\mathcal{A})$. The statement (33) of Theorem 3 then results from the above machinery for double complexes, applied to the double complex $(P_{\bullet} \otimes_{\mathcal{A}} \mathfrak{Q})_{\bullet}$.

We construct the exact sequence $P_{\bullet} \rightarrow X \rightarrow 0$ just mentioned, using an iterative process.

Set $X_0 = X$ and fix a family of homogeneous generators $([x_i])_{i \in I_0}$ of $H(X_0) \in \text{Mod}(H(\mathcal{A}))$. Homogeneity means that x_i has a homogeneous degree $n_i \in \mathbb{N}$, so that $x_i \in X_{0,n_i} \cap \ker d_{X_0}$. Notice that the number of generators needs not be finite in any way, so that one can in particular choose as generators all the homogeneous elements of $H(X_0) = \bigoplus_n H_n(X_0)$. Next we define a $\text{Mod}(\mathcal{A})$ -morphism

$$q_0 : \mathcal{A} \otimes_{\mathcal{O}} \bigoplus_{i \in I_0} S_i^{n_i} \rightarrow X_0 .$$

According to Lemma 2 such a morphism is uniquely defined by its values $q_0(1_{\mathcal{A}} \otimes \mathbf{1}_i[n_i]) \in X_{0,n_i} \cap \ker d_{X_0}$, in particular by the choice of representatives x_i of the generating homology classes $[x_i]$. Any trivial cofibration - fibration decomposition of q_0 leads to a trivial cofibration $\iota_0 : \mathcal{A} \otimes_{\mathcal{O}} S_0 \xrightarrow{\sim} P_0$ and a fibration $\pi_0 : P_0 \rightarrow X_0$ of $\text{Mod}(\mathcal{A})$ (where S_0 is of course a compact notation for $\mathcal{A} \otimes_{\mathcal{O}} \bigoplus_{i \in I_0} S_i^{n_i}$). First, the weak equivalence ι_0 of $\text{Mod}(\mathcal{A})$ induces an isomorphism $H(\iota_0)$ of $\text{Mod}(H(\mathcal{A}))$. Since, due to Lemma 3, we have also a $\text{Mod}(H(\mathcal{A}))$ -isomorphism

$$H(\mathcal{A} \otimes_{\mathcal{O}} S_0) \simeq H(\mathcal{A}) \otimes_{\mathcal{O}} S_0 ,$$

we see that $H(P_0)$ is isomorphic as $H(\mathcal{A})$ -module to the free $H(\mathcal{A})$ -module $H(\mathcal{A}) \otimes_{\mathcal{O}} S_0$. Moreover, we already mentioned that modules of the type $\mathcal{A} \otimes_{\mathcal{O}} \bigoplus_{i \in I_0} S_i^{n_i}$ are Sullivan and therefore cofibrant \mathcal{A} -modules. Hence, in the composite

$$0 \rightarrowtail \mathcal{A} \otimes_{\mathcal{O}} \bigoplus_{i \in I_0} S_i^{n_i} \xrightarrow{\iota_0} P_0$$

is a cofibration and P_0 is a cofibrant \mathcal{A} -module. Second, the $\text{Mod}(\mathcal{A})$ -morphism π_0 is also a fibration in DGDM , hence it is surjective in positive degrees $n > 0$. On the other hand, the induced $\text{Mod}(H(\mathcal{A}))$ -morphism $H(\pi_0) : H(P_0) \rightarrow H(X_0)$ is surjective, since any homology class $[x] \in H(X_0)$ can be expressed as finite combination of generators, so that

$$[x] = \sum_i [a_i] \triangleleft [x_i] = \sum_i [a_i \triangleleft x_i] = H(q_0)[\sum_i a_i \otimes \mathbf{1}_i[n_i]] = H(\pi_0)(H(i_0)([\sum_i a_i \otimes \mathbf{1}_i[n_i]])) .$$

One easily checks that the $\text{Mod}(\mathcal{A})$ -morphism $\pi_0 : P_0 \rightarrow X_0$, which is surjective in positive degrees and surjective in homology, is actually surjective in all degrees.

In the previous paragraph, we started from a module $X_0 \in \text{Mod}(\mathcal{A})$ and constructed a cofibrant module $P_0 \in \text{Mod}(\mathcal{A})$ whose homology $H(P_0)$ is a free $H(\mathcal{A})$ -module, as well as a morphism $\pi_0 : P_0 \rightarrow X_0$ that is surjective and induces a surjective morphism in homology. We now set

$$X_1 := \ker(\pi_0 : P_0 \rightarrow X_0) \xhookrightarrow{k_1} P_0 ,$$

where $\xhookrightarrow{k_1}$ is the canonical inclusion of the \mathcal{A} -submodule X_1 into the \mathcal{A} -module P_0 , and we iterate the process of the previous paragraph. This way we get $\text{Mod}(\mathcal{A})$ -morphisms

$$\cdots \hookrightarrow P_2 \xrightarrow{\pi_2} X_2 \hookrightarrow P_1 \xrightarrow{\pi_1} X_1 \hookrightarrow P_0 \xrightarrow{\pi_0} X_0 \rightarrow 0 .$$

The $\text{Mod}(\mathcal{A})$ -sequences

$$0 \rightarrow X_{n+1} \xhookrightarrow{k_{n+1}} P_n \xrightarrow{\pi_n} X_n \rightarrow 0 \tag{35}$$

$(n \in \mathbb{N})$ and

$$\cdots \rightarrow P_2 \xrightarrow{\tilde{\pi}_2} P_1 \xrightarrow{\tilde{\pi}_1} P_0 \xrightarrow{\pi_0} X \rightarrow 0 \tag{36}$$

are obviously exact.

As mentioned earlier, we have to check whether the cofibrant resolution P_{\bullet} of $X = X_0$ in $\text{Mod}(\mathcal{A})$ (i.e., resolution made of cofibrant \mathcal{A} -modules) induces a projective resolution $H(P_{\bullet})$ of $H(X)$ in $\text{Mod}(H(\mathcal{A}))$. When applying the homology functor $H : \text{Mod}(\mathcal{A}) \rightarrow \text{Mod}(H(\mathcal{A}))$ to the sequences (35) and (36), we get sequences

$$0 \longrightarrow H(X_{n+1}) \xrightarrow{H(k_{n+1})} H(P_n) \xrightarrow{H(\pi_n)} H(X_n) \longrightarrow 0 \tag{37}$$

and

$$\dots \longrightarrow H(P_2) \xrightarrow{H(\tilde{\pi}_2)} H(P_1) \xrightarrow{H(\tilde{\pi}_1)} H(P_0) \xrightarrow{H(\pi_0)} H(X) \longrightarrow 0. \quad (38)$$

As (35) is also a short exact sequence in $\mathbf{Ch}_+(\mathcal{DM})$, it induces an exact \mathbf{GDM} -triangle in homology. Since $H(\pi_n)$ is surjective, the connecting homomorphism vanishes and the morphism $H(k_{n+1})$ is injective (it is actually the canonical inclusion), so that the short $\mathbf{Mod}(H(\mathcal{A}))$ -sequence (37) is exact. It follows that the long $\mathbf{Mod}(H(\mathcal{A}))$ -sequence (38) is exact. Indeed, since

$$H(\tilde{\pi}_{n+1}) = H(k_{n+1}) \circ H(\pi_{n+1}),$$

the preceding properties imply that

$$\ker H(\tilde{\pi}_{n+1}) = \ker H(\pi_{n+1}) = H(X_{n+2}) = \text{im } H(\pi_{n+2}) = \text{im } H(\tilde{\pi}_{n+2}).$$

Hence $H(P_\bullet)$ is a free (hence projective) resolution of $H(X)$ in $\mathbf{Mod}(H(\mathcal{A}))$.

We continue to follow the procedure described above and consider the first quadrant double complex $(P_p \otimes_{\mathcal{A}} \mathfrak{Q})_q \in \mathcal{DM}$ in the abelian category of \mathcal{D} -modules (index q refers to the grading of the \mathcal{A} -module $P_p \otimes_{\mathcal{A}} \mathfrak{Q} \in \mathbf{DGDM}$). The horizontal differential

$$d_{pq}^h : (P_p \otimes_{\mathcal{A}} \mathfrak{Q})_q \rightarrow (P_{p-1} \otimes_{\mathcal{A}} \mathfrak{Q})_q$$

is the \mathcal{D} -linear map $(\tilde{\pi}_p)_q$ given by the chain map

$$\bar{\pi}_p := \tilde{\pi}_p \otimes \text{id}_{\mathfrak{Q}}.$$

The map d^h actually squares to zero due to the exactness of (36). The vertical differential

$$d_{pq}^v : (P_p \otimes_{\mathcal{A}} \mathfrak{Q})_q \rightarrow (P_p \otimes_{\mathcal{A}} \mathfrak{Q})_{q-1}$$

is the \mathcal{D} -linear map $(d_p^\otimes)_q$ given by the differential

$$d_p^\otimes := d_{P_p} \otimes \text{id} + \text{id} \otimes d_{\mathfrak{Q}}$$

of the \mathcal{A} -module $P_p \otimes_{\mathcal{A}} \mathfrak{Q} \in \mathbf{DGDM}$. This way we actually get a double complex, i.e.,

$$d_{p-1,q}^v d_{pq}^h = d_{p,q-1}^h d_{pq}^v,$$

as $\tilde{\pi}_p$ is a chain map.

The second page of the horizontal spectral sequence is

$${}^h E_{pq}^2 = H_p^h(H_q^v((P_\bullet \otimes_{\mathcal{A}} \mathfrak{Q})_\bullet)) = H_q^v((P_p \otimes_{\mathcal{A}} \mathfrak{Q})_\bullet) \cap \ker d_{\sharp}^h / d_{\sharp}^h H_q^v((P_{p+1} \otimes_{\mathcal{A}} \mathfrak{Q})_\bullet),$$

where d_{\sharp}^h is the differential induced in homology by d^h . Since d^v is d^{\otimes} and P_i is a cofibrant \mathcal{A} -module whose homology is a free $H(\mathcal{A})$ -module, there is a \mathcal{DM} -isomorphism

$$H_q^v((P_i \otimes_{\mathcal{A}} \mathfrak{Q})_{\bullet}) \cong (H(P_i) \otimes_{H(\mathcal{A})} H(\mathfrak{Q}))_q ,$$

in view of Proposition 6. We thus get

$${}^h E_{pq}^2 = H_p^h((H(P_{\bullet}) \otimes_{H(\mathcal{A})} H(\mathfrak{Q}))_q) = \text{Tor}_p^{H(\mathcal{A})}(H(M), H(N))_q ,$$

as $d_{\sharp}^h = \tilde{\pi}_{\sharp} \otimes \text{id}$ when read through the previous isomorphism and as $H(X) \cong H(M)$ and $H(\mathfrak{Q}) \cong H(N)$.

The second sheet of the vertical spectral sequence is

$${}^v E_{pq}^2 = H_p^v(H_q^h((P_{\bullet} \otimes_{\mathcal{A}} \mathfrak{Q})_{\bullet})) = H_q^h((P_{\bullet} \otimes_{\mathcal{A}} \mathfrak{Q})_p) \cap \ker d_{\sharp}^v / d_{\sharp}^v H_q^h((P_{\bullet} \otimes_{\mathcal{A}} \mathfrak{Q})_{p+1}) .$$

The horizontal homology $H_q^h((P_{\bullet} \otimes_{\mathcal{A}} \mathfrak{Q})_{\ell})$ is the homology of the differential $d^h = \tilde{\pi} \otimes \text{id}$, so that in its computation the differential $d^v = d_P \otimes \text{id} + \text{id} \otimes d_{\mathfrak{Q}}$ is irrelevant, only the graded \mathcal{D} -module structure of

$$P_k \otimes_{\mathcal{A}} \mathfrak{Q} = P_k \otimes_{\mathcal{A}} (\mathcal{A} \otimes_{\mathcal{O}} V) \in \mathbf{Mod}_{\mathbf{DGDM}}(\mathcal{A}) \subset \mathbf{GDM}$$

matters. We can therefore ignore the lowering differential of $\mathfrak{Q} = \mathcal{A} \otimes_{\mathcal{O}} V$ and even replace it by the differential $d_{\mathcal{A}} \otimes \text{id}$, thus viewing V as the corresponding direct sum S of spheres (this direct sum is a differential graded \mathcal{D} -module with vanishing differential). By Proposition 5 we now get

$$P_k \otimes_{\mathcal{A}} \mathfrak{Q} \cong P_k \otimes_{\mathcal{O}} S \in \mathbf{GDM} .$$

Since $\tilde{\pi}_k$ is \mathcal{A} -linear, we get an isomorphism

$$((P_{\bullet} \otimes_{\mathcal{A}} \mathfrak{Q})_{\ell}, (\tilde{\pi}_{\bullet} \otimes \text{id}_{\mathfrak{Q}})_{\ell}) \cong ((P_{\bullet} \otimes_{\mathcal{O}} S)_{\ell}, (\tilde{\pi}_{\bullet} \otimes \text{id}_S)_{\ell})$$

of chain complexes of \mathcal{D} -modules, so that

$$H_q^h((P_{\bullet} \otimes_{\mathcal{A}} \mathfrak{Q})_{\ell}) \cong H_q((P_{\bullet} \otimes_{\mathcal{O}} S)_{\ell}) = \bigoplus_{r+s=\ell} H_q(P_{\bullet,r} \otimes_{\mathcal{O}} S(s))$$

in \mathcal{DM} . The complex

$$P_{\bullet,r} \otimes_{\mathcal{O}} S(s) = \bigoplus_k P_{k,r} \otimes_{\mathcal{O}} S(s)$$

with differential $(\tilde{\pi}_{\bullet})_r \otimes \text{id}_{S(s)}$ is the tensor product of the complex $P_{\bullet,r}$ with differential $(\tilde{\pi}_{\bullet})_r$ and the complex $S(s)$ concentrated in degree zero with zero differential. It suffices to compute

its homology as homology of a tensor product of chain complexes of \mathcal{O} -modules. We already mentioned earlier that $S(s)$, $d(S(s)) = 0$, $H_0(S(s)) = S(s)$ and $H_{\sigma>0}(S(s)) = 0$ are all \mathcal{O} -flat. From Künneth's theorem it therefore follows that

$$\begin{aligned} H_q^h((P_\bullet \otimes_{\mathcal{A}} \mathfrak{Q})_\bullet) &\cong \bigoplus_{\ell} \bigoplus_{r+s=\ell} \bigoplus_{\rho+\sigma=q} H_\rho(P_{\bullet,r}) \otimes_{\mathcal{O}} H_\sigma(S(s)) = H_q(P_\bullet) \otimes_{\mathcal{O}} S \cong \\ H_q(P_\bullet) \otimes_{\mathcal{A}} (\mathcal{A} \otimes_{\mathcal{O}} S) &\cong H_q(P_\bullet) \otimes_{\mathcal{A}} (\mathcal{A} \otimes_{\mathcal{O}} V) = H_q(P_\bullet) \otimes_{\mathcal{A}} \mathfrak{Q}, \end{aligned}$$

where the isomorphisms are isomorphisms of graded \mathcal{D} -modules. However, the differential on the LHS is that induced by $d^v = d_P \otimes \text{id} + \text{id} \otimes d_{\mathfrak{Q}}$ and the same is true for the differential on the RHS. In other words, the third isomorphism resets the correct vertical differential, so that the LHS and the RHS are isomorphic differential graded \mathcal{D} -modules and have therefore isomorphic vertical homologies. Since P_\bullet is a resolution of X in the abelian category $\text{Mod}(\mathcal{A})$, we have $H_{q>0}(P_\bullet) = 0$ and $H_0(P_\bullet) \cong X = QM$ in $\text{Mod}(\mathcal{A})$. Hence

$${}^v E_{pq}^2 = \{0\} \quad (q > 0)$$

and

$${}^v E_{p0}^2 \cong H_p(QM \otimes_{\mathcal{A}} \mathfrak{Q}) \cong H_p(M \otimes_{\mathcal{A}}^{\mathbb{L}} N),$$

due to Equation (15) of Theorem 2. The vertical spectral sequence thus collapses at its second page and

$$H_n(\text{Tot}(C)) \cong {}^v E_{n0}^2 \cong H_n(M \otimes_{\mathcal{A}}^{\mathbb{L}} N)$$

(see text below Equation (34)), so that

$${}^h E_{pq}^2 = \text{Tor}_p^{H(\mathcal{A})}(H(M), H(N))_q \Rightarrow H_{p+q}(M \otimes_{\mathcal{A}}^{\mathbb{L}} N),$$

as announced.

Remark 7. Edge homomorphisms of the first quadrant double complexes' spectral sequences

$$G_p^v H_p(\text{Tot } C_{\bullet\bullet}) \rightarrow H_p^v H_0^h(C_{\bullet\bullet}) \quad \text{and} \quad H_0^h H_p^v(C_{\bullet\bullet}) \rightarrow G_0^h H_p \text{Tot}(C_{\bullet\bullet})$$

in any Abelian category \mathbf{A} are natural. More precisely, they are the $C_{\bullet\bullet}$ -components of natural transformations

$$G_p^v H_p \circ \text{Tot} \Rightarrow H_p^v H_0^h \quad \text{and} \quad H_0^h H_p^v \Rightarrow G_0^h H_p \circ \text{Tot}$$

between functors from the category of first quadrant double complexes in \mathbf{A} to the category \mathbf{A} . From here, it can be shown that the edge homomorphisms of the Tor spectral sequence

$$(HM \otimes_{H\mathcal{A}} HN)_q = {}^h E_{0q}^2 \rightarrow H_q(M \otimes_{\mathcal{A}}^{\mathbb{L}} N)$$

are the N -components of a natural transformation

$$(HM \otimes_{H\mathcal{A}} H-)_q \Rightarrow H_q(M \otimes_{\mathcal{A}}^{\mathbb{L}} -).$$

6 Exactness of the derived tensor product

Given $M \in \text{Mod}(\mathcal{A})$, limit-preserving property of the tensor product $-\otimes_A M$ is not invariant under weak equivalences, and should be replaced by a homotopy invariant notion. In [33], the property that the derived tensor product $-\otimes_A^{\mathbb{L}} M$ preserves homotopy pullbacks is recognized as the correct notion of flatness. The rest of the paper is dedicated to the implementation of this property in our homotopical \mathcal{D} -algebraic framework. It is worth noting that if M is flat in the later derived sense, Tor spectral sequence collapses on the second page, yielding a natural isomorphism

$$(HM \otimes_{H\mathcal{A}} HN)_q \cong H_q(M \otimes_{\mathcal{A}}^{\mathbb{L}} N).$$

This is the content of the Corollary 2.

6.1 Homotopy pullbacks and flat modules

As detailed in the introduction, the requirement for the derived tensor product to preserve homotopy pullbacks is ambiguous to a certain extent. The present subsection gives it a precise meaning, applicable to any monoidal model category in which the tensoring with cofibrant objects preserves weak equivalences.

Denote by $S = \{\bullet \rightarrow \bullet \leftarrow \bullet\}$ the small category with three objects $\{c, d, b\}$, and two non-identity morphisms $b \rightarrow d$ and $c \rightarrow d$. Given any model category \mathbb{M} , the functor category $\text{Fun}(S, \mathbb{M})$ – the category of cospan diagrams in \mathbb{M} – has three model structures with objectwise weak equivalences, in which fibrant objects are objectwise fibrant diagrams for which respectively $b \rightarrow d$, $c \rightarrow d$, and both $b \rightarrow d$, $c \rightarrow d$ are fibrations [16]. The limit (pullback) functor is right Quillen for all three model structures, and the corresponding right derived functors, i.e. homotopy pullbacks, agree. Moreover, given a commutative square

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & & \downarrow \\ C & \longrightarrow & D \end{array}$$

in \mathbb{M} , whether the universal map $A \rightarrow \text{Lim}(R(B \rightarrow D \leftarrow C))$ from A to the homotopy pullback of the cospan $B \rightarrow D \leftarrow C$ is a weak equivalence is independent of the chosen model structure or the chosen fibrant replacement R in $\text{Fun}(S, \mathbb{M})$. Commutative squares for which those maps are indeed weak equivalences are referred to as model squares, and denoted by $ABCD$. The vertex A is called a generalized representative of the homotopy pullback $B \times_D^h C$. In the

literature, model squares are also called homotopy fiber squares [17], and homotopy pullback squares [22].

Generally, we say that a functor $F : \mathbb{M} \rightarrow \mathbb{N}$ preserves model squares if the F -image of any model square in \mathbb{M} is a model square in \mathbb{N} . If there is a natural weak equivalence $\eta : F \Rightarrow F'$ between two functors from \mathbb{M} to \mathbb{N} , then by [14, Proposition 2] F preserves model squares if and only if F' does.

We now turn our attention from general functors to the derived tensor product. To begin with, there is a certain ambiguity in its definition. Denoting by Q the cofibrant F -replacement functor, and by C a local cofibrant replacement, the derived tensor product

$$- \otimes_{\mathcal{A}}^{\mathbb{L}} M : \text{Ho}(\text{Mod}(\mathcal{A})) \rightarrow \text{Ho}(\text{Mod}(\mathcal{A}))$$

can be equivalently defined as the derived functor of any of the three functors

$$Q - \otimes_{\mathcal{A}} CM : N \mapsto QN \otimes_{\mathcal{A}} CM, \quad Q - \otimes_{\mathcal{A}} M : N \mapsto QN \otimes_{\mathcal{A}} M, \quad - \otimes_{\mathcal{A}} CM : N \mapsto N \otimes_{\mathcal{A}} CM, \quad (39)$$

which all respect weak equivalences 2.

Any one of them preserves model squares if any other does, due to the natural weak equivalences

$$\begin{aligned} q \otimes_{\mathcal{A}} CM : Q - \otimes_{\mathcal{A}} CM &\Rightarrow - \otimes_{\mathcal{A}} CM, \\ Q - \otimes_{\mathcal{A}} CM : Q - \otimes_{\mathcal{A}} CM &\Rightarrow Q - \otimes_{\mathcal{A}} M. \end{aligned}$$

Similarly, whether the functors in question preserve model squares is independent of the chosen cofibrant replacements. Indeed, if Q_1, Q_2 are two different cofibrant F -replacement functors, and C, C' two different local cofibrant replacements, one gets following chains of natural weak equivalences:

$$\begin{aligned} - \otimes_{\mathcal{A}} CM &\Leftarrow Q_1 - \otimes_{\mathcal{A}} CM \Rightarrow Q_1 - \otimes_{\mathcal{A}} M \Leftarrow Q_1 - \otimes_{\mathcal{A}} C' M \Rightarrow - \otimes_{\mathcal{A}} C' M, \\ - \otimes_{\mathcal{A}} CM &\Leftarrow Q_2 - \otimes_{\mathcal{A}} CM \Rightarrow Q_2 - \otimes_{\mathcal{A}} M \Leftarrow Q_2 - \otimes_{\mathcal{A}} C' M. \end{aligned}$$

Finally, we are prepared to define flat modules.

Definition 3. *The derived tensor product $- \otimes_{\mathcal{A}}^{\mathbb{L}} M$ is said to preserve model squares if any, and hence all the functors in 39 preserve model squares. In this situation, we say that M is a flat \mathcal{A} -module.*

To verify that tensoring with an \mathcal{A} -module preserves all model squares is not a simple task. Luckily, for the derived tensor product to preserve model squares it suffices that it

preserves homotopy fiber sequences. For this, we first recall the later concept, as defined in [14], together with the related notions which are relevant for this paper: mapping fiber, loop object (shift), and the connecting homomorphism. Definitions slightly deviate from the standard references [30], [18].

6.2 Homotopy fiber sequences and flat modules

The purpose of this subsection is to prove that an \mathcal{A} -module M is flat if and only if the derived tensor product with M preserves homotopy fiber sequences. The argumentation used in the proof of the general result (Theorem 7) is applicable to any pointed model category in which suspension is the right inverse to the looping. We begin by recalling the notion of a homotopy fiber sequence in $\mathbf{Mod}(\mathcal{A})$.

For us, a homotopy fiber sequence in a pointed model category \mathbb{M} , denoted by $A \rightarrow B \rightarrow D$, is a model square

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & & \downarrow \\ C & \longrightarrow & D, \end{array} \tag{40}$$

in which the unique map from C to the terminal/initial object is a weak equivalence. Canonical examples are fibrations together with their kernels, and the pullbacks of arbitrary maps $f : B \rightarrow D$ along the based path object (dual cone) of D ([14, Sections 4, 6]).

To give an elegant description of the based path object it is useful to consider unbounded chain complexes as well. A chain complex X of \mathcal{D} -modules which is not necessarily concentrated in non-negative degrees will be called an unbounded \mathcal{A} -module if it is equipped with a morphism $\mathcal{A} \otimes X \rightarrow X$ in the category of unbounded dg \mathcal{D} -modules, such that the usual associativity and unitality diagrams commute. The category of unbounded \mathcal{A} -modules will be denoted by $\mathbf{Mod}(\mathcal{A})^{\text{ub}}$. $\mathbf{Mod}(\mathcal{A})$ is its full subcategory. Given $X \in \mathbf{Mod}(\mathcal{A})^{\text{ub}}$, its good truncation, denoted by $X^{\geq 0}$, is its subcomplex

$$\dots \xrightarrow{d_3} D_2 \xrightarrow{d_2} D_1 \xrightarrow{d_1} \text{Ker}(d_0).$$

Being closed under the \mathcal{A} -action, $X^{\geq 0}$ is an object in $\mathbf{Mod}(\mathcal{A})$. It is straightforwardly verified that the obtained functor $(-)^{\geq 0} : \mathbf{Mod}(\mathcal{A})^{\text{ub}} \rightarrow \mathbf{Mod}(\mathcal{A})$ is right adjoint to the inclusion $\iota : \mathbf{Mod}(\mathcal{A}) \rightarrow \mathbf{Mod}(\mathcal{A})^{\text{ub}}$.

Notice that the truncation is a right Quillen functor as it respects quasi-isomorphisms, and sends degreewise surjective maps to fibrations in $\mathbf{Mod}(\mathcal{A})$ – the maps surjective in strictly positive degrees.

Given $D \in \text{Mod}(\mathcal{A})^{\text{ub}}$, its unbounded based path object $\text{Path}_0^{\text{ub}} D$ is its (-1) – shifted cone, explicitly,

$$(\text{Path}_0^{\text{ub}} D)_n = D_n \oplus D_{n+1},$$

$$d = \begin{pmatrix} d_D & 0 \\ -\text{id}_D & -d_D \end{pmatrix}.$$

For the explicit description of the \mathcal{A} -action see the Remark 8.

Denote by π_D^{ub} the degreewise projection

$$\text{Path}_0^{\text{ub}} D \rightarrow D, \quad D_n \oplus D_{n+1} \rightarrow D_n$$

Given a morphism $f : B \rightarrow D$ in $\text{Mod}(\mathcal{A})^{\text{ub}}$, the pullback

$$\begin{array}{ccc} K_f^{\text{ub}} & \xrightarrow{\pi_f^{\text{ub}}} & B \\ \downarrow p_f^{\text{ub}} & & \downarrow f \\ \text{Path}_0^{\text{ub}} D & \xrightarrow{\pi_D^{\text{ub}}} & D, \end{array} \quad (41)$$

will be called the unbounded fiber of f . It equals the (-1) shifted mapping cone. The (-1) -shift of D ,

$$D[-1]_n = D_{n+1}, \quad d_n^{D[-1]} = -d_{n+1}^D$$

can be seen as the kernel of $\pi_D^{\text{ub}} : \text{Path}_0^{\text{ub}} D \rightarrow D$, and the connecting homomorphism $\delta_f^{\text{ub}} : D[-1] \rightarrow K_f^{\text{ub}}$ is the pullback's universal map

$$\begin{array}{ccccc} D[-1] & \xrightarrow{\quad 0 \quad} & & & \\ \delta_f^{\text{ub}} \dashv & \nwarrow & K_f^{\text{ub}} & \xrightarrow{\pi_f^{\text{ub}}} & B \\ \text{Ker} \searrow & & \downarrow p_f^{\text{ub}} & & \downarrow f \\ & & \text{Path}_0^{\text{ub}} D & \xrightarrow{\pi_D^{\text{ub}}} & D. \end{array} \quad (42)$$

Assume now $f : B \rightarrow D$ is a morphism in $\text{Mod}(\mathcal{A})$. By a slight abuse of notation, we denote the map $\iota(f)$ in $\text{Mod}(\mathcal{A})^{\text{ub}}$ also by f . Let

$$\begin{array}{ccccc}
& & 0 & & \\
& \swarrow \delta_f & \curvearrowright & \searrow \pi_f & \\
\Omega D & & K_f & \xrightarrow{\pi_f} & B \\
\downarrow \text{Ker} & & \downarrow p_f & & \downarrow f \\
\text{Path}_0 D & \xrightarrow{\pi_D} & D. & &
\end{array} \tag{43}$$

be the image of Diagram (42) under truncation.

As truncation is right adjoint, the commutative square $K_f, B, \text{Path}_0 D, D$ is a pullback. As $\text{Path}_0 D$ is acyclic, and π_D is a fibration, it is even a homotopy fiber sequence. Finally, homotopy fiber sequences induce long exact sequences in homology, in this example

$$\dots \xrightarrow{H_1 f} H_1(D) \xrightarrow{H_0 \delta_f} H_0(K_f) \xrightarrow{H_0 p_f} H_0(B) \xrightarrow{H_0 f} H_0(D).$$

Looping functor Ω , defined as the composition of (-1) -shift and truncation has the left adjoint suspension functor Σ , which equals the 1 -shift. Explicitly, it is defined on objects as

$$\Sigma D_n = D_{n-1}, \quad d_n^{\Sigma D} = -d_{n-1}^D,$$

and on morphisms as

$$\Sigma f_n = f_{n-1}.$$

Since Ω preserves fibrations and weak equivalences, $\Sigma \dashv \Omega$ is even a Quillen adjunction. Unlike with unbounded chain complexes, where 1 -shift is the two sided inverse of (-1) -shift, Σ is only a right inverse to Ω .

Remark 8. There is a subtlety involved when determining the correct \mathcal{A} -action on $\text{Path}_0^{ub} D$ (which induces the \mathcal{A} -action on both $\text{Path}_0 D$ and ΩD). \mathcal{A} being graded commutative, the notions of left and right module structures on D are equivalent, and related by $m \triangleright a = (-1)^{am} a \triangleleft m$ (Remark 6). A reasonable way to define the \mathcal{A} -action on $\text{Path}_0^{ub} D$ is from the \mathcal{A} -action on D , in the decomposition $\text{Path}_0^{ub} D = D \oplus D[-1]$. A priori, there is a choice: either one takes the left induced action (coming from the term-wise left \mathcal{A} -action), or the right induced action. The two will agree on the first summand, but differ by a sign on the second. However, only the right induced action is compatible with the differential. Concretely,

for $m \in (\text{Path}_0 D)_n$, and $a \in \mathcal{A}$, denoting by \cdot the \mathcal{A} -action on D , \mathcal{A} -action on $(\text{Path}_0 D)_n$, denoted by \triangleleft , is given by

$$a \triangleleft m = \begin{cases} a \cdot m, & m \in D_n \\ (-1)^a a \cdot m, & m \in D_{n+1}. \end{cases}$$

Consequently, \mathcal{A} -action on the (-1) -shift $D[-1]$, and its truncation ΩD is given by $a \triangleleft m = (-1)^a a \cdot m$. To assure that unit and counit of $\Sigma \dashv \Omega$ adjunction respect the \mathcal{A} -action, the same formula defines the \mathcal{A} -action on the (1) -shift ΣD .

With all the relevant notions illuminated, we move on to showing that $-\otimes_A^L M$ preserves all model squares if it preserves all homotopy fiber sequences. Again, there is a certain ambiguity in what it means for the derived tensor product to preserve homotopy fiber sequences.

In general, we say that a functor $F : \mathbf{M} \rightarrow \mathbf{N}$ preserves homotopy fiber sequences if the F -image of any homotopy fiber sequence in \mathbf{M} (diagram 40) is a homotopy fiber sequence in \mathbf{N} . If there is a natural weak equivalence $\eta : F \Rightarrow F'$ between two functors from \mathbf{M} to \mathbf{N} , then by [14, Proposition 2] F preserves homotopy fiber sequences if and only if F' does. Thus, the following definition makes sense:

Definition 4. *The derived tensor product $-\otimes_A^L M$ is said to preserve homotopy fiber sequences if any, and hence all the functors in 39 preserve homotopy fiber sequences.*

Similarly, as detailed in the previous section, any of the functors 39 representing $-\otimes_A^L M$ preserves model squares whenever any other does. Consequently, to show that $-\otimes_A^L M$ preserves all model squares if it preserves all homotopy fiber sequence, it suffices to prove the statement for the functor $-\otimes_A CM$, or equivalently, that for any cofibrant object $M \in \text{Mod}(\mathcal{A})$, the functor $-\otimes_A M$ preserves all model squares if it preserves all homotopy fiber sequences.

Proposition 7. *A functor $F : \text{Mod}(\mathcal{A}) \rightarrow \text{Mod}(\mathcal{A})$ which preserves weak equivalences preserves all model squares if it preserves all homotopy fiber sequences.*

Proof. Let

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & & \downarrow f \\ C & \xrightarrow{g} & D \end{array} \tag{44}$$

be a model square in $\text{Mod}(\mathcal{A})$. Given any DGDM -morphism $\phi : X \rightarrow Y$, the map $\Omega\phi$ fits in a model square

$$\begin{array}{ccc} \Omega X & \longrightarrow & \text{Path}_0 X \\ \downarrow \Omega\phi & & \downarrow \\ \Omega Y & \longrightarrow & K_\phi \end{array} \quad (45)$$

(see the diagram 49 in [14]). Σ being the right inverse of Ω , the model squares 44 and 45 (for $\phi = \Sigma f$) can be pasted into

$$\begin{array}{ccccc} A & \longrightarrow & B & \longrightarrow & \text{Path}_0 \Sigma B \\ \downarrow & & \downarrow f & & \downarrow \\ C & \xrightarrow{g} & D & \longrightarrow & K_{\Sigma f}. \end{array}$$

By the pasting law [16, Proposition 8], the total square is a model square as well. It is in fact a homotopy fiber sequence, as $\text{Path}_0 \Sigma B$ is acyclic. Applying the functor F which by the assumption preserves homotopy fiber sequences, we get a diagram

$$\begin{array}{ccccc} FA & \longrightarrow & FB & \longrightarrow & F \text{Path}_0 \Sigma B \\ \downarrow & & \downarrow Ff & & \downarrow \\ FC & \xrightarrow{Fg} & FD & \longrightarrow & FK_{\Sigma f} \end{array}$$

whose right-hand-side square and total square are homotopy fiber sequences, especially model squares. By the pasting law we conclude that the left-hand-side square is a model square as well. \square

Corollary 1. $M \in \text{Mod}(\mathcal{A})$ is flat if and only if $- \otimes_{\mathcal{A}}^{\mathbb{L}} M$ preserves homotopy fiber sequences.

7 A characterization of flat modules

As an application of the developed Tor spectral sequence, we characterize flatness as strong flatness. The characterization is specific to the contexts such as differential non-negatively graded or simplicial modules. For unbounded chains, strong flatness is a strictly stronger property (see [33, Chapter 2.3]). The characterization relies on the explicit constructions in the section 6.2. We begin with the definition of strongly flat modules.

Definition 5. 1. Let $\mathcal{A} \in \text{DGDA}$. A module $M \in \text{Mod}(\mathcal{A})$ is **strong**, if the natural $\text{Mod}_{\mathcal{D}\mathcal{M}}(H_0(\mathcal{A}))$ -morphism

$$\phi_{\bullet, \mathcal{A}, M} : H_{\bullet}(\mathcal{A}) \otimes_{H_0(\mathcal{A})} H_0(M) \rightarrow H_{\bullet}(M), \quad [a] \otimes [m_0] \mapsto [a \cdot m_0] \quad (46)$$

is an isomorphism, i.e., if all $\text{Mod}_{\mathcal{D}\mathcal{M}}(H_0(\mathcal{A}))$ -morphisms $\phi_{k, \mathcal{A}, M}$ are isomorphisms.

2. $M \in \text{Mod}(\mathcal{A})$ is **strongly flat** if it is strong, and if $H_0(M)$ is a flat $H_0(\mathcal{A})$ -module in the classical sense.

Before moving forward with the proof that strongly flat modules are equivalently flat, we prove that the isomorphism (46) is well defined.

Let $\mathcal{A} \in \mathcal{D}\mathcal{A}$. In order to define a $\text{Mod}_{\mathcal{D}\mathcal{M}}(\mathcal{A})$ -morphism (an \mathcal{A} - and \mathcal{D} -linear map) $\phi : M' \otimes_{\mathcal{A}} M'' \rightarrow M$, one starts defining an \mathcal{O} -bilinear map $\varphi : M' \times M'' \rightarrow M$, hence, an \mathcal{O} -linear map $\varphi : M' \otimes_{\mathcal{O}} M'' \rightarrow M$. One then checks that φ is a $\mathcal{D}\mathcal{M}$ -morphism, i.e., is \mathcal{D} -linear, or, equivalently, is linear for the action ∇_{θ} of vector fields θ . Recall that, by definition,

$$\nabla_{\theta}(m' \otimes m'') = (\nabla_{\theta}m') \otimes m'' + m' \otimes (\nabla_{\theta}m'').$$

Denote by $\kappa : M' \otimes_{\mathcal{O}} M'' \rightarrow M' \otimes_{\mathcal{A}} M''$ coequalizer's universal map. If, for $a \in \mathcal{A}$,

$$\varphi((a \cdot m') \otimes m'') = \varphi(m' \otimes (a \cdot m'')) ,$$

it follows from the universal property of a coequalizer, that there is a unique \mathcal{D} -linear map $\phi : M' \otimes_{\mathcal{A}} M'' \rightarrow M$, such that $\phi \circ \kappa = \varphi$. Finally, if φ is \mathcal{A} -bilinear on $M' \times M''$, then ϕ is \mathcal{A} -linear on $M' \otimes_{\mathcal{A}} M''$. Indeed,

$$\varphi(m', m'') = \varphi(m' \otimes m'') = \phi(\kappa(m' \otimes m'')) = \phi(m' \otimes_{\mathcal{A}} m'').$$

We now come back to Definition 5. The tensor product in (46) makes sense since $H_0(\mathcal{A}) \in \mathcal{D}\mathcal{A}$ and $H_k(\mathcal{A}), H_0(M) \in \text{Mod}_{\mathcal{D}\mathcal{M}}(H_0(\mathcal{A}))$. In order to define the morphism $\phi_{k, \mathcal{A}, M}$ (we will write ϕ), we apply the just detailed method to the preceding \mathcal{D} -algebra and modules in $\mathcal{D}\mathcal{M}$ over that algebra. Let φ be the map

$$\varphi : H_k(\mathcal{A}) \times H_0(M) \ni ([a_k], [m_0]) \mapsto [\nu(a_k \otimes m_0)] \in H_k(M) ,$$

where the DGDM -morphism $\nu : \mathcal{A} \otimes_{\mathcal{O}} M \rightarrow M$ is the action of \mathcal{A} on M . It is easy to check that φ is well-defined on homology classes. In view of the $\mathcal{D}\mathcal{A}$ -morphism

$$\mathcal{O} \ni f \mapsto f \cdot [1_{\mathcal{A}}] = [f \cdot 1_{\mathcal{A}}] \in H_0(\mathcal{A}) ,$$

if φ is $H_0(\mathcal{A})$ -bilinear, it is in particular \mathcal{O} -bilinear. As for $H_0(\mathcal{A})$ -bilinearity, taking into account that the action $*$ of $H_0(\mathcal{A})$ on $H_k(\mathcal{A})$ is induced by the multiplication $*$ in \mathcal{A} , and that the action ν of $H_0(\mathcal{A})$ on $H_k(M)$ is induced by the action ν of \mathcal{A} on M , we get

$$\varphi([a_0] * [a_k], [m_0]) = \varphi([a_0 * a_k], [m_0]) = [\nu((a_0 * a_k) \otimes m_0)],$$

$$\nu([a_0] \otimes \varphi([a_k], [m_0])) = \nu([a_0] \otimes [\nu(a_k \otimes m_0)]) = [\nu(a_0 \otimes \nu(a_k \otimes m_0))] = [\nu((a_0 * a_k) \otimes m_0)],$$

and

$$\varphi([a_k], \nu([a_0] \otimes [m_0])) = \varphi([a_k], [\nu(a_0 \otimes m_0)]) = [\nu(a_k \otimes \nu(a_0 \otimes m_0))] = [\nu((a_k * a_0) \otimes m_0)],$$

so that φ is actually $H_0(\mathcal{A})$ -bilinear and thus \mathcal{O} -bilinear. We now check linearity of

$$\varphi : H_k(\mathcal{A}) \otimes_{\mathcal{O}} H_0(M) \rightarrow H_k(M)$$

with respect to the action ∇_θ by vector fields θ . As the \mathcal{D} -action on homology is induced by the \mathcal{D} -action on the underlying complex, we have

$$\begin{aligned} \varphi(\nabla_\theta([a_k] \otimes [m_0])) &= \varphi((\nabla_\theta[a_k]) \otimes [m_0]) + [a_k] \otimes (\nabla_\theta[m_0]) = \\ \varphi([\nabla_\theta a_k] \otimes [m_0]) + \varphi([a_k] \otimes [\nabla_\theta m_0]) &= [\nu((\nabla_\theta a_k) \otimes m_0)] + [\nu(a_k \otimes (\nabla_\theta m_0))] = \\ [\nu(\nabla_\theta(a_k \otimes m_0))] &= \nabla_\theta[\nu(a_k \otimes m_0)] = \nabla_\theta \varphi([a_k] \otimes [m_0]). \end{aligned}$$

In view of what has been said above, it follows that φ induces a unique $\mathbf{Mod}_{\mathcal{DM}}(H_0(\mathcal{A}))$ -morphism $\phi : H_k(\mathcal{A}) \otimes_{H_0(\mathcal{A})} H_0(M) \rightarrow H_k(M)$.

7.1 Strongly flat modules are flat

Notice that both flatness and strong flatness are invariant under the isomorphisms in the homotopy category. This allows us to assume without the loss of generality that M is cofibrant, so that $- \otimes_{\mathcal{A}} M$ represents the derived tensor product $- \otimes_{\mathcal{A}}^{\mathbb{L}} M$.

Assume that M is a strongly flat \mathcal{A} -module. To prove that M is flat, it suffices to show that the functor $- \otimes_{\mathcal{A}} M$ preserves homotopy fiber sequences. By [14, Corollary 7], we further reduce to showing that for any map $f : B \rightarrow D$ of \mathcal{A} -modules, the image of the homotopy fiber sequence

$$\begin{array}{ccc} K_f & \xrightarrow{\pi_f} & B \\ \downarrow p_f & & \downarrow f \\ \mathrm{Path}_0 D & \xrightarrow{\pi_D} & D, \end{array} \tag{47}$$

under the functor $- \otimes_{\mathcal{A}} M$ is still a homotopy fiber sequence.

Let $f : B \rightarrow D$ be a morphism in $\text{Mod}(\mathcal{A})$. Unbounded \mathcal{A} -modules $D[\pm 1] \otimes M$ and $(D \otimes M)[\pm 1]$ are equal, since the equality of graded modules

$$(D[\pm 1] \otimes M)_n = \bigoplus_{k+l=n} (D_{k\pm 1} \otimes M_l) = \bigoplus_{k+l=n\pm 1} (D_k \otimes M_l) = (D \otimes M)[\pm 1]_n$$

respects both the differential and the \mathcal{A} -action. As right \mathcal{A} -actions on $D[\pm 1]$ and D coincide (Remark 8),

$$D[\pm 1] \otimes_{\mathcal{A}} M = (D \otimes_{\mathcal{A}} M)[\pm 1]. \quad (48)$$

On the other side, the tensor products $M \otimes_{\mathcal{A}} D[\pm 1]$ and $(M \otimes_{\mathcal{A}} D)[\pm 1]$ not equal, but can still be identified via the isomorphism $m \otimes d \mapsto (-1)^m m \otimes d$.

The isomorphism of graded \mathcal{A} -modules

$$(\text{Path}_0^{ub} D) \otimes_{\mathcal{A}} M = (D \oplus D[-1]) \otimes_{\mathcal{A}} M \rightarrow D \otimes_{\mathcal{A}} M \oplus D[-1] \otimes_{\mathcal{A}} M = \text{Path}_0^{ub}(D \otimes_{\mathcal{A}} M)$$

also respects the differential. The obtained isomorphism in $\text{Mod}^{ub}(\mathcal{A})$ is

$$\phi^{ub} : (\text{Path}_0^{ub} D) \otimes_{\mathcal{A}} M \xrightarrow{\cong} \text{Path}_0^{ub}(D \otimes_{\mathcal{A}} M), \quad (d + d^{-1}) \otimes m \mapsto d \otimes m + d^{-1} \otimes m,$$

for $d \in D$, $d^{-1} \in D[-1]$, $m \in M$.

Denote by $K^{\otimes, ub}$ the pullback

$$(B \otimes_{\mathcal{A}} M) \prod_{D \otimes_{\mathcal{A}} M} \text{Path}_0^{ub}(D \otimes_{\mathcal{A}} M),$$

and by K^{\otimes} its good truncation. Denote by ϵ the counit of the $\iota \vdash (-)^{\geq 0}$ adjunction, explicitly the natural map

$$\begin{array}{ccccccccccc} \dots & \xrightarrow{d_3} & D_2 & \xrightarrow{d_2} & D_1 & \xrightarrow{d_1} & \text{Ker}(d_0) & \longrightarrow & 0 & \longrightarrow & \dots \\ & & \parallel & & \parallel & & \downarrow & & \downarrow & & \\ \dots & \xrightarrow{d_3} & D_2 & \xrightarrow{d_2} & D_1 & \xrightarrow{d_1} & D_0 & \xrightarrow{d_0} & D_{-1} & \xrightarrow{d_{-1}} & \dots \end{array} \quad (49)$$

As $(\text{Path}_0 D) \otimes_{\mathcal{A}} M$ is non-negatively graded, the composition

$$(\text{Path}_0 D) \otimes_{\mathcal{A}} M \xrightarrow{\epsilon \otimes M} (\text{Path}_0^{ub} D) \otimes_{\mathcal{A}} M \xrightarrow{\phi^{ub}} \text{Path}_0^{ub}(D \otimes_{\mathcal{A}} M)$$

factors through $\text{Path}_0(D \otimes_{\mathcal{A}} M)$. Let

$$\phi : (\text{Path}_0 D) \otimes_{\mathcal{A}} M \rightarrow \text{Path}_0(D \otimes_{\mathcal{A}} M)$$

be the resulting map.

The following commutative diagram, in which the dashed arrows are universal maps of the pullback, is central. For notation, see Diagrams (42) and (43).

$$\begin{array}{ccccccc}
 \Omega D \otimes_{\mathcal{A}} M & \xrightarrow{\quad} & 0 & & & & \\
 \downarrow \delta_f \otimes M & \dashrightarrow \psi_{\Omega} & \downarrow & \downarrow & \downarrow & & \\
 K_f \otimes_{\mathcal{A}} M & \xrightarrow{\quad} & \Omega(D \otimes_{\mathcal{A}} M) & \xrightarrow{\epsilon} & (D \otimes_{\mathcal{A}} M)[-1] & \xrightarrow{\quad} & 0 \\
 \downarrow p_f \otimes M & \dashrightarrow \psi_K & \downarrow \delta_{f \otimes M} & \downarrow \pi_{f \otimes M} & \downarrow \delta_{f \otimes M}^{ub} & \downarrow & \downarrow f \otimes M \\
 (\text{Path}_0 D) \otimes_{\mathcal{A}} M & \xrightarrow{\quad} & K^{\otimes} & \xrightarrow{\epsilon} & K^{\otimes, ub} & \xrightarrow{\pi_{f \otimes M}^{ub}} & B \otimes_{\mathcal{A}} M \\
 \downarrow \phi & \searrow & \downarrow p_{f \otimes M} & \downarrow \pi_{D \otimes M} & \downarrow p_{f \otimes M}^{ub} & \downarrow \pi_{D \otimes M}^{ub} & \downarrow \\
 \text{Path}_0(D \otimes_{\mathcal{A}} M) & \xrightarrow{\epsilon} & \text{Path}_0^{ub}(D \otimes_{\mathcal{A}} M) & \xrightarrow{\phi^{ub}} & \text{Path}_0^{ub}(D \otimes_{\mathcal{A}} M) & \xrightarrow{\pi_{D \otimes M}^{ub}} & D \otimes_{\mathcal{A}} M
 \end{array} \tag{50}$$

To prove that $- \otimes M$ preserves homotopy fiber sequences, it suffices to show that ψ_K is a quasi-isomorphism. The composition $p_f \circ \delta_f$ is the inclusion of the kernel of π_D (Diagram 43). Likewise, $p_{f \otimes M}^{ub} \circ \delta_{f \otimes M}^{ub}$ is inclusion of the kernel of $\pi_{D \otimes M}^{ub}$. Finally, the composition $\epsilon \circ \psi_{\Omega}$ equals the natural map

$$\epsilon \otimes M : \Omega D \otimes_{\mathcal{A}} M \rightarrow D[-1] \otimes_{\mathcal{A}} M = (D \otimes_{\mathcal{A}} M)[-1],$$

as, in view of kernel's universal property, it is the unique map for which the square

$$\begin{array}{ccccc}
 \Omega D \otimes_{\mathcal{A}} M & \xrightarrow{\epsilon \circ \psi_{\Omega}} & D[-1] \otimes_{\mathcal{A}} M & \xrightarrow{\quad} & (D \otimes_{\mathcal{A}} M)[-1] \\
 \downarrow (p_f \otimes M) \circ (\delta_f \otimes M) & = \epsilon \otimes M & \downarrow p_{f \otimes M}^{ub} \circ \delta_{f \otimes M}^{ub} & = \ker(\pi_D^{ub}) \otimes M & \downarrow \ker(\pi_{D \otimes M}^{ub}) \\
 \text{Path}_0 D \otimes_{\mathcal{A}} M & \xrightarrow{\epsilon \otimes M} & \text{Path}_0^{ub} D \otimes_{\mathcal{A}} M & \xrightarrow{\phi^{ub}} & \text{Path}_0^{ub}(D \otimes_{\mathcal{A}} M)
 \end{array}$$

commutes. The left-hand side commutes due to the naturality of ϵ . The commutativity of the right-hand side follows from the explicit description of involved maps.

Strongness of M , together with the Proposition 5, gives the isomorphisms

$$H_{\bullet} D \otimes_{H_0 \mathcal{A}} H_0 M \cong H_{\bullet} D \otimes_{H_{\bullet} \mathcal{A}} (H_{\bullet} \mathcal{A} \otimes_{H_0 \mathcal{A}} H_0 M) \cong H_{\bullet} D \otimes_{H_{\bullet} \mathcal{A}} H_{\bullet} M, \tag{51}$$

natural in D . As $H_0 M$ is a flat $H_0 \mathcal{A}$ -module, $H_\bullet M$ is a flat graded $H_\bullet \mathcal{A}$ -module, and the Tor spectral sequence

$$\mathrm{Tor}_p^{H_\bullet \mathcal{A}}(H_\bullet D, H_\bullet M)_q \Rightarrow H_{p+q}(D \otimes_{\mathcal{A}} M)$$

collapses on the second page. Consequently, the natural edge homomorphism

$$H_\bullet D \otimes_{H_\bullet \mathcal{A}} H_\bullet M \rightarrow H_\bullet(D \otimes_{\mathcal{A}} M) \quad (52)$$

is an isomorphism. Composing with 51, we get the isomorphism

$$H_\bullet D \otimes_{H_0 \mathcal{A}} H_0 M \rightarrow H_\bullet(D \otimes_{\mathcal{A}} M), \quad (53)$$

natural in D . Applying to ϵ , we conclude that

$$H_k(\epsilon \otimes M) : H_k(\Omega D \otimes_{\mathcal{A}} M) \rightarrow H_k(D[-1] \otimes_{\mathcal{A}} M)$$

is an isomorphism for $k \geq 0$. A word of caution is in order: since $\epsilon : \Omega D \rightarrow D[-1]$ is not a map of complexes concentrated in non-negative degrees, to be completely rigorous, before using the natural isomorphism (53), one should apply the 1-shift.

Being the good truncation of $\epsilon \otimes M$, ψ_Ω is a quasi-isomorphism, natural in D . In the diagram

$$\begin{array}{ccccccccccc} H_{\bullet+1}(B) \otimes_{H_0(A)} H_0(M) & \rightarrow & H_{\bullet+1}(D) \otimes_{H_0(A)} H_0(M) & \rightarrow & H_\bullet(K_f) \otimes_{H_0(A)} H_0(M) & \rightarrow & H_\bullet(B) \otimes_{H_0(A)} H_0(M) & \rightarrow & H_\bullet(D) \otimes_{H_0(A)} H_0(M) & & \\ \downarrow \cong & & \downarrow \cong \\ H_\bullet(\Omega B \otimes_{\mathcal{A}} M) & \longrightarrow & H_\bullet(\Omega D \otimes_{\mathcal{A}} M) & \longrightarrow & H_\bullet(K_f \otimes_{\mathcal{A}} M) & \longrightarrow & H_\bullet(B \otimes_{\mathcal{A}} M) & \longrightarrow & H_\bullet(D \otimes_{\mathcal{A}} M) & & \\ \downarrow H_\bullet(\psi_\Omega) & & \downarrow H_\bullet(\psi_\Omega) & & \downarrow H_\bullet(\psi_K) & & \parallel & & \parallel & & \\ H_\bullet \Omega(B \otimes_{\mathcal{A}} M) & \longrightarrow & H_\bullet \Omega(D \otimes_{\mathcal{A}} M) & \longrightarrow & H_\bullet(K^\otimes) & \longrightarrow & H_\bullet(B \otimes_{\mathcal{A}} M) & \longrightarrow & H_\bullet(D \otimes_{\mathcal{A}} M) & & \end{array}$$

the bottom line is the exact sequence in homology associated to the homotopy fiber sequence

$$K^\otimes \rightarrow B \otimes_{\mathcal{A}} M \rightarrow D \otimes_{\mathcal{A}} M.$$

The top line is the long exact sequence in homology associated to the homotopy fiber sequence

$$K_f \rightarrow B \rightarrow D,$$

tensored with the flat $H_0(A)$ -module $H_0(M)$. Since ψ_Ω is a quasi-isomorphism, 5-lemma implies that ψ_K is a quasi-isomorphism as well.

7.2 Postnikov towers

The proof that all flat modules are strongly flat uses the machinery of Postnikov towers, which is – for the category $\text{Mod}(\mathcal{A})$ – recalled in this section.

For given $M \in \text{Mod}(\mathcal{A})$, set $M_{\leq n}$ to be the complex

$$\dots \rightarrow 0 \rightarrow \text{Im}(d_{n+1}) \hookrightarrow M_n \xrightarrow{d_n} M_{n-1} \xrightarrow{d_{n-1}} \dots \xrightarrow{d_1} M_0. \quad (54)$$

Degreewise surjection $\phi_n : M \rightarrow M_{\leq n}$ is defined in degrees $\leq n$ as identity, and in degree $n+1$ as d_{n+1} . Degreewise surjections $p_n : M_{\leq n} \rightarrow M_{\leq n-1}$ are defined alike. $M_{\leq n}$ is an \mathcal{A} -module, with the \mathcal{A} -action defined as follows:

- For $a \in A$ and $m \in M_k, k \leq n$, we set $a \cdot m$ to be the image of $a \cdot m \in M$ under ϕ_n .
- For $dm \in \text{Im}(d_{n+1})$, $a \cdot dm$ is set to zero unless $a \in A_0$, and for $a \in A_0$ we have $a \cdot dm = d(a \cdot m) \in \text{Im}(d_{n+1})$.

Maps p_n and ϕ_n are compatible in the sense that $p_n \circ \phi_n = \phi_{n-1}$, and are easily checked to be morphisms of \mathcal{A} -modules. In fact, being degreewise surjective, the maps are fibrations.

Definition 6. *Postnikov tower of $M \in \text{Mod}(\mathcal{A})$ is the inverse system*

$$\dots \rightarrow M_{\leq n} \xrightarrow{p_n} M_{\leq n-1} \xrightarrow{p_{n-1}} \dots \xrightarrow{p_2} M_{\leq 1} \xrightarrow{p_1} M_{\leq 0},$$

together with the sequence of compatible maps $\phi_n : M \rightarrow M_{\leq n}$.

Indeed, so-defined Postnikov towers satisfy the defining properties of the more familiar topological concept:

Proposition 8.

1. $H_k(\phi_n) : H_k(M) \rightarrow H_k(M_{\leq n})$ is an isomorphism for $k \leq n$;
2. $H_k(M_{\leq n}) = 0$ for $k > n$;
3. the homotopy fiber of the fibration $M_{\leq n} \rightarrow M_{\leq n-1}$ is $H_n(M)$, viewed as a complex concentrated in degree n .

Proof. The first two statements are obvious. For the third, it suffices to notice that the homotopy fiber in question coincides with the kernel of p_n (on the nose), as the map in question is a fibration, and all objects in $\text{Mod}(\mathcal{A})$ are fibrant. \square

As with CW complexes, M can be recovered as the homotopy limit of the inverse system $(M_{\leq n})_{n \in \mathbb{N}}$. Namely, given any model category \mathbb{M} , the functor category $\text{Fun}(\mathbb{N}^{\text{op}}, \mathbb{M})$ has an injective model structure whose weak equivalences and cofibrations are objectwise [18, Theorem 5.3.1.]. Fibrant objects are sequences of fibrations between fibrant objects in \mathbb{M} . Especially, Postnikov tower $\dots M_{\leq n} \twoheadrightarrow M_{\leq n-1} \twoheadrightarrow \dots \twoheadrightarrow M_{\leq 0}$ is a fibrant object in $\text{Fun}(\mathbb{N}^{\text{op}}, \text{Mod}(\mathcal{A}))$, so that its homotopy limit coincides with its projective limit on the nose, which is clearly M .

7.3 Flat modules are strongly flat

Given a morphism $f : \mathcal{A} \rightarrow \mathcal{B}$ in DGDA , any \mathcal{B} -module has an induced \mathcal{A} -module structure defined as $a \cdot m := f(a) \cdot m$. For $\mathcal{A} \in \text{DGDA}$, both \mathcal{A}_0 , and $H_0\mathcal{A}$ are commutative \mathcal{D} -algebras, and can be viewed as differential graded \mathcal{D} -algebras concentrated in degree zero. Due to DGDA morphisms $\mathcal{A}_0 \hookrightarrow \mathcal{A} \twoheadrightarrow H_0\mathcal{A}$, any $H_0\mathcal{A}$ -module is also an \mathcal{A} -module, and any \mathcal{A} -module is also an \mathcal{A}_0 -module. These facts will be used repeatedly throughout this subsection. For $D \in \text{Mod}(\mathcal{A})$, we will denote by $\pi_0 : D \rightarrow H_0(D)$ the natural map

$$\begin{array}{ccccccc} \dots & \xrightarrow{d_3} & D_2 & \xrightarrow{d_2} & D_1 & \xrightarrow{d_1} & D_0 \\ & & \downarrow & & \downarrow & & \downarrow \\ \dots & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & H_0(D) \end{array} \tag{55}$$

Suppose that M is a flat \mathcal{A} -module. As in the subsection 7.1, we assume M to be cofibrant. Any exact sequence

$$0 \rightarrow N \hookrightarrow P$$

of $H_0\mathcal{A}$ -modules is a homotopy fiber sequence in $\text{Mod}(\mathcal{A})$, since any map between complexes concentrated in degree zero is a fibration. Since M is flat,

$$0 \rightarrow N \otimes_{\mathcal{A}} M \rightarrow P \otimes_{\mathcal{A}} M \tag{56}$$

is a homotopy fiber sequence as well. From the associated long exact sequence in homology, we conclude that

$$0 \rightarrow H_0(N \otimes_{\mathcal{A}} M) \rightarrow H_0(P \otimes_{\mathcal{A}} M) \tag{57}$$

is an exact sequence of $H_0\mathcal{A}$ -modules.

Lemma 1. *For $N \in \text{Mod}(H_0\mathcal{A})$, $M \in \text{Mod}(\mathcal{A})$, there exists a natural isomorphism*

$$\phi : H_0(N \otimes_{\mathcal{A}} M) \rightarrow N \otimes_{H_0\mathcal{A}} H_0 M, \quad [n \otimes m_0] \mapsto n \otimes [m_0]$$

of $H_0\mathcal{A}$ -modules.

From here and (57), it follows that H_0M is a flat $H_0\mathcal{A}$ -module.

Proof. For $\mathcal{A} \in \mathsf{DGDA}$, and $P, Q \in \mathsf{Mod}(\mathcal{A})$, the tensor product $P \otimes_{\mathcal{A}} Q$ is the quotient of $P \otimes Q$ by the graded submodule \mathcal{I}_A generated by the homogeneous elements $p \cdot a \otimes q - p \otimes a \cdot q$, together with the induced differential, and the \mathcal{A} -action induced by the left \mathcal{A} -action on P . For $P, Q \in \mathsf{Mod}(\mathcal{B})$, the inclusion $\mathcal{I}_A \subseteq \mathcal{I}_B$ of submodules in $P \otimes Q$ induces the quotient map $P \otimes_{\mathcal{A}} Q \rightarrow P \otimes_{\mathcal{B}} Q$ in $\mathsf{Mod}(\mathcal{B})$.

For $m \in M$, $a \in \mathcal{A}$, and $n \in N$, $n \cdot a \otimes m - n \otimes a \cdot m$ is a homogeneous element of degree zero in $M \otimes N$ if and only if both m and a are of degree zero. Thus, in degree zero, the projection $p : N \otimes_{\mathcal{A}_0} M \rightarrow N \otimes_{\mathcal{A}} M$ is isomorphism. As, p is degreewise surjective, H_0p is an isomorphism of $H_0\mathcal{A}$ -modules. Due to right exactness of the tensor product, the universal map

$$\begin{aligned} H_0(N \otimes_{\mathcal{A}_0} M) &= \text{coker}(N \otimes_{\mathcal{A}_0} M_1 \xrightarrow{\text{id} \otimes d} N \otimes_{\mathcal{A}_0} M_0) \\ &\rightarrow N \otimes_{\mathcal{A}_0} \text{coker}(M_1 \xrightarrow{d} M_0) = N \otimes_{\mathcal{A}_0} H_0M, \\ &[n \otimes m_0] \mapsto n \otimes [m_0] \end{aligned}$$

is an isomorphism. Composing with inverse of H_0p , we get another isomorphism

$$H_0(N \otimes_{\mathcal{A}} M) \rightarrow N \otimes_{\mathcal{A}_0} H_0M, \quad [n \otimes m_0] \mapsto n \otimes [m_0]. \quad (58)$$

As the map $\mathcal{A}_0 \rightarrow H_0(\mathcal{A})$ is surjective, the quotient map

$$N \otimes_{\mathcal{A}_0} H_0M \rightarrow N \otimes_{H_0\mathcal{A}} H_0M$$

is an isomorphism as well. Composing with (58), we get the desired result. \square

We now show that that M is a strong \mathcal{A} -module.

Proposition 9. *Let $M, N \in \mathsf{Mod}(\mathcal{A})$, with M flat and cofibrant. There exists a natural isomorphism of graded $H_0(\mathcal{A})$ -modules*

$$\phi_{\bullet, N, M} : H_{\bullet}(N) \otimes_{H_0(\mathcal{A})} H_0(M) \rightarrow H_{\bullet}(N \otimes_{\mathcal{A}} M), \quad [n] \otimes [m_0] \mapsto [n \otimes m_0]. \quad (59)$$

Applying to $N = \mathcal{A}$, we find that M is a strong \mathcal{A} -module.

Proof. Taking $N = 0$, from the long exact sequence in homology associated to the fiber sequence 56, we conclude that $H_i(P \otimes_{\mathcal{A}} M) = 0$, for any $H_0\mathcal{A}$ -module P and $i > 0$. Thus, the map

$$\pi_0 : P \otimes_{\mathcal{A}} M \rightarrow H_0(P \otimes_{\mathcal{A}} M) \quad (60)$$

is a weak equivalence. Since (n) -shifts preserve weak equivalences, for any $H_0(A)$ -module P , and $n > 0$, the map

$$P[n] \otimes_{\mathcal{A}} M = (P \otimes_{\mathcal{A}} M)[n] \xrightarrow{(\phi \circ \pi_0)[n]} (P \otimes_{H_0\mathcal{A}} H_0 M)[n] = P[n] \otimes_{H_0\mathcal{A}} H_0 M \quad (61)$$

which is in degree n given by

$$p \otimes m_0 \mapsto p \otimes [m_0]$$

is a weak equivalence as well.

Notice that for any $n \in \mathbb{N}$, and $i < n$,

$$H_i(\phi_n \otimes \text{id}_M) : H_i(N \otimes_{\mathcal{A}} M) \rightarrow H_i(N_{\leq n} \otimes_{\mathcal{A}} M) \quad (62)$$

is an isomorphism. Indeed, in degrees $\leq n$, ϕ_n is identity. The same goes for $\phi_n \otimes \text{id}_M$, and hence for the induced map in homology up to degree $n - 1$.

Consequently, it suffices to prove that for all $n \in \mathbb{N}$, the map $\phi_{\bullet, N_{\leq n}, M}$, is an isomorphism. We proceed by induction.

For $n = 0$, the projection $\pi_0 : N_{\leq 0} \rightarrow H_0(N_{\leq 0})$ is a weak equivalence. As M is a cofibrant \mathcal{A} -module, $- \otimes_{\mathcal{A}} M$ preserves weak equivalences. Consequently, the composition

$$N_{\leq 0} \otimes_{\mathcal{A}} M \xrightarrow{\pi_0 \otimes M} H_0(N_{\leq 0}) \otimes_{\mathcal{A}} M \xrightarrow{\phi \circ \pi_0} H_0(N_{\leq 0}) \otimes_{H_0\mathcal{A}} H_0 M$$

is a weak equivalence as well, and the induced isomorphism in homology is exactly the inverse of (59).

Suppose now that $\phi_{\bullet, N_{\leq n-1}, M}$ is isomorphism for an $n \in \mathbb{N}$. Kernel of the fibration $p_n : N_{\leq n} \rightarrow N_{\leq n-1}$ is the complex

$$\text{Ker } p_n = (\dots \rightarrow 0 \rightarrow \text{im}(d_{n+1}^N) \hookrightarrow \ker(d_n^N) \rightarrow 0 \rightarrow \dots)$$

concentrated in degrees n and $n + 1$. There is the evident weak equivalence $\pi_n : \text{Ker } p_n \rightarrow H_n(N)[n]$. As M is flat and cofibrant,

$$\text{Ker } p_n \otimes_{\mathcal{A}} M \xrightarrow{\text{ker}(p_n) \otimes M} N_{\leq n} \otimes_{\mathcal{A}} M \xrightarrow{p_n \otimes M} N_{\leq n-1} \otimes_{\mathcal{A}} M$$

is a homotopy fiber sequence. From the associated long exact sequence in homology, we read the following:

- For $0 < i < n$, $H_i(p_n \otimes M)$ is an isomorphism. It now follows from the induction hypothesis that $\phi_{i, N_{\leq n}, M}$ is isomorphism. Degree $i = 0$ is subtle as the long exact sequence in homology confirms only that $H_0(p_n \otimes M)$ is injective. The map is surjective as p_n is degreewise surjective, and functors $-\otimes_{\mathcal{A}} M : \mathbf{Mod}(\mathcal{A}) \rightarrow \mathbf{Mod}(\mathcal{A})$, $H_0 : \mathbf{Mod}(\mathcal{A}) \rightarrow \mathbf{Mod}(H_0\mathcal{A})$ are both left adjoint, thus right exact. Functor $-\otimes_{\mathcal{A}} M$ is left adjoint as $\mathbf{Mod}(\mathcal{A})$ is a closed monoidal category, and H_0 is the left adjoint of the inclusion functor $\mathbf{Mod}(H_0\mathcal{A}) \hookrightarrow \mathbf{Mod}(\mathcal{A})$.
- $H_n(\ker p_n \otimes M)$ is an isomorphism. Let ψ denote the composed quasi-isomorphism

$$\text{Ker } p_n \otimes_{\mathcal{A}} M \xrightarrow{\pi_n \otimes M} H_n(N)[n] \otimes_{\mathcal{A}} M \xrightarrow{(\phi \circ \pi_0)[n]} H_n(N)[n] \otimes_{H_0\mathcal{A}} H_0 M.$$

It is straightforwardly verified that $\phi_{n, N_{\leq n}, M} = H_n(\ker p_n \otimes M) \circ H_n(\psi)^{-1}$, hence isomorphism.

- For $i > m$, $H_i(N_{\leq n} \otimes_{\mathcal{A}} M) = 0$, and $\phi_{n, N_{\leq n}, M} : 0 \rightarrow 0$ is trivially isomorphism.

□

Finally, we have shown that notions of flatness and strong flatness are equivalent. The story of flat modules summarizes as:

Theorem 4. *For $M \in \mathbf{Mod}(\mathcal{A})$, the following properties are equivalent:*

1. M is flat,
2. the derived tensor product $-\otimes_{\mathcal{A}}^{\mathbb{L}} M$ preserves homotopy fiber sequences,
3. M is strongly flat.

Corollary 2. *Given $M, N \in \mathbf{Mod}(\mathcal{A})$, if M is a flat \mathcal{A} -module, the above Tor spectral sequence collapses on the second page, yielding a natural isomorphism*

$$(HM \otimes_{H\mathcal{A}} HN)_q \cong H_q(M \otimes_{\mathcal{A}}^{\mathbb{L}} N).$$

Proof. Since M is flat it is also strongly flat. For strongly flat modules, the above property is proven in the subsection 7.1. □

8 Outlook

This text is part of a program which aims to establish homotopical algebraic geometry over differential operators as a natural framework for partial differential equations and their symmetries [5, 25]. For this we have to show in particular that the triplet $(\mathbf{DGDM}, \mathbf{DGDM}, \mathbf{DGDA})$ together with étale coverings and smooth morphisms, is not only a homotopical algebraic but even a homotopical algebraic geometric context. This includes proving that, in our specific environment, flat (resp., étale) morphisms are the same as strongly flat (resp., strongly étale). The proofs require that Quillen's Tor spectral sequence be valid in the \mathcal{D} -geometric setting. The latter has been proved in this work, and we expect to use it to complete the proof that solid concepts of derived stack and geometric derived stack do exist in our homotopical \mathcal{D} -geometric setting. Viewed from a broader perspective, the present work is also part of an effort to strengthen the role of the functor-of-points as a fundamental approach in derived algebraic geometry [32, 26] and colored supergeometry [8, 6, 7, 28].

References

- [1] Ardizzoni A. The category of modules over a monoidal category: Abelian or not? *Ann. Univ. Ferrara*, 50:167–185, 2004.
- [2] Behrend K, Liao H Y, Xu P. Derived Differentiable Manifolds. arXiv 2006.01376, 2020.
- [3] Beilinson A and Drinfeld V. *Chiral algebras*. Colloquium Publications, volume 51, AMS, 2004.
- [4] Bruce A, Ibarguengoytia E, Poncin N. Linear \mathbb{Z}_n^2 -manifolds and linear actions. *SIGMA*, 17(060), 2021.
- [5] Costello K and Gwilliam O. *Factorization Algebras in Quantum Field Theory, volume 1*. Cambridge University Press, 2016.
- [6] Covolo T, Grabowski J, Poncin N. The category of \mathbb{Z}_2^n -supermanifolds. *J. Math. Phys.*, 57(7), 2016.
- [7] Covolo T, Kwok S, Poncin N. Local forms of morphisms of colored supermanifolds. *J. Geom. Phys.*, 168, 2021.
- [8] Covolo T, Ovsienko V, Poncin N. Higher trace and Berezinian of matrices over a Clifford algebra. *J. Geom. Phys.*, 62(12):2294–2319, 2012.

- [9] Di Brino G, Pištalo D, Poncin N. Koszul-Tate resolutions as cofibrant replacements of algebras over differential operators. *J. Homotopy Relat. Struct.*, 13(4):793–846, 2018.
- [10] Di Brino G, Pištalo D, Poncin N. Homotopical algebraic context over differential operators. *J. Homotopy Relat. Struct.*, 14(1):293–347, 2018.
- [11] Elmendorf A D, Kříž I, Mandell M A, May J P. Rings, modules, and algebras in stable homotopy theory. *Math. surv. monogr.*, 47, 2007.
- [12] Dwyer W G and Spaliński J. Homotopy theories and model categories. In James I M, editor, *Handbook of algebraic topology*, chapter 2, pages 73–126. Elsevier, Amsterdam, 1995.
- [13] Gaitsgory D, Rozenblyum N. Crystals and \mathbb{D} -modules. *Pure appl. math.*, 10:57–154, 2014.
- [14] Govzmann A, Pištalo D, Poncin N. A new approach to model categorical homotopy fiber sequences. *Dissert. Math.*, 581, 2022.
- [15] Govzmann A, Pištalo D, Poncin N. Comparison theorems for Kan, faintly universal and strongly universal derived functors. *Surv. Math. & Appl.*, 17:397 – 429, 2022.
- [16] Govzmann A, Pištalo D, Poncin N. Indeterminacies and models of homotopy limits. *Theory Appl. Categ.*, 38, 2022.
- [17] Hirschhorn P S. *Model categories and their localizations*. Mathematical Surveys and Monographs, volume 99, AMS, 2003.
- [18] Hovey M. *Model categories*. Mathematical Surveys and Monographs, volume 63, AMS, 1999.
- [19] Joyce D. An introduction to d-manifolds and derived differential geometry. In P. Thomas R Newstead P Brambila-Paz L and García-Prada 0, editors, *Moduli Spaces (volume 411 of London Math. Soc. Lecture Note Ser.)*, chapter 5, page 230–281. Cambridge University Press, Cambridge, 2014.
- [20] Kashiwara M. *\mathcal{D} -modules and Microlocal Calculus*. Transl. Math. Monographs, volume 217, AMS, 2002.
- [21] Kříž I, May J P. Operads, algebras, modules and motives. *Astérisque*, 233, 1995.

- [22] Lurie J. *Higher Topos Theory*. AM-170, Annals of Mathematics Studies, Princeton University Press, 2009.
- [23] Lurie J. Higher algebra, 2017.
- [24] Mac Lane S. *Categories for the working mathematician (second edition)*. Graduate Texts in Mathematics, volume 5, Springer-Verlag, 1998.
- [25] Paugam F. Histories and observables in covariant field theory. *J. Geom. Phys.*, 61(9):1675–1702, 2011.
- [26] Paugam F. *Towards the mathematics of quantum Field theory*. A Series of Modern Surveys in Mathematics, volume 59, Springer, 2014.
- [27] Pištalo D and Poncin N. Koszul-Tate resolutions and Sullivan models. *Dissert. Math.*, 531, 2018.
- [28] Poncin N. Towards integration on colored supermanifolds. *Banach Center Publ.*, 110:201–217, 2016.
- [29] Porta M. GAGA theorems in derived complex geometry. *J. Algebraic Geom.*, 28:519–565, 2019.
- [30] Quillen D. *Homotopical algebra*. Lecture Notes in Mathematics, volume 43, Springer-Verlag, 1967.
- [31] Spivak D I. Derived smooth manifolds. *Duke Math. J.*, 153(1):55–128, 2010.
- [32] Toën B and Vezzosi G. Homotopical algebraic geometry I, Topos theory. *Adv. Math.*, 193(2):257–372, 2005.
- [33] Toën B and Vezzosi G. Homotopical algebraic geometry II, Geometric stacks and applications. *Mem. Amer. Math. Soc.*, 193(902), 2008.
- [34] van Nigtevecht S. The K-theory cochains of H-spaces and height 1 chromatic homotopy theory. arXiv 2209.07311, 2022.
- [35] Vinogradov A M. *Cohomological analysis of Partial Differential Equations and Secondary Calculus*. Transl. Math. Monographs, volume 204, AMS, 2001.
- [36] Weibel C A. *An introduction to homological algebra*. Cambridge Studies in Advanced Mathematics, volume 38, Cambridge University Press, 1994.

email: alisa.govzmann@uni.lu; *email:* damjan.pistalo@uni.lu; *email:* norbert.poncin@uni.lu.