

# Derived categories of flips and cubic hypersurfaces

Pieter Belmans      Lie Fu      Theo Raedschelders

February 13, 2020

*dedicated to the memory of Tom Nevins*

## Abstract

A classical result of Bondal–Orlov states that a standard flip in birational geometry gives rise to a fully faithful functor between derived categories of coherent sheaves. We complete their embedding into a semiorthogonal decomposition by describing the complement. As an application, we can lift the “quadratic Fano correspondence” (due to Galkin–Shinder) in the Grothendieck ring of varieties between a smooth cubic hypersurface, its Fano variety of lines, and its Hilbert square, to a semiorthogonal decomposition.

We also show that the Hilbert square of a cubic hypersurface of dimension at least 3 is again a Fano variety, so in particular the Fano variety of lines on a cubic hypersurface is a Fano visitor. The most interesting case is that of a cubic fourfold, where this exhibits the first higher-dimensional hyperkähler variety as a Fano visitor.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>A fully faithfulness criterion for EZ-type functors</b>	<b>5</b>
<b>3</b>	<b>A semiorthogonal decomposition for standard flips</b>	<b>9</b>
<b>4</b>	<b>Application: quadratic Fano correspondence for cubic hypersurfaces</b>	<b>16</b>
<b>5</b>	<b>Hilbert squares of low-degree hypersurfaces</b>	<b>21</b>
<b>A</b>	<b>Short proof of theorem A(ii) for <math>\ell = 1</math></b>	<b>27</b>

# 1 Introduction

## Derived categories of flips

The (conjectural) interaction between birational geometry and derived categories is largely based on the *DK-hypothesis* [22]. This hypothesis predicts that K-equivalent varieties  $X$  and  $Y$  are D-equivalent, i.e. their derived categories of coherent sheaves are equivalent as triangulated categories. Here K-equivalence means the existence of a diagram

$$(1) \quad \begin{array}{ccc} & Z & \\ f \swarrow & & \searrow g \\ X & & Y \end{array}$$

of birational morphisms between smooth projective varieties, such that  $f^*K_X \sim_{\text{lin}} g^*K_Y$ . If on the other hand we have a K-inequality, i.e.  $f^*K_X + D \sim_{\text{lin}} g^*K_Y$  for some effective divisor  $D$  on  $Z$ , then it predicts the existence of a fully faithful functor  $\mathbf{D}^b(Y) \hookrightarrow \mathbf{D}^b(X)$ .

We will study the case of a specific K-inequality, where we complete the fully faithful functor predicted by the DK-hypothesis into a semiorthogonal decomposition, i.e. we explicitly describe the complement. The special instance of a birational transformation we are interested in is that of a (standard) flip. We recall the setup: let  $F$  be a smooth projective variety defined over an algebraically closed field  $k$  of characteristic zero. Let  $k$  and  $\ell$  be two positive integers. Let  $\mathcal{V}$  be a vector bundle of rank  $k + 1$  on  $F$ , denote  $Z := \mathbb{P}(\mathcal{V})$  and let  $\pi : Z \rightarrow F$  be the associated projective bundle.

Let  $X$  be a smooth projective variety containing  $Z$  as a closed subvariety such that the restriction of the normal bundle  $N_{Z/X}$  to each fiber of  $\pi$  is isomorphic to  $\mathcal{O}(-1)^{\oplus \ell+1}$ . In other words,  $N_{Z/X} \simeq \mathcal{O}_\pi(-1) \otimes \pi^*\mathcal{V}'$  for some vector bundle  $\mathcal{V}'$  of rank  $\ell + 1$  on  $F$ .

Let  $\tau : \tilde{X} \rightarrow X$  be the blow up of  $X$  along the smooth center  $Z$ , then the exceptional divisor  $E$  is isomorphic to  $\mathbb{P}(\mathcal{V}) \times_F \mathbb{P}(\mathcal{V}')$  with normal bundle  $N_{E/\tilde{X}} \simeq \mathcal{O}(-1, -1)$ . Therefore we can contract  $E$  along the other direction,  $E \rightarrow Z' := \mathbb{P}(\mathcal{V}')$ , which identifies  $E$  as the projectivization of the vector bundle  $\pi'^*(\mathcal{V}')$ , where  $\pi' : Z' \rightarrow F$  is the natural projection. We summarize the situation in the following diagram.

$$(2) \quad \begin{array}{ccccc} & & E & & \\ & & \downarrow j & & \\ & p \swarrow & & \searrow p' & \\ & & \tilde{X} & & \\ & \tau \swarrow & & \searrow \tau' & \\ Z \xrightarrow{i} X & \xrightarrow{\phi} & X' & \xleftarrow{i'} & Z' \\ & \pi \searrow & & \swarrow \pi' & \\ & & F & & \end{array}$$

Here the birational transform  $\phi$  is called a *standard flip* (resp. *standard flop* when  $k = \ell$ ) while the inner triangle is its resolution; the upper two trapezoids are blowup diagrams and the outer square is cartesian. Observe that  $\pi$  and  $p'$  are  $\mathbb{P}^k$ -bundles while  $\pi'$  and  $p$  are  $\mathbb{P}^\ell$ -bundles. We will refer to (2) as a *standard flip diagram*.

By symmetry, we are free to assume that  $k \geq \ell$ . According to the general conjecture on derived categories under birational transformations discussed above, the derived category of  $X'$  is expected to be “smaller” than that of  $X$ . Bondal–Orlov established this in the above setting of standard flips [12, theorem 3.6].

**Theorem 1** (Bondal–Orlov). Assume a standard flip diagram (2) is given, with  $k \geq \ell$ . The functor

$$(3) \quad \mathbf{R}\tau_* \circ \mathbf{L}\tau'^*: \mathbf{D}^b(X') \rightarrow \mathbf{D}^b(X)$$

is fully faithful. Moreover, if  $k = \ell$ , this functor is an equivalence of triangulated categories.

The first goal of this paper is to identify the complement of  $\mathbf{D}^b(X')$  inside  $\mathbf{D}^b(X)$ , thus completing theorem 1 into a semiorthogonal decomposition. This specific question was stated in [37, remarque 4.5], and the following theorem accomplishes this.

**Theorem A.** Assume a standard flip diagram (2) is given, with  $k > \ell$ .

(i) For every integer  $m$ , the functor

$$(4) \quad \Phi_m: \mathbf{D}^b(F) \rightarrow \mathbf{D}^b(X): \mathcal{E} \mapsto i_*(\pi^*(\mathcal{E}) \otimes \mathcal{O}_\pi(m)),$$

is fully faithful.

(ii) We have the following semiorthogonal decomposition of  $\mathbf{D}^b(X)$ :

$$(5) \quad \mathbf{D}^b(X) = \langle \Phi_{-k+\ell}(\mathbf{D}^b(F)), \dots, \Phi_{-1}(\mathbf{D}^b(F)), \mathbf{R}\tau_* \circ \mathbf{L}\tau'^* \mathbf{D}^b(X') \rangle.$$

In fact, part (i) of theorem A follows from a more general fully faithfulness criterion for EZ-type functors which might be of independent interest, see proposition 3.

For the application in theorem B below we only need theorem A(ii) for  $\ell = 1$ . In this case a more elegant proof is possible, which moreover gives the slightly stronger result that compares the category  $\mathbf{D}^b(X)$  and the components of (5) as subcategories in  $\mathbf{D}^b(\tilde{X})$ , and not only after applying  $\mathbf{R}\tau_*$ . This will be discussed in appendix A.

Note also that the case  $\ell = 0$  of theorem A is exactly Orlov’s blowup formula [35, theorem 4.3].

More recently, [1] and [21] discusses the behavior of derived categories under flops (but not flips), and [20] discusses the behavior of Chow groups under standard flips. Kawamata has proven analogous results in the toric and toroidal case, see [23, theorem 6.1], [24, theorem 1] and [25, theorem 1].

**Remark 2.** By mutating the first few terms of (5) to the far right, we get a series of similar semiorthogonal decompositions: for any integer  $0 \leq m \leq k - \ell$ , we have

$$(6) \quad \mathbf{D}^b(X) = \langle \Phi_{-m}(\mathbf{D}^b(F)), \dots, \Phi_{-1}(\mathbf{D}^b(F)), \mathbf{R}\tau_* \circ \mathbf{L}\tau'^*(\mathbf{D}^b(X')), \Phi_0(\mathbf{D}^b(F)), \dots, \Phi_{k-\ell-m-1}(\mathbf{D}^b(F)) \rangle.$$

## Smooth cubic hypersurfaces: quadratic Fano correspondence

As an illustration of the usefulness of theorem A, we will consider a smooth cubic hypersurface  $Y \subset \mathbb{P}^{n+1}$ . Building upon insights of Galkin–Shinder [15] and Voisin

[43], we show in corollary 15 that the *quadratic Fano correspondence*

$$(7) \quad \begin{array}{ccc} P_2 & \hookrightarrow & Y^{[2]} \\ \downarrow & & \\ F(Y) & & \end{array}$$

obtained from the universal family (or *Fano correspondence*)

$$(8) \quad \begin{array}{ccc} P & \longrightarrow & Y \\ \downarrow & & \\ F(Y) & & \end{array}$$

by taking the relative Hilbert square of  $P \rightarrow Y$  gives a standard flip diagram where  $X$  is the Hilbert square of  $Y$  and  $F$  the Fano variety  $F(Y)$  of lines on  $Y$ . The role of  $X'$  is played by a certain  $\mathbb{P}^n$ -bundle over  $Y$ . For more on the terminology and notation, one is referred to section 4. An excellent reference for more background on cubic hypersurfaces is [18].

This brings us to the second main result.

**Theorem B.** Let  $Y \subset \mathbb{P}^{n+1}$  be a smooth cubic hypersurface. Let  $F(Y)$  be its Fano variety of lines and  $Y^{[2]}$  be its Hilbert square. Then there is a semiorthogonal decomposition

$$(9) \quad \mathbf{D}^b(Y^{[2]}) = \langle \mathbf{D}^b(F(Y)), \underbrace{\mathbf{D}^b(Y), \dots, \mathbf{D}^b(Y)}_{n+1 \text{ copies}} \rangle.$$

For the precise form of the functors one is referred to section 4.

This is a derived categorical version, of a comparison between the Fano variety of lines and the Hilbert square of a cubic hypersurface, which was known to hold in various contexts, such as cohomology, classes in the Grothendieck ring of varieties, or (rational) Chow motives. In section 4 we discuss these.

## Fano visitors

Using theorem B we can study the Fano visitor problem for the Fano variety of lines on a cubic hypersurface. Recall that a smooth projective variety  $X$  is said to be a *Fano visitor* if there exists a smooth projective Fano variety  $Y$  and a fully faithful functor  $\mathbf{D}^b(X) \hookrightarrow \mathbf{D}^b(Y)$ . In this case we call  $Y$  a *Fano host* for  $X$ .

Bondal raised the question whether every smooth projective variety is a Fano visitor. A positive answer would imply that (additive invariants of) derived categories of Fano varieties are as complicated as those of arbitrary varieties. This question has been addressed in [27, 8, 6, 34, 30, 14] for various families of varieties, in particular it is known that curves, Enriques surfaces and complete intersections are Fano visitors.

To study the Fano visitor problem for Fano varieties of lines<sup>1</sup>, we prove the following result.

---

<sup>1</sup>To avoid any confusion with other usages of the name Fano, we will always write “Fano variety of lines” in full when we refer to  $F(Y)$ , where  $Y$  is a cubic hypersurface.

**Theorem C.** Let  $n \geq 3$ . Let  $Y$  be  $\mathbb{P}^n$ , or a smooth quadric (resp. cubic) hypersurface in  $\mathbb{P}^{n+1}$ . Then  $Y^{[2]}$  is Fano.

This is in contrast to the case of smooth projective surface  $S$ , in which case the Hilbert scheme  $S^{[m]}$  of  $m$  points is never a Fano variety. We immediately obtain the following corollary to theorem B and theorem C.

**Corollary D.** Let  $Y$  be a smooth cubic hypersurface of dimension  $n = 3, 4$ . Then the Fano variety of lines  $F(Y)$  is a Fano visitor with Fano host  $D^b(Y^{[2]})$ .

This is the first construction (to our knowledge) of a Fano host for a surface of general type which is not a complete intersection or a product of curves (for  $n = 3$ ), and the first construction of a Fano host for a (higher-dimensional) hyperkähler variety (for  $n = 4$ ). For  $n \geq 5$  the Fano variety of lines is itself a Fano variety, so the Fano visitor problem is trivial.

**Conventions** Throughout we will assume that  $\mathbf{k}$  is an algebraically closed field of characteristic 0.

**Acknowledgements** We want to thank Sergey Galkin, Daniel Huybrechts, Robert Laterveer, Will Sawin and Mingmin Shen for helpful discussions.

The results in this article were conceived during the “Problems and recent developments in hyperkähler geometry” research seminar, organised by Daniel Huybrechts, and we heartily thank him.

The first author is partially supported by a postdoctoral fellowship from the Research Foundation—Flanders (FWO). The third author is supported by a postdoctoral fellowship from the Research Foundation—Flanders (FWO).

## 2 A fully faithfulness criterion for EZ-type functors

Let  $F, Z$  and  $X$  be smooth projective varieties. Assume there is a smooth proper morphism  $\pi: Z \rightarrow F$  and a closed immersion  $i: Z \rightarrow X$  as in the following diagram.

$$(10) \quad \begin{array}{ccc} Z & \xhookrightarrow{i} & X \\ \downarrow \pi & & \\ F & & \end{array}$$

Let  $c$  be the codimension of  $Z$  in  $X$ . The following result is inspired by [16, Theorem 2.1], in the formulation of [1, Theorem 1.3]. The terminology *EZ-type* is taken from [16], where the diagram (10) is denoted with  $E$  instead of  $F$ .

**Proposition 3.** Assume that for every fiber  $P$  of  $\pi$  and all integers  $p, q$  with  $p + q > 0$ ,

$$(11) \quad H^p(P, \bigwedge^q \mathcal{N}') = 0,$$

where  $\mathcal{N}' := N_{Z/X}|_P$  is the restriction of the normal bundle of  $Z$  in  $X$  to  $P$ . Then for every line bundle  $\mathcal{L}$  on  $Z$ , the functor

$$(12) \quad \Phi: D^b(F) \rightarrow D^b(X) : \mathcal{E} \mapsto i_*(\pi^*(\mathcal{E}) \otimes \mathcal{L}),$$

is fully faithful.

Its proof is an application of the Bondal–Orlov criterion for fully faithfulness [12, Theorem 1.1], which we will quickly recall.

**Proposition 4** (Bondal–Orlov criterion). Let  $X$  and  $Y$  be smooth projective varieties, and  $\mathcal{E}$  an object in  $\mathbf{D}^b(X \times Y)$ . The corresponding Fourier–Mukai functor  $\Phi_{\mathcal{E}}: \mathbf{D}^b(X) \rightarrow \mathbf{D}^b(Y)$  is fully faithful if and only if for all closed points  $x, x' \in X$  we have that

- (i)  $\mathrm{Hom}_Y(\Phi_{\mathcal{E}}(\mathcal{O}_x), \Phi_{\mathcal{E}}(\mathcal{O}_x)) \cong \mathbf{k}$ ;
- (ii)  $\mathrm{Hom}_Y(\Phi_{\mathcal{E}}(\mathcal{O}_x), \Phi_{\mathcal{E}}(\mathcal{O}_x)[m]) \cong 0$  for all  $m \notin [0, \dim X]$ ;
- (iii)  $\mathrm{Hom}_Y(\Phi_{\mathcal{E}}(\mathcal{O}_x), \Phi_{\mathcal{E}}(\mathcal{O}_{x'})[m]) \cong 0$  for all  $m \in \mathbb{Z}$  and  $x \neq x'$ ;

where  $\mathcal{O}_x$  denotes the skyscraper sheaf at  $x$ .

*Proof of proposition 3.* For a (closed) point  $x$  in  $F$ , denote by  $P := \pi^{-1}(x)$  the fiber. We denote  $j: P \hookrightarrow X$  for the closed immersion obtained by composition with  $i$ . Then,  $\Phi(\mathcal{O}_x) = i_*(\mathcal{O}_P \otimes \mathcal{L}) = j_*(\mathcal{L}|_P)$ .

Let us first check (iii) of proposition 4. Let  $P'$  denote the fiber  $\pi^{-1}(x')$  for  $x \neq x'$  in  $F$ . Since  $P \cap P' = \emptyset$ , the objects  $j_*(\mathcal{L}|_P)$  and  $j'_*(\mathcal{L}|_{P'})$  have disjoint support, hence they are completely orthogonal.

To check (i) and (ii), we have by adjunction that

$$(13) \quad \mathrm{Hom}_X(j_*(\mathcal{L}|_P), j_*(\mathcal{L}|_P)[m]) \cong \mathrm{Hom}_P(\mathcal{L}|_P, j^! \circ j_*(\mathcal{L}|_P)[m]).$$

By [17, Corollary 11.2 and Proposition 11.8], the cohomology sheaves of the complex  $\mathbf{L}j^* \circ j_*(\mathcal{L}|_P)$  are given by

$$(14) \quad \mathcal{H}^{-s}(\mathbf{L}j^* \circ j_*(\mathcal{L}|_P)) \cong \mathcal{L}|_P \otimes \bigwedge^s \mathcal{N}^\vee \text{ for } s = 0, \dots, c + \dim F,$$

where  $\mathcal{N} := \mathbf{N}_{P/X}$  is the normal bundle of  $P$  in  $X$ . As  $j^!$  and  $\mathbf{L}j^*$  are related by Grothendieck duality, we have that

$$(15) \quad j^! \circ j_*(\mathcal{L}|_P) = \mathbf{L}j^* \circ j_*(\mathcal{L}|_P) \otimes \omega_j[\dim P - \dim X] = \mathbf{L}j^* \circ j_*(\mathcal{L}|_P) \otimes \det(\mathcal{N})[-\dim F - c]$$

and this object has cohomology sheaves

$$(16) \quad \mathcal{H}^s(j^! \circ j_*(\mathcal{L}|_P)) \cong \mathcal{L}|_P \otimes \bigwedge^s \mathcal{N} \text{ for } s = 0, \dots, c + \dim F.$$

We can compute the right-hand side of (13) via the hypercohomology spectral sequence and we obtain the spectral sequence

$$(17) \quad E_2^{p,q} = \mathrm{H}^p(P, \mathcal{H}^q(j^! \circ j_*(\mathcal{L}|_P) \otimes \mathcal{L}^{-1}|_P)) = \mathrm{H}^p(P, \bigwedge^q \mathcal{N}) \Rightarrow \mathrm{Hom}_X(j_*(\mathcal{L}|_P), j_*(\mathcal{L}|_P)[p+q]).$$

Hence (i), and (ii) for  $m < 0$  follow immediately.

To check (ii) for  $m > \dim F$ , consider the following identification of the short exact sequence of normal bundles

$$(18) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathbf{N}_{P/Z} & \longrightarrow & \mathbf{N}_{P/X} & \longrightarrow & \mathbf{N}_{Z/X}|_P \longrightarrow 0 \\ & & \parallel & & \parallel & & \parallel \\ 0 & \longrightarrow & \mathcal{O}_P^{\oplus \dim F} & \longrightarrow & \mathcal{N} & \longrightarrow & \mathcal{N}' \longrightarrow 0 \end{array}$$

as the first term is isomorphic to  $\mathcal{O}_P \otimes_k T_x F$ . Therefore, for any positive integer  $q$ , the bundle  $\wedge^q \mathcal{N}$  is a successive extension of direct sums of exterior powers of  $\mathcal{N}'$ . From the vanishing hypothesis, we see that  $H^p(P, \wedge^q \mathcal{N}) = 0$  for all  $p + q > \dim F$ . As a result,  $E_\infty^{p,q} = 0$  for all  $p + q > \dim F$ , and we are done.  $\square$

## 2.1 Some applications

We will briefly discuss a few situations where this criterion applies, before proceeding with the setting of standard flips in section 3.

**Orlov’s blowup formula** The easiest setting where one can apply proposition 3 is that of the usual blowup of a smooth projective variety  $X$  in a smooth center  $Z$  of codimension  $c \geq 2$ , i.e. we consider the diagram

$$(19) \quad \begin{array}{ccc} E & \xhookrightarrow{i} & \mathrm{Bl}_Z X \\ \downarrow \pi & & \\ Z & & \end{array}$$

where  $i$  denotes the inclusion of the exceptional divisor  $E$ . Then the vanishing condition (11) is clearly satisfied as we have the identification  $\mathcal{N}' \cong \mathcal{O}_P(-1)$  using the notation of proposition 3, and we see that the functor

$$(20) \quad \Phi_m : \mathbf{D}^b(Z) \rightarrow \mathbf{D}^b(\mathrm{Bl}_Z X) : \mathcal{E} \mapsto i_*(\pi^*(\mathcal{E}) \otimes \mathcal{O}_E(mE))$$

is fully faithful for every  $m \in \mathbb{Z}$ , recovering [35, assertion 4.2(i)], which is one of the ingredients for Orlov’s blowup formula.

**Krug–Ploog–Sosna’s cyclic quotient singularities** More interestingly, proposition 3 also applies to the “singular” blowups of [28], for cyclic quotient singularities. The setting, using the notation of op. cit., is as follows. Let  $Y$  denote a smooth quasiprojective variety with an action of a cyclic group  $G \cong \mu_m$ . Then the fixed point locus  $S \subset Y$  (of codimension  $n$ ) is smooth and the quotient  $Y/G$  has rational singularities. We further assume that only the isotropy groups 1 and  $G$  occur, and that a generator of  $G$  acts on the normal bundle  $N_{S/Y}$  by multiplication with a fixed primitive  $m$ th root of unity.

In this case there is a diagram

$$(21) \quad \begin{array}{ccc} Z := \mathbb{P}(N_{S/Y}) & \xhookrightarrow{i} & X := \mathrm{Bl}_S(Y/G) \cong (\mathrm{Bl}_S Y)/G \\ \downarrow \pi & & \\ S & & \end{array}$$

of smooth projective varieties. Here  $Y/G$  has cyclic quotient singularities, and the “singular blowup”  $\mathrm{Bl}_S(Y/G)$  is a resolution of singularities.

By [28, lemma 4.10(ii) and (vi)], we have that

$$(22) \quad N_{Z/X}|_P \cong \mathcal{O}_{\mathbb{P}^{n-1}}(-m)$$

for every fiber  $P$  of  $\pi$ , so proposition 3 applies as soon as  $n > m$  and one obtains fully faithful functors

$$(23) \quad \Theta_\beta: \mathbf{D}^b(S) \rightarrow \mathbf{D}^b(\mathrm{Bl}_S(Y/G)) : \mathcal{E} \mapsto i_*(\pi^*(\mathcal{E}) \otimes \mathcal{O}_\pi(\beta)),$$

for  $\beta \in \mathbb{Z}$ , as in [28, theorem 4.1(ii)], which is one of the ingredients for the Krug–Ploog–Sosna semiorthogonal decomposition.

A particularly relevant example for this paper is given by taking  $Y := S \times S$  with  $S$  a smooth projective variety, and  $G = \mu_2$  acting via transposition; in this case  $X \cong S^{[2]}$ .

**Proposition 5** (Krug–Ploog–Sosna). Let  $S$  be a smooth projective variety of dimension  $n \geq 2$ . Then there exists a semiorthogonal decomposition

$$(24) \quad \mathbf{D}^b(S^{[2]}) = \langle \mathbf{D}_{\mu_2}^b(S^2), \underbrace{\mathbf{D}^b(S), \dots, \mathbf{D}^b(S)}_{n-2 \text{ copies}} \rangle.$$

**Cayley’s trick** A third application is Cayley’s trick, which gives a relationship between a complete intersection and a canonically associated hypersurface in a projective bundle. This relationship can be studied at different levels, and the name “Cayley’s trick” originates from [19] where it was used to study cohomology. In [27] Cayley’s trick was used on the level of derived categories to study Fano visitors as in section 5.3, and we will now explain how to obtain part of their semiorthogonal decomposition.

Let  $\mathcal{E}$  denote a vector bundle of rank  $r$  on a smooth variety  $Y$ , and  $s \in H^0(Y, \mathcal{E})$  a regular section with smooth projective zero locus  $Z(s)$ . By the isomorphism

$$(25) \quad H^0(Y, \mathcal{E}) \cong H^0(\mathbb{P}(\mathcal{E}^\vee), \mathcal{O}_{\mathbb{P}(\mathcal{E}^\vee)}(1)),$$

the section  $s$  also defines a section  $f_s \in H^0(\mathbb{P}(\mathcal{E}^\vee), \mathcal{O}_{\mathbb{P}(\mathcal{E}^\vee)}(1))$  and we denote  $Z(f_s)$  its zero locus, which defines a hypersurface in  $\mathbb{P}(\mathcal{E}^\vee)$ . In particular, there is a cartesian diagram

$$(26) \quad \begin{array}{ccc} Z := \mathbb{P}(\mathbf{N}_{Z(s)/Y}) & \xleftarrow{i} & X := Z(f_s) \\ \downarrow \pi & \square & \downarrow \\ Z(s) & \xleftarrow{\quad} & Y \end{array}$$

where  $X \rightarrow Y$  is a  $\mathbb{P}^{r-2}$ -bundle over  $Y \setminus Z(s)$  and the restriction  $\pi$  is a  $\mathbb{P}^{r-1}$ -bundle. In this case, the normal bundle is

$$(27) \quad \mathbf{N}_{Z/X}|_P \cong \Omega_{\mathbb{P}^{r-1}}(1),$$

for a fiber  $P \cong \mathbb{P}^{r-1}$  of  $\pi$ , and proposition 3 applies by Bott vanishing. This yields fully faithful functors

$$(28) \quad \Phi: \mathbf{D}^b(Z(s)) \rightarrow \mathbf{D}^b(Z(f_s)) : \mathcal{E} \mapsto i_*(\pi^*(\mathcal{E}) \otimes \mathcal{L}),$$

for every line bundle  $\mathcal{L}$  on  $\mathbb{P}(\mathbf{N}_{Z(s)/Y})$ , recovering part of [36, proposition 2.10].



### 3 A semiorthogonal decomposition for standard flips

In this section we prove theorem A. Part (i) follows immediately from proposition 3.

The proof of part (ii) of theorem A is more interesting, and is very much inspired by Kuznetsov's homological projective duality [31] and its interpretation by Thomas [41]. In the case  $\ell = 1$ , a slightly different (and shorter) proof of a stronger statement is given in appendix A.

The semiorthogonal decomposition (5) will be established in several steps:

1. The fully faithfulness of the functor  $\Phi_m$  on  $\mathbf{D}^b(F)$  follows from proposition 3.
2. The fully faithfulness of the functor  $\mathbf{R}\tau_* \circ \mathbf{L}\tau'^*$  on  $\mathbf{D}^b(X')$  is Bondal–Orlov's theorem, see theorem 1 and also remark 10.
3. The semiorthogonality of the subcategories in (5) is the combination of proposition 8 and proposition 9.
4. The completeness of the semiorthogonal decomposition (5) is corollary 12.

Let us first introduce some notation. Assume we are given a standard flip diagram (2). For integers  $a, b \in \mathbb{Z}$ , denote by

$$(29) \quad \mathcal{O}(a, b) := p^* \mathcal{O}_\pi(a) \otimes p'^* \mathcal{O}_{\pi'}(b),$$

which is a line bundle on  $E$ . We then define the following triangulated subcategory of  $\mathbf{D}^b(\tilde{X})$ :

$$(30) \quad \mathcal{A}(a, b) := j_*(p^* \circ \pi^* \mathbf{D}^b(F) \otimes \mathcal{O}(a, b)).$$

We note first that the components

$$(31) \quad \Phi_m(\mathbf{D}^b(F)) := i_*(\pi^*(\mathbf{D}^b(F)) \otimes \mathcal{O}_\pi(m)),$$

for  $m \in \mathbb{Z}$  appearing in the decomposition (5) can be expressed in terms of the categories just defined, by the following lemma.

**Lemma 6.** For every  $m \in \mathbb{Z}$  we have the identification of subcategories

$$(32) \quad \Phi_m(\mathbf{D}^b(F)) = \mathbf{R}\tau_* \mathcal{A}(m, 0)$$

in  $\mathbf{D}^b(X)$ .

*Proof.* This follows from the computation

$$(33) \quad \begin{aligned} \mathbf{R}\tau_* \mathcal{A}(m, 0) &= \mathbf{R}\tau_* \circ j_*(p^* \circ \pi^* \mathbf{D}^b(F) \otimes p^* \mathcal{O}_\pi(m)) \\ &= i_* \circ \mathbf{R}p_* \circ p^*(\pi^* \mathbf{D}^b(F) \otimes \mathcal{O}_\pi(m)) \\ &= i_*(\pi^* \mathbf{D}^b(F) \otimes \mathcal{O}_\pi(m)) \\ &= \Phi_m(\mathbf{D}^b(F)). \end{aligned}$$

□

For convenience, we also define the following subcategories which are equivalent to  $\mathbf{D}^b(Z')$  and  $\mathbf{D}^b(Z)$  respectively.

$$(34) \quad \begin{aligned} \mathcal{A}(a, \star) &:= j_*(p^* \mathcal{O}_\pi(a) \otimes p'^* \mathbf{D}^b(Z')), \\ \mathcal{A}(\star, b) &:= j_*(p^* \mathbf{D}^b(Z) \otimes p'^* \mathcal{O}_{\pi'}(b)). \end{aligned}$$

By Orlov's projective bundle formula [35, theorem 2.6], there are the following semiorthogonal decompositions for every  $m \in \mathbb{Z}$ :

$$(35) \quad \mathcal{A}(a, \star) = \langle \mathcal{A}(a, m-l), \mathcal{A}(a, m-l+1), \dots, \mathcal{A}(a, m) \rangle.$$

Similarly we have the decompositions

$$(36) \quad \mathcal{A}(\star, b) = \langle \mathcal{A}(m-k, b), \mathcal{A}(m-k+1, b), \dots, \mathcal{A}(m, b) \rangle.$$

Applying Orlov's blowup formula [35, theorem 4.3] to the blowup  $\tau: \tilde{X} \rightarrow X$ , we have the semiorthogonal decomposition:

$$(37) \quad \mathbf{D}^b(\tilde{X}) = \langle \mathcal{A}(\star, -l), \mathcal{A}(\star, -l+1), \dots, \mathcal{A}(\star, -1), \mathbf{L}\tau^* \mathbf{D}^b(X) \rangle.$$

Similarly, applying the blowup formula to  $\tau': \tilde{X} \rightarrow X$ , we have

$$(38) \quad \mathbf{D}^b(\tilde{X}) = \langle \mathcal{A}(-k, \star), \mathcal{A}(-k+1, \star), \dots, \mathcal{A}(-1, \star), \mathbf{L}\tau'^* \mathbf{D}^b(X') \rangle.$$

Inserting appropriately chosen (35) and (36) into (37) and (38) we get different semiorthogonal decompositions of  $\mathbf{D}^b(\tilde{X})$ . The main theme of the proof is to compare them. The following vanishing result plays a central role in the argument, and is similar to [41, lemma 4.3], but we are not in the setting of homological projective duality context.

**Lemma 7.** Let  $a_1, a_2, b_1, b_2$  be integers. Then  $\mathbf{RHom}_{\tilde{X}}(\mathcal{A}(a_1, b_1), \mathcal{A}(a_2, b_2)) = 0$  in any of the following cases:

- $1 \leq a_1 - a_2 \leq k - 1$ ,
- $1 \leq b_1 - b_2 \leq \ell - 1$ ,
- $a_1 - a_2 = k$  and  $0 \leq b_1 - b_2 \leq \ell - 1$ ,
- $b_1 - b_2 = \ell$  and  $0 \leq a_1 - a_2 \leq k - 1$ .

*Proof.* For arbitrary  $\mathcal{F}_1, \mathcal{F}_2 \in \mathbf{D}^b(F)$ , by adjunction,

$$(39) \quad \begin{aligned} &\mathbf{RHom}(j_*(p^* \circ \pi^* \mathcal{F}_1 \otimes \mathcal{O}(a_1, b_1)), j_*(p^* \circ \pi^* \mathcal{F}_2 \otimes \mathcal{O}(a_2, b_2))) \\ &= \mathbf{RHom}(\mathbf{L}j^* \circ j_*(p^* \circ \pi^* \mathcal{F}_1 \otimes \mathcal{O}(a_1, b_1)), p^* \circ \pi^* \mathcal{F}_2 \otimes \mathcal{O}(a_2, b_2)) \end{aligned}$$

By [17, Corollary 11.4(ii)], there is a distinguished triangle:

$$(40) \quad p^* \circ \pi^* \mathcal{F}_1 \otimes \mathcal{O}(a_1, b_1) \otimes \mathcal{O}_E(-E)[1] \rightarrow \mathbf{L}j^* \circ j_*(p^* \circ \pi^* \mathcal{F}_1 \otimes \mathcal{O}(a_1, b_1)) \rightarrow p^* \circ \pi^* \mathcal{F}_1 \otimes \mathcal{O}(a_1, b_1) \xrightarrow{+1}$$

Since  $\mathcal{O}_E(E) = \mathcal{O}(-1, -1)$ , this triangle reduces to

$$(41)$$

$$p^* \circ \pi^* \mathcal{F}_1 \otimes \mathcal{O}(a_1+1, b_1+1)[1] \rightarrow \mathbf{L}j^* \circ j_* (p^* \pi^* \mathcal{F}_1 \otimes \mathcal{O}(a_1, b_1)) \rightarrow p^* \circ \pi^* \mathcal{F}_1 \otimes \mathcal{O}(a_1, b_1) \xrightarrow{+1}$$

Applying the functor  $\mathbf{R}\mathrm{Hom}(-, p^* \circ \pi^* \mathcal{F}_2 \otimes \mathcal{O}(a_2, b_2))$ , we see that (the right-hand side of) (39) is the cone of the morphism

(42)

$$\mathbf{R}\mathrm{Hom}(p^* \circ \pi^* \mathcal{F}_1 \otimes \mathcal{O}(a_1+1, b_1+1)[2], p^* \circ \pi^* \mathcal{F}_2 \otimes \mathcal{O}(a_2, b_2)) \xrightarrow{f} \mathbf{R}\mathrm{Hom}(p^* \circ \pi^* \mathcal{F}_1 \otimes \mathcal{O}(a_1, b_1), p^* \circ \pi^* \mathcal{F}_2 \otimes \mathcal{O}(a_2, b_2))$$

or equivalently the cone of

(43)

$$\mathbf{R}\mathrm{Hom}(p^* \circ \pi^* \mathcal{F}_1[2], p^* \circ \pi^* \mathcal{F}_2 \otimes \mathcal{O}(a_2-a_1-1, b_2-b_1-1)) \xrightarrow{f} \mathbf{R}\mathrm{Hom}(p^* \circ \pi^* \mathcal{F}_1, p^* \circ \pi^* \mathcal{F}_2 \otimes \mathcal{O}(a_2-a_1, b_2-b_1)).$$

Now note that for all  $a, b \in \mathbb{Z}$ ,

$$\begin{aligned} & \mathbf{R}\mathrm{Hom}(p^* \circ \pi^* \mathcal{F}_1, p^* \circ \pi^* \mathcal{F}_2 \otimes \mathcal{O}(a, b)) \\ &= \mathbf{R}\mathrm{Hom}(p^* \circ \pi^* \mathcal{F}_1, p^* \circ \pi^* \mathcal{F}_2 \otimes p^* \mathcal{O}_\pi(a) \otimes p'^* \mathcal{O}_{\pi'}(b)) \\ (44) \quad &= \mathbf{R}\mathrm{Hom}(\pi^* \mathcal{F}_1, \pi^* \mathcal{F}_2 \otimes \mathcal{O}_\pi(a) \otimes \mathbf{R}p_* \circ p'^* \mathcal{O}_{\pi'}(b)) \\ &= \mathbf{R}\mathrm{Hom}(\pi^* \mathcal{F}_1, \pi^* \mathcal{F}_2 \otimes \mathcal{O}_\pi(a) \otimes \pi^* \circ \mathbf{R}\pi'_* \mathcal{O}_{\pi'}(b)) \\ &= \mathbf{R}\mathrm{Hom}(\mathcal{F}_1, \mathcal{F}_2 \otimes \mathbf{R}\pi_* \mathcal{O}_\pi(a) \otimes \mathbf{R}\pi'_* \mathcal{O}_{\pi'}(b)), \end{aligned}$$

which vanishes when  $a \in [-k, -1]$  or  $b \in [-\ell, -1]$ . Therefore, in each of the cases in the statement, both the source and the target of  $f$  vanish, hence also the cone.  $\square$

### 3.1 Semiorthogonality

We will first prove that the subcategories in (5) are semiorthogonal. As e.g. in the proof of Orlov's blowup formula, this falls apart into two statements:

- (i) the semiorthogonality of the subcategories  $\Phi_{-k+\ell}(\mathbf{D}^b(F)), \dots, \Phi_{-1}(\mathbf{D}^b(F))$ ;
- (ii) the semiorthogonality of the pair  $\Phi_m(\mathbf{D}^b(F)), \mathbf{R}\tau_* \circ \mathbf{L}\tau'^* \mathbf{D}^b(X')$  for  $m = -k+\ell, \dots, -1$ .

Let us first prove (i).

**Proposition 8.** If  $m' < m$  are two integers such that  $m - m' < k - \ell$ , then for all  $\mathcal{E}, \mathcal{F} \in \mathbf{D}^b(F)$ , we have

$$(45) \quad \mathbf{R}\mathrm{Hom}_X(\Phi_m(\mathcal{E}), \Phi_{m'}(\mathcal{F})) = 0.$$

*Proof.* It suffices to prove this for  $\mathcal{E}$  and  $\mathcal{F}$  coherent sheaves, by the standard dévissage argument for bounded complexes of coherent sheaves. By adjunction,

$$(46) \quad \begin{aligned} \mathbf{R}\mathrm{Hom}(\Phi_m(\mathcal{E}), \Phi_{m'}(\mathcal{F})) &\cong \mathbf{R}\mathrm{Hom}(i_* (\pi^* \mathcal{E} \otimes \mathcal{O}_\pi(m)), i_* (\pi^* \mathcal{F} \otimes \mathcal{O}_\pi(m'))) \\ &\cong \mathbf{R}\mathrm{Hom}\left(\pi^* \mathcal{E} \otimes \mathcal{O}_\pi(m), i^! \circ i_* (\pi^* \mathcal{F} \otimes \mathcal{O}_\pi(m'))\right). \end{aligned}$$

To show this space vanishes, it suffices to show that

$$(47) \quad \mathbf{R}\mathrm{Hom}\left(\pi^* \mathcal{E} \otimes \mathcal{O}_\pi(m), \mathcal{H}^q(i^! \circ i_* (\pi^* \mathcal{F} \otimes \mathcal{O}_\pi(m')))[*]\right) = 0,$$

for all  $q \in \mathbb{Z}$ . By [1, lemma 1.4], the Fourier–Mukai kernel of  $i^! \circ i_* : \mathbf{D}^b(Z) \rightarrow \mathbf{D}^b(Z)$  has cohomology sheaves  $\mathcal{H}^q = \Delta_{Z,*}(\wedge^q \mathcal{N})$ , where  $\mathcal{N} := \mathbf{N}_{Z/X} \cong \mathcal{O}_\pi(-1) \otimes \pi^*(\mathcal{V}')$  for some vector bundle  $\mathcal{V}'$  on  $F$  of rank  $\ell + 1$  by the standing hypothesis. We then find:

$$\begin{aligned}
& \mathrm{RHom}\left(\pi^* \mathcal{E} \otimes \mathcal{O}_\pi(m), \mathcal{H}^q(i^! \circ i_*(\pi^* \mathcal{F} \otimes \mathcal{O}_\pi(m'))[*])\right) \\
(48) \quad & \cong \mathrm{RHom}\left(\pi^* \mathcal{E}, \pi^* \mathcal{F} \otimes \mathcal{O}_\pi(m' - m) \otimes \wedge^q \mathcal{N}[*]\right) \\
& \cong \mathrm{RHom}\left(\pi^* \mathcal{E}, \pi^* \mathcal{F} \otimes \mathcal{O}_\pi(m' - m) \otimes \mathcal{O}_\pi(-q) \otimes \wedge^q \pi^*(\mathcal{V}')[*]\right) \\
& \cong \mathrm{RHom}\left(\pi^* \mathcal{E}, \pi^*(\mathcal{F} \otimes \wedge^q \mathcal{V}') \otimes \mathcal{O}_\pi(m' - m - q)[*]\right)
\end{aligned}$$

As  $\mathrm{rk} \mathcal{V}' = \ell + 1$ , this space is zero except possibly for  $0 \leq q \leq \ell + 1$ . Combined with the assumption that  $m' - m \in [\ell - k + 1, -1]$ , we see that  $m' - m - q \in [-k, -1]$ .

Now  $\pi^* \mathcal{E} \in \pi^* \mathbf{D}^b(F)$  whilst  $\pi^*(\mathcal{F} \otimes \wedge^q \mathcal{V}') \otimes \mathcal{O}_\pi(m' - m - q) \in \pi^* \mathbf{D}^b(F) \otimes \mathcal{O}_\pi(m' - m - q)$ , so using the semiorthogonality in Orlov’s projective bundle formula, we deduce that (47) vanishes, and the proposition follows.  $\square$

Let us show the remaining semiorthogonality. We use an argument from the original proof of theorem 1 of Bondal–Orlov from [12, theorem 3.6]. For the convenience of the reader, we give a complete proof.

**Proposition 9.** For all integers  $m \in [-k + \ell, -1]$ , and for all  $\mathcal{F} \in \mathbf{D}^b(X')$  and  $\mathcal{G} \in \mathbf{D}^b(F)$ , we have

$$(49) \quad \mathrm{RHom}_X(\mathbf{R}\tau_* \circ \mathbf{L}\tau'^*(\mathcal{F}), \Phi_m(\mathcal{G})) = 0.$$

*Proof.* By lemma 6, we are to show that  $\mathrm{RHom}_X(\mathbf{R}\tau_* \circ \mathbf{L}\tau'^* \mathbf{D}^b(X'), \mathbf{R}\tau_* \mathcal{A}(m, 0)) = 0$ . By adjunction, it is enough to show that

$$(50) \quad \mathrm{RHom}_{\tilde{X}}(\mathbf{L}\tau^* \circ \mathbf{R}\tau_* \circ \mathbf{L}\tau'^* \mathbf{D}^b(X'), \mathcal{A}(m, 0)) = 0.$$

Take any  $\mathcal{E} \in \mathbf{L}\tau'^* \mathbf{D}^b(X')$ , viewed as an object in  $\mathbf{D}^b(\tilde{X})$ . Then  $\mathbf{L}\tau^* \circ \mathbf{R}\tau_*(\mathcal{E})$  is its right projection to the component  $\mathbf{L}\tau^* \mathbf{D}^b(X)$  with respect to the semiorthogonal decomposition (37).

Using (36), we will use the following version of (37):


$$\begin{aligned}
(51) \quad \mathbf{D}^b(\tilde{X}) = & \langle \mathcal{A}(-k, -\ell), \mathcal{A}(-k + 1, -\ell), \dots, \mathcal{A}(-1, -\ell), \mathcal{A}(0, -\ell); \\
& \mathcal{A}(-k + 1, -\ell + 1), \dots, \mathcal{A}(0, -\ell + 1), \mathcal{A}(1, -\ell + 1); \\
& \dots; \\
& \mathcal{A}(-k + \ell - 1, -1), \mathcal{A}(-k + \ell, -1), \dots, \mathcal{A}(\ell - 1, -1); \\
& \mathbf{L}\tau^* \mathbf{D}^b(X) \rangle.
\end{aligned}$$

In figure 1a we make the order of terms  $\mathcal{A}(i, j)$  explicit in the case of  $k = 6$  and  $\ell = 4$ .


The virtue of this decomposition is that, thanks to lemma 7, among all the components appearing above, there are no Hom’s from  $\mathcal{A}(a, b)$  to  $\mathcal{A}(a', b')$  with  $a \geq 0$  and  $a' < 0$ . Therefore, we can rearrange the order of the components to obtain the following semiorthogonal decomposition

$$(52) \quad \mathbf{D}^b(\tilde{X}) = \langle \mathcal{D}_1, \mathcal{D}_2, \mathbf{L}\tau^* \mathbf{D}^b(X) \rangle,$$

Figure 1: Visualising the mutation of the semiorthogonal decomposition in (51), for  $k = 6$  and  $\ell = 4$ .



(a) Before the mutation



(b) After the mutation, with the grouping according to (53)

where

$$\begin{aligned}
 \mathcal{D}_1 &:= \langle \mathcal{A}(-k, -\ell), \dots, \mathcal{A}(-1, -\ell); \mathcal{A}(-k+1, -\ell+1), \dots, \mathcal{A}(-1, -\ell+1); \\
 (53) \quad &\dots; \mathcal{A}(-k+\ell-1, -1), \dots, \mathcal{A}(-1, -1) \rangle, \\
 \mathcal{D}_2 &:= \langle \mathcal{A}(0, -\ell); \mathcal{A}(0, -\ell+1), \mathcal{A}(1, -\ell+1); \dots; \mathcal{A}(0, -1), \dots, \mathcal{A}(\ell-1, -1) \rangle.
 \end{aligned}$$

See figure 1b for the order of the terms  $\mathcal{A}(i, j)$  after the mutation for the case  $k = 6$  and  $\ell = 4$ .

Since  $\mathcal{D}_1 \subset \langle \mathcal{A}(-k, \star), \dots, \mathcal{A}(-1, \star) \rangle$ , using the semiorthogonality of (38), we see that  $\mathcal{E} \in {}^\perp \mathcal{D}_1 = \langle \mathcal{D}_2, \mathbf{L}\tau^* \mathbf{D}^b(X) \rangle$ .

We deduce that there is an object  $\mathcal{E}' \in \mathcal{D}_2$  fitting into the following distinguished triangle

$$(54) \quad \mathbf{L}\tau^* \circ \mathbf{R}\tau_* \mathcal{E} \rightarrow \mathcal{E} \rightarrow \mathcal{E}' \xrightarrow{+1}.$$

Therefore, to show that  $\mathbf{R}\mathrm{Hom}_{\tilde{\mathcal{X}}}(\mathbf{L}\tau^* \circ \mathbf{R}\tau_* \mathcal{E}, \mathcal{A}(m, 0)) = 0$  for all  $m \in [-k + \ell, -1]$ , it suffices to show that there is no morphism from  $\mathcal{E}$  or  $\mathcal{E}'$  to  $\mathcal{A}(m, 0)$ .

For the vanishing of  $\mathrm{RHom}_X(\mathcal{E}, \mathcal{A}(m, 0))$ , it suffices to apply the semiorthogonality from (38).

For the second vanishing, since  $\mathcal{E}' \in \mathcal{D}_2$ , to see that  $\mathrm{RHom}_X(\mathcal{E}', \mathcal{A}(m, 0)) = 0$ , it is enough to check that

$$(55) \quad \mathrm{RHom}_X(\mathcal{A}(a, b), \mathcal{A}(m, 0)) = 0$$

for any  $(a, b)$  with  $\ell - 1 \geq a \geq 0$  and  $-1 \geq b \geq a - \ell$ . But this holds by lemma 7, since  $a - m \leq \ell - 1 - (-k + \ell) = k - 1$ .

The vanishing (50) is proved and so is the proposition.  $\square$

**Remark 10** (Bondal–Orlov’s fully faithfulness). In the literature the proof of theorem 1 is given only in the case that  $F$  is a point (see e.g. [17, proposition 11.23], or the original [12, theorem 3.6]), but fully faithfulness of the functor  $\mathbf{R}\tau_* \circ \mathbf{L}\tau'^*$  in the general case follows with hardly any extra work. Indeed, by adjunction and the fully faithfulness of  $\mathbf{L}\tau'^*$ , we are reduced to show that for all  $\mathcal{E}, \mathcal{F} \in \mathbf{L}\tau'^* \mathbf{D}^b(X')$ , the natural map  $\mathrm{Hom}(\mathcal{E}, \mathcal{F}) \rightarrow \mathrm{Hom}(\mathbf{L}\tau^* \circ \mathbf{R}\tau_* \mathcal{E}, \mathcal{F})$  is an isomorphism.

Keeping the notation from the proof of proposition 9, it is enough to show that  $\mathrm{Hom}(\mathcal{E}', \mathcal{F}) = 0$ . By adjunction, this amounts to the vanishing  $\mathbf{R}\tau'_*(\mathcal{A}(a, b) \otimes \omega_{\tau'}^\vee) = 0$  for all  $(a, b)$  with  $\ell - 1 \geq a \geq 0$  and  $-1 \geq b \geq a - \ell$ , which follows again from lemma 7. We stress that this is the original proof of Bondal–Orlov.

### 3.2 Generation

We follow Thomas’ presentation in [41] of Kuznetsov’s argument for “mutation-cancellation”, which is initially in the context of homological projective duality [31]. This will be used to show that the sequence of subcategories in (5), shown to be semiorthogonal in the previous section, moreover generate the category  $\mathbf{D}^b(X)$ . The main result of this section is the following.

**Proposition 11.** For all  $m \in [-k, -k + \ell - 1]$ , the subcategory  $\mathcal{A}(m, 0)$  is in the triangulated subcategory of  $\mathbf{D}^b(\tilde{X})$  generated by the following subcategories:

$$(56) \quad \mathcal{A}(\star, -\ell), \dots, \mathcal{A}(\star, -1); \mathcal{A}(-k + \ell, 0), \dots, \mathcal{A}(-1, 0), \mathbf{L}\tau'^* \mathbf{D}^b(X').$$

Assuming this proposition, let us deduce the generation part in theorem A(ii).

**Corollary 12.** The triangulated category  $\mathbf{D}^b(X)$  is generated by the subcategories

$$(57) \quad \Phi_{-k+\ell}(\mathbf{D}^b(F)), \dots, \Phi_{-1}(\mathbf{D}^b(F)), \mathbf{R}\tau_* \circ \mathbf{L}\tau'^*(\mathbf{D}^b(X'))$$

*Proof.* Let  $\mathcal{C} \subset \mathbf{D}^b(\tilde{X})$  be the triangulated subcategory generated by the subcategories in (56). By definition,  $\mathcal{C}$  contains  $\mathcal{A}(a, b)$  for all  $a \in [-k, -1]$  and  $b \in [-\ell, -1]$ .

Thanks to proposition 11,  $\mathcal{C}$  also contains  $\mathcal{A}(-k, 0), \dots, \mathcal{A}(-1, 0)$ . Therefore,  $\mathcal{C}$  contains  $\mathcal{A}(a, b)$  for all  $a \in [-k, -1]$  and  $b \in [-\ell, 0]$ . By (35) (with  $m = 0$ ),  $\mathcal{C}$  contains  $\mathcal{A}(a, \star)$  for all  $a \in [-k, -1]$ . By (38), we deduce that in fact

$$(58) \quad \mathcal{C} = \mathbf{D}^b(\tilde{X}),$$

i.e.  $\mathbf{D}^b(\tilde{X})$  is generated by the subcategories in (56). Applying the projection functor  $\mathbf{R}\tau_*$ , one finds that  $\mathbf{D}^b(X)$  is generated by the subcategories

$$(59) \quad \mathbf{R}\tau_*\mathcal{A}(\star, -l), \dots, \mathbf{R}\tau_*\mathcal{A}(\star, -1); \mathbf{R}\tau_*\mathcal{A}(-k+l, 0), \dots, \mathbf{R}\tau_*\mathcal{A}(-1, 0), \mathbf{R}\tau_*\circ\mathbf{L}\tau'^* \mathbf{D}^b(X').$$

However, the first  $\ell$  terms are zero by (37), and the next  $k-\ell$  terms are in fact  $\Phi_{-k+\ell}(\mathbf{D}^b(F)), \dots, \Phi_{-1}(\mathbf{D}^b(F))$  by lemma 6. We get the desired generation.  $\square$

*Proof of proposition 11.* Let  $\mathcal{C} \subset \mathbf{D}^b(\tilde{X})$  be the triangulated subcategory generated by (56).

We proceed by descending induction on  $m$ . The induction hypothesis is that

$$(60) \quad \mathcal{A}(m+1, 0), \dots, \mathcal{A}(-1, 0) \subset \mathcal{C}.$$

For all  $a \in [m+1, -1]$ , we have that

$$(61) \quad \mathcal{A}(a, -\ell), \dots, \mathcal{A}(a, -1) \subset \mathcal{C}.$$

Combined with the induction hypothesis that  $\mathcal{A}(a, 0) \subset \mathcal{C}$  and with (35), we have

$$(62) \quad \mathcal{A}(a, \star) \subset \mathcal{C} \text{ for all } a \in [m+1, -1].$$

Recall that similar to remark 2, but now for the classical case of Orlov's blowup formula, we have also the semiorthogonal decomposition

$$(63) \quad \mathbf{D}^b(\tilde{X}) = \langle \mathcal{A}(m+1, \star), \dots, \mathcal{A}(-1, \star), \mathbf{L}\tau'^* \mathbf{D}^b(X'), \mathcal{A}(0, \star), \dots, \mathcal{A}(m+k, \star) \rangle.$$

Now for any object  $\mathcal{E} \in \mathcal{A}(m, 0)$ , there is a distinguished triangle

$$(64) \quad \mathcal{E}' \rightarrow \mathcal{E} \rightarrow \mathcal{E}'' \xrightarrow{+1},$$

with

$$(65) \quad \mathcal{E}' \in \langle \mathcal{A}(0, \star), \dots, \mathcal{A}(m+k, \star) \rangle$$

$$(66) \quad \mathcal{E}'' \in \langle \mathcal{A}(m+1, \star), \dots, \mathcal{A}(-1, \star), \mathbf{L}\tau'^* \mathbf{D}^b(X') \rangle.$$

By (62),  $\mathcal{E}'' \in \mathcal{C}$ . Hence to show that  $\mathcal{E} \in \mathcal{C}$ , it suffices to show that  $\mathcal{E}' \in \mathcal{C}$ .

We expand (65) using the projective bundle formula (35) as

$$(67) \quad \mathcal{E}' \in \langle \mathcal{A}(0, -\ell-1), \dots, \mathcal{A}(0, -1); \mathcal{A}(1, -\ell-1), \dots, \mathcal{A}(1, -1); \dots; \mathcal{A}(m+k, -\ell-1), \dots, \mathcal{A}(m+k, -1) \rangle.$$

For any  $(a, b)$  with  $a \in [0, m+k]$  and  $b \in [-\ell-1, -1]$ , let us denote by  $\mathcal{E}(a, b)$  the projection of  $\mathcal{E}'$  in  $\mathcal{A}(a, b)$ . We will show by induction that

$$(68) \quad \mathcal{E}(a, b) = 0 \text{ for any } a \in [0, m+k] \text{ and } b \in [-\ell-1, -\ell-1+a].$$

The induction is with the descending order on  $a$  and, for fixed  $a$ , with ascending order on  $b$ . Assuming that  $\mathcal{E}(a', b') = 0$  for all  $m+k \geq a' > a$ , or  $a' = a$  and  $-\ell-1 \leq b' < b$ , let us show  $\mathcal{E}(a, b) = 0$ . We have that

$$(69) \quad \mathbf{R}\mathrm{Hom}_X(\mathcal{E}'', \mathcal{A}(a-k, b+1)) = 0.$$

Indeed, on one hand,  $\mathcal{E}'' \in \langle \mathcal{A}(m+1, \star), \dots, \mathcal{A}(-1, \star), \mathbf{L}\tau^{*} \mathbf{D}^b(X') \rangle$ ; on the other hand,  $\mathcal{A}(a-k, b+1) \in \langle \mathcal{A}(-k, \star), \dots, \mathcal{A}(m, \star) \rangle$ , since  $a-k \in [-k, m]$ . The vanishing follows from the semiorthogonality in (38).

The second required vanishing result is that

$$(70) \quad \mathbf{RHom}_X(\mathcal{E}, \mathcal{A}(a-k, b+1)) = 0.$$

Indeed,  $\mathcal{E} \in \mathcal{A}(m, 0)$  and the vanishing follows from lemma 7. To see this, observe that we always have  $a-k \leq m$ . If  $a-k < m$ , it is fine by lemma 7. If  $a-k = m$ , then  $b+1 \in [-\ell, -\ell + m + k] \subset [-\ell, -1]$ , which is again covered by lemma 7.

Consequently,  $\mathbf{RHom}(\mathcal{E}', \mathcal{A}(a-k, b+1)) = 0$ . However, by lemma 7, a non-zero morphism from  $\mathcal{E}(a', b')$  to  $\mathcal{A}(a-k, b+1)$  only exists when  $a' > a$ , or  $a' = a$  and  $b' \leq b$ . But by the induction hypothesis,  $\mathcal{E}(a', b') = 0$  for all  $a' > a$ , or  $a' = a$  and  $b' < b$ . Therefore, we deduce that

$$(71) \quad \mathbf{RHom}_X(\mathcal{E}(a, b), \mathcal{A}(a-k, b+1)) = 0.$$

Now write  $\mathcal{E}(a, b) = j_*(p^* \circ \pi^* \mathcal{F} \otimes \mathcal{O}(a, b))$  for some  $\mathcal{F} \in \mathbf{D}^b(F)$ . We define

$$(72) \quad \mathcal{G} := j_*(p^* \circ \pi^* \mathcal{F} \otimes \mathcal{O}(a-k, b+1)[k+1]),$$

which is in  $\mathcal{A}(a-k, b+1)$ . Making use of the triangle (40) and

$$(73) \quad \mathbf{RHom}_E(p^* \circ \pi^* \mathcal{F} \otimes \mathcal{O}(a, b), p^* \circ \pi^* \mathcal{F} \otimes \mathcal{O}(a-k, b+1)[k+1]) = 0,$$

we find

$$(74) \quad \begin{aligned} \mathbf{RHom}(\mathcal{E}(a, b), \mathcal{G}) &\cong \mathbf{RHom}_E(\mathbf{L}j^* \circ j_*(p^* \circ \pi^* \mathcal{F} \otimes \mathcal{O}(a, b)), p^* \circ \pi^* \mathcal{F} \otimes \mathcal{O}(a-k, b+1)[k+1]) \\ &\cong \mathbf{RHom}_E(p^* \circ \pi^* \mathcal{F} \otimes \mathcal{O}(a+1, b+1)[1], p^* \circ \pi^* \mathcal{F} \otimes \mathcal{O}(a-k, b+1)[k+1]) \end{aligned}$$

which contains the element  $\psi := \text{id}_{p^* \circ \pi^* \mathcal{F}} \otimes p^*(\alpha) \otimes p^{*k}(\text{id}_{\mathcal{O}_{\pi^{-1}(b+1)}})$ , where  $\alpha$  is a non-zero element in

$$(75) \quad \begin{aligned} \mathbf{RHom}(\mathcal{O}_\pi(a+1), \mathcal{O}_\pi(a-k)[k]) &\cong \mathbf{H}^k(Z, \mathcal{O}_\pi(-k-1)) \\ &\cong \mathbf{H}^0(F, \mathbf{R}^k \pi_* \mathcal{O}_\pi(-k-1)) \\ &\cong \mathbf{k}. \end{aligned}$$

Then (71) implies that  $\psi$  must be zero. This is only possible if  $p^* \circ \pi^* \mathcal{F}$  is itself zero. Hence  $\mathcal{E}(a, b) = 0$ . The induction for (68) is complete.

In particular, the components  $\mathcal{E}(a, -\ell-1)$  of  $\mathcal{E}'$  vanish for  $a \in [0, m+k]$ . Therefore  $\mathcal{E}' \in \langle \mathcal{A}(\star, -\ell), \dots, \mathcal{A}(\star, -1) \rangle$ , hence is contained in  $\mathcal{C}$ . As we see above, this implies  $\mathcal{E} \in \mathcal{C}$ . In other words,  $\mathcal{A}(m, 0) \subset \mathcal{C}$ . The induction on  $m$  is complete and the proposition is proved.  $\square$

## 4 Application: quadratic Fano correspondence for cubic hypersurfaces

In this section we show there is a standard flip diagram associated to a smooth cubic hypersurface, so that theorem B follows from theorem A. In corollary 18, we use this



result and [20, theorem 3.4] to deduce a corresponding isomorphism of Chow motives, generalizing [32, theorem 5].

We now recall the setup for the *quadratic Fano correspondence* for cubic hypersurfaces. Let  $n \geq 2$ , and let  $Y \subset \mathbb{P}^{n+1}$  be a smooth cubic hypersurface. Denote  $F := F(Y)$  the Fano variety of lines on  $Y$  and  $Y^{[2]}$  the Hilbert scheme of two points on  $Y$ , which will be discussed more explicitly in section 5. Galkin–Shinder [15, §5] and Voisin [43] established the existence of the following diagram:

$$(76) \quad \begin{array}{ccccc} & & E & & \\ & & \downarrow j & & \\ & & D & & \\ & p \swarrow & & \searrow p' & \\ P_2 & \xrightarrow{i} & Y^{[2]} & \xrightarrow{\phi} & P_Y & \xleftarrow{i'} & P \\ & \searrow \pi & & \swarrow \tau' & & \swarrow \pi' & \\ & & F & & & & \end{array}$$

In this diagram, we use the following notation:

- $\pi' : P \rightarrow F$  is the universal  $\mathbb{P}^1$ -bundle;
- $\pi : P_2 := P^{[2]/F} \rightarrow F$  is the relative Hilbert square of  $\pi'$ , which is a  $\mathbb{P}^2$ -bundle;
- $P_Y := \mathbb{P}(\mathbb{T}_{\mathbb{P}^{n+1}}|_Y) \rightarrow Y$  is the projectivization of the restriction to  $Y$  of the tangent bundle of the projective space  $\mathbb{P}^{n+1}$ , which is a  $\mathbb{P}^n$ -bundle over  $Y$ , and  $P_Y$  parametrises a point on  $Y$  together with a line in  $\mathbb{P}^{n+1}$  passing through it;
- $D := \{(L, z, x) \in \text{Gr}(2, n+2) \times Y^{[2]} \times Y \mid z \subset L, x \in L, \text{ and if } L \not\subset Y, z+x = Y \cap L\}$ , where  $\text{Gr}(2, n+2) = \text{Gr}(\mathbb{P}^1, \mathbb{P}^{n+1})$ ;
- given a point  $(L, z, x)$  of  $D$ , the morphism  $\tau$  maps it to  $z \in Y^{[2]}$  while the morphism  $\tau'$  maps it to  $(L, x) \in P_Y$ ;
- $E := \{(L, z, x) \in D \mid L \subset Y\}$  is a divisor in  $D$ ;
- $\phi$  is the Galkin–Shinder map [15, (5.3)] which sends  $z \in Y^{[2]}$  to  $(L, x) \in P_Y$ , where  $L$  is the line determined (i.e. spanned) by  $z$ , and  $x$  is the residual intersection point of  $L$  with  $Y$ , i.e.  $L \cap Y = z + x$ , when  $L$  is not completely contained in  $Y$ ;
- by Voisin [43, proposition 2.9] the upper two trapezoids are blowup diagrams, and in particular all varieties appearing above are smooth and projective;
- the outer square is cartesian, and observe that  $\pi$  and  $p'$  are  $\mathbb{P}^2$ -bundles while  $\pi'$  and  $p$  are  $\mathbb{P}^1$ -bundles.

Denote by  $\mathcal{S}$  the restriction of the tautological rank-2 bundle over the Grassmannian  $\text{Gr}(2, n+2)$  to  $F$ . Note that since  $\mathcal{S}$  is of rank 2, there is a canonical  $F$ -isomorphism  $\mathbb{P}(\mathcal{S}) \simeq \mathbb{P}(\mathcal{S}^\vee)$ .

In corollary 15 we will show that this diagram is in fact a standard flip diagram, as in (2).

**Lemma 13.** The natural morphism  $i : P_2 \rightarrow Y^{[2]}$  is a closed immersion.

*Proof.* Let us describe more carefully the morphism  $i$ . Let  $[L] \in F$  be a point representing (by abuse of notation) a line  $L \subset Y$ . Then for two distinct points  $x, y \in L$ , the morphism  $i$  sends  $x + y$  to the length-2 subscheme  $\{x, y\} \in Y^{[2]}$ ; while for the double point  $2x \in L^{[2]}$ , it sends it to the length-2 subscheme of  $Y$  supported on  $x$  with the tangent direction given by the 1-dimensional subspace  $T_x L \subset T_x Y$ .

Now observe that if  $L_1 \neq L_2$  are two distinct points in  $F$ , then  $L_1^{[2]} \cap L_2^{[2]} = \emptyset$ . To see this, we discuss the two possibilities:

- if  $L_1 \cap L_2 = \emptyset$ , then this is immediate;
- if  $L_1 \cap L_2 = \{z\}$ , then points on  $L_1^{[2]}$  and  $L_2^{[2]}$  whose support contains  $z$  are distinguished by the support of the second point, or the tangent direction.

□

**Lemma 14.** We have the following descriptions for the normal bundles of  $i$  and  $i'$ :

- (i)  $N_{P_2/Y^{[2]}} \cong \mathcal{O}_\pi(-1) \otimes \pi^*(\mathcal{S})$ .
- (ii)  $N_{P/P_Y} \cong \mathcal{O}_{\pi'}(-1) \otimes \pi'^*(\text{Sym}^2 \mathcal{S})$ .

*Proof.* We will use the construction of Voisin in the proof of [43, Proposition 2.9], which we recall now. Denote  $\text{Gr} := \text{Gr}(\mathbb{P}^1, \mathbb{P}^{n+1})$ . Let  $\sigma: Q \rightarrow \text{Gr}$  be the universal  $\mathbb{P}^1$ -bundle associated to the tautological rank-2 vector bundle.

For (i), pulling back the  $\mathbb{P}^1$ -bundle using the natural morphism  $Y^{[2]} \rightarrow \text{Gr}$ , we obtain the morphism  $\sigma_1: Q_1 \rightarrow Y^{[2]}$ . Denote the evaluation morphism  $e_1: Q_1 \rightarrow \mathbb{P}$ . Then the preimage  $e_1^{-1}(Y)$  is a divisor in  $Q_1$  of degree 3 over  $Y^{[2]}$ , with one component  $D_1$  being the universal subscheme  $D_1$  and the other component exactly  $D$ . We can summarise the notation as

$$(77) \quad \begin{array}{ccccc} & & Y & \hookrightarrow & \mathbb{P} \\ & & \uparrow & & \uparrow e_1 \\ D_1 \cup D & \longleftarrow & e_1^{-1}(Y) & \hookrightarrow & Q_1 & \longrightarrow & Q \\ & & & & \downarrow \sigma_1 & & \downarrow \sigma \\ & & & & Y^{[2]} & \longrightarrow & \text{Gr} \end{array} .$$

Then  $P_2$  is the zero set of a regular section of the rank-2 vector bundle  $\sigma_{1,*}(\mathcal{O}_{\sigma_1}(3) \otimes \mathcal{O}_{Q_1}(-D_1))$ . Therefore the normal bundle of  $P_2$  inside  $Y^{[2]}$  is its restriction to  $P_2$ , which is

$$(78) \quad \text{pr}_{1,*}(\text{pr}_2^* \mathcal{O}_\omega(3) \otimes \mathcal{O}(-\mathcal{U}))$$

where  $\mathcal{U}$  is the universal subscheme (i.e. the restriction of  $D_1$ ) in the diagram

$$(79) \quad \begin{array}{ccc} \mathcal{U} & \hookrightarrow & P_2 \times_F \mathbb{P}(\mathcal{S}^\vee) \xrightarrow{\text{pr}_2} \mathbb{P}(\mathcal{S}^\vee) \\ & \searrow & \downarrow \text{pr}_1 \qquad \qquad \downarrow \omega \\ & & P_2 \xrightarrow{\pi} F \end{array}$$

Let us compute this vector bundle (78).

First, for any  $[L] \in F$ , the bundle (78) restricted to  $\pi^{-1}(L) \cong L^{[2]}$  is given by

$$(80) \quad \text{pr}_{1,*} (\text{pr}_2^* \mathcal{O}_L(3) \otimes \mathcal{O}_{L^{[2]} \times L}(-U))$$

where  $U$  is the universal subscheme (the fiber of  $D_1$  over  $L$ ) in the diagram

$$(81) \quad \begin{array}{ccc} U & \hookrightarrow & L^{[2]} \times L \xrightarrow{\text{pr}_2} L \\ & \searrow & \downarrow \text{pr}_1 \\ & & L^{[2]} \end{array}$$

and  $U \cong L \times L$ , so that  $U \rightarrow L^{[2]}$  is the quotient by the involution and  $U \rightarrow L$  is the projection on the first factor. We obtain that

$$(82) \quad \mathcal{O}_{L^{[2]} \times L}(U) \cong \mathcal{O}_{\mathbb{P}^2 \times \mathbb{P}^1}(1, 2).$$

Indeed, in the Chow ring of  $\mathbb{P}^2 \times \mathbb{P}^1$  we compute that  $U \cdot (\{z\} \times L) = 2$  for  $z \in L^{[2]}$ , and we have that  $U|_{L^{[2]} \times \{x\}} \cong \mathcal{O}_{L^{[2]}}(1)$  for  $x \in L$ .

Next, globalizing the computation, we have that  $\mathcal{O}(\mathcal{U}) \cong \text{pr}_1^* \mathcal{O}_\pi(1) \otimes \text{pr}_2^* \mathcal{O}_\omega(2)$ . In fact, via the identification  $P_2 = \mathbb{P}(\text{Sym}^2 \mathcal{S}) \rightarrow F$ ,  $\mathcal{U}$  corresponds to the canonical element in

$$(83) \quad H^0(P_2 \times_F \mathbb{P}(\mathcal{S}^\vee), \text{pr}_1^* \mathcal{O}_\pi(1) \otimes \text{pr}_2^* \mathcal{O}_\omega(2)) \cong H^0(F, \text{Sym}^2 \mathcal{S}^\vee \otimes \text{Sym}^2 \mathcal{S}).$$

As a result, we can compute the normal bundle via the identification (78) as

$$(84) \quad \begin{aligned} \mathbf{N}_{P_2/Y^{[2]}} &\cong \text{pr}_{1,*} (\text{pr}_2^* \mathcal{O}_\omega(3) \otimes \mathcal{O}(-\mathcal{U})) \\ &\cong \text{pr}_{1,*} (\text{pr}_1^* \mathcal{O}_\pi(-1) \otimes \text{pr}_2^* \mathcal{O}_\omega(1)) \\ &\cong \mathcal{O}_\pi(-1) \otimes \text{pr}_{1,*} (\text{pr}_2^* \mathcal{O}_\omega(1)) \\ &\cong \mathcal{O}_\pi(-1) \otimes \pi^* (\omega_* \mathcal{O}_\omega(1)) \\ &\cong \mathcal{O}_\pi(-1) \otimes \pi^* \mathcal{S}. \end{aligned}$$

Similarly for (ii), the base change of  $\sigma : Q \rightarrow \text{Gr}$  by the natural morphism  $P_Y \rightarrow \text{Gr}$  is a  $\mathbb{P}^1$ -bundle  $\sigma_2 : Q_2 \rightarrow P_Y$ , which is endowed with a canonical section. Denote the evaluation morphism  $e_2 : Q_2 \rightarrow \mathbb{P}$ . Then the preimage  $e_2^{-1}(Y)$  is a divisor in  $Q_2$  of degree 3 over  $P_Y$ , with one component  $D_2$  being the image of the canonical section of  $\sigma_2$  and the other component exactly  $D$ . We can summarise the notation as

$$(85) \quad \begin{array}{ccccc} Y & \hookrightarrow & \mathbb{P} & & \\ \uparrow & & e_2 \uparrow & & \\ D_2 \cup D & \longleftarrow & e_2^{-1}(Y) & \hookrightarrow & Q_2 \longrightarrow Q \\ & & \downarrow \sigma_2 & & \downarrow \sigma \\ & & P_Y & \longrightarrow & \text{Gr} \end{array}$$

Then  $P$  is the zero set of a regular section of the rank-3 vector bundle  $\sigma_{2,*}(\mathcal{O}_{\sigma_2}(3) \otimes \mathcal{O}_{Q_2}(-D_2))$ . Therefore the normal bundle of  $P$  inside  $P_Y$  is its restriction to  $P$ , which is

$$(86) \quad \text{pr}_{1,*} (\text{pr}_2^* \mathcal{O}_\omega(3) \otimes \mathcal{O}(-\mathcal{J}))$$

where  $\mathcal{J}$  is the incidence subscheme (i.e. the restriction of  $D_2$ ) in the diagram

$$(87) \quad \begin{array}{ccccc} \mathcal{J} & \hookrightarrow & P \times_F \mathbb{P}(\mathcal{S}^\vee) & \xrightarrow{\text{pr}_2} & \mathbb{P}(\mathcal{S}^\vee) \\ & \searrow & \downarrow \text{pr}_1 & & \downarrow \omega \\ & & P & \xrightarrow{\pi'} & F \end{array}$$

As before we compute that  $\mathcal{O}(\mathcal{J}) \cong \text{pr}_1^* \mathcal{O}_{\pi'}(1) \otimes \text{pr}_2^* \mathcal{O}_\omega(1)$ . As a result, we can compute the normal bundle via the identification (86) as

$$(88) \quad \begin{aligned} N_{P/P_Y} &\cong \text{pr}_{1,*} (\text{pr}_2^* \mathcal{O}_\omega(3) \otimes \mathcal{O}(-\mathcal{J})) \\ &\cong \text{pr}_{1,*} (\text{pr}_1^* \mathcal{O}_{\pi'}(-1) \otimes \text{pr}_2^* \mathcal{O}_\omega(2)) \\ &\cong \mathcal{O}_{\pi'}(-1) \otimes \text{pr}_{1,*} (\text{pr}_2^* \mathcal{O}_\omega(2)) \\ &\cong \mathcal{O}_{\pi'}(-1) \otimes \pi'^* (\omega_* \mathcal{O}_\omega(2)) \\ &\cong \mathcal{O}_{\pi'}(-1) \otimes \pi'^* \text{Sym}^2 \mathcal{S}. \end{aligned}$$

and we are done.  $\square$

**Corollary 15.** The diagram (76) is a standard flip diagram.

*Proof.* Indeed, thanks to lemma 14, we take in the notation of (2),  $X := Y^{[2]}$ ,  $X' := P_Y$ ,  $Z := P_2$ ,  $Z' := P$ ,  $D := \tilde{X}$ ,  $k = 2$ ,  $\ell = 1$ ,  $\mathcal{V} = \text{Sym}^2 \mathcal{S}$  and  $\mathcal{V}' = \mathcal{S}$ .  $\square$

Hence we obtain the following.

*Proof of theorem B.* By corollary 15, we can apply theorem A to the diagram (76) to get semiorthogonal decompositions

$$(89) \quad \begin{aligned} \mathbf{D}^b(Y^{[2]}) &= \langle \Phi_{-1}(\mathbf{D}^b(F)), \mathbf{R}\tau_* \circ \mathbf{L}\tau'^* \mathbf{D}^b(P_Y) \rangle \\ &= \langle \mathbf{R}\tau_* \circ \mathbf{L}\tau'^* \mathbf{D}^b(P_Y), \Phi_0(\mathbf{D}^b(F)) \rangle \end{aligned}$$

where  $\Phi_m(-) = i_*(\pi^*(-) \otimes \mathcal{O}_\pi(m))$ :  $\mathbf{D}^b(F) \rightarrow \mathbf{D}^b(Y^{[2]})$  is the fully faithful functor from (4).

We can now further decompose the  $\mathbb{P}^n$ -bundle  $P_Y \rightarrow Y$  using the projective bundle formula, and conclude.  $\square$

**Remark 16** (Grothendieck ring of categories). The  $Y$ - $\mathbf{F}(Y)$ -relation from [15, theorem 5.1] is the equality

$$(90) \quad [Y^{[2]}] = [\mathbb{P}^n][Y] + \mathbb{L}^2[\mathbf{F}(Y)]$$

in the Grothendieck ring of varieties  $K_0(\text{Var}/\mathbf{k})$ , for a cubic hypersurface  $Y \hookrightarrow \mathbb{P}^{n+1}$ .

Taking the motivic measure in the Grothendieck ring of dg categories  $K_0(\text{dgCat}/\mathbf{k})$  from [11] gives the relation

$$(91) \quad [\mathbf{D}^b(Y^{[2]})] = (n+1)[\mathbf{D}^b(Y)] + [\mathbf{D}^b(\mathbf{F}(Y))],$$

which is suggestive of a semiorthogonal decomposition of the form (9), and theorem B indeed provides this.

**Remark 17** (An isomorphism of Chow motives). Using Jiang’s analysis of the Chow theory of a standard flip diagram in [20], we can obtain the following improvement of Laterveer’s result in [32, theorem 5] to integral coefficients. For a smooth scheme  $X$  over some ground field  $\mathbf{k}$ , we write  $\mathfrak{h}(X) = (X, \Delta_X)$  for the motive of  $X$  in the rigid tensor category of Chow motives over  $\mathbf{k}$ , see [33]. By  $\mathfrak{h}(X)(-i)$  we denote the motive  $\mathfrak{h}(X) \otimes \mathbb{L}^i$ , where  $\mathbb{L} = (\mathbb{P}^1, [\mathbb{P}^1 \times \{0\}])$  is the Lefschetz motive.

**Corollary 18.** For an arbitrary base field  $\mathbf{k}$ , there is an isomorphism of Chow motives (with integral coefficients) over  $\mathbf{k}$ :

$$(92) \quad \mathfrak{h}(F)(-2) \oplus \bigoplus_{i=0}^n \mathfrak{h}(Y)(-i) \xrightarrow{\cong} \mathfrak{h}(Y^{[2]}),$$

where the isomorphism is given by

$$(93) \quad \left( i_* \circ \pi^*, \bigoplus_{i=0}^n \tau_* \circ \tau'^* \circ \cdot c_1(\mathcal{O}_u(1))^i \circ u^* \right)$$

where  $u: P_Y \rightarrow Y$  the  $\mathbb{P}^n$ -bundle and the other notation is as in (76).

*Proof.* It suffices to combine corollary 15 with [20, corollary 3.8].  $\square$

## 5 Hilbert squares of low-degree hypersurfaces

Let  $X$  be a smooth projective variety of dimension  $n = \dim X$ . Then its Hilbert square (which is short-hand for the Hilbert scheme of length-2 subschemes)  $X^{[2]}$  is a smooth projective variety of dimension  $2n$  [13, theorem 3.0.1]. Throughout we will assume that  $n \geq 2$ , with the main focus being on  $n \geq 3$ . For the benefit of the reader we will recall its relevant properties.

It has the following explicit description:

$$(94) \quad \begin{array}{ccc} & X^{[2]} & \\ & q \uparrow & \\ \text{Bl}_\Delta(X \times X) & \longleftarrow & E \\ \tau \downarrow & \square & \downarrow \\ X \times X & \xleftarrow{\Delta} & X \end{array}$$

where the square is the cartesian blowup square, and  $q$  is the quotient by the  $\mathbb{Z}/2\mathbb{Z}$ -action on  $\text{Bl}_\Delta(X \times X)$ .

We can consider  $q$  as a double cover, and we need to describe it more explicitly. We have that

$$(95) \quad \text{Pic}(X^{[2]}) \cong \text{Pic}(X) \oplus \mathbb{Z}[\delta]$$

for the divisor  $\delta$  which is half of the divisor  $D$  on  $X^{[2]}$  parametrising non-reduced length-2 subschemes. The double cover is branched along  $D$ , and we have the follow-

ing picture

$$(96) \quad \begin{array}{ccccc} E & \hookrightarrow & 2E & \hookrightarrow & \mathrm{Bl}_\Delta(X \times X) \\ & \searrow \cong & \downarrow & \square & \downarrow q \\ & & D & \hookrightarrow & X^{[2]} \end{array}$$

where  $2E$  is the second infinitesimal neighbourhood of the exceptional divisor  $E$ .

By [3, lemma I.17.1] we have that the description as a double cover gives the identifications

$$(97) \quad \omega_{\mathrm{Bl}_\Delta(X \times X)} \cong q^*(\omega_{X^{[2]}} \otimes \mathcal{O}_{X^{[2]}}(\delta))$$

$$(98) \quad q^*(\mathcal{O}_{X^{[2]}}(\delta)) \cong \mathcal{O}_{\mathrm{Bl}_\Delta(X \times X)}(E)$$

whilst the description as a blowup gives the identification

$$(99) \quad \omega_{\mathrm{Bl}_\Delta(X \times X)} \cong \tau^*(\omega_{X \times X}) \otimes \mathcal{O}_{\mathrm{Bl}_\Delta(X \times X)}((n-1)E).$$

The Hilbert square is a resolution of singularities, via the Hilbert–Chow morphism  $X^{[2]} \rightarrow \mathrm{Sym}^2 X$ , sending a length-2 subscheme to the cycle it is supported on. The resolution is crepant if  $n = 2$ , but fails to be crepant for  $n \geq 3$ . To see this, observe that the Hilbert–Chow morphism fits in the commutative diagram

$$(100) \quad \begin{array}{ccc} \mathrm{Bl}_\Delta(X \times X) & \xrightarrow{\tau} & X \times X \\ q \downarrow & & \downarrow \\ X^{[2]} & \longrightarrow & \mathrm{Sym}^2 X \end{array}$$

which allows us to compare the canonical bundles. We conclude that it is crepant if and only if the multiplicity of the contribution of  $\delta$  in the double cover agrees with that of the exceptional divisor in the blowup, which happens if and only if  $n = 2$ .

We will now use this description of the Hilbert square to prove that in some cases the Hilbert square of a Fano variety is again Fano. This possibility was suggested in [38], and later used in [39, lemma 2.9], for  $X = \mathbb{P}^n$  where  $n \geq 3$ . We will amplify this result to also cover low-degree hypersurfaces.

## 5.1 Projective space

We will recall the setup and proof of [39, lemma 2.9]. We will use the notation of (94) for  $X = \mathbb{P}^n$ . Consider the morphism

$$(101) \quad \varphi: \mathrm{Bl}_\Delta(\mathbb{P}^n \times \mathbb{P}^n) \rightarrow \mathbb{P}^n \times \mathbb{P}^n \times \mathrm{Gr}(2, n+1)$$

which is the product of the blowup  $\tau$  and the morphism which sends a point on the blowup to the line in  $\mathbb{P}^n$  spanned by two distinct points or a point together with a tangent direction, which gives a point on the Grassmannian  $\mathrm{Gr}(2, n+1)$ . It is quasi-finite by construction (its fibres are either empty or singletons), hence by [40, tag 02LS] it is actually finite. By Zariski’s main theorem we can conclude it is a closed immersion.

We have that  $\text{Pic}(\mathbb{P}^n \times \mathbb{P}^n \times \text{Gr}(2, n+1)) \cong \mathbb{Z}^{\oplus 3}$  and we will denote

$$(102) \quad \mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^n \times \text{Gr}(2, n+1)}(i, j, k) := \mathcal{O}_{\mathbb{P}^n}(i) \boxtimes \mathcal{O}_{\mathbb{P}^n}(j) \boxtimes \mathcal{O}_{\text{Gr}(2, n+1)}(k),$$

here  $\mathcal{O}_{\text{Gr}(2, n+1)}(1)$  is the Plücker line bundle. The line bundles  $\mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^n \times \text{Gr}(2, n+1)}(i, j, k)$  are ample if and only if  $i, j, k \geq 1$ , and because  $\varphi$  is finite (so in particular affine) ampleness is preserved under pullback.

We have the identifications

$$(103) \quad \begin{aligned} \varphi^*(\mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^n \times \text{Gr}(2, n+1)}(1, 1, 0)) &\cong \tau^*(\mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^n}(1, 1)) \\ \varphi^*(\mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^n \times \text{Gr}(2, n+1)}(0, 0, 1)) &\cong \tau^*(\mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^n}(1, 1)) \otimes \mathcal{O}_{\text{Bl}_\Delta(\mathbb{P}^n \times \mathbb{P}^n)}(-E), \end{aligned}$$

where the second isomorphism is obtained by considering Plücker coordinates in terms of bilinear forms (so as sections of  $\mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^n}(1, 1)$ ) which are non-vanishing on the diagonal. We then obtain the following proposition, which is also [39, lemma 2.9].

**Proposition 19** (Sawin). Let  $n \geq 3$ . Then  $\mathbb{P}^{n[2]}$  is Fano.

*Proof.* By (97) and (98) we have the identification

$$(104) \quad \begin{aligned} q^*(\omega_{\mathbb{P}^{n[2]}}^\vee) &\cong \omega_{\text{Bl}_\Delta(\mathbb{P}^n \times \mathbb{P}^n)}^\vee \otimes q^*(\mathcal{O}_{\mathbb{P}^{n[2]}}(\delta)) \\ &\cong \omega_{\text{Bl}_\Delta(\mathbb{P}^n \times \mathbb{P}^n)}^\vee \otimes \mathcal{O}_{\text{Bl}_\Delta(\mathbb{P}^n \times \mathbb{P}^n)}(E). \end{aligned}$$

By (99) we can further rewrite this as

$$(105) \quad \cong \tau^*(\mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^n}(n+1, n+1)) \otimes \mathcal{O}_{\text{Bl}_\Delta(\mathbb{P}^n \times \mathbb{P}^n)}((2-n)E).$$

But using (103) we can realise this line bundle as

$$(106) \quad \cong \varphi^*(\mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^n \times \text{Gr}(2, n+1)}(3, 3, n-2))$$

and hence it is ample if and only if  $n \geq 3$ .

To conclude, by [40, tag 0B5V] we know that  $\omega_{\mathbb{P}^{n[2]}}^\vee$  is ample if and only if  $q^*(\omega_{\mathbb{P}^{n[2]}}^\vee)$  is ample, and we are done.  $\square$

## 5.2 Quadrics and cubic hypersurfaces

We will now bootstrap from the result in proposition 19 to show that low-degree hypersurfaces again give rise to Hilbert squares which are Fano.

We will extend the notation of (94) for  $X \hookrightarrow \mathbb{P}^{n+1}$  a hypersurface of degree  $d = 2, 3$  by decorating the morphisms involving  $\mathbb{P}^{n+1}$ . By functoriality of the Hilbert scheme construction we obtain the following diagram.

$$(107) \quad \begin{array}{ccccccc} & & X^{[2]} & \hookrightarrow & \mathbb{P}^{n+1[2]} & & \\ & & \uparrow q & & \uparrow q' & & \\ E & \hookrightarrow & \text{Bl}_\Delta(X \times X) & \xrightarrow{t} & \text{Bl}_\Delta(\mathbb{P}^{n+1} \times \mathbb{P}^{n+1}) & \longleftarrow & E' \\ & \downarrow & \tau \downarrow & & \downarrow \tau' & & \downarrow \\ X & \xrightarrow{\Delta} & X \times X & \hookrightarrow & \mathbb{P}^{n+1} \times \mathbb{P}^{n+1} & \xleftarrow{\Delta'} & \mathbb{P}^{n+1} \end{array} .$$

**Proposition 20.** Let  $n \geq 3$  and  $d = 1, 2, 3$ . Let  $X \subseteq \mathbb{P}^{n+1}$  be a hypersurface of degree  $d$ . Then  $X^{[2]}$  is Fano.

*Proof.* Observe that we have  $\text{Pic}(X \times X) \cong \text{Pic } X \oplus \text{Pic } X$ , and we will write  $\mathcal{O}_{X \times X}(i, j) := \mathcal{O}_X(i) \boxtimes \mathcal{O}_X(j)$ .

By the adjunction formula we have  $\omega_X \cong \mathcal{O}_X(-n-2+d)$ . As in the proof of proposition 19, using (97), (98) and (99) we have the isomorphism

$$(108) \quad q^*(\omega_{X^{[2]}}) \cong \tau^*(\mathcal{O}_{X \times X}(-n-2+d, -n-2+d)) \otimes \mathcal{O}_{\text{Bl}_\Delta(X \times X)}((n-2)E).$$

Now, by the blowup closure lemma from [42, lemma 22.2.6] we get that the exceptional divisor  $E$  of  $\text{Bl}_\Delta(X \times X)$  is the restriction of the exceptional divisor  $E'$  on  $\text{Bl}_{\Delta'}(\mathbb{P}^{n+1} \times \mathbb{P}^{n+1})$ , so we have an isomorphism

$$(109) \quad \begin{aligned} & \tau^*(\mathcal{O}_{X \times X}(n-2, n-2)) \otimes \mathcal{O}_{\text{Bl}_\Delta(X \times X)}((2-n)E) \\ & \cong i^*(\tau'^*(\mathcal{O}_{\mathbb{P}^{n+1} \times \mathbb{P}^{n+1}}(n-2, n-2)) \otimes \mathcal{O}_{\text{Bl}_{\Delta'}(\mathbb{P}^{n+1} \times \mathbb{P}^{n+1})}((2-n)E')). \end{aligned}$$

Next we consider the composition of the morphism  $\varphi$  (now for  $\mathbb{P}^{n+1}$ ) with the inclusion  $\iota$ , which allows us to further rewrite this line bundle as

$$(110) \quad \cong i^* \circ \varphi^*(\mathcal{O}_{\mathbb{P}^{n+1} \times \mathbb{P}^{n+1} \times \text{Gr}(2, n+2)}(0, 0, n-2)).$$

Combining this with the isomorphism

$$(111) \quad \tau^*(\mathcal{O}_{X \times X}(4-d, 4-d)) \cong i^* \circ \varphi^*(\mathcal{O}_{\mathbb{P}^{n+1} \times \mathbb{P}^{n+1} \times \text{Gr}(2, n+2)}(4-d, 4-d, 0))$$

we obtain after dualising the isomorphism

$$(112) \quad q^*(\omega_{X^{[2]}}^\vee) \cong i^* \circ \varphi^*(\mathcal{O}_{\mathbb{P}^{n+1} \times \mathbb{P}^{n+1} \times \text{Gr}(2, n+2)}(4-d, 4-d, n-2)).$$

Hence we see that  $q^*(\omega_{X^{[2]}}^\vee)$  is ample as soon as  $d \leq 3$  and  $n \geq 3$ , because  $\varphi \circ \iota$  is a again finite morphism.

We can now conclude as in the proof of proposition 19, as by [40, tag 0B5V] we know that  $\omega_{X^{[2]}}^\vee$  is ample if and only if  $q^*(\omega_{X^{[2]}}^\vee)$  is ample.  $\square$

**Remark 21** (Geometric properties of the Hilbert squares). By [5, theorem C] we can understand the deformation theory of these Fano Hilbert squares. We have that  $\mathbb{P}^n$  and  $Q^n$  are rigid, and therefore so are  $\mathbb{P}^{n[2]}$  and  $Q^{n[2]}$ . Cubic hypersurfaces of dimension  $n$  on the other hand come in a family of dimension  $\binom{n+2}{3}$ , and so do their Hilbert squares.

Finally, by [7, theorem 4] we have that the automorphism group of  $\mathbb{P}^{n[2]}$  is isomorphic to that  $\mathbb{P}^n$ , i.e. is given by  $\text{PGL}_{n+1}$ . It would be interesting to understand the automorphism groups of Hilbert squares (and especially those which are again Fano) more generally.

### 5.3 Application: the Fano variety of lines is a Fano visitor

As discussed in the introduction, the *Fano visitor problem* for a smooth projective variety  $X$  asks for the construction of a fully faithful functor  $\text{D}^b(X) \hookrightarrow \text{D}^b(Y)$  into the derived category of a smooth projective Fano variety  $Y$ . In this case we call  $X$  a *Fano*



visitor and  $Y$  the *Fano host*. We will study this for the Fano variety of lines on a cubic hypersurface.

Now let  $Y \subset \mathbb{P}^{n+1}$  be a smooth cubic hypersurface. By theorem B we have that  $\mathbf{D}^b(F(Y))$  is an admissible subcategory of  $\mathbf{D}^b(Y^{[2]})$ , whilst  $Y^{[2]}$  is Fano if  $n \geq 3$  by theorem C. This proves corollary D. In fact, if  $n \geq 5$  then  $F(Y)$  is itself a Fano variety, which is why we are only interested in  $n = 3, 4$  for the Fano visitor problem [2, proposition 1.8].

**Cubic threefolds** If  $n = 3$ , then  $F(Y)$  is a smooth projective surface of general type [2, proposition 1.21], whose Hodge diamond is

$$(113) \quad \begin{array}{cccc} & & 1 & \\ & 5 & & 5 \\ 10 & & 25 & & 10 \\ & 5 & & 5 & \\ & & 1 & & \end{array} .$$

As far as we know this is the first construction of a Fano host for a surface of general type which is not a complete intersection or a product of curves.

Observe that  $\omega_{F(Y)} \cong \mathcal{O}_{F(Y)}(1)$  is a very ample line bundle (as it is the restriction of  $\mathcal{O}_{\mathbb{P}^{\binom{n+2}{2}-1}}(1)$  along the closed immersions  $F(Y) \hookrightarrow \text{Gr}(2, n+2) \hookrightarrow \mathbb{P}^{\binom{n+2}{2}-1}$ ), hence by [26, corollary 1.5] we have that  $\mathbf{D}^b(F(Y))$  is indecomposable.

**Cubic fourfolds** If  $n = 4$ , then the Fano variety of lines on a cubic fourfold is a 4-dimensional hyperkähler variety, deformation equivalent to the Hilbert square of a K3 surface [4]. Hence we have constructed Fano hosts for the complete 20-dimensional family of hyperkählers arising from Fano varieties for cubic fourfolds. As  $\mathbf{D}^b(F(Y))$  is Calabi–Yau, it is indecomposable.

It is still an interesting question to find a Fano host for other 20-dimensional families of hyperkähler 4-folds deformation equivalent to the Hilbert square of a K3 surface.

**Higher dimensions** If  $n \geq 5$ , then the Fano variety of lines is itself a Fano variety, and the interesting question is to find a natural semiorthogonal decomposition for it, rather than embed it in the derived category of a Fano variety. We do not address this here.

**Cubic surfaces** If  $n = 2$ , then  $Y^{[2]}$  is not a Fano variety, because the Hilbert–Chow morphism is a crepant resolution of singularities of  $\text{Sym}^2 Y$ , but it is log Fano [9, corollary 3]. Theorem B still applies though, and we obtain a semiorthogonal decomposition

$$(114) \quad \mathbf{D}^b(Y^{[2]}) = \langle \mathbf{D}^b(F(Y)), \mathbf{D}^b(Y), \mathbf{D}^b(Y), \mathbf{D}^b(Y) \rangle,$$

which can be refined into a full exceptional collection of length 54.

As  $F(Y)$  consists of 27 points, we obtain 27 completely orthogonal objects in  $\mathbf{D}^b(Y^{[2]})$ . These are the structure sheaves of the  $\mathbb{P}^2$ 's embedded in  $Y^{[2]}$  which parametrise 2 points on each of the 27 lines. Alternatively, consider the natural morphism  $Y^{[2]} \rightarrow \text{Gr}(2, 4)$

sending 2 points to the line they span in  $\mathbb{P}^3$ . This morphism is generically finite of degree  $\binom{3}{2} = 3$  onto its image, and the 27 completely orthogonal objects correspond to the locus where the morphism is not finite.

## A Short proof of theorem A(ii) for $\ell = 1$

In this appendix we give a short proof of theorem A(ii) when  $\ell = 1$ . This proof has the added benefit of comparing the category  $\mathbf{D}^b(X)$  to the components of (5) as subcategories in  $\mathbf{D}^b(\tilde{X})$ , and not only after applying  $\mathbf{R}\tau_*$ .

Assume we are given a standard flip diagram (2) with  $\ell = 1$ . We recall some of the notation introduced in section 3. Let  $a, b \in \mathbb{Z}$ , then we denote by  $\mathcal{O}(a, b) := p^* \mathcal{O}_\pi(a) \otimes p'^* \mathcal{O}_{\pi'}(b)$ , which is a line bundle on  $E$ . We will consider the following triangulated subcategory of  $\mathbf{D}^b(\tilde{X})$ :

$$(115) \quad \begin{aligned} \mathcal{A}(a, b) &:= j_* (p^* \circ \pi^* (\mathbf{D}^b(F) \otimes \mathcal{O}(a, b))) \\ &= j_* (p'^* \circ \pi'^* (\mathbf{D}^b(F) \otimes \mathcal{O}(a, b))). \end{aligned}$$

As  $\tilde{X}$  is the blow up of  $X$  along  $Z$  (which has codimension 2 by assumption), Orlov's blowup formula [35, theorem 4.3] gives a semiorthogonal decomposition

$$(116) \quad \mathbf{D}^b(\tilde{X}) = \langle j_* (p^* (\mathbf{D}^b(Z)) \otimes \mathcal{O}_E(E)), \mathbf{L}\tau^* \mathbf{D}^b(X) \rangle.$$

Using the fact that  $Z$  is a  $\mathbb{P}^k$ -bundle over  $F$  by (2), the first component in (116) can be further decomposed using Orlov's projective bundle formula [35, theorem 2.6] using our notation as

$$(117) \quad j_* (p^* (\mathbf{D}^b(Z)) \otimes \mathcal{O}_E(E)) = \langle \mathcal{A}(-k, -1), \mathcal{A}(-k+1, -1), \dots, \mathcal{A}(0, -1) \rangle.$$

Combining (116) with (117), we have the following semiorthogonal decomposition:

$$(118) \quad \mathbf{D}^b(\tilde{X}) = \langle \mathcal{A}(-k, -1), \mathcal{A}(-k+1, -1), \dots, \mathcal{A}(0, -1), \mathbf{L}\tau^* \mathbf{D}^b(X) \rangle.$$

Here all the components except the last one are equivalent to  $\mathbf{D}^b(F)$  via the defining functors.

Similarly, using the fact that  $\tilde{X}$  is also the blowup of  $X'$  along  $Z'$  and that  $Z'$  is a  $\mathbb{P}^1$ -bundle over  $F$ , we get a second semiorthogonal decomposition:

$$(119) \quad \mathbf{D}^b(\tilde{X}) = \langle \mathcal{A}(-k, -1), \mathcal{A}(-k, 0), \mathcal{A}(-k+1, -1), \mathcal{A}(-k+1, 0), \dots, \mathcal{A}(-1, -1), \mathcal{A}(-1, 0), \mathbf{L}\tau'^* \mathbf{D}^b(X') \rangle.$$

Again all the components of (119) except the last one are equivalent to  $\mathbf{D}^b(F)$ .

To compare (118) and (119), we perform a sequence of mutations on (119).

First we mutate  $\mathcal{A}(-k, 0)$  to the left. The resulting semiorthogonal decomposition is

$$(120) \quad \mathbf{D}^b(\tilde{X}) = \langle \mathcal{A}(-k, -2), \mathcal{A}(-k, -1), \mathcal{A}(-k+1, -1), \mathcal{A}(-k+1, 0), \dots, \mathcal{A}(-1, -1), \mathcal{A}(-1, 0), \mathbf{L}\tau'^* (\mathbf{D}^b(X')) \rangle.$$

Next we mutate  $\mathcal{A}(-k, -2)$  to the far right. By [10, Proposition 3.6], the resulting decomposition is

$$(121)$$

$$\mathbf{D}^b(\tilde{X}) = \langle \mathcal{A}(-k, -1), \mathcal{A}(-k+1, -1), \mathcal{A}(-k+1, 0), \dots, \mathcal{A}(-1, -1), \mathcal{A}(-1, 0), \mathbf{L}\tau'^* \mathbf{D}^b(X'), \mathbb{S}_{\tilde{X}}^{-1} \mathcal{A}(-k, -2) \rangle,$$

where  $\mathbb{S}_{\tilde{X}} = -\otimes \omega_{\tilde{X}}[\dim \tilde{X}]$  is the Serre functor of  $\mathbf{D}^b(\tilde{X})$ . We can easily compute  $\mathbb{S}_{\tilde{X}}^{-1} \mathcal{A}(-k, -2)$  as follows

$$\begin{aligned} \mathbb{S}_{\tilde{X}}^{-1} \mathcal{A}(-k, -2) &= \mathcal{A}(-k, -2) \otimes \omega_{\tilde{X}}^\vee \\ &= j_*(p^* \circ \pi^*(\mathbf{D}^b(F)) \otimes \mathcal{O}(-k, -2)) \otimes \omega_{\tilde{X}}^\vee \\ &= j_*(p^* \circ \pi^*(\mathbf{D}^b(F)) \otimes \mathcal{O}(-k, -2) \otimes \omega_{\tilde{X}}|_E^\vee) \\ (122) \quad &= j_*(p^* \circ \pi^*(\mathbf{D}^b(F)) \otimes \mathcal{O}(-k, -2) \otimes \omega_E^\vee \otimes \mathcal{O}_E(E)) \\ &= j_*(p^* \circ \pi^*(\mathbf{D}^b(F)) \otimes \mathcal{O}(-k, -2) \otimes \mathcal{O}(k+1, 2) \otimes \mathcal{O}(-1, -1)) \\ &= j_*(p^* \circ \pi^*(\mathbf{D}^b(F)) \otimes \mathcal{O}(0, -1)) \\ &= \mathcal{A}(0, -1). \end{aligned}$$

The result of the mutation is therefore

$$(123) \quad \mathbf{D}^b(\tilde{X}) = \langle \mathcal{A}(-k, -1), \mathcal{A}(-k+1, -1), \mathcal{A}(-k+1, 0), \dots, \mathcal{A}(-1, -1), \mathcal{A}(-1, 0), \mathbf{L}\tau'^*(\mathbf{D}^b(X')), \mathcal{A}(0, -1) \rangle,$$

Then for each  $m = -k+1, \dots, -2$ , we right mutate the component  $\mathcal{A}(m, 0)$  through each of the categories  $\mathcal{A}(m, -1), \mathcal{A}(m+1, -1), \dots, \mathcal{A}(-1, -1)$ . Lacking control over the resulting category, we denote the result for now as

$$(124) \quad \begin{aligned} \mathbf{D}^b(\tilde{X}) &= \langle \mathcal{A}(-k, -1), \mathcal{A}(-k+1, -1), \dots, \mathcal{A}(-1, -1), \\ &\mathcal{B}(-k+1, 0), \dots, \mathcal{B}(-2, 0), \mathcal{A}(-1, 0), \mathbf{L}\tau'^* \mathbf{D}^b(X'), \mathcal{A}(0, -1) \rangle, \end{aligned}$$

where for all  $-k+1 \leq m \leq -2$ , we denote

$$(125) \quad \begin{aligned} \mathcal{B}(m, 0) &:= \mathbf{R}_{\langle \mathcal{A}(m, -1), \mathcal{A}(m+1, -1), \dots, \mathcal{A}(-1, -1) \rangle} \mathcal{A}(m, 0) \\ &= \mathbf{R}_{\mathcal{A}(-1, -1)} \circ \dots \circ \mathbf{R}_{\mathcal{A}(m+1, -1)} \circ \mathbf{R}_{\mathcal{A}(m, -1)} \mathcal{A}(m, 0). \end{aligned}$$

The identification of the mutation through the subcategory generated by a sequence of subcategories and the composition of mutations follows from e.g. [29, lemma 2.2(i)].

Finally we right mutate the subcategory  $\langle \mathcal{B}(-k+1, 0), \dots, \mathcal{B}(-2, 0), \mathcal{A}(-1, 0), \mathbf{L}\tau'^* \mathbf{D}^b(X') \rangle$  through  $\mathcal{A}(0, -1)$ . The result is

$$(126) \quad \begin{aligned} \mathbf{D}^b(\tilde{X}) &= \langle \mathcal{A}(-k, -1), \mathcal{A}(-k+1, -1), \dots, \mathcal{A}(-1, -1), \mathcal{A}(0, -1), \\ &\mathbf{R}_{\mathcal{A}(0, -1)} \mathcal{B}(-k+1, 0), \dots, \mathbf{R}_{\mathcal{A}(0, -1)} \mathcal{B}(-2, 0), \mathbf{R}_{\mathcal{A}(0, -1)} \mathcal{A}(-1, 0), \mathbf{R}_{\mathcal{A}(0, -1)} \mathbf{L}\tau'^* \mathbf{D}^b(X') \rangle. \end{aligned}$$

As mutations induce equivalences between the original and the mutated components, in the previous decomposition, each component except the last one is equivalent to  $\mathbf{D}^b(F)$ .

Comparing the semiorthogonal decomposition (118) with the mutated semiorthogonal decomposition (126) obtained in the last step, we deduce the following equality of subcategories of  $\mathbf{D}^b(\tilde{X})$ :

$$(127)$$

$$\mathbf{L}\tau^* \mathbf{D}^b(X) = \langle \mathbf{R}_{\mathcal{A}(0,-1)} \mathcal{B}(-k+1, 0), \dots, \mathbf{R}_{\mathcal{A}(0,-1)} \mathcal{B}(-2, 0), \mathbf{R}_{\mathcal{A}(0,-1)} \mathcal{A}(-1, 0), \mathbf{R}_{\mathcal{A}(0,-1)} \mathbf{L}\tau'^* \mathbf{D}^b(X') \rangle.$$

By construction, each component except the last one is equivalent to  $\mathbf{D}^b(F)$ .

By the fully faithfulness of  $\mathbf{L}\tau^* : \mathbf{D}^b(X) \rightarrow \mathbf{D}^b(\tilde{X})$ , its right adjoint  $\mathbf{R}\tau_* : \mathbf{D}^b(\tilde{X}) \rightarrow \mathbf{D}^b(X)$  induces an equivalence between  $\mathbf{L}\tau^* \mathbf{D}^b(X)$  and  $\mathbf{D}^b(X)$ . Applying this equivalence to (127), we get a semiorthogonal decomposition

(128)

$$\mathbf{D}^b(X) = \langle \mathbf{R}\tau_* \mathbf{R}_{\mathcal{A}(0,-1)} \mathcal{B}(-k+1, 0), \dots, \mathbf{R}\tau_* \mathbf{R}_{\mathcal{A}(0,-1)} \mathcal{B}(-2, 0), \mathbf{R}\tau_* \mathbf{R}_{\mathcal{A}(0,-1)} \mathcal{A}(-1, 0), \mathbf{R}\tau_* \mathbf{R}_{\mathcal{A}(0,-1)} \mathbf{L}\tau'^* \mathbf{D}^b(X') \rangle.$$

It remains to compute the components appearing in this decomposition. We do this via the following trivial lemma, which says that if a functor annihilates a subcategory, then the mutation through this subcategory does not change the image by this functor.

**Lemma 22.** Let  $\mathcal{A}$  be an admissible triangulated subcategory of a triangulated subcategory  $\mathcal{T}$ . If  $\Psi : \mathcal{T} \rightarrow \mathcal{T}'$  is a triangulated functor to another triangulated category such that  $\Psi(\mathcal{A}) = 0$ , then

$$(129) \quad \Psi(\mathcal{A}^\perp) = \Psi(\mathcal{T}) = \Psi(\mathcal{A}^\perp).$$

Note that  ${}^\perp \mathcal{A} = \mathbf{R}_{\mathcal{A}} \mathcal{A}^\perp$ .

Now, for any integer  $m$ , we claim that the functor  $\mathbf{R}\tau_*$  annihilates the subcategory  $\mathcal{A}(m, -1)$ . Indeed, for any  $\mathcal{E} \in \mathbf{D}^b(F)$  we have

(130)

$$\begin{aligned} \mathbf{R}\tau_* \circ j_* (p^* \circ \pi^* (\mathcal{E}) \otimes \mathcal{O}(m, -1)) &= \mathbf{R}\tau_* \circ j_* (p^* \circ \pi^* (\mathcal{E}) \otimes p^* \mathcal{O}_\pi(m) \otimes p'^* \mathcal{O}_{\pi'}(-1)) \\ &= i_* \circ \mathbf{R}p_* (p^* \circ \pi^* (\mathcal{E}) \otimes \mathcal{O}_\pi(m)) \otimes p'^* \mathcal{O}_{\pi'}(-1) \\ &= i_* (\pi^* (\mathcal{E}) \otimes \mathcal{O}_\pi(m) \otimes \mathbf{R}p_* \circ p'^* (\mathcal{O}_{\pi'}(-1))) \end{aligned}$$

However by the base change formula we have that

$$(131) \quad \mathbf{R}p_* \circ p'^* (\mathcal{O}_{\pi'}(-1)) = \pi^* \circ \mathbf{R}\pi'_* \mathcal{O}_{\pi'}(-1) = 0$$

which proves the necessary vanishing.

Applying lemma 22 to the functor  $\mathbf{R}\tau_*$ , we see that a mutation through any category of the form  $\mathcal{A}(m, -1)$  with  $m \in \mathbb{Z}$  does not change the image under  $\mathbf{R}\tau_*$ . In particular we have

$$(132) \quad \mathbf{R}\tau_* \mathbf{R}_{\mathcal{A}(0,-1)} \mathcal{B}(m, 0) = \mathbf{R}\tau_* \mathcal{B}(m, 0) = \mathbf{R}\tau_* \mathcal{A}(m, 0).$$

Therefore, the semiorthogonal decomposition (128) is nothing else but the following:

$$(133) \quad \mathbf{D}^b(X) = \langle \mathbf{R}\tau_* \mathcal{A}(-k+1, 0), \dots, \mathbf{R}\tau_* \mathcal{A}(-2, 0), \mathbf{R}\tau_* \mathcal{A}(-1, 0), \mathbf{R}\tau_* \circ \mathbf{L}\tau'^* \mathbf{D}^b(X') \rangle.$$

Finally, since

$$(134) \quad \begin{aligned} \mathbf{R}\tau_* \circ j_* (p^* \circ \pi^* \mathbf{D}^b(F) \otimes \mathcal{O}(n, 0)) &= i_* \circ \mathbf{R}p_* (p^* \circ \pi^* \mathbf{D}^b(F) \otimes p^* \mathcal{O}_\pi(n)) \\ &= i_* (\pi^* \mathbf{D}^b(F) \otimes \mathcal{O}_\pi(n)) \\ &= \Phi_n(\mathbf{D}^b(F)), \end{aligned}$$

where  $\Phi_n$  is defined in (4), the decomposition (133) is exactly the one in theorem A(ii).

## References

- [1] Nicolas Addington, Will Donovan, and Ciaran Meachan. Mukai flops and  $\mathbb{P}$ -twists. *J. Reine Angew. Math.*, 748:227–240, 2019.
- [2] Allen B. Altman and Steven L. Kleiman. Foundations of the theory of Fano schemes. *Compositio Math.*, 34(1):3–47, 1977.
- [3] Wolf P. Barth, Klaus Hulek, Chris A. M. Peters, and Antonius Van de Ven. *Compact complex surfaces*, volume 4 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, second edition, 2004.
- [4] Arnaud Beauville and Ron Donagi. La variété des droites d’une hypersurface cubique de dimension 4. *C. R. Acad. Sci. Paris Sér. I Math.*, 301(14):703–706, 1985.
- [5] Pieter Belmans, Lie Fu, and Theo Raedschelders. Hilbert squares: derived categories and deformations. *Selecta Math. (N.S.)*, 25(3):Art. 37,32, 2019.
- [6] Pieter Belmans and Swarnava Mukhopadhyay. Admissible subcategories in derived categories of moduli of vector bundles on curves. *Adv. Math.*, 351:653–675, 2019.
- [7] Pieter Belmans, Georg Oberdieck, and J yrge Vold Rennemo. Automorphisms of Hilbert schemes of points on surfaces. <https://arxiv.org/abs/1907.07064v2>.
- [8] Marcello Bernardara, Michele Bolognesi, and Daniele Faenzi. Homological projective duality for determinantal varieties. *Adv. Math.*, 296:181–209, 2016.
- [9] Aaron Bertram and Izzet Coskun. The birational geometry of the Hilbert scheme of points on surfaces. In *Birational geometry, rational curves, and arithmetic*, Simons Symp., pages 15–55. Springer, Cham, 2013.
- [10] Alexey Bondal and Mikhail Kapranov. Representable functors, Serre functors, and reconstructions. *Izv. Akad. Nauk SSSR Ser. Mat.*, 53(6):1183–1205,1337, 1989.
- [11] Alexey Bondal, Michael Larsen, and Valery Lunts. Grothendieck ring of pretriangulated categories. *Int. Math. Res. Not.*, (29):1461–1495, 2004.
- [12] Alexey Bondal and Dmitri Orlov. Semiorthogonal decomposition for algebraic varieties, 1995. <https://arxiv.org/abs/alg-geom/9506012>.
- [13] Jan Cheah. Cellular decompositions for nested Hilbert schemes of points. *Pacific J. Math.*, 183(1):39–90, 1998.
- [14] Anton Fonarev and Alexander Kuznetsov. Derived categories of curves as components of Fano manifolds. *J. Lond. Math. Soc. (2)*, 97(1):24–46, 2018.
- [15] Sergey Galkin and Evgeny Shinder. The fano variety of lines and rationality problem for a cubic hypersurface. <https://arxiv.org/abs/1405.5154v2>.
- [16] R. Paul Horja. Derived category automorphisms from mirror symmetry. *Duke Math. J.*, 127(1):1–34, 2005.

- [17] Daniel Huybrechts. *Fourier-Mukai transforms in algebraic geometry*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, Oxford, 2006.
- [18] Daniel Huybrechts. The geometry of cubic hypersurfaces, <http://www.math.uni-bonn.de/~huybrech/>. book-in-progress, see.
- [19] Atanas Iliev and Laurent Manivel. Fano manifolds of Calabi-Yau Hodge type. *J. Pure Appl. Algebra*, 219(6):2225–2244, 2015.
- [20] Qingyuan Jiang. On the Chow theory of projectivization. <https://arxiv.org/abs/1910.06730v1>.
- [21] Qingyuan Jiang and Naichung Conan Leung. Derived category of projectivization and flops. <https://arxiv.org/abs/1811.12525v1>.
- [22] Yujiro Kawamata.  $D$ -equivalence and  $K$ -equivalence. *J. Differential Geom.*, 61(1):147–171, 2002.
- [23] Yujiro Kawamata. Derived categories of toric varieties. *Michigan Math. J.*, 54(3):517–535, 2006.
- [24] Yujiro Kawamata. Derived categories of toric varieties II. *Michigan Math. J.*, 62(2):353–363, 2013.
- [25] Yujiro Kawamata. Derived categories of toric varieties III. *Eur. J. Math.*, 2(1):196–207, 2016.
- [26] Kotaro Kawatani and Shinnosuke Okawa. Nonexistence of semiorthogonal decompositions and sections of the canonical bundle. <https://arxiv.org/abs/1508.00682v2>.
- [27] Young-Hoon Kiem, In-Kyun Kim, Hwayoung Lee, and Kyoung-Seog Lee. All complete intersection varieties are Fano visitors. *Adv. Math.*, 311:649–661, 2017.
- [28] Andreas Krug, David Ploog, and Pawel Sosna. Derived categories of resolutions of cyclic quotient singularities. *Q. J. Math.*, 69(2):509–548, 2018.
- [29] Alexander Kuznetsov. Calabi-Yau and fractional Calabi-Yau categories. *J. Reine Angew. Math.*, 753:239–267, 2019.
- [30] Alexander Kuznetsov. Embedding derived categories of an Enriques surface into derived categories of Fano varieties. *Izv. Ross. Akad. Nauk Ser. Mat.*, 83(3):127–132, 2019.
- [31] Alexander Kuznetsov. Homological projective duality. *Publ. Math. Inst. Hautes Études Sci.*, (105):157–220, 2007.
- [32] Robert Laterveer. A remark on the motive of the Fano variety of lines of a cubic. *Ann. Math. Qué.*, 41(1):141–154, 2017.
- [33] Jacob P. Murre, Jan Nagel, and Chris A. M. Peters. *Lectures on the theory of pure motives*, volume 61 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2013.
- [34] Mudumbai S. Narasimhan. Derived categories of moduli spaces of vector bundles on curves. *J. Geom. Phys.*, 122:53–58, 2017.

- [35] Dmitri Orlov. Projective bundles, monoidal transformations, and derived categories of coherent sheaves. *Izv. Ross. Akad. Nauk Ser. Mat.*, 56(4):852–862, 1992.
- [36] Dmitri Orlov. Triangulated categories of singularities, and equivalences between Landau-Ginzburg models. *Mat. Sb.*, 197(12):117–132, 2006.
- [37] Raphaël Rouquier. Catégories dérivées et géométrie birationnelle (d’après Bondal, Orlov, Bridgeland, Kawamata et al.). Number 307, pages Exp. No. 946,viii,283–307. 2006. Séminaire Bourbaki. Vol. 2004/2005.
- [38] Will Sawin. Can free rational curves lift to ramified covers of Fano varieties? MathOverflow. (version: 2018-11-14).
- [39] Will Sawin. Freeness alone is insufficient for Manin-Peyre. <https://arxiv.org/abs/2001.06078v1>.
- [40] The Stacks project authors. The Stacks project, 2019.
- [41] Richard P. Thomas. Notes on homological projective duality. In *Algebraic geometry: Salt Lake City 2015*, volume 97 of *Proc. Sympos. Pure Math.*, pages 585–609. Amer. Math. Soc., Providence, RI, 2018.
- [42] Ravi Vakil. *The Rising Sea: Foundation of algebraic geometry*, 2017.
- [43] Claire Voisin. On the universal  $CH_0$  group of cubic hypersurfaces. *J. Eur. Math. Soc. (JEMS)*, 19(6):1619–1653, 2017.

Pieter Belmans, pbelmans@math.uni-bonn.de  
 UNIVERSITÄT BONN, ENDENICHER ALLEE 60, 53115 BONN, GERMANY  
 UNIVERSITEIT ANTWERPEN, MIDDELHEIMLAAN 1, 2020 ANTWERPEN, BELGIUM  
 UNIVERSITEIT HASSELT, AGORALAAN, 3590 DIEPENBEEK, BELGIUM

Lie Fu, fu@math.univ-lyon1.fr  
 UNIVERSITÉ CLAUDE BERNARD LYON 1, 43 BD DU 11 NOVEMBRE 1918, 69622 VILLE-URBANNE CEDEX, FRANCE  
 RADBOUD UNIVERSITY, HEYENDAALSEWEG 135, 6525 AJ NIJMEGEN, NETHERLANDS

Theo Raedschelders, theo.raedschelders@vub.be  
 VRIJE UNIVERSITEIT BRUSSEL, PLEINLAAN 2, 1050 BRUSSELS, BELGIUM