Projectivity of the moduli space of vector bundles on a curve

Jarod Alper Pieter Belmans Daniel Bragg Jason Liang Tuomas Tajakka

June 14, 2022

Abstract

We discuss the projectivity of the moduli space of semistable vector bundles on a curve of genus $g \ge 2$. This is a classical result from the 1960s, obtained using geometric invariant theory. We outline a modern approach that combines the recent machinery of good moduli spaces with determinantal line bundle techniques. The crucial step producing an ample line bundle follows an argument by Faltings with improvements by Esteves–Popa. We hope to promote this approach as a blueprint for other projectivity arguments.

Dedicated to the memory of Conjeevaram Srirangachari Seshadri

In [17], Kollár took a modern approach to the construction of the moduli space $\overline{\mathrm{M}}_g$ of stable curves of genus $g \geq 2$ as a projective variety, avoiding methods from geometric invariant theory. The approach can be summarized in three steps.

- (i). Prove that the stack of stable curves is a proper Deligne–Mumford stack $\overline{\mathcal{M}}_{g}$ [7].
- (ii). Use the Keel–Mori theorem to show that $\overline{\mathcal{M}}_g$ has a coarse moduli space $\overline{\mathcal{M}}_g \to \overline{\mathcal{M}}_g$, where $\overline{\mathcal{M}}_g$ is a proper algebraic space [15].
- (iii). Prove that some line bundle on $\overline{\mathcal{M}}_g$ descends to an *ample* line bundle on $\overline{\mathcal{M}}_g$ [17, 6].

In [5] the projectivity of $\overline{\mathrm{M}}_g$ was established by giving a stack-theoretic treatment of Kollár's paper.

We take a similar approach to constructing the moduli space $M_X^{ss}(r, d)$ of semistable vector bundles of rank r and degree d on a smooth, proper curve Xas a projective variety. There are again three steps in the proof.

(i). Prove that $\mathcal{M}_X^{ss}(r,d)$ is a universally closed algebraic stack of finite type [19].

(ii). Use an analogue of the Keel–Mori theorem to show that there exists a good moduli space $\mathcal{M}_X^{ss}(r,d) \to \mathcal{M}_X^{ss}(r,d)$ [2], where $\mathcal{M}_X^{ss}(r,d)$ is a proper algebraic space.

Unlike the case of $\overline{\mathcal{M}}_g$, the stack $\mathcal{M}_X^{\mathrm{ss}}(r, d)$ is never Deligne–Mumford and the morphism to $\mathcal{M}_X^{\mathrm{ss}}(r, d)$ will identify nonisomorphic vector bundles, introducing the notion of S-equivalence.

(iii). Prove that a suitable determinantal line bundle on $\mathcal{M}_X^{ss}(r, d)$ descends to an ample line bundle on $\mathcal{M}_X^{ss}(r, d)$.

In Sections 1–3 we define and study the moduli stack $\mathcal{M}_X^{ss}(r, d)$ and establish steps (i) and (ii). The main points are summarized in Theorems 2.17 and 3.12.

In Section 5 we discuss step (iii) and give a detailed exposition of the construction of an ample line bundle on $M_X^{ss}(r, d)$ using techniques from Esteves–Popa [9]. This method is more recent than the first GIT-free construction due to Faltings [10], which has received expository accounts by Seshadri (and Nori) in [22] and for a special case by Hein in [13]. All these constructions build upon the notion of determinantal line bundles, which we introduce in Section 4.

Because the existence results for good moduli spaces from [2] are currently restricted to characteristic 0, we make that restriction as well.

1 The moduli stack of all vector bundles

Let k be an algebraically closed field of characteristic 0, and let X be a smooth, projective, connected curve over k.

We denote by $\underline{\operatorname{Coh}}_X$ the stack of all coherent sheaf on X. For integers r > 0and d, we let $\mathcal{M}_X(r, d)$ denote the stack parameterizing vector bundles on X of rank r and degree d. An object of $\mathcal{M}_X(r, d)$ over a scheme S is a vector bundle \mathcal{E} on $X \times S$ such that for every geometric point $\operatorname{Spec} K \to S$, the restriction $\mathcal{E}|_{X \times \operatorname{Spec} K}$ has rank r and degree d.

Proposition 1.1. The stack $\mathcal{M}_X(r,d)$ is an algebraic stack, locally of finite type over k with affine diagonal.

Proof. The stack $\underline{\operatorname{Coh}}_X$ is algebraic by [23, Tag 09DS]. Given a map $T \to \underline{\operatorname{Coh}}_X$ corresponding to a sheaf \mathcal{E} on $T \times X$ flat over T, the locus $Z \subseteq T \times X$ where \mathcal{E} is not locally free is closed, so $U = T \setminus \operatorname{pr}_1(Z) \subseteq T$ is open by properness, showing that the substack $\mathcal{M} \subset \underline{\operatorname{Coh}}_X$ parameterizing vector bundles on X is an open substack. The Hilbert polynomial P_E of a coherent sheaf E on X is determined by its rank and degree, and by the Riemann–Roch theorem it is given by

$$P_E(n) = \operatorname{rk}(E)n + \deg(E) + \operatorname{rk}(E)(1-g).$$
(1)

It is locally constant in flat families, so by the discussion in [23, Tag 0DNF] there is an open and closed substack $\mathcal{M}_X(r, d) \subset \mathcal{M}$ parameterizing vector bundles of rank r and degree d.

Since \underline{Coh}_X is locally of finite type over k by [23, Tag 0DLZ], and has affine diagonal by [23, Tag 0DLY], so does $\mathcal{M}_X(r, d)$.

There is a universal vector bundle \mathcal{E}_{univ} on $X \times \mathcal{M}_X(r, d)$. Specializing to the rank-1 case, we obtain the Picard stack $\mathcal{P}ic_X^d$ parameterizing line bundles of degree d on X. In this case the universal bundle is called the Poincaré bundle and denoted \mathcal{P} . The line bundle det (\mathcal{E}_{univ}) on $X \times \mathcal{M}_X(r, d)$ induces a determinant morphism

$$\det\colon \mathcal{M}_X(r,d)\to \mathcal{P}\mathrm{ic}_X^d,$$

with the property that $(\mathrm{id}_X \times \mathrm{det})^* \mathcal{P} \cong \mathrm{det}(\mathcal{E}_{\mathrm{univ}}).$

Let L be a line bundle on X of degree d corresponding to a closed point of Pic_X^d , and let [L]: Spec $k \to \operatorname{Pic}_X^d$ be the corresponding morphism. The algebraic stack parameterizing vector bundles with fixed determinant L is defined as the fiber product

$$\begin{array}{cccc}
\mathcal{M}_X(r,L) & \longrightarrow & \mathcal{M}_X(r,d) \\
& & & & & \downarrow & & \\
& & & & \downarrow & & \downarrow & \\
& & & & & \downarrow & \downarrow & \\
& & & & & & \downarrow & \\
& & & & & & \downarrow & \\
& & & & & & \downarrow & \\
& & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & \downarrow & & \downarrow & & \downarrow & \\
& & & & & & \downarrow & & & \downarrow & \\
& & & & & & & \downarrow & \downarrow & \downarrow & \downarrow & & \downarrow &$$

Explicitly, an object of $\mathcal{M}_X(r,L)$ over a k-scheme S is a pair (\mathcal{E},φ) , where \mathcal{E} is a vector bundle on $X \times S$ with rank r and degree d on the fibers, and $\varphi: \det \mathcal{E} \xrightarrow{\sim} L|_{X \times S}$ is an isomorphism.

An alternative approach to fixing the determinant uses the residual gerbe $\mathbb{BG}_{\mathrm{m}} \hookrightarrow \mathcal{P}\mathrm{ic}_X^d$ instead. We compare these in Example 3.2.

Proposition 1.2. The stack $\mathcal{M}_X(r, L)$ is an algebraic stack, locally of finite type over k with affine diagonal.

Proof. This follows by applying [23, Tag 04TF] for the algebraicity and [23, Tag 06FU] for being locally of finite type. To see that $\mathcal{M}_X(r, L)$ has affine diagonal, we use the fact that det: $\mathcal{M}_X(r, d) \to \mathcal{P}ic_X^d$ has affine diagonal (because both stacks do) and that having affine diagonal is stable under base change.

Let \mathscr{X} be an algebraic stack. Let $x_0 \in \mathscr{K}(k)$ be a k-point. By [23, Tag 07WZ], the tangent space $T_{\mathscr{X},x_0}$ to \mathscr{X} at x_0 is the set of isomorphism classes of objects in the groupoid $\mathscr{X}_{x_0}(k[\varepsilon]/(\varepsilon^2))$, where \mathscr{X}_{x_0} denotes the slice category. Using deformation theory one can prove the following result about the moduli stacks.

Proposition 1.3.

- (i). The algebraic stacks $\mathcal{M}_X(r,d)$ and $\mathcal{M}_X(r,L)$ are smooth over k.
- (ii). If E is a vector bundle of rank r and φ : $L \cong \det E$ is an isomorphism, then there are identifications of the Zariski tangent space

$$T_{\mathcal{M}_{X}(r,d),[E]} \cong \operatorname{Ext}^{1}_{X}(E,E), \tag{3}$$

$$T_{\mathcal{M}_{X}(r,L),[(E,\varphi)]} \cong \ker \left(\operatorname{Ext}_{X}^{1}(E,E) \xrightarrow{\operatorname{tr}} \operatorname{H}^{1}(X,\mathcal{O}_{X}) \right).$$
(4)

Remark 1.4. Let *E* be a *simple* vector bundle of rank *r* and degree *d*, that is, dim Hom_{*X*}(*E*, *E*) = 1. We will see in Lemma 2.5 that stable bundles are simple. By the Riemann–Roch theorem, we have

$$\dim \operatorname{Ext}_X^1(E, E) = \dim \operatorname{Hom}_X(E, E) - \chi(X, E^{\vee} \otimes E)$$
$$= 1 - \deg(E^{\vee} \otimes E) - \operatorname{rk}(E^{\vee} \otimes E)(1 - g)$$
$$= r^2(g - 1) + 1.$$

Next, note that since we are working over a field of characteristic 0, the trace map $E^{\vee} \otimes E \to \mathcal{O}_X$ is split by the section $\frac{1}{d} \operatorname{id}_E \colon \mathcal{O}_X \to E^{\vee} \otimes E \cong \operatorname{Hom}_X(E, E)$. Thus, the surjection $\operatorname{Ext}^1_X(E, E) \to \operatorname{H}^1(X, \mathcal{O}_X)$ in (4) is split, and so its kernel has dimension

$$\dim \operatorname{Ext}_X^1(E, E) - \dim \operatorname{H}^1(X, \mathcal{O}_X) = r^2(g-1) + 1 - g$$
$$= (r^2 - 1)(g - 1).$$

2 Semistability

In order to obtain a more well-behaved geometric object, we will restrict ourselves to an open substack $\mathcal{M}_X^{ss}(r,d)$ of the stack $\mathcal{M}_X(r,d)$. This substack parameterizes the *semistable* vector bundles. The Jordan–Hölder and Harder–Narasimhan filtrations explain how it is sufficient to consider these vector bundles in order to describe all vector bundles on X:

- the Harder–Narasimhan filtration allows one to uniquely break up any vector bundles into semistable constituents;
- the Jordan–Hölder filtration allows one to break up a semistable vector bundle into *stable* bundles.

The essential properties of semistability, which will be formalized in Theorem 2.17, are

- openness: $\mathcal{M}_X^{ss}(r,d)$ is an open substack of $\mathcal{M}_X(r,d)$;
- boundedness: $\mathcal{M}_X^{ss}(r, d)$ is a quasicompact;
- irreducibility of the semistable locus.

For more information, in particular on the boundedness of semistable sheaves in higher dimensions and in arbitrary characteristic, see [11].

2.1 Definition and basic properties

Definition 2.1. Given a nonzero vector bundle E on X, we define the *slope* of E to be the rational number

$$\mu(E) := \frac{\deg(E)}{\operatorname{rk}(E)}.$$

Recall that on a smooth curve, any subsheaf of a locally free sheaf is locally free.

Definition 2.2. A vector bundle E on X is

- semistable if, for every nonzero subsheaf $E' \subset E$, we have $\mu(E') \leq \mu(E)$,
- stable if it is semistable and the only subsheaf $E' \subset E$ with $\mu(E') = \mu(E)$ is E' = E,
- *polystable* if it is isomorphic to a finite direct sum $\bigoplus_i E_i$, where each E_i is stable with $\mu(E_i) = \mu(E)$.

Below are some basic properties of semistable vector bundles.

Lemma 2.3. For a line bundle L, a vector bundle E is (semi)stable if and only if $E \otimes L$ is (semi)stable.

Proof. Note that, for any vector bundle E, we have

$$\deg(E \otimes L) = \deg(E) + \operatorname{rk}(E) \cdot \deg(L).$$

Therefore, by definition of the slope,

$$\mu(E \otimes L) = \mu(E) + \deg(L).$$

Hence, for any subsheaf $E' \subset E$, we have

$$\mu(E) - \mu(E') = \mu(E \otimes L) - \mu(E' \otimes L).$$

By definition, E is (semi)stable if and only if for all $E' \subset E$, the above quantity is positive (resp. non-negative), which holds if and only if $E \otimes L$ is (semi)stable. \Box

Lemma 2.4. If E and F are semistable vector bundles with $\mu(E) > \mu(F)$, then $\operatorname{Hom}_X(E,F) = 0$.

Proof. Suppose that on the contrary $E \to F$ is a nonzero map, and let K and I be its kernel and image respectively. Since E is semistable, $\mu(K) \leq \mu(E)$. Hence,

$$\mu(I) \operatorname{rk}(I) = \mu(E) \operatorname{rk}(E) - \mu(K) \operatorname{rk}(K)$$
$$\geq \mu(E) \operatorname{rk}(E) - \mu(E) \operatorname{rk}(K)$$
$$= \mu(E) \operatorname{rk}(I)$$

so we see that $\mu(I) \ge \mu(E) > \mu(F)$. But now I cannot be a subsheaf of F because F is semistable.

The following is proved similarly.

Lemma 2.5. Any nonzero endomorphism of a stable bundle E is invertible. In particular, as k is algebraically closed we have $\operatorname{End}_X(E) \cong k$.

We will use the following lemma in the proof of Proposition 2.15 to establish boundedness.

Lemma 2.6. If E is a semistable vector bundle on X and $\mu(E) > 2g - 1$, then E is globally generated and $H^1(X, E) = 0$.

Proof. Let $x \in X(k)$ be a closed point. Let \mathcal{O}_x denote the skyscraper sheaf at x. By taking the sheaf cohomology of the short exact sequence

$$0 \to E \otimes \mathcal{O}_X(-x) \to E \to \mathcal{O}_x^{\oplus \operatorname{rk}(E)} \to 0,$$

we see that it suffices to show that $\mathrm{H}^1(X, E \otimes \mathcal{O}_X(-x)) = 0$. By Serre duality, this is equivalent to $\mathrm{Hom}_X(E \otimes \mathcal{O}_X(-x), \omega_X) = 0$. Now, $\mu(E \otimes \mathcal{O}_X(-x)) = \mu(E) - 1 > 2g - 2 = \mu(\omega_X)$ so the result follows by Lemma 2.4 since ω_X is semistable as a line bundle.

2.2 Jordan–Hölder and Harder–Narasimhan filtrations

Every semistable vector bundle E on X admits a Jordan-Hölder filtration

$$0 = E_0 \subset E_1 \subset E_2 \subset \cdots \subset E_k = E$$

where each subquotient $G_i = E_i/E_{i-1}$ is a *stable* vector bundle with $\mu(G_i) = \mu(E)$. This filtration is not unique, but the list of subquotients is unique by the following result. See [20] for a proof.

Proposition 2.7 (Jordan–Hölder filtration). Let E be a semistable vector bundle on X. Suppose that

$$0 = E_0 \subset E_1 \subset E_2 \subset \cdots \subset E_k = E$$

and

$$0 = E'_0 \subset E'_1 \subset E'_2 \subset \dots \subset E'_\ell = E$$

are two Jordan-Hölder filtrations with associated graded bundles $G_i = F_i/F_{i-1}$ and $G'_i = F'_i/F'_{i-1}$. We have $\ell = k$, and there exists a permutation $\sigma \in \text{Sym}_k$ such that $G'_i \cong G_{\sigma(i)}$.

Hence the associated graded $\operatorname{gr}_E = \bigoplus_{i=1}^k G_i$ of a semistable bundle E is well defined up to isomorphism. This allows us to make the following definition, where the S refers to Seshadri.

Definition 2.8. Semistable vector bundles E and E' of the same rank and degree are called *S*-equivalent if their associated graded vector bundles are isomorphic.

We can think of the Jordan–Hölder filtration as stating that semistable vector bundles are built out of stable vector bundles of the same slope. Meanwhile, the result below tells us that *all* vector bundles are built out of semistable bundles in a *unique* way. See again [20] for a proof. **Proposition 2.9** (Harder–Narasimhan filtration). Let F be a vector bundle on X. There exists a unique filtration of F by subbundles

 $0 = F_0 \subset F_1 \subset F_2 \subset \cdots \subset F_k = F$

such that $G_i = F_i/F_{i-1}$ is semistable for each i and

$$\mu(G_1) > \mu(G_2) > \cdots > \mu(G_k).$$

This filtration is called the Harder–Narasimhan filtration.

Example 2.10 $(X = \mathbb{P}^1)$. By a theorem of Birkhoff and Grothendieck, vector bundles on $X = \mathbb{P}^1$ split as direct sums of line bundles. Writing $E \cong \bigoplus_{i=1}^k \mathcal{O}_{\mathbb{P}^1}(d_i)^{\oplus m_i}$ with $d_1 > d_2 > \cdots > d_k$, the associated graded bundles of the Harder–Narasimhan filtration are $G_i = \mathcal{O}_{\mathbb{P}^1}(d_i)^{\oplus m_i}$ and their slopes are $\mu(G_i) = d_i$. In particular E is semistable if and only if k = 1, and stable if and only if it is a line bundle.

2.3 Openness of semistability

We can now show that semistability is an open property in families. We will use the results from [23, Tag 0DM1] on Quot, where we will denote $\operatorname{Quot}_{\mathcal{E}/X/B}^{P(t)}$ the object from [23, Tag 0DP6] for a numerical polynomial P(t), a morphism $f: X \to B$ of schemes, and a quasicoherent sheaf \mathcal{E} on X.

Proposition 2.11 (Openness of semistability). Let S be a k-scheme and \mathcal{E} be a vector bundle of rank r and relative degree d on $X \times S \to S$. The points $p \in S$ for which \mathcal{E}_p is semistable form an open subscheme.

Proof. By standard approximation techniques we can assume that S is of finite type over k. We may also assume that S is connected. We will show that the locus of points $s \in S$ for which \mathcal{E}_s is *not semistable* is closed by expressing it as a finite union of images of proper morphisms. Because our family is flat, the Hilbert polynomial $P_{\mathcal{E}_s}(t)$ is independent of $p \in S$.

By definition, \mathcal{E}_s is not semistable if and only if there exists some subsheaf $E' \hookrightarrow \mathcal{E}_s$ with $\mu(E') > d/r$. Considering the cokernel of $E' \hookrightarrow \mathcal{E}_s$, we see that \mathcal{E}_s is not semistable if and only if \mathcal{E}_s has a quotient with Hilbert polynomial $P_{\mathcal{E}_s}(t) - P_{E'}(t)$ for some E' with $\mu(E') > d/r$. In other words, the set of $s \in S$ such that \mathcal{E}_s is not semistable is the union of images of relative Quot schemes $Quot_{\mathcal{E}/X \times S/S}^{Q(t)}$ for the projection $X \times S \to S$, where Q(t) ranges over the set

$$\{Q(t) = \mathcal{P}_{\mathcal{E}_s}(t) - \mathcal{P}_{E'}(t) \mid \exists s \in S, E' \subset \mathcal{E}_s \text{ with } \mu(E') > d/r\}.$$

Relative Quot schemes are proper by [23, Tag 0DPC], so to show that the nonsemistable locus is closed, it suffices to show that the above set of polynomials is finite.

The Hilbert polynomial $P_{E'}(t)$ is determined by the rank and degree of E'. Being a subsheaf of E there are only finitely many possibilities for the rank of E'. The requirement $\mu(E') > d/r$ puts a lower bound on the degree of E'. It remains to show that the degrees of all subsheaves of \mathcal{E}_s as s varies over S are bounded above.

To see this, choose a finite morphism $X \to \mathbb{P}^1$ and consider the induced morphism $f: X \times S \to \mathbb{P}^1 \times S$. If $E' \subset \mathcal{E}_p$ is a subsheaf, then $f_*E' \subset f_*\mathcal{E}_p$ is a subsheaf.

According to Example 2.10, vector bundles on \mathbb{P}^1 split as direct sums of line bundles. Only finitely many splitting types occur among the $f_*\mathcal{E}_s$ for $s \in S$, since otherwise $h^0(\mathbb{P}^1, f_*\mathcal{E}_s)$ would achieve infinitely many values, which is impossible by semicontinuity. Moreover, for each vector bundle on \mathbb{P}^1 , the set of Euler characteristics of all its subsheaves is bounded above.

Because f is finite, $\chi(X, f_*E') = \chi(X, E')$. It follows that the Euler characteristics of all subsheaves $E' \subset \mathcal{E}_s$ are bounded above and, hence, the degrees of all subsheaves $E' \subset \mathcal{E}_s$ are bounded above.

2.4 Boundedness and irreducibility

To obtain a finite-type moduli space, the objects parameterized by the moduli problem must be bounded in the following sense.

Definition 2.12 (Boundedness). A collection of isomorphism classes of vector bundles on X is called *bounded* if there exists a scheme S of finite type and a vector bundle \mathcal{F} on $X \times S \to S$ such that every isomorphism class in the collection is represented by \mathcal{F}_s for some k-point $s \in S$.

When ranging over a bounded collection of isomorphism classes, the set of values of any discrete algebraic invariant must be finite. However, as soon as the rank is at least 2, fixing the rank and degree is not sufficient to obtain a bounded family.

Example 2.13 (An unbounded family). Let $x \in X$ be a point, and consider the collection $\{F_n\}$ of rank $r \ge 2$, degree d vector bundles

$$F_n = \mathcal{O}_X(-nx) \oplus \mathcal{O}_X((n+d)x) \oplus \mathcal{O}_X^{\oplus (r-2)}.$$

The range of values of $h^1(X, F_n)$ is unbounded in this family. Hence, the collection of all isomorphism classes of vector bundles of fixed rank $r \ge 2$ and degree d cannot be bounded.

One important source of bounded families comes from quotients of a fixed vector bundle.

Example 2.14 (Boundedness of quotients). Fix a vector bundle F on X and a polynomial $P(t) \in \mathbb{Z}[t]$. The Quot scheme $\operatorname{Quot}_{F/X/\operatorname{Spec} k}^{P(t)}$ parameterizing quotients of F with Hilbert polynomial P(t) is of finite type by [23, Tag 0DP9]. Therefore, the collection of isomorphism classes of vector bundles that appear as a quotient of F with Hilbert polynomial P(t) is bounded: restricting the universal quotient sheaf on $X \times \operatorname{Quot}_{F/X/\operatorname{Spec} k}^{P(t)}$ to the open locus where the quotient is locally free provides a family of vector bundles realizing each isomorphism class. Example 2.14 helps us see that semistable bundles of fixed rank and degree are bounded.

Proposition 2.15 (Boundedness of semistability). The collection of isomorphism classes of semistable vector bundles of rank r and degree d on X is bounded.

Proof. Let $x \in X$ be a k-point and let n be an integer such that d/r + n > 2g - 1. For every semistable vector bundle E, the vector bundle $E \otimes \mathcal{O}_X(nx)$ is semistable by Lemma 2.3 and has slope greater than 2g - 1. Hence, by Lemma 2.6, the vector bundle $E \otimes \mathcal{O}_X(nx)$ is globally generated and $h^1(X, E \otimes \mathcal{O}_X(nx)) = 0$. In other words, $E \otimes \mathcal{O}_X(nx)$ is a quotient of $V \otimes \mathcal{O}_X$ where V is a vector space of dimension $h^0(X, E \otimes \mathcal{O}_X(nx))$, and this dimension is independent of E. It follows that every semistable bundle E of rank r and degree d arises as a quotient

$$V \otimes \mathcal{O}_X(-nx) \to E.$$

Boundedness of this family now follows from Example 2.14 with $F = V \otimes \mathcal{O}_X(-nx)$ and P(t) = d + r(t+1-g).

Proposition 2.16 (Irreducibility of the moduli stack). The moduli stack $\mathcal{M}_X^{ss}(r, d)$ is irreducible. The same is true for $\mathcal{M}_X^{ss}(r, L)$.

Proof. For a closed point $x \in X(k)$, Lemma 2.3 says that E is semistable if and only if $E \otimes \mathcal{O}_X(x)$ is semistable. This gives rise to an isomorphism $\mathcal{M}_X^{ss}(r,d) \to \mathcal{M}_X^{ss}(r,d+r)$. Therefore, it suffices to prove the claim when d > r(2g-1). In other words, we may assume that the slope is at least 2g-1, and so all semistable bundles of our fixed rank and degree are globally generated by Lemma 2.6.

We will construct an irreducible variety V^{ss} that maps to $\mathcal{M}_X^{ss}(r, d)$ surjectively. Because V^{ss} is a scheme the morphism is automatically representable, and so we get an induced surjection on the level of topological spaces by [23, Tag 04XI]. But then the topological space is irreducible, and hence so is the stack. Our space V^{ss} will parameterize semistable extensions of a degree-d line bundle by a rank-(r-1) trivial bundle. For a fixed line bundle L, the extensions

$$0 \to \mathcal{O}_X^{r-1} \to E \to L \to 0 \tag{5}$$

are parameterized by $\operatorname{Ext}_X^1(L, \mathcal{O}_X^{r-1}) = \operatorname{H}^1(X, L^{\vee})^{r-1}$. To parameterize such extensions of all degree d line bundles, let \mathcal{L} be a Poincaré line bundle on $X \times \operatorname{Pic}_X^d$ and let $\pi \colon X \times \operatorname{Pic}_X^d \to \operatorname{Pic}_X^d$ be the projection. By the theorem on cohomology and base change, the fiber of the vector bundle $V = (\operatorname{R}^1 \pi_* \mathcal{L}^{\vee})^{r-1}$ over $L \in \operatorname{Pic}_X^d$ is the space of extensions $\operatorname{H}^1(X, L^{\vee})^{r-1}$, and there is a universal extension on $X \times V$. By Proposition 2.11, the locus V^{ss} where the universal extension is semistable is an open subvariety, so we obtain a map $V^{\operatorname{ss}} \to \mathcal{M}_X^{\operatorname{ss}}(r, d)$. Clearly V is irreducible, hence so is the open subvariety V^{ss} .

To show that this map is *surjective*, we must show that every semistable bundle E arises as an extension as in (5). By the first paragraph of the proof, E is

globally generated. For such an E, we claim there exists a subspace $\Gamma \subset \mathrm{H}^0(X, E)$ of dimension r-1, so that the evaluation map

$$\Gamma \otimes \mathcal{O}_X \to E \tag{6}$$

has rank r-1 at all points of X. To see this, we use a dimension count. At each point $x \in X$, we consider the map of vector spaces $\mathrm{H}^0(X, E) \to E(x)$, where E(x) denotes the fiber of E at x. We wish to select a Γ in the Grassmannian $\mathrm{Gr}(r-1, \mathrm{H}^0(X, E))$ which does not meet the kernel nontrivially. Since the kernel has codimension r, the linear spaces meeting it form a codimension-2 Schubert variety in $\mathrm{Gr}(r-1, \mathrm{H}^0(X, E))$. Varying x over X, we see that the collection of Γ 's where (6) has rank less than r-1 has codimension at least 1. Hence, there exists some Γ where (6) has rank r-1. For such Γ , we obtain an exact sequence

$$0 \to \Gamma \otimes \mathcal{O}_X \to E \to \det(E) \to 0.$$

This shows that E occurs as an extension of a degree d line bundle by a rank-(r-1) trivial bundle. Hence, E is in the image of $V^{ss} \to \mathcal{M}_X^{ss}(r, d)$.

The proof for $\mathcal{M}_X^{\mathrm{ss}}(r, L)$ is similar, using an open subscheme of $\mathrm{Ext}_X^1(L, \mathcal{O}_X^{r-1})$ surjecting onto the stack.

We summarize the results of this section as follows.

Theorem 2.17. The moduli stacks $\mathcal{M}_X^{ss}(r, d)$ and $\mathcal{M}_X^{ss}(r, L)$ are irreducible algebraic stacks, of finite type over k with affine diagonal. They are moreover smooth, and their Zariski tangent spaces are given in Proposition 1.3.

Remark 2.18. One can show that if the genus of X is at least 2, then for any integer r > 0 and line bundle $L \in \operatorname{Pic}_X$ there exist stable vector bundles of rank r and determinant L; see, e.g., [21, Lemma 4.3]. Hence the open substacks $\mathcal{M}_X^{\mathrm{s}}(r,d) \subseteq \mathcal{M}_X^{\mathrm{ss}}(r,d)$ and $\mathcal{M}_X^{\mathrm{s}}(r,L) \subseteq \mathcal{M}_X^{\mathrm{ss}}(r,L)$ of stable bundles are nonempty and dense.

3 Good moduli spaces

To study the properties of an algebraic stack, we often want to find the best possible approximation as a scheme or algebraic space. For a Deligne–Mumford stack such a procedure is given by the Keel–Mori theorem, producing a coarse moduli space. For an algebraic stack the first author introduced the notion of a good moduli space in [1] and in [2] criteria to verify its existence were established. The goal of this section is to briefly discuss how $\mathcal{M}_X^{ss}(r,d)$ fits into this general framework.

The following example explains why the stack $\mathcal{M}_X^{\mathrm{ss}}(r,d)$ is not Deligne–Mumford and the Keel–Mori theorem does not apply.

Example 3.1. The automorphism group of any semistable vector bundle contains a canonical copy of \mathbb{G}_m acting by scalars, and moreover $\operatorname{Aut}(F) = \mathbb{G}_m$ when F is a stable bundle by Lemma 2.5. In particular, the stack $\mathcal{M}_X^{ss}(r, d)$ is not Deligne–Mumford, as the automorphism groups of all points are infinite.

A prototypical example of a strictly semistable vector bundle of rank 2 and degree 0 is the polystable bundle $F = \mathcal{O}_X \oplus \mathcal{O}_X$. Now $\operatorname{Aut}(F) = \operatorname{GL}_2$, so the dimension of the automorphism group can jump up when going from the stable locus to the strictly semistable locus.

The automorphism groups do not have to be reductive either. If F fits in a non-split exact sequence

$$0 \to \mathcal{O}_X \xrightarrow{i} F \xrightarrow{q} \mathcal{O}_X \to 0,$$

then the map $(a, b) \mapsto a \cdot \mathrm{id}_F + ab \cdot i \circ q$ gives an an isomorphism $\mathbb{G}_m \times \mathbb{G}_a \cong \mathrm{Aut}(F)$. Such extensions are parameterized by the projective space

$$\mathbb{P}\mathrm{Ext}^1_X(\mathcal{O}_X,\mathcal{O}_X)\cong\mathbb{P}^{g-1}$$

Example 3.2. For $\mathcal{M}_X^{ss}(r, L)$ as defined in (2) one has generically a finite stabilizer, because for a *stable* vector bundle with fixed determinant the automorphisms need to preserve the isomorphism, and hence are isomorphic to μ_r .

If instead we took the fiber product along the closed substack $\mathbb{B}\mathbb{G}_{\mathrm{m}} \hookrightarrow \mathcal{P}\mathrm{ic}_X^d$ then the resulting closed substack would have the same stabilizers as $\mathcal{M}_X^{\mathrm{ss}}(r, d)$, and in particular they would never be finite.

Observe that, with the first definition of $\mathcal{M}_X^{ss}(r, L)$, if gcd(r, deg L) = 1then there are no strictly semistable vector bundles, and $\mathcal{M}_X^{ss}(r, L)$ is Deligne– Mumford. In this case $\mathcal{M}_X^{ss}(r, L)$ is a μ_r -gerbe over its coarse moduli space. With the second definition the stack is rather a \mathbb{G}_m -gerbe over this coarse moduli space.

Example 3.3. The moduli stack $\mathcal{M}_X^{\mathrm{ss}}(r, d)$ is *not* separated, despite its many other nice properties. Indeed, if it were separated, its diagonal would be proper. But the diagonal is affine, hence it would be finite, and so the stabilizers would be finite as they are the fibers of the diagonal. This contradicts the previous example. Likewise, the moduli stack $\mathcal{M}_X^{\mathrm{ss}}(r, L)$ is *not* separated if the strictly semistable locus is nonempty.

Definition 3.4. Let $f: \mathscr{X} \to X$ be a quasi-compact and quasi-separated morphism over k, where \mathscr{X} is an algebraic stack and X is an algebraic space. We say that f is a *good moduli space* if

- (i). f is cohomologically affine, i.e., f_* : Qcoh $\mathscr{X} \to \operatorname{Qcoh} X$ is exact;
- (ii). the morphism $\mathcal{O}_X \to f_*\mathcal{O}_{\mathscr{X}}$ is an isomorphism.

We now summarize some of the fundamental properties of good moduli spaces that will be relevant for us. See [1] for a complete statement.

Theorem 3.5. Let \mathscr{X} be an algebraic stack of finite type over k, and let $f: \mathscr{X} \to X$ be a good moduli space.

- (i). The map f is surjective, universally closed, and initial for maps to algebraic spaces.
- (ii). The space X is of finite type over k.
- (iii). The map f identifies two k-points x, y: Spec k → X if and only if the closures of {x} and {y} in |X| intersect. In particular, f induces a bijection on closed points.
- (iv). The pullback f^* induces an equivalence of categories between vector bundles on X and those vector bundles \mathcal{F} on \mathscr{X} such that the stabilizer of any closed point $x: \operatorname{Spec} k \to \mathscr{X}$ acts trivially on $x^* \mathcal{F}$.

The following existence theorem for good moduli spaces is a specialized version of [2, Theorem A], in the form in which we will apply it. The terms "S-complete" and " Θ -reductive" in the statement refer to certain valuative criteria that we will discuss below.

Theorem 3.6. Let \mathscr{X} be an algebraic stack of finite type over k with affine diagonal. There exists a good moduli space $\mathscr{X} \to X$ where X is a separated algebraic space over k if and only if

- (i). \mathscr{X} is S-complete, and
- (ii). \mathscr{X} is Θ -reductive.

Moreover, X is proper if and only if \mathscr{X} is universally closed, or equivalently satisfies the existence part of the valuative criterion for properness.

3.1 Langton's semistable reduction theorem

We first discuss the last part of Theorem 3.6. Since the stacks $\mathcal{M}_X^{\mathrm{ss}}(r, d)$ and $\mathcal{M}_X^{\mathrm{ss}}(r, L)$ are quasicompact and have affine diagonal, hence are quasiseparated, it follows from [23, Tags 0CLW and 0CLX] that universal closedness is equivalent to the existence part of the valuative criterion for properness.

The valuative criterion states the following. If R is a discrete valuation ring R with residue field κ and fraction field K, then for every 2-commutative diagram

$$\begin{array}{ccc} \operatorname{Spec} K & \longrightarrow & \mathscr{X} \\ & & & \downarrow \\ & & & \downarrow \\ & \operatorname{Spec} R & \longrightarrow & \operatorname{Spec} k \end{array}$$

there exists a field extension K'/K and a discrete valuation ring $R' \subseteq K'$ dominating R, such that there exists a morphism $\operatorname{Spec} R' \to \mathscr{X}$ making the diagram



2-commutative.

It turns out that for $\mathcal{M}_X^{ss}(r, d)$ and $\mathcal{M}_X^{ss}(r, L)$ it is not necessary to consider field extensions of K, by the following classical theorem of Langton [19].

Theorem 3.7. Let E be a semistable vector bundle of rank r and degree don X_K . There exists a vector bundle \mathcal{E} on X_R such that $\mathcal{E}_K \cong E$ and \mathcal{E}_{κ} is semistable. Thus, the stacks $\mathcal{M}_X^{ss}(r, d)$ and $\mathcal{M}_X^{ss}(r, L)$ are universally closed.

According to Example 3.3, the stack $\mathcal{M}_X^{ss}(r, d)$ is not separated, and there will in general be many extensions of E to X_R . However, for any two extensions, the central fibers will be *S*-equivalent, as we will see in Theorem 3.12, justifying the notion of S-equivalence. In particular, over the stable locus $\mathcal{M}_X^s(r, d)$ any two extensions are isomorphic, although not uniquely.

3.2 Θ -reductivity and S-completeness

We now turn to the more contemporary ingredients of Theorem 3.6. Let R denote a discrete valuation ring with field of fractions K and residue field k, and let π be a uniformizer. We define the quotient stacks

- $\Theta_R = [\operatorname{Spec} R[t]/\mathbb{G}_m]$ where the action has weight -1,
- $\overline{\operatorname{ST}}_R := [(\operatorname{Spec} R[s,t]/(st-\pi))/\mathbb{G}_m]$ where the action has weight 1 on s and -1 on t.

Both Θ -reductivity and S-completeness are lifting criteria for *codimension-2* punctures in Θ_R and \overline{ST}_R . The notion of Θ -reductivity is encoded by the lifting criterion in the left diagram, whereas S-completeness is the lifting criterion in the right diagram:

The notion of Θ -reductivity was introduced in [12] and that of S-completeness in [2, Section 3.5]. If \mathscr{X} is an algebraic stack with affine diagonal, then in either criterion, a lift is automatically unique by [2, Proposition 3.17].

Each of the two lifting criteria encodes the extension of a certain type of filtration over the puncture, which we will now describe.

3.2.1 Θ -reductivity

We let $0 \in \Theta_R$ be the unique closed point defined by the vanishing of t and the uniformizer $\pi \in R$. Observe that $\Theta_R \setminus 0 = \Theta_K \cup_{\operatorname{Spec} K} \operatorname{Spec} R$. We have the

following two cartesian diagrams which describe the geometry of Θ_R :



Here the maps on the left are open embeddings and on the right are closed embeddings.

A morphism $\rho: \Theta \to \underline{\operatorname{Coh}}_X$ corresponds to the data of a coherent sheaf E on X and a filtration $0 = E_0 \subset E_1 \subset \cdots \subset E_n = E$. Under this correspondence, $\rho(1) = E$ and $\rho(0) = \bigoplus_i E_{i+1}/E_i$ is the associated graded. The map $\rho: \Theta \to \underline{\operatorname{Coh}}_X$ factors through $\mathcal{M}_X(r,d)$ (resp. $\mathcal{M}_X^{\mathrm{ss}}(r,d)$) if and only if E is a vector bundle (resp. semistable vector bundle) of rank r and degree d and each factor E_{i+1}/E_i is a vector bundle (resp. semistable vector bundle). There are analogous descriptions for morphisms from Θ_A for a k-algebra A.

Under this correspondence, the Θ -reductivity of these stacks translates into a valuative criterion for filtrations. For $\mathcal{M}_X^{ss}(r,d)$, Θ -reductivity translates into the following: for every discrete valuation ring R with fraction field Kand semistable vector bundle E on X_R of rank r and degree d, any filtration $0 = G_0 \subset G_1 \subset \cdots \subset G_n = E_K$ of the generic fiber where each factor G_{i+1}/G_i is semistable of slope $\mu = d/r$ extends to a filtration $0 = E_0 \subset E_1 \subset \cdots \subset E_n = E$ where each factor E_{i+1}/E_i is a family of semistable vector bundles on X over R. Note that the factors G_{i+1}/G_i are semistable of slope d/r for all i if and only if G_i is semistable of slope d/r for all i if and only if $\mu(G_i) = d/r$ for all i.

Proposition 3.8. The stacks \underline{Coh}_X , $\mathcal{M}_X^{ss}(r, d)$, and $\mathcal{M}_X^{ss}(r, L)$ are Θ -reductive.

Proof. First consider \underline{Coh}_X . Let E be an R-flat coherent sheaf on X_R and let $0 = G_0 \subset G_1 \subset \cdots \subset G_n = E_K$ be a filtration of the generic fiber. By descending induction starting with j = n, suppose that we have constructed a filtration $E_j \subset E_{j+1} \subset \cdots \subset E_n = E$ extending G_{\bullet} . Since the relative Quot scheme of quotients of E_j with the same Hilbert polynomial as G_j/G_{j-1} is proper over R, there is a unique subsheaf $E_{j-1} \subset E_j$ with $(E_{j-1})_K = G_{j-1}$ and E_j/E_{j-1} flat over R. This gives the next step in the filtration E_{\bullet} .

To see that $\mathcal{M}_X^{ss}(r, d)$ is Θ -reductive, suppose that E is a family of semistable vector bundles on X_R and G_{\bullet} is a filtration of E_K such that the sheaves G_i are semistable vector bundles with $\mu(G_i) = d/r$. By the previous paragraph, G_{\bullet} extends to E_{\bullet} over R, and we must show that in the sheaves $(E_i)_k$ are semistable as well. Since each E_i is flat over R, we have $\mu((E_i)_k) = \mu(G_i) = d/r$, and so $(E_i)_k$ is a subsheaf of the semistable vector bundle E_k of the same slope, hence a semistable vector bundle itself. The same argument shows that $\mathcal{M}_X^{ss}(r, L)$ is Θ -reductive.

3.2.2 S-completeness

The stack $\overline{\mathrm{ST}}_R$ can be viewed as a local model of the quotient $[\mathbb{A}^2/\mathbb{G}_m]$ where \mathbb{A}^2 has coordinates s and t with weights 1 and -1; indeed, $\overline{\mathrm{ST}}_R$ is the base change of the good moduli space $[\mathbb{A}^2/\mathbb{G}_m] \to \operatorname{Spec} k[st]$ along $\operatorname{Spec} R \to \operatorname{Spec} k[st]$ defined by $st \mapsto \pi$.

We let $0 \in \overline{\mathrm{ST}}_R$ be the unique closed point defined by the vanishing of s and t. Observe that $\overline{\mathrm{ST}}_R \setminus 0$ is the non-separated union $\operatorname{Spec} R \cup_{\operatorname{Spec} K} \operatorname{Spec} R$. We have the following two cartesian diagrams which describe the geometry of $\overline{\mathrm{ST}}_R$:



Here the maps on the left are open embeddings and those on the right are closed embeddings.

Remark 3.9. If G is a linear algebraic group over k, then BG is S-complete if and only if G is reductive (see [2, Proposition 3.45 and Remark 3.46]). Moreover, as S-completeness is preserved under closed substacks, it follows that every closed point (corresponding to a polystable object) in an S-complete algebraic stack with affine diagonal has a *reductive stabilizer*.

We will see in Lemma 3.11 below that the closed points of $\mathcal{M}_X^{ss}(r,d)$ correspond to polystable vector bundles, so although some k-points of $\mathcal{M}_X^{ss}(r,d)$ may have nonreductive stabilizers as in Example 3.1, such points necessarily correspond to non-polystable bundles.

Proposition 3.10. The stacks \underline{Coh}_X , $\mathcal{M}_X^{ss}(r, d)$, and $\mathcal{M}_X^{ss}(r, L)$ are S-complete.

We omit the proof of this proposition, which can be proven along the same lines as Proposition 3.8 using the interpretation of morphisms from $\overline{\mathrm{ST}}_R$ to $\mathcal{M}_X^{\mathrm{ss}}(r,d)$ from [2, Remark 3.36].

We have summarized in Table 1 the properties of $\mathcal{M}_X^{ss}(r,d)$, $\mathcal{M}_X(r,d)$, and \underline{Coh}_X discussed in the first three sections.

3.3 Existence of good moduli spaces

In this subsection we will establish the existence and basic properties of good moduli spaces for the stacks $\mathcal{M}_X^{ss}(r,d)$ and $\mathcal{M}_X^{ss}(r,L)$, but before doing so we make an observation regarding their topology.

Lemma 3.11.

 (i). If E is a semistable vector bundle, then the corresponding k-point [E] ∈ M^{ss}_X(r,d)(k) contains the point [gr_E] in its closure, where gr_E is the graded bundle associated to a Jordan–Hölder filtration.

Property	$\mathcal{M}_X^{\mathrm{ss}}(r,d)$	$\mathcal{M}_X(r,d)$	$\underline{\mathrm{Coh}}_X$
affine diagonal	yes	yes, Proposition 1.1	yes, [23, Tag 0DLY]
locally of finite type	yes	yes, Proposition 1.1	yes, [23, Tag 0DLZ]
quasicompact	yes, Proposition 2.15	no, Example 2.13	no
Θ -reductive	yes, Proposition 3.8	no	yes, Proposition 3.8
S-complete	yes, Proposition 3.10	no	yes, Proposition 3.10
existence part of valuative criterion	yes	yes	yes
separated	no, Example 3.3	no	no
existence of good moduli space	yes, Theorem 3.12	no	no

Table 1: Properties of the stacks $\mathcal{M}_X^{ss}(r,d) \subset \mathcal{M}_X(r,d) \subset \underline{\mathrm{Coh}}_X$

(ii). A point $[E] \in \mathcal{M}_X^{ss}(r,d)(k)$ is closed if and only if E is polystable.

The same holds for the stack $\mathcal{M}_X^{ss}(r, L)$.

Proof. (i). If E is semistable but not stable, there exists a non-split extension $0 \to E' \to E \to E'' \to 0$ of semistable vector bundles of the same slope. Let \mathcal{E} be the universal family over the affine line in $\operatorname{Ext}^1_X(E'', E')$ spanned by this extension, so that \mathcal{E} is a family of semistable vector bundles on X parameterized by \mathbb{A}^1 such that $\mathcal{E}_t \cong E$ if $t \neq 0$ and $\mathcal{E}_0 \cong E' \oplus E''$. The resulting map

$$[\mathcal{E}]: \mathbb{A}^1 \to \mathcal{M}_X^{\mathrm{ss}}(r, d)$$

satisfies $t \mapsto [E]$ if $t \neq 0$ and $0 \mapsto [E' \oplus E'']$. It follows that $[E' \oplus E'']$ is contained in the closure of [E]. Iterating this construction for E' and E'' shows that $[\operatorname{gr}_E]$ is in the closure of [E].

(ii). For a contradiction, suppose E is a polystable vector bundle such that [E] is not closed, and let [F] be in its closure. By (i), $[gr_F]$ is in the closure of [F] and, since no two points can be in the closure of each other, we must have $E \not\cong gr_F$.

On the other hand, if E_i is stable with the same slope as E, then E_i appears as a direct summand of E with multiplicity $\hom_X(E_i, E) = \dim_k \operatorname{Hom}_X(E_i, E)$ and similarly for gr_F .

For any E_i , the function $\hom_X(E_i, -)$ is upper semicontinuous in the second variable, since $[\operatorname{gr}_F]$ is in the closure of [E], so we have $\hom_X(E_i, E) \leq$

 $\hom_X(E_i, \operatorname{gr}_F)$. This means that any stable summand of E appears in gr_F with at least the same multiplicity. But E and F have the same rank, so we must have $E \cong \operatorname{gr}_F$, a contradiction. Thus, [E] is closed.

Theorem 3.12.

(i). There exist good moduli spaces

$$\mathcal{M}_X^{\mathrm{ss}}(r,d) \to \mathrm{M}_X^{\mathrm{ss}}(r,d)$$
$$\mathcal{M}_X^{\mathrm{ss}}(r,L) \to \mathrm{M}_X^{\mathrm{ss}}(r,L).$$

- (ii). These good moduli space maps induce bijections between the k-points of the good moduli space and S-equivalence classes of semistable sheaves.
- (iii). The good moduli spaces $M_X^{ss}(r, d)$ and $M_X^{ss}(r, L)$ are irreducible, proper algebraic spaces of dimensions

$$\dim \mathcal{M}_X^{ss}(r,d) = r^2(g-1) + 1,$$
$$\dim \mathcal{M}_X^{ss}(r,L) = (r^2 - 1)(g-1).$$

Proof.

- (i). This follows from Propositions 3.8 and 3.10 and the first part of Theorem 3.6.
- (ii). By Theorem 3.5(iii), two k-points $[E], [E'] \in \mathcal{M}_X^{ss}(r, d)$ map to the same point in $\mathcal{M}_X^{ss}(r, d)$ if and only if the closures of $\{[E]\}$ and $\{[E']\}$ in $\mathcal{M}_X^{ss}(r, d)$ intersect. On the one hand, if E is any semistable vector bundle, then by Lemma 3.11(i) [E] contains $[gr_E]$ in its closure, so both points map to the same point in $\mathcal{M}_X^{ss}(r, d)$. On the other hand, if E and E' are polystable and nonisomorphic, then by Lemma 3.11(ii), the corresponding points in $\mathcal{M}_X^{ss}(r, d)$ are closed and distinct, hence map to distinct points in $\mathcal{M}_X^{ss}(r, d)$.
- (iii). The stacks are irreducible by Proposition 2.16 and the good moduli space maps are surjective by Theorem 3.5(i), so the good moduli spaces are irreducible as well. They are proper by Proposition 3.7 and the second part of Theorem 3.6.

To compute the dimensions, one can show using (ii) and Lemma 2.5 that over the stable locus $\mathcal{M}_X^{s}(r,d) \subseteq \mathcal{M}_X^{ss}(r,d)$ the good moduli space map is a \mathbb{G}_m -gerbe over its image $\mathrm{M}_X^{s}(r,d)$, and similarly for $\mathcal{M}_X^{ss}(r,L)$. Since the stable loci are moreover nonempty by Remark 2.18, the dimensions now follow from Proposition 1.3 and Remark 1.4.

4 Determinantal line bundles

Let X be a smooth, projective and connected curve over k. For an algebraic stack $\mathscr S$ over k we consider the diagram



In this setting we can freely apply cohomology and base change arguments, because p is representable by schemes. If \mathcal{E} is a vector bundle on $X \times \mathscr{S}$, then the derived direct image $\mathbf{R}p_*\mathcal{E}$ is a perfect complex on \mathscr{S} with amplitude in [0,1]. In fact, something stronger is true. We will use the construction of [14, Proposition 2.1.10]. Let \mathcal{E} be a vector bundle on $X \times \mathcal{M}_X(r, d)$. Then there exists a short exact sequence

$$0 \to \mathcal{E}^{-1} \to \mathcal{E}^0 \to \mathcal{E} \to 0$$

of vector bundles such that $\mathbb{R}^0 p_* \mathcal{E}^j = 0$ and $K^j := \mathbb{R}^1 \mathcal{E}^{j+1}$ is locally free for j = 0, 1. In particular, we have a quasi-isomorphism $K^{\bullet} := [K^0 \to K^1] \to \mathbb{R} p_* \mathcal{E}$. We make the following definition.

Definition 4.1 (Determinants and sections). If \mathcal{E} is a vector bundle on $X \times \mathscr{S}$ and $[K^0 \to K^1]$ is a two-term complex of locally free sheaves such that $\mathbf{R}p_*\mathcal{E} \sim [K^0 \to K^1]$, we define the line bundle

$$\det \mathbf{R} p_* \mathcal{E} := \det(K^0) \otimes \det(K^1)^{\vee}.$$

If $\operatorname{rk}(\mathbf{R}p_*\mathcal{E}) = 0$, then $\operatorname{rk} K^0 = \operatorname{rk} K^1$ and the dual $(\det \mathbf{R}p_*\mathcal{E})^{\vee}$ is equipped with a section, locally given by the determinant of the map $K^0 \to K^1$.

As described in [23, Tag 0FJI], the definition of det $\mathbf{R}p_*\mathcal{E}$ is independent of the choice of quasi-isomorphism with a two-term complex.

We will apply this construction in the case $\mathscr{S} = \mathcal{M}_X(r, d)$ and $\mathcal{E} = \mathcal{E}_{\text{univ}} \otimes q^* V$, where $\mathcal{E}_{\text{univ}}$ is the universal vector bundle on $X \times \mathcal{M}_X(r, d)$ and V is a vector bundle on X. The universal vector bundle exists tautologically because we are working with the moduli stack. Consider the diagram



Proposition 4.2 (Determinant is multiplicative). If

$$0 \to \mathcal{E}' \to \mathcal{E} \to \mathcal{E}'' \to 0,$$

is an exact sequence of vector bundles on $X \times \mathcal{M}_X(r, d)$, then

$$\det \mathbf{R} p_* \mathcal{E} = (\det \mathbf{R} p_* \mathcal{E}') \otimes (\det \mathbf{R} p_* \mathcal{E}'')$$

Proof. As before, we can find a two-term complex $\mathcal{E}^{\bullet} = [\mathcal{E}^{-1} \to \mathcal{E}^{0}]$ supported in degrees [-1, 0] and a quasi-isomorphism $\mathcal{E}^{\bullet} \to \mathcal{E}$ with the properties described above. We choose \mathcal{E}'^{\bullet} and \mathcal{E}''^{\bullet} similarly. In fact, it follows from the construction in the proof of [14, Proposition 2.1.10] that we may choose these resolutions compatibly, so that we have a short exact sequence

$$0 \to \mathcal{E}^{\prime \bullet} \to \mathcal{E}^{\bullet} \to \mathcal{E}^{\prime \prime \bullet} \to 0$$

of complexes, compatible with the quasi-isomorphisms to the members in our original exact sequence. Taking cohomology, we find a short exact sequence

$$0 \to K'^{\bullet} \to K^{\bullet} \to K''^{\bullet} \to 0$$

of complexes of locally free sheaves on $\mathcal{M}_X(r,d)$. The result follows from the multiplicativity of the determinant in short exact sequences of locally free sheaves.

Definition 4.3 (Determinantal line bundles and sections). For a vector bundle V on X, we define the *determinantal line bundle*

$$\mathcal{L}_V := (\det \mathbf{R} p_*(q^*V \otimes \mathcal{E}_{\mathrm{univ}}))^{\vee}$$

on $\mathcal{M}_X(r,d)$ associated to V. If $\chi(X, V \otimes E) = 0$ for all $[E] \in \mathcal{M}_X(r,d)$, or equivalently by the Riemann–Roch theorem

$$d \operatorname{rk}(V) + r \operatorname{deg}(V) + r \operatorname{rk}(V)(1 - g) = 0,$$
(10)

then the rank of $\mathbf{R}p_*(q^*V \otimes \mathcal{E}_{univ})$ is zero and we define the section $s_V \in \Gamma(\mathcal{M}_X(r,d),\mathcal{L}_V)$ as in Definition 4.1.

Remark 4.4. Since $\mathbf{R}p_*(q^*V \otimes \mathcal{E}_{univ})$ is perfect, its construction commutes with base change. In particular, its restriction to a k-point $[E] \in \mathcal{M}_X(r, d)$ is identified with the two-term complex $\mathbf{R}\Gamma(X, E \otimes V)$. If moreover $\chi(X, V \otimes E) =$ $h^0(X, V \otimes E) - h^1(X, V \otimes E) = 0$, we see that indeed $\operatorname{rk}(\mathbf{R}p_*(q^*V \otimes \mathcal{E}_{univ})) = 0$.

In order to state some basic properties of the construction in Definition 4.3, we recall that, by the universal property of the Picard stack $\mathcal{P}ic_X^d$, the determinant $\det(\mathcal{E}_{univ})$ of the universal vector bundle on $X \times \mathcal{M}_X(r, d)$ induces a morphism

$$\det\colon \mathcal{M}_X(r,d)\to \mathcal{P}\mathrm{ic}_X^d$$

such that $(\det \times \operatorname{id}_X)^* \mathscr{P} \cong \det(\mathcal{E}_{\operatorname{univ}})$, where \mathscr{P} is the Poincaré bundle on $\operatorname{Pic}_X^d \times X$.

Proposition 4.5 (Properties of the determinantal line bundle). *The following hold:*

(i). The assignment $V \mapsto \mathcal{L}_V$ induces a group homomorphism

$$\mathrm{K}_{0}(X) \to \mathrm{Pic}(\mathcal{M}_{X}(r,d)).$$
 (11)

Consequently, the isomorphism class of the line bundle \mathcal{L}_V depends only on $\operatorname{rk}(V)$ and $\det(V)$.

(ii). If V and W are vector bundles with equal rank and degree, then there exists a line bundle \mathcal{N} on Pic_X^d such that

$$\mathcal{L}_W \cong \mathcal{L}_V \otimes \det^* \mathcal{N}.$$

Proof. An exact sequence $0 \to V_1 \to V_2 \to V_3 \to 0$ of vector bundles on X induces an exact sequence of vector bundles

$$0 \to q^* V_1 \otimes \mathcal{E}_{\text{univ}} \to q^* V_2 \otimes \mathcal{E}_{\text{univ}} \to q^* V_3 \otimes \mathcal{E}_{\text{univ}} \to 0,$$

which by Proposition 4.2 induces an isomorphism

$$\mathcal{L}_{V_2}\cong \mathcal{L}_{V_1}\otimes \mathcal{L}_{V_3}.$$

Thus, we have a group homomorphism $K_0(X) \to \operatorname{Pic}(\mathcal{M}_X(r,d))$. The second statement follows from the isomorphism $K_0(X) \cong \mathbb{Z} \otimes \operatorname{Pic}(X)$ given by $V \mapsto (\operatorname{rk}(V), \det(V))$. This proves (i).

By (i), the isomorphism type of \mathcal{L}_V depends only on the rank r_V and determinant of V, so we may assume that $V = \mathcal{O}_X^{\oplus r_V - 1} \oplus \mathcal{O}_X(D)$ for some divisor D on X, and similarly for W. Moreover, writing $D = D_1 - D_2$ as a difference of effective divisors and using standard exact sequences, we see that $[\mathcal{O}_X(D)] = [\mathcal{O}_X] + [\mathcal{O}_{D_1}] - [\mathcal{O}_{D_2}]$ in $K_0(X)$. This implies that the classes of Vand W in $K_0(X)$ differ only by the class of a divisor of degree 0. Thus, by the additivity of the determinantal construction, it suffices to prove that

$$\det \mathbf{R}p_*(\mathcal{E}_{\mathrm{univ}} \otimes q^*\mathcal{O}_x) \cong \det^*\mathcal{N}$$

for some line bundle \mathcal{N}' on $\mathcal{P}ic_X^d$, where $x \in X$ is a closed point. If we view $\mathcal{E}_x = \mathcal{E}_{univ}|_{\mathcal{M}_X^{ss}(r,d) \times \{x\}}$ and $\mathcal{P}_x = \mathcal{P}|_{\mathcal{P}ic_X^d \times \{x\}}$ as sheaves on $\mathcal{M}_X^{ss}(r,d)$ and $\mathcal{P}ic_X^d$ respectively, then we have

$$\det \mathbf{R}p_*(\mathcal{E}_{\mathrm{univ}} \otimes q^*\mathcal{O}_x) = \det \mathcal{E}_x = \det^* \mathcal{P}_x.$$

This proves (ii).

Definition 4.6. Let *E* be a vector bundle on *X*. We say that *E* is *cohomology-free* if $h^0(X, E) = h^1(X, E) = 0$.

The following proposition relates cohomology-freeness to the properties of sections of determinantal line bundles. It will be an essential tool in the ampleness proof.

Proposition 4.7. If $\chi(X, E \otimes V) = 0$ for all $[E] \in \mathcal{M}_X(r, d)$, then the following are equivalent:

- the section $s_V \in \Gamma(\mathcal{M}_X(r,d),\mathcal{L}_V)$ is nonzero at a vector bundle $[E] \in \mathcal{M}_X(r,d);$
- $E \otimes V$ is cohomology-free.

Proof. In the setup of Definition 4.3 the morphism of line bundles det $j: \det(K_0) \to \det(K_1)$ is nonzero at the point $[E] \in \mathcal{M}_X(r, d)$ if and only if the morphism of vector bundles $j: K_0 \to K_1$ is an isomorphism at [E]. Since the derived direct image sheaf is the cohomology of this complex, this occurs if and only if $h^0(X, E \otimes V) = h^1(X, E \otimes V) = 0$.

Remark 4.8. We emphasize that while \mathcal{L}_V only depends on rk(V) and det(V), the section s_V does depend on V itself. We will leverage this fact to produce enough sections of \mathcal{L}_V to establish ampleness. Notice also that, under the assumption $\chi(X, E \otimes V) = 0$, the vanishing of $H^0(X, E \otimes V)$ is equivalent to the vanishing of $H^1(X, E \otimes V)$.

We now specialize the construction to the open substack $\mathcal{M}_X^{ss}(r,d) \subseteq \mathcal{M}_X(r,d)$. As explained in the introduction, our goal is to prove that the proper algebraic space $\mathcal{M}_X^{ss}(r,d)$ obtained in Theorem 3.12 is actually a projective scheme. For this we need to produce an ample line bundle on it, and the determinantal line bundle we have constructed a priori lives on $\mathcal{M}_X^{ss}(r,d)$. To remedy this with the following.

Proposition 4.9. The determinantal line bundle \mathcal{L}_V of Proposition 4.7 associated to a vector bundle V descends to $M_X^{ss}(r,d)$, that is, there exists a unique line bundle $L_V \in \text{Pic}(M_X^{ss}(r,d))$ such that $\mathcal{L}_V \cong \phi^* L_V$.

Proof. By Theorem 3.5(iv), we must show that stabilizers of $\mathcal{M}_X^{ss}(r,d)$ act trivially on the fibers of \mathcal{L}_V . By Theorem 3.12(ii), the closed points of $\mathcal{M}_X^{ss}(r,d)$ correspond to polystable bundles, and for a polystable bundle $E = \bigoplus_{j=1}^n E_j^{\oplus m_j}$, where the E_i are pairwise nonisomorphic, we have $\operatorname{Aut}(E) \cong \operatorname{GL}_{m_1} \times \cdots \times \operatorname{GL}_{m_n}$. The fiber of $\mathcal{L}_V|_{[E]}$ is identified with

$$\det \mathbf{R}\Gamma(X, E \otimes V) = \prod_{i=1}^{n} (\det \mathrm{H}^{i}(X, E \otimes V))^{\otimes (-1)^{i}}$$

An element $(g_1, \ldots, g_n) \in \operatorname{Aut}(E)$ acts on

$$\det \mathrm{H}^{i}(X, E \otimes V) \cong \bigotimes_{j=1}^{n} \left(\det \mathrm{H}^{i}(X, E_{j} \otimes V) \right)^{\otimes m_{j}}$$

by multiplication with $\prod_{j=1}^{n} \det(g_1)^{\dim \operatorname{H}^i(X, E_j \otimes V)}$, and thus on $\mathcal{L}_V|_{[E]}$ by multiplication with $\prod_{j=1}^{n} \det(g_j)^{\chi(X, E_j \otimes V)}$. But each E_j has slope equal to d/r, so by the assumption on V we have $\chi(X, E_j \otimes V) = 0$, and so the action is trivial. \Box

Corollary 4.10. For a line bundle L of degree d on X, the restriction of the determinantal line bundle L_V to $M_X^{ss}(r, L)$ only depends on the rank and degree of V.

Proof. We have commuting diagrams

$$\begin{aligned} \mathcal{M}_X^{\mathrm{ss}}(r,L) & \longrightarrow \mathcal{M}_X^{\mathrm{ss}}(r,d) & \qquad \mathcal{M}_X^{\mathrm{ss}}(r,L) & \longrightarrow \mathcal{M}_X^{\mathrm{ss}}(r,d) \\ & \downarrow & \Box & \downarrow_{\mathrm{det}} & \mathrm{and} & \downarrow & \downarrow \\ & \mathrm{Spec}\,k & \stackrel{[L]}{\longrightarrow} \mathcal{P}\mathrm{ic}_X^d & \qquad \mathrm{M}_X^{\mathrm{ss}}(r,L) & \longleftrightarrow & \mathrm{M}_X^{\mathrm{ss}}(r,d). \end{aligned}$$

where in the second square, both vertical arrows are good moduli space maps. If V and W are vector bundles of equal rank and degree on X, both satisfying condition (10), then by Proposition 4.7(ii), there exists a line bundle \mathcal{N} on $\mathcal{P}ic_X^d$ such that $\mathcal{L}_W \cong \mathcal{L}_V \otimes \det^* \mathcal{N}$. The left diagram shows that, restricting to $\mathcal{M}_X^{ss}(r, L)$, this isomorphism becomes $\mathcal{L}_W \cong \mathcal{L}_V$. The right diagram now shows that the restrictions of \mathcal{L}_V and \mathcal{L}_W to $\mathcal{M}_X^{ss}(r, L)$ become isomorphic after pulling back to $\mathcal{M}_X^{ss}(r, L)$, so the restrictions must be isomorphic to begin with, by Theorem 3.5(iv).

Using geometric invariant theory, Drézet–Narasimhan gave in [8, Théorèmes B, C] the following description of the Picard groups of the good moduli spaces.

Theorem 4.11 (Drézet-Narasimhan). There exist isomorphisms

$$\operatorname{Pic}(\operatorname{M}_{X}^{\operatorname{ss}}(r,d)) \cong \operatorname{Pic}(\operatorname{Pic}_{X}^{d}) \oplus \mathbb{Z},$$

$$\operatorname{Pic}(\operatorname{M}_{X}^{\operatorname{ss}}(r,L)) \cong \mathbb{Z}.$$
(12)

In the second line \mathbb{Z} is generated by the determinantal line bundle L_V , where V is chosen to be of minimal rank.

This explains how the isomorphism type of the line bundle L_V on the good moduli space depends on the invariants of the vector bundle V satisfying $\chi(X, E \otimes V) = 0$, building upon Proposition 4.5.

5 Projectivity

In this section we will prove the following result.

Theorem 5.1. Let X be a smooth projective curve of genus $g \ge 2$. The good moduli spaces $M_X^{ss}(r, d)$ and $M_X^{ss}(r, L)$ are projective varieties of dimension $r^2(g-1)+1$ and $(r^2-1)(g-1)$ respectively.

We begin by reducing the projectivity of $M_X^{ss}(r, d)$ to that of $M_X^{ss}(r, L)$. The determinant morphism

$$\det\colon \mathcal{M}_X^{\mathrm{ss}}(r,d)\to \mathcal{P}\mathrm{ic}_X^d$$

discussed above descends to a map

det:
$$M_X^{ss}(r, d) \to \operatorname{Pic}_X^d$$

of good moduli spaces whose fiber over $[L] \in \operatorname{Pic}_X^d$ is the moduli space $\operatorname{M}_X^{\operatorname{ss}}(r, L)$. It is a classical fact that Pic_X^d is a projective variety isomorphic to the Jacobian of X. Moreover, since $\operatorname{M}_X^{\operatorname{ss}}(r, d)$ is a proper algebraic space, to conclude that it is a projective scheme, it suffices by [23, Tag 0D36] to produce a line bundle on it whose restriction to each fiber $M_X^{ss}(r, L)$ is ample. We will show that the determinantal line bundle L_V will work for an arbitrary vector bundle Vsatisfying condition (10) below.

From now on, we will fix a line bundle L of degree d on X and focus on the moduli space $M_X^{ss}(r, L)$.

Setup Denote h = gcd(r, d) and set $r_1 = r/h$ and $d_1 = d/h$. Let V be a vector bundle on X, with invariants

$$rk(V) = mr_1, \quad deg(V) = mr_1(g-1) - md_1$$
 (13)

for some $m \geq 1$. By the Riemann–Roch theorem this ensures that $\chi(X, V \otimes E) = 0$ for a vector bundle E of rank r and degree d. Hence by Section 4 the associated determinantal line bundle L_V on $M_X^{ss}(r, L)$ depends only on $\operatorname{rk}(V)$ and $\operatorname{deg}(V)$, and comes with a section s_V , depending on V, such that $s_V([E]) \neq 0$ if and only if $V \otimes E$ is cohomology-free.

We will prove the theorem using the following two propositions.

Proposition 5.2. Let E be a stable vector bundle of rank r and degree d. For large enough m, a general vector bundle V with

$$rk(V) = mr_1, \quad deg(V) = mr_1(g-1) - md_1$$

satisfies

$$\mathrm{H}^{0}(X, V \otimes E) = \mathrm{H}^{1}(X, V \otimes E) = 0.$$
(14)

In particular, the determinantal line bundle L_V on $M_X^{ss}(r, L)$ is semiample.

Remark 5.3. In the approach of Faltings this statement (albeit for $M_X^{ss}(r, d)$) corresponds to [10, Theorem I.2] (note that op. cit. is concerned about the more complicated notion of stable Higgs bundles) and [22, Lemma 3.1], but the method of proof will be different.

Proposition 5.4. Let E and E' be polystable vector bundles of rank r and degree d such that E has a stable summand that is not a summand of E'. There exists an integer m > 0 and a stable vector bundle V of rank mr_1 and degree $mr_1(g-1) - md_1$ such that

$$\mathrm{H}^{0}(X, V \otimes E) \neq 0, \quad \mathrm{H}^{0}(X, V \otimes E') = 0.$$

$$(15)$$

In particular, the sections s_V of the determinantal line bundle L_V on $M_X^{ss}(r, L)$ separate points determined by polystable bundles for which at least one stable summand is different.

Remark 5.5. It is not true that for any two polystable bundles E and E' we can find a vector bundle V of the appropriate rank and degree for which $\operatorname{H}^0(X, V \otimes E) \neq 0$ and $\operatorname{H}^0(X, V \otimes E') = 0$. Namely, consider two polystable vector bundles

$$E = E_1 \oplus E_1 \oplus E_2$$
 and $E' = E_1 \oplus E_2 \oplus E_2$,

where E_1 and E_2 are nonisomorphic stable vector bundles of the same rank and degree. However, it can be shown that sections of the form s_V span the space of global sections of L_V for $m \gg 0$ and induce a closed embedding $M_X^{ss}(r, d) \hookrightarrow \mathbb{P}^N$; see [3, Corollary 7.15].

Remark 5.6. The method of Faltings (see [10, Theorem I.4] and [22, Lemma 4.2]) takes a different approach at this point. He shows that, given a map $C \to \mathcal{M}_X^{ss}(r,d)$ from a smooth projective curve C corresponding to a family \mathcal{E} of semistable vector bundles on X, the degree of the determinantal line bundle \mathcal{L}_V is nonnegative and is zero if and only if all the fibers $\{\mathcal{E}_t\}_{t\in C}$ are S-equivalent. This allows one to conclude that the morphism $M_X^{ss}(r,d) \to \mathbb{P}^N$ has finite fibers, and then the argument can proceed as in the proof of Theorem 5.1. Interestingly, Faltings's argument uses stability of vector bundles on the curve C.

Proof of Theorem 5.1. By Proposition 5.2 the determinantal line bundle \mathcal{L}_V associated to the appropriate choice of V is semiample, so choose some power for which it is basepoint-free and obtain a morphism $\phi: \mathrm{M}^{\mathrm{ss}}_X(r, d) \to \mathbb{P}^N$ for some N.

By Proposition 5.4 the morphism is actually quasi-finite. Indeed, for a given polystable bundle $E = E_1^{\oplus r_1} \oplus \cdots \oplus E_s^{\oplus r_s}$ where the E_1, \ldots, E_s are nonisomorphic stable bundles, there are only finitely many other polystable bundles whose stable summands are precisely E_1, \ldots, E_s , and for any other polystable bundle E', there exists a section s_V separating E and E'.

Since $M_X^{ss}(r, d)$ is a proper algebraic space, the morphism $M_X^{ss}(r, d) \to \mathbb{P}^N$ is proper by [23, Tag 04NX] and hence finite by [23, Tag 0A4X]. But then it is also affine, hence representable, so $M_X^{ss}(r, d)$ is a scheme. Finally, pulling back an ample line bundle on \mathbb{P}^N along the affine morphism produces an ample line bundle on $M_X^{ss}(r, d)$, by [23, Tag 0892].

We are left with proving Propositions 5.2 and 5.4. Their proofs will be based on a dimension-counting argument inspired by [9] (see also [18, Section 5] for similar methods), where the authors actually prove a more refined result that yields an effective bound on the rank of V in the theorem. To get started, we will need the following lemma, discussed in [4, Remark 4.2].

Lemma 5.7. If $\{F_{\alpha}\}_{\alpha \in \Lambda}$ is a bounded family of vector bundles of rank r on X, there exists a scheme S of finite type over k of dimension at most $r^2(g-1)+1$ and a vector bundle \mathcal{F} on $S \times X$ such that each F_{α} appears as the fiber of \mathcal{F} over some point $s \in S$.

In fact, the stable bundles of rank r and degree d can be parameterized by a smooth variety of dimension $r^2(g-1)+1$, and any bounded family of non-stable bundles by a variety of dimension at most $r^2(g-1)$.

The following lemma will be the technical heart for the proof of both propositions. We will apply it for $\epsilon = -1, 0, 1$, and think of it as a "fudge factor" perturbing the equality $\chi(X, E \otimes V) = 0$ for $\epsilon = 0$.

Lemma 5.8. Let E be a vector bundle of rank r and degree d on X, and let ϵ be an integer. Let m be a sufficiently large integer and V a general stable vector

bundle with

$$rk(V) = mr_1, \quad deg(V) = mr_1(g-1) - md_1 + \epsilon.$$

(i). There exist no maps $V^{\vee} \to E$ whose image F has slope $\mu(F) < \mu(E)$.

(ii). If moreover E is stable and $\epsilon \leq 0$, there exist no nonzero maps $V^{\vee} \to E$.

Proof. We can assume that rk(V) > r. Note that V^{\vee} satisfies

$$\operatorname{rk}(V^{\vee}) = mr_1, \quad \deg(V^{\vee}) = mr_1(1-g) + md_1 - \epsilon.$$

If $V^{\vee} \to E$ is a map with image F, then, since V^{\vee} is stable, we must have

$$\mu(F)>\mu(V^{\vee})=\mu(E)-g+1-\frac{\epsilon}{mr_1}\geq \mu(E)-g+1-\epsilon.$$

Thus, there are only finitely many options for the slope of the image F such that $\mu(F) \leq \mu(E)$. Since moreover we must have $1 \leq \operatorname{rk}(F) \leq r$, there are only finitely many options for the rank and degree of F.

To prove (i), fix integers $1 \le k \le r$ and l with $l/k < \mu(E)$. We will find an upper bound for the dimension of a variety that parameterizes all stable bundles V such that there exists a map $V^{\vee} \to E$ whose image F has rank k and degree l. Any such V^{\vee} fits into an exact sequence

where G satisfies

$$\operatorname{rk}(G) = mr_1 - k, \quad \deg(G) = mr_1(1 - g) + md_1 - \epsilon - l.$$

The possible sheaves F that can appear in (16) form a bounded family since they are all subsheaves of E of fixed rank and degree. By Lemma 5.7 they are parameterized by a scheme U_1 of finite type over k with

$$\dim \mathcal{U}_1 \le k^2(g-1) + 1.$$

Similarly, since the family of stable bundles of fixed rank and degree is bounded, the sheaves G that can appear in (16) form a bounded family and so by Lemma 5.7 are parameterized by a scheme U_2 with

$$\dim \mathcal{U}_2 \le (mr_1 - k)^2 (g - 1) + 1.$$

Note that since V^{\vee} in (16) is stable, we must have $\operatorname{Hom}_X(F, G) = 0$, because for any nonzero map $F \to G$ the composition

$$V^{\vee} \twoheadrightarrow F \to G \hookrightarrow V^{\vee}$$

would give a nonzero endomorphism of V^{\vee} that is not an isomorphism. Now, for a fixed F and G, the possible V^{\vee} appearing in (16) are parameterized by an open subset of $\mathbb{P}\text{Ext}^1_X(F,G)$. By the Riemann–Roch theorem we have

$$\dim \operatorname{Ext}_{X}^{1}(F,G) = -\chi(X, F^{\vee} \otimes G) = -\deg(F^{\vee} \otimes G) - \operatorname{rk}(F^{\vee} \otimes G)(1-g) = \deg(F)\operatorname{rk}(G) - \operatorname{rk}(F) \deg(G) + \operatorname{rk}(F)\operatorname{rk}(G)(g-1) = l(mr_{1}-k) - k(mr_{1}(1-g) + md_{1}-\epsilon - l) + k(mr_{1}-k)(g-1) = (lr_{1}-kd_{1})m + 2kr_{1}(g-1)m - k^{2}(g-1) + k\epsilon.$$

Note that by our assumption on k and l, we have

$$lr_1 - kd_1 = kr_1\left(\frac{l}{k} - \frac{d_1}{r_1}\right) = kr_1(\mu(F) - \mu(E)) < 0.$$

Thus, the possible V^{\vee} that could appear are parameterized by an open subset of a projective bundle \mathcal{P} over an open subset $\mathcal{U} \subseteq \mathcal{U}_1 \times \mathcal{U}_2$ of dimension

$$\dim \mathcal{P} \le \dim \mathcal{U}_1 + \dim \mathcal{U}_2 + \dim \operatorname{Ext}_X^1(F, G) - 1$$

$$\le k^2(g-1) + 1 + (mr_1 - k)^2(g-1) + 1$$

$$+ kr_1(\mu(F) - \mu(E))m + 2kr_1(1-g)m - k^2(g-1) - k\epsilon - 1$$

$$= (mr_1)^2(g-1) + kr_1(\mu(F) - \mu(E))m + k^2(g-1) + k\epsilon + 1.$$

Since the coefficient of m in the last expression is negative, we see that for large enough m, we have dim $\mathcal{P} < (mr_1)^2(g-1) + 1$, where by Theorem 3.12 the right-hand side is the dimension of the moduli space of stable bundles of rank mr_1 and fixed degree. Thus, a general stable bundle V cannot appear in an extension of the form (16). This proves (i).

To prove (ii), assume that E is stable and $\epsilon \leq 0$. By (i), if V is a general stable vector bundle of the given rank and degree and m is sufficiently large, there are no maps $V^{\vee} \to E$ whose image has slope less than $\mu(E)$. Thus, since E is stable, any nonzero map must be surjective, and so we are led to estimate the dimension of a variety parameterizing all stable bundles V such that there exists a short exact sequence

$$0 \to G \to V^{\vee} \to E \to 0.$$

Such V^{\vee} are parameterized by a projective bundle \mathcal{P} over a scheme \mathcal{U}_2 , where \mathcal{U}_2 parameterizes bundles G with

$$rk(G) = mr_1 - r, \quad deg(G) = mr_1(1 - g) + md_1 - \epsilon - d$$

and again $\operatorname{Hom}_X(E,G) = 0$ since V^{\vee} is stable. The fiber of \mathcal{P} over $[G] \in \mathcal{U}_2$ is $\operatorname{\mathbb{P}Ext}^1_X(E,G)$, and the dimension calculations from above apply, so we have

$$\dim \mathcal{P} \le \dim \mathcal{U}_2 + \dim \operatorname{Ext}_X^1(E, G) - 1$$

= $(mr_1 - r)^2(g - 1) + 1 + 2rr_1(g - 1)m - r^2(g - 1) + r\epsilon - 1$
= $(mr_1)^2(g - 1) + r\epsilon$.

Since $\epsilon \leq 0$, we have dim $\mathcal{P} < (mr_1)^2(g-1) + 1$, where the left-hand side is the dimension of the moduli space of stable bundles of rank mr_1 by Theorem 3.12.

Proof of Proposition 5.2. Applying Lemma 5.8 with $\epsilon = 0$ to a stable vector bundle E produces an open subset in the relevant moduli space such that any V in the open subset satisfies the first claim of the proposition.

Now let E be a polystable vector bundle corresponding to a point in $M_X^{ss}(r, L)$. Taking the intersection of the open subsets for all the stable summands of E produces a nonempty open subset where one can find a vector bundle V such that $H^i(X, E \otimes V) = 0$ for i = 0, 1. Hence the characterization from Proposition 4.7 shows that the associated section s_V does not vanish at $[E] \in M_X^{ss}(r, L)$.

Lemma 5.9. Let E_0, E_1, \ldots, E_n be stable vector bundles of slope $\mu = r/d$. Denote the rank of E_i by k_i . For m sufficiently large, a general stable vector bundle V with

$$rk(V) = mr_1$$
 and $deg(V) = mr_1(g-1) - md_1 + 1$

satisfies the following:

- (i) dim Hom_X(V^{\vee}, E_i) = k_i , or equivalently Ext¹_X(V^{\vee}, E_i) = 0;
- (ii) for $0 \leq i \leq n$, every nonzero map $V^{\vee} \to E_i$ is surjective;
- (iii) for $1 \leq i \leq n$, for all nonzero maps $V^{\vee} \to E_0$ and $V^{\vee} \to E_i$, the induced map $V^{\vee} \to E_0 \oplus E_i$ is surjective.

Proof. Note first that $\mu = d_1/r_1 = \deg(E_i)/k_i$. We have

$$\dim \operatorname{Hom}_X(V^{\vee}, E_i) = \operatorname{h}^0(V \otimes E_i) = \chi(X, V \otimes E_i) + \operatorname{h}^1(X, V \otimes E_i),$$

and by the Riemann-Roch theorem,

$$\chi(X, V \otimes E_i) = \deg(V \otimes E_i) + \operatorname{rk}(V \otimes E_i)(1-g)$$

= deg(V) rk(E_i) + rk(V) deg(E_i) + rk(V) rk(E_i)(1-g)
= (mr_1(g-1) - md_1 + 1)k_i + mr_1k_i\mu + mr_1k_i(1-g)
= k_i.

To obtain (i) we must show that for a general stable V we have $H^1(X, V \otimes E_i) = 0$ for $0 \le i \le n$. By Serre duality, we have

$$h^{1}(X, V \otimes E_{i}) = h^{0}(X, V^{\vee} \otimes \omega_{X} \otimes E_{i}^{\vee}) = \dim \operatorname{Hom}_{X}(V \otimes \omega_{X}^{\vee}, E_{i}^{\vee}).$$

Now $V^{\vee} \otimes \omega_X$ is a general stable bundle of rank mr_1 and degree

$$\deg(V^{\vee} \otimes \omega_X) = -\deg(V) + \operatorname{rk}(V) \deg(\omega_X) = mr_1(1-g) + md_1 - 1 + mr_1(2g-2) = mr_1(g-1) + md_1 - 1,$$

and E_i^{\vee} is a stable bundle of slope $-d_1/r_1$, so we may apply Lemma 5.8(ii) with $\epsilon = -1$ to each E_i^{\vee} to conclude that, for sufficiently large m and general stable V, we have $\operatorname{Hom}_X(V \otimes \omega_X^{\vee}, E_i^{\vee}) = 0$ for each i. This gives (i).

By applying Lemma 5.8(i) with $\epsilon = 1$ to each E_i and by increasing m if necessary, we may assume that for general stable V the image of every nonzero map $V^{\vee} \to E_i$ has slope equal to μ , and, since E_i is stable, such a map must be surjective. This gives (ii).

Finally, by applying Lemma 5.8(i) to each $E_0 \oplus E_i$ for $1 \le i \le n$ and increasing m if necessary, we may assume that for general stable V, every nonzero map $V^{\vee} \to E_0 \oplus E_i$ has image with slope equal to μ . For such a V and two nonzero maps $V^{\vee} \to E_0$ and $V^{\vee} \to E_i$, the image $F \subseteq E_0 \oplus E_i$ of the induced map $V^{\vee} \to E_0 \oplus E_i$ must thus have slope $\mu(F) = \mu$, and moreover F surjects onto both summands. But the only nonzero subsheaves of $E_0 \oplus E_i$ of slope μ are E_0 , E_i , and $E_0 \oplus E_i$ and, since E_0 and E_i are by assumption nonisomorphic, we must have $F = E_0 \oplus E_i$. This gives (iii).

Before we give the proof of Proposition 5.4 we introduce a method for constructing new vector bundles out of old ones.

Definition 5.10. Let E be a vector bundle on X and let $x \in X(k)$ be a closed point. Denote by \mathcal{O}_x the skyscraper sheaf at x and E(x) the fiber of E at x. Let $\alpha \in \operatorname{Hom}_X(E, \mathcal{O}_x) = E(x)^{\vee}$ be a nonzero morphism. The *elementary* transformation of E at x associated to α is the vector bundle E' defined as the kernel of α , so that we have the short exact sequence

$$0 \to E' \to E \stackrel{\alpha}{\to} \mathcal{O}_x \to 0. \tag{17}$$

The kernel of α does not change under rescaling, so elementary transformations of E at x correspond to closed points of $\mathbb{P}E(x)^{\vee}$.

If $\phi: F \to E$ is a morphism, then $\operatorname{im} \phi(x)$ is a subspace of E(x), while ker $\alpha(x)$ is a hyperplane in E(x), which we will denote by H. If $\operatorname{im} \phi(x) \subseteq H$ then the composition $F \xrightarrow{\phi} E \xrightarrow{\alpha} \mathcal{O}_x$ vanishes and so $F \to E$ factors through E'. Conversely, if $\operatorname{im} \phi(x)$ is not contained in H then $F \to E$ does not factor through E'.

Proof of Proposition 5.4. Let E_1, \ldots, E_n be the stable summands of E', and let E_0 be a stable summand of E not isomorphic to any of the E_i . As before, let k_i denote the rank of E_i . It suffices to find a vector bundle F such that

$$\mathrm{H}^{0}(X, F \otimes E_{0}) \neq 0$$
 but $\mathrm{H}^{0}(X, F \otimes E_{i}) = 0$ for $i = 1, \ldots, n$.

Let V be a stable vector bundle satisfying the conditions of Lemma 5.9, and fix a nonzero map $\phi: E_0^{\vee} \to V$. The goal is to find a subsheaf $F \subset V$ of degree $\deg(F) = \deg(V) - 1$ such that

- ϕ factors through F, but
- for i = 1, ..., n, no nonzero map $E_i^{\vee} \to V$ factors through F.

Let Q denote the cokernel of ϕ . It is locally free since by Lemma 5.9(ii) ϕ is the dual of a surjective map $V^{\vee} \to E_0$. Note that, for any nonzero map $\psi_i \colon E_i^{\vee} \to V$, the induced map $E_0^{\vee} \oplus E_i^{\vee} \to V$ is dual to a surjection by Lemma 5.9(iii) and hence injective. The inclusions

$$E_0^{\vee} \hookrightarrow E_0^{\vee} \oplus E_i^{\vee} \stackrel{(\phi,\psi_i)}{\hookrightarrow} V$$

induce a short exact sequence

$$0 \to E_i^{\vee} \xrightarrow{\overline{\psi}_i} Q \to V/E_0^{\vee} \oplus E_i^{\vee} \to 0,$$

and so the composition $\overline{\psi}_i : E_i^{\vee} \to V \to Q$ is injective with locally free cokernel, from which it follows that the restriction of $\overline{\psi}_i$ to a fiber is also injective.

Let $x \in X$ be a closed point, let Q(x) and V(x) denote the fibers of Qand V at x respectively, and let $\operatorname{Grass}(k_i, Q(x))$ denote the Grassmannian of k_i -dimensional subspaces of the vector space Q(x). Let $M_i \subseteq \operatorname{Grass}(k_i, Q(x))$ denote the image of the morphism

$$\mathbb{P}\operatorname{Hom}_X(E_i^{\vee}, V) \to \operatorname{Grass}(k_i, Q(x))$$

that sends a nonzero map ψ_i to its image under the composition

$$E_i^{\vee} \xrightarrow{\psi_i} Q \to Q(x).$$

This morphism is well defined, because in the previous paragraph we established that $\overline{\psi}_i(x)$ is injective. We have dim $M_i \leq k_i - 1$ by Lemma 5.9(i).

For a hyperplane $H \subseteq Q(x)$, let $Z_{H,i} \subseteq \operatorname{Grass}(k_i, Q(x))$ denote the Schubert variety parameterizing subspaces contained in H. The codimension of $Z_{H,i}$ in $\operatorname{Grass}(k_i, Q(x))$ is k_i , and so the expected dimension of the intersection $M_i \cap Z_{H,i}$ is -1. Thus, by the Bertini–Kleiman theorem [16, Corollary 4(i)] we obtain for the general hyperplane H that $M_i \cap Z_{H,i}$ is empty for $i = 1, \ldots, n$. Let us fix one such hyperplane H.

Let F denote the elementary transformation of V at x defined by the subspace H, i.e., it is the kernel of the morphism $V \to \mathcal{O}_x$ determined by the composition

$$V \to V(x) \to Q(x) \to Q(x)/H.$$

By construction $\phi \colon E_0^{\vee} \to V$ factors through F, and so

$$\mathrm{H}^{0}(X, E_{0} \otimes F) = \mathrm{Hom}_{X}(E_{0}^{\vee}, F) \neq 0.$$

On the other hand, no nonzero map $E_i^\vee \to V$ factors through F, as the composition

$$E_i^{\vee}(x) \xrightarrow{\psi_i(x)} V(x) \to Q(x)/H$$

is nonzero by the choice of H. Thus,

$$\mathrm{H}^{0}(X, E_{i} \otimes F) = \mathrm{Hom}_{X}(E_{i}^{\vee}, F) = 0 \quad \text{for} \quad i = 1, \dots, n.$$

Since $\operatorname{rk}(F) = \operatorname{rk}(V)$ and $\operatorname{deg}(F) = \operatorname{deg}(V) - 1$, this completes the proof. \Box

Acknowledgements We thank Hannah Larson and Shizhang Li for being part of our group during SPONGE, and the many interesting conversations during and after the workshop. We are grateful for the referee's comments. PB and TT would also like to thank Chiara Damiolini, Hans Franzen, Vicky Hoskins, and Svetlana Makarova for interesting discussions on a GIT-free proof of the (quasi-)projectivity of moduli of semistable quiver representations. JA was partially supported by NSF grants DMS-1801976 and DMS-2100088. DB was supported by NSF Postdoctoral Research Fellowship DMS-1902875.

References

- Jarod Alper. Good moduli spaces for Artin stacks. Ann. Inst. Fourier (Grenoble), 63(6):2349-2402, 2013.
- [2] Jarod Alper, Daniel Halpern-Leistner, and Jochen Heinloth. Existence of moduli spaces for algebraic stacks, 2019. arXiv:1812.01128v3.
- [3] Luis Álvarez Cónsul and Alastair King. A functorial construction of moduli of sheaves. *Invent. Math.*, 168(3):613–666, 2007.
- [4] Leticia Brambila-Paz, Iwona Grzegorczyk, and Peter Newstead. Geography of Brill–Noether loci for small slopes. J. Algebraic Geom., 6(4):645–669, 1997.
- [5] Raymond Cheng, Carl Lian, and Takumi Murayama. Projectivity of the moduli of curves. In *Stacks Project Expository Collection*, number 480 in London Mathematical Society Lecture Note Series, chapter 1. Cambridge University Press, 2022.
- [6] Maurizio Cornalba. On the projectivity of the moduli spaces of curves. J. Reine Angew. Math., 443:11–20, 1993.
- [7] Pierre Deligne and David Mumford. The irreducibility of the space of curves of given genus. Inst. Hautes Études Sci. Publ. Math., (36):75–109, 1969.
- [8] Jean-Marc Drézet and M. S. Narasimhan. Groupe de Picard des variétés de modules de fibrés semi-stables sur les courbes algébriques. *Invent. Math.*, 97(1):53–94, 1989.
- [9] Eduardo Esteves and Mihnea Popa. Effective very ampleness for generalized theta divisors. Duke Math. J., 123(3):429–444, 2004.
- [10] Gerd Faltings. Stable G-bundles and projective connections. J. Algebraic Geom., 2(3):507–568, 1993.
- [11] Haoyang Guo, Sanal Shivaprasad, Dylan Spence, and Yueqiao Wu. Boundedness of semistable sheaves. In *Stacks Project Expository Collection*, number 480 in London Mathematical Society Lecture Note Series, chapter 4. Cambridge University Press, 2022.
- [12] Daniel Halpern-Leistner. On the structure of instability in moduli theory, 2018. arXiv:1411.0627v4.
- [13] Georg Hein. Faltings' construction of the moduli space of vector bundles on a smooth projective curve. In Affine flag manifolds and principal bundles, Trends Math., pages 91–122. Birkhäuser/Springer Basel AG, Basel, 2010.

- [14] Daniel Huybrechts and Manfred Lehn. The geometry of moduli spaces of sheaves. Cambridge Mathematical Library. Cambridge University Press, second edition, 2010.
- [15] Seán Keel and Shigefumi Mori. Quotients by groupoids. Ann. Math., 145(1):193– 213, 1997.
- [16] Steven L. Kleiman. The transversality of a general translate. Compositio Math., 28:287–297, 1974.
- [17] János Kollár. Projectivity of complete moduli. J. Differential Geom., 32(1):235– 268, 1990.
- [18] Herbert Lange. Zur Klassifikation von Regelmannigfaltigkeiten. Math. Ann., 262(4):447–459, 1983.
- [19] Stacy G. Langton. Valuative criteria for families of vector bundles on algebraic varieties. Ann. Math., 101:88–110, 1975.
- [20] Joseph Le Potier. Lectures on vector bundles, volume 54 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1997. Translated by A. Maciocia.
- [21] M. S. Narasimhan and S. Ramanan. Moduli of vector bundles on a compact Riemann surface. Ann. Math., 89:14–51, 1969. doi:10.2307/1970807.
- [22] C. S. Seshadri. Vector bundles on curves. In *Linear algebraic groups and their representations (Los Angeles, CA, 1992)*, volume 153 of *Contemp. Math.*, pages 163–200. American Mathematical Society, 1993.
- [23] The Stacks Project Authors. The Stacks project. https://stacks.math.columbia. edu.

Department of Mathematics, University of Washington, Box 354350, C-138 Padelford, Seattle, WA 98195-4350, United States

Department of Mathematics, University of Luxembourg, 6, Avenue de la Fonte, L-4364 Esch-sur-Alzette, Luxembourg

Department of Mathematics, University of California, Berkeley, 970 Evans Hall, Berkeley, CA 94720-3840, United States

Department of Mathematics, University of Michigan, 530 Church Street, Ann Arbor, MI 48109, United States

Matematiska institutionen, Stockholm Universitet, Kräftriket 5A, 114 19 Stockholm, Sweden

alperQuw.edu, pieter.belmansQuni.lu, braggdanQberkeley.edu, liangjiQumich.edu, tuomas.tajakkaQmath.su.se