

Journal Pre-proof

Distribution of psychological instability among surgeons

Anna L. Kalywis, Robin Samuel, Felix Scholtes, Gilles Reuter, Martin N. Stienen, Erich Seifritz, Werner Surbeck



PII: S1878-8750(23)00461-8

DOI: <https://doi.org/10.1016/j.wneu.2023.03.137>

Reference: WNEU 20401

To appear in: *World Neurosurgery*

Received Date: 16 December 2022

Revised Date: 29 March 2023

Accepted Date: 30 March 2023

Please cite this article as: Kalywis AL, Samuel R, Scholtes F, Reuter G, Stienen MN, Seifritz E, Surbeck W, Distribution of psychological instability among surgeons, *World Neurosurgery* (2023), doi: <https://doi.org/10.1016/j.wneu.2023.03.137>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 The Author(s). Published by Elsevier Inc.

Distribution of psychological instability among surgeons

Anna L. Kalywis^{1,2}, Robin Samuel³, Felix Scholtes⁴, Gilles Reuter⁵, Martin N. Stienen⁶,
Erich Seifritz¹, Werner Surbeck¹

¹) Department of Psychiatry, Psychotherapy and Psychosomatics, Psychiatric Hospital of the University of Zurich, Zurich, Switzerland

²) Department of Consultation-Liaison-Psychiatry and Psychosomatic Medicine, University Hospital, Zurich, University of Zurich, Zurich, Switzerland.

³) Department of Social Sciences, University of Luxembourg, Esch-sur-Alzette, Luxembourg

⁴) Department of Neuroanatomy, Faculty of Medicine, Université de Liège, Liège, Belgium

⁵) Department of Neurosurgery, Faculty of Medicine, Université de Liège, Liège, Belgium

⁶) Department of Neurosurgery, Cantonal Hospital St. Gallen, St. Gallen, Switzerland

Correspondence to: Werner Surbeck, MD PhD; Department of Psychiatry, Psychotherapy and Psychosomatics, Psychiatric Hospital of the University of Zurich, Lenggstrasse 31 8032 Zurich, Switzerland; Tel: +41 (0)44 296 73 86; Email: surbeck.werner@gmail.com

Short title: Distribution of psychological instability among surgeons

Keywords: medicine, surgery, neuroticism, medical error, second victim, physicians' wellbeing

Compliance with Ethical Requirements: The study was submitted to the institutional review board of the Canton St.Gallen, Switzerland (EKSG 16/020) and the "Comité d'Ethique Hospitalo-Facultaire Universitaire de Liège" (2016/74). Both estimated that it did not fall under the legislation for research involving human beings and that the collected anonymous data did not require any consent beyond the deliberate participation.

Conflicts of interest: No author has any conflict of interest with regard to its content.

Manuscript word count (without title page and references): 2496

Abstract

Background: High emotional instability, i.e. neuroticism, is associated with poor mental health. Conversely, traumatic experiences may increase neuroticism. Stressful experiences such as complications are common in the surgical profession, with neurosurgeons being particularly affected. We compared the personality trait neuroticism between physicians in a prospective cross-sectional study.

Methods: We used an online survey using the Ten-Item Personality Inventory, an internationally validated measure of the five-factor model of personality dimensions. It was distributed to board-certified physicians, residents, and medical students in several European countries and Canada ($n = 5,148$). Multivariate linear regression was used to model differences between surgeons, non-surgeons and specialties with occasional surgical interventions with respect to neuroticism, adjusting for gender, age, age squared and their interactions, then testing equality of parameters of adjusted predictions separately and jointly using Wald tests.

Results: With an expected variability within disciplines, average levels of neuroticism are lower in surgeons than non-surgeons, especially in the first part of their career. However, the course of neuroticism across age follows a quadratic pattern, i.e., an increase after the initial decrease. The acceleration of neuroticism with age is specifically significant in surgeons. Levels of neuroticism are lowest towards mid-career, but exhibit a strong secondary increase towards the end of the surgeons' career. This pattern seems driven by neurosurgeons.

Conclusion: Despite initially lower levels of neuroticism, surgeons suffer a stronger increase of neuroticism together with age. Since, beyond well-being, neuroticism influences professional performance and healthcare systems costs, explanatory studies are mandatory to enlighten causes of this burden.

Introduction

“Every surgeon carries within himself a small cemetery, where from time to time he goes to pray – a place of bitterness and regret, where he must look for an explanation for his failures.” René Leriche, La philosophie de la chirurgie, 1951”¹

Mental health differs between medical practitioners and the general population and suicide risk is increased in physicians²⁻⁴. Emotional distress in practitioners not only has negative effects on the individual, it also weigh on patient care and even on the health care systems^{5, 6}.

Distress can result from the occurrence of adverse clinical events, especially individual errors and surgical incidents, such as complications with demonstrated negative impacts on health and wellbeing⁷⁻¹⁰. Emotional responses of health care professionals to adverse clinical events with intense professional suffering and possible long-term impact thus create a “second victim”^{11, 12}. Health care professionals are exposed to such experiences over their careers^{6, 13}, especially in the surgical field^{12, 14} and the impact of complications can be devastating, especially in neurosurgery.

Neuroticism, the opposite of emotional stability, is one of the “higher-order factors” used to describe personality in the classic five-factor model (FFM)¹⁵⁻¹⁷. It reflects a tendency to experience negative emotions such as irritability, anxiety and sadness. It has been identified as a strong health predictor^{18, 19}. It is associated with depression and other psychopathological states²⁰⁻²², with burnout and psychological distress²³⁻²⁵ as well as perceived job stress²⁶. Generally, neuroticism appears to decrease somewhat with age²⁷⁻²⁹, even if the literature is not perfectly consistent on this subject³⁰. Deteriorated quality of life and stressful life events may increase neuroticism³¹⁻³⁵. Stressful life events might also alter the normative development trajectory, so that expected age-related reduction of neuroticism does not occur³⁶⁻³⁸.

We hypothesized that, contrary to the general population, in medical professionals, neuroticism increases with age, as adverse experiences accumulate over time. Second, we hypothesized that surgeons and especially neurosurgeons, who face more adverse

events that are directly causal to their actions, may be at a higher risk compared to physicians in non-surgical specialties.

Methods

Survey design and distribution

We used data collected by means of an online survey (prospective, cross-sectional study design; for details see ³⁹). The survey was conducted in Austria, Belgium, Canada, Germany and Switzerland collecting respondents' age, gender, primary language, level of training (board certified physician, resident, or medical student), as well as the (intended) medical specialty. Data was collected from February 12, 2016 to May 12, 2016. The study was submitted to the institutional review board of the Canton St.Gallen, Switzerland (EKSG 16/020) and the "Comité d'Ethique Hospitalo-Facultaire Universitaire de Liège" (2016/74). Both estimated that it did not fall under the legislation for research involving human beings and that the collected anonymous data did not require any consent beyond the deliberate participation.

Study variables

Our key outcome variable was neuroticism, measured by the Ten Item Personality Inventory (TIPI), an internationally validated short measure of the five-factor model (FFM). ⁴⁰ The latter comprehensively describes personality using five higher-order factors: agreeableness, conscientiousness, openness to experience, neuroticism, and extraversion. ¹⁵ For assessing neuroticism, participants rate the extent to which they agree (1 = "Disagree strongly" to 7 = "Agree strongly") with the statements of being anxious and easily upset as well as being calm and emotionally stable (reversed item). ⁴⁰ The score for neuroticism is the average of both items. Despite less precise estimation of the FFM than with more complex and time-consuming tools, its results have been shown to converge with other widely used FFM measures in self-report, observer report and peer report, test–retest reliability, patterns of predicted external correlates, and convergence between self and observer ratings. ⁴¹

We used raw and unstandardized test scores and included age and gender in our models to adjust for these factors. Respondents were categorized based on the discipline they stated to be non-surgeons (non-surgical disciplines with clinical patient contact), surgeons, or “occasionals” (specialties with clinical patient contact and occasional invasive interventions). Supplementary table 1 provides an overview of the assignment of subspecialties to these three categories.

Statistical Analysis

Continuous data are reported as mean and standard deviation (SD) and were compared using two-sample unpaired t-tests. Potential bias due to unequal variances and unequal sample sizes was accounted for by Satterthwaite approximation.

Multivariate linear regression was used to model differences between non-surgeons, surgeons, neurosurgeons, and occasionals with respect to neuroticism. We adjusted for gender and age and included age squared to allow for a change in the association of age and levels of neuroticism as we expect. To analyse differences between non-surgeons, surgeons, neurosurgeons, and occasionals, marginal effects were calculated, and their differences tested using Wald tests. All statistical tests were two-sided and p values < 0.05 were considered statistically significant. All analysis were performed in Stata, version 17.0 (StataCorp LP, College Station, Texas, USA).

Analysis Sample

5,148 respondents completed the survey (Stienen, Scholtes et al. 2018). Excluding specialties without direct clinical contact to patients (as outlined under *study variables*), that is, clinical pathologists, forensic pathologists, microbiologists, pathologists, and radiologists leaves us with a final analytical sample of 4,678 cases, with 2,200 non-surgeons (age 35.4 ± 11.7 ; 55.7% female), 1,035 surgeons (age 34.9 ± 11.2 ; 41.3% female), and 1,443 occasionals (age 36.9 ± 13.1 ; 60.4% female).

Results

On average, surgeons score lower on neuroticism (1.5) than non-surgeons (1.9) and occasionals (1.9), yielding a statistically significant difference between groups as determined by one-way ANOVA ($F(2, 4,675) = 40.06, p < .001$). Multivariate linear regression suggests that these differences remain when adjusting for age and gender (Model 1, Table 1). Model 1 further provides evidence that males score on average lower on neuroticism than females do and that there is a negative association with age. Model 2 shows that these findings hold when allowing for an interaction between speciality and age. Model 3 further allows this interaction to vary over gender, Model 4 allows for a quadratic age pattern.

Overall, all models support a linear negative association of age with levels of neuroticism. Model 4 suggests that this decline is more pronounced for surgeons. Furthermore, Model 4 provides evidence that there is a significant quadratic pattern of age with respect to neuroticism for surgeons.

Plotting marginal effects derived from Model 4 (Figure 1) we see that overall levels of neuroticism decline with age, but increase again at some point following a u-shaped pattern. This pattern is most pronounced for surgeons, with the quadratic age terms being jointly significantly different between non-surgeons, surgeons, and occasionals ($F(2, 4,660) = 3.38, p = .034$).

In a sub analysis, we find that the prominent pattern for surgeons was particularly driven by neurosurgeons (see supplementary table 2) with the quadratic age pattern being jointly significantly different between neurosurgeons, surgeons, non-surgeons, and occasionals ($F(3, 4,654) = 2.75, p = .041$).

Discussion

The present cross-sectional observational study indicates a distinct pattern of the personality facet neuroticism within the medical field, across age and gender. It shows significant differences between surgical and other medical specialties.

As described before ^{39, 42}, internists score, on average, higher on neuroticism than surgeons, including after adjustment for gender, especially in the first part of their career. It is plausible that relatively low baseline neuroticism helps surgeons to make incisive and irreversible decisions with less anxiety in critical situations. Lower neuroticism scores among surgeons are probably correlated with personality traits and non-technical skills inherent to the choice of pursuing a career in surgery ⁴³⁻⁴⁵.

The current results also replicate findings that females score higher on neuroticism than males ²⁷. Furthermore, we found that both female and male internist's degree of neuroticism differs from that of female and male surgeons', respectively. This strengthens the notion of a personality trait difference, specifically neuroticism, in surgeons vs non-surgeons, independently of gender differences.

To the best of our knowledge, no previous data about neuroticism in clinicians of different disciplines across age has been published. In the general population, mostly decreasing patterns were described in longitudinal ^{28, 29, 46}, as well as cross-sectional studies ^{47, 48}. There is some previous data showing higher levels of neuroticism in older surgeons compared to younger ones ⁴⁹. We hypothesize that the impact of repeated adverse work-related events, accumulating over time in a surgical career, increase neuroticism in physicians with time.

Our results indeed provide evidence for a significant quadratic pattern of age in respect to neuroticism in all disciplines, with a clearer pronouncement in surgeons. This prominent pattern seem especially driven by neurosurgeons. While surgeons start with lower levels of neuroticism compared with other medicals, older surgeons show increased levels of neuroticism. This pattern is not statistically significant in physicians with occasional interventional tasks and non-surgeons. This is possibly explained by the intensity of work-related negative impacts through directly experienced adverse events. Indeed, surgical incidents have been shown to generate significant negative personal and professional impact ^{8, 10, 50}. Surgeons must take greater individual risks and are thus more prone to concrete, immediate and severe adverse work related events ⁵¹. They are mostly directly involved in adverse events and often attribute these to their own capabilities ⁵⁰ and thus

tend to feel responsible and blame themselves⁵². If, in addition to these general aspects, the extent of the damage caused by the operated organ is taken into account, neurosurgeons seem to be particularly affected. Balch and Shanafelt⁵³ reported nearly 9% of surgeons who perceive to have made a recent major medical error, and these score higher on suicide ideation, depression and burnout⁵³. According to Han and colleagues⁵⁴ 80% of the studied surgeons reported at least one intraoperative adverse event in the past year, and intraoperative events only represent one, though very immediately apparent, type of surgical complication. An unwritten code of social norms and expectations — including the value attributed to “not complaining” or keeping emotions from interfering with one’s work⁵³ — as well as convictions to be more resilient in comparison to non-surgeons, reflect a unique overall surgical attitude. This might lead to silent suffering and not seeking adequate support^{10, 55, 56}. Competitive and non-supportive working cultures seem to hinder support seeking and might aggravate impacts of adverse events⁵⁴. Thus: being more prone to, and possibly more affected by these kinds of negative events add strength to our hypothesis that negative work-related incidents affect well-being and performance not only directly¹⁰, but through progressively increasing neuroticism over a career, with further negative collateral impact.

Further longitudinal investigations could strengthen this hypothesis, help understand the impacts of negative incidents and design strategies to lower these impacts. Indeed, in this perspective, neuroticism’s significant economic burden⁵⁷ and personal negative effect⁵⁸, as well as negative impact on job performance⁵⁹ highlight the importance of further investigation of these findings. We concur with the assumption of Ullström⁶⁰ that psychological and institutional support are likely to lower the negative impacts of adverse events. Additionally, a change of attitudes towards medical errors and incidents could evolve: errors can be analyzed from an structural instead of an individual perspective⁶¹; learning from medical errors can be strengthened as a coping mechanism⁶². This type of changed attitude could improve not only the direct well-being of physicians, but also their performance and, ultimately, overall patient outcomes. In addition, complications are clearly underreported in the medical literature at present. It is somewhat easier to publish a series of successful cases than to report complications. Upgrading the reporting of

complications could be beneficial to the surgeon, the patients, and the entire health care system ⁶³. Previous work has already underlined the benefit of perceived social support on neuroticism ⁶⁴ as well as favorable psychotherapeutic treatment effects on neuroticism ^{65, 66}. In short, adequate support might lower perceived stress ¹² and prevent increase of neuroticism with its associated negative implications.

Limitations

Our analytical strategy aims to provide a description of cross-sectional data but does not allow for a causal interpretation. For the limited number of recorded baseline variables, we were unable to adjust for other factors, e.g., at the individual and job level. We could not consider possible changing personality characteristics of physicians or changing profiles concerning the chosen discipline over time. We also could not observe traumatic events during the professional career, as this information was not available. Another limitation is the self-selection of respondents into survey participation. This is also true for neurosurgeons, whose numbers might be considered too small to reliably determine their relative importance for some of the observed associations. All in all, the external validity stays unclear. Nevertheless, the findings of this study are of tremendous importance and give ground for further investigation with the ultimate aim to improve practitioners' wellbeing and patient care.

Conclusion

Surgeons show higher emotional stability than their medical colleagues and the general population at the beginning of their careers. First, emotional stability further increases, but then neuroticism rises above initial levels with advancing age. This finding is alarming, even if taking into account the current study limitations and its descriptive nature. Are surgeons not only carrying their “inner cemetery”, as Leriche described, but also paying a high personal cost for their work? The hypothesis that traumatic or stressful work experiences contribute to this finding needs to be tested in further investigations. Meanwhile, we suggest that easily accessible and systematic psychological support throughout the career, as well as a change of attitudes towards medical errors and

incidents are likely to show benefits, without expected negative impacts, regarding physicians' mental health and work quality.

Acknowledgements

The authors thank the journals "Deutsches Ärzteblatt, Germany", "Schweizer Ärztezeitung, Switzerland", and "Le Quotidien du Médecin, France" to help distribute the survey among their readership. We thank all participating specialists, residents and medical students for their time and trust.

Contributors

Those persons listed as authors on the manuscript have made substantial contributions to the conception or design of the work, and have been involved in acquiring, analyzing or interpreting the data for the work. They all have been active in drafting or revising the manuscript for important intellectual content, which is the basis of the current article. Concept: ALK, RS, ES, WS. Data collection: MNS, FS, WS. Data analysis: RS, ALK, WS. Article drafting: ALK, WS, FS, RS, GR. Critical revision: all authors.

References

1. Leriche R. *La philosophie de la chirurgie*: Flammarion; 1951.
2. Agerbo E, Gunnell D, Bonde JP, Mortensen PB, Nordentoft M. Suicide and occupation: the impact of socio-economic, demographic and psychiatric differences. *Psychol Med*. 2007;37(8): 1131-1140. <https://doi.org/10.1017/S0033291707000487>.
3. Shanafelt TD, Balch CM, Dyrbye L, et al. Special report: suicidal ideation among American surgeons. *Arch Surg*. 2011;146(1): 54-62. <https://doi.org/10.1001/archsurg.2010.292>.
4. Dutheil F, Aubert C, Pereira B, et al. Suicide among physicians and health-care workers: A systematic review and meta-analysis. *PLoS One*. 2019;14(12): e0226361. <https://doi.org/10.1371/journal.pone.0226361>.
5. Wallace JE, Lemaire JB, Ghali WA. Physician wellness: a missing quality indicator. *Lancet*. 2009;374(9702): 1714-1721. [https://doi.org/10.1016/S0140-6736\(09\)61424-0](https://doi.org/10.1016/S0140-6736(09)61424-0).
6. Hu YY, Fix ML, Hevelone ND, et al. Physicians' needs in coping with emotional stressors: the case for peer support. *Arch Surg*. 2012;147(3): 212-217. <https://doi.org/10.1001/archsurg.2011.312>.
7. Waterman AD, Garbutt J, Hazel E, et al. The emotional impact of medical errors on practicing physicians in the United States and Canada. *Jt Comm J Qual Patient Saf*. 2007;33(8): 467-476. [https://doi.org/10.1016/s1553-7250\(07\)33050-x](https://doi.org/10.1016/s1553-7250(07)33050-x).

8. Pinto A, Faiz O, Bicknell C, Vincent C. Surgical complications and their implications for surgeons' well-being. *Br J Surg*. 2013;100(13): 1748-1755. <https://doi.org/10.1002/bjs.9308>.
9. Serou N, Sahota L, Husband AK, et al. Systematic review of psychological, emotional and behavioural impacts of surgical incidents on operating theatre staff. *BJS Open*. 2017;1(4): 106-113. <https://doi.org/10.1002/bjs5.21>.
10. Serou N, Slight SP, Husband AK, Forrest SP, Slight RD. Surgical incidents and their impact on operating theatre staff: qualitative study. *BJS Open*. 2021;5(2). <https://doi.org/10.1093/bjsopen/zraa007>.
11. Wu AW. Medical error: the second victim. The doctor who makes the mistake needs help too. *BMJ*. 2000;320(7237): 726-727. <https://doi.org/10.1136/bmj.320.7237.726>.
12. Seys D, Wu AW, Van Gerven E, et al. Health care professionals as second victims after adverse events: a systematic review. *Eval Health Prof*. 2013;36(2): 135-162. <https://doi.org/10.1177/0163278712458918>.
13. Vincent C, Neale G, Woloshynowych M. Adverse events in British hospitals: preliminary retrospective record review. *BMJ*. 2001;322(7285): 517-519. <https://doi.org/10.1136/bmj.322.7285.517>.
14. Thomas EJ, Studdert DM, Burstin HR, et al. Incidence and types of adverse events and negligent care in Utah and Colorado. *Med Care*. 2000;38(3): 261-271. <https://doi.org/10.1097/00005650-200003000-00003>.
15. Digman JM. Personality Structure - Emergence of the 5-Factor Model. *Annual Review of Psychology*. 1990;41: 417-440. <https://doi.org/DOI 10.1146/annurev.ps.41.020190.002221>.
16. Tupes EC, Christal RE. Recurrent personality factors based on trait ratings. *J Pers*. 1992;60(2): 225-251. <https://doi.org/10.1111/j.1467-6494.1992.tb00973.x>.
17. McCrae RR, John OP. An introduction to the five-factor model and its applications. *J Pers*. 1992;60(2): 175-215. <https://doi.org/10.1111/j.1467-6494.1992.tb00970.x>.
18. Lahey BB. Public health significance of neuroticism. *Am Psychol*. 2009;64(4): 241-256. <https://doi.org/10.1037/a0015309>.
19. Montoliu T, Pulopulos MM, Puig-Perez S, Hidalgo V, Salvador A. Mediation of perceived stress and cortisol in the association between neuroticism and global cognition in older adults: A longitudinal study. *Stress Health*. 2021. <https://doi.org/10.1002/smi.3088>.
20. Kotov R, Gamez W, Schmidt F, Watson D. Linking "big" personality traits to anxiety, depressive, and substance use disorders: a meta-analysis. *Psychol Bull*. 2010;136(5): 768-821. <https://doi.org/10.1037/a0020327>.
21. Ormel J, Jeronimus BF, Kotov R, et al. Neuroticism and common mental disorders: meaning and utility of a complex relationship. *Clin Psychol Rev*. 2013;33(5): 686-697. <https://doi.org/10.1016/j.cpr.2013.04.003>.
22. Jeronimus BF, Kotov R, Riese H, Ormel J. Neuroticism's prospective association with mental disorders halves after adjustment for baseline symptoms and psychiatric history, but the adjusted association hardly decays with time: a meta-analysis on 59 longitudinal/prospective studies with 443 313 participants. *Psychol Med*. 2016;46(14): 2883-2906. <https://doi.org/10.1017/S0033291716001653>.
23. Ormel J, Wohlfarth T. How neuroticism, long-term difficulties, and life situation change influence psychological distress: a longitudinal model. *J Pers Soc Psychol*. 1991;60(5): 744-755. <https://doi.org/10.1037//0022-3514.60.5.744>.
24. Ormel J, Rosmalen J, Farmer A. Neuroticism: a non-informative marker of vulnerability to psychopathology. *Soc Psychiatry Psychiatr Epidemiol*. 2004;39(11): 906-912. <https://doi.org/10.1007/s00127-004-0873-y>.
25. Divinakumar KJ, Bhat PS, Prakash J, Srivastava K. Personality traits and its correlation to burnout in female nurses. *Ind Psychiatry J*. 2019;28(1): 24-28. https://doi.org/10.4103/ipj.ipj_52_19.

26. Gramstad TO, Gjestad R, Haver B. Personality traits predict job stress, depression and anxiety among junior physicians. *BMC Med Educ.* 2013;13: 150. <https://doi.org/10.1186/1472-6920-13-150>.
27. Costa PT, Terracciano A, McCrae RR. Gender differences in personality traits across cultures: robust and surprising findings. *J Pers Soc Psychol.* 2001;81(2): 322-331. <https://doi.org/10.1037/0022-3514.81.2.322>.
28. Roberts BW, Walton KE, Viechtbauer W. Patterns of mean-level change in personality traits across the life course: a meta-analysis of longitudinal studies. *Psychol Bull.* 2006;132(1): 1-25. <https://doi.org/10.1037/0033-2909.132.1.1>.
29. Rantanen J, Metsapelto RL, Feldt T, Pulkkinen L, Kokko K. Long-term stability in the Big Five personality traits in adulthood. *Scand J Psychol.* 2007;48(6): 511-518. <https://doi.org/10.1111/j.1467-9450.2007.00609.x>.
30. Costa PT, Jr., McCrae RR, Lockenhoff CE. Personality Across the Life Span. *Annu Rev Psychol.* 2019;70: 423-448. <https://doi.org/10.1146/annurev-psych-010418-103244>.
31. Specht J, Egloff B, Schmukle SC. Stability and change of personality across the life course: the impact of age and major life events on mean-level and rank-order stability of the Big Five. *J Pers Soc Psychol.* 2011;101(4): 862-882. <https://doi.org/10.1037/a0024950>.
32. Jeronimus BF, Ormel J, Aleman A, Penninx BW, Riese H. Negative and positive life events are associated with small but lasting change in neuroticism. *Psychol Med.* 2013;43(11): 2403-2415. <https://doi.org/10.1017/S0033291713000159>.
33. Jeronimus BF, Riese H, Sanderman R, Ormel J. Mutual reinforcement between neuroticism and life experiences: a five-wave, 16-year study to test reciprocal causation. *J Pers Soc Psychol.* 2014;107(4): 751-764. <https://doi.org/10.1037/a0037009>.
34. Goldstein BL, Perlman G, Eaton NR, Kotov R, Klein DN. Testing explanatory models of the interplay between depression, neuroticism, and stressful life events: a dynamic trait-stress generation approach. *Psychol Med.* 2020;50(16): 2780-2789. <https://doi.org/10.1017/S0033291719002927>.
35. Metts A, Yarrington J, Enders C, et al. Reciprocal effects of neuroticism and life stress in adolescence. *J Affect Disord.* 2021;281: 247-255. <https://doi.org/10.1016/j.jad.2020.12.016>.
36. Mroczek DK, Spiro A, 3rd. Modeling intraindividual change in personality traits: findings from the normative aging study. *J Gerontol B Psychol Sci Soc Sci.* 2003;58(3): P153-165. <https://doi.org/10.1093/geronb/58.3.p153>.
37. Lockenhoff CE, Terracciano A, Patriciu NS, Eaton WW, Costa PT, Jr. Self-reported extremely adverse life events and longitudinal changes in five-factor model personality traits in an urban sample. *J Trauma Stress.* 2009;22(1): 53-59. <https://doi.org/10.1002/jts.20385>.
38. Sutin AR, Costa PT, Jr., Wethington E, Eaton W. Turning points and lessons learned: stressful life events and personality trait development across middle adulthood. *Psychol Aging.* 2010;25(3): 524-533. <https://doi.org/10.1037/a0018751>.
39. Stienen MN, Scholtes F, Samuel R, Weil A, Weyerbrock A, Surbeck W. Different but similar: personality traits of surgeons and internists-results of a cross-sectional observational study. *BMJ Open.* 2018;8(7): e021310. <https://doi.org/10.1136/bmjopen-2017-021310>.
40. Gosling SD, Rentfrow PJ, Swann WB. A very brief measure of the Big-Five personality domains. *J Res Pers.* 2003;37(6): 504-528. [https://doi.org/10.1016/S0092-6566\(03\)00046-1](https://doi.org/10.1016/S0092-6566(03)00046-1).
41. Muck PM, Hell B, Gosling SD. Construct validation of a short five-factor model instrument - A self-peer study on the German adaptation of the Ten-Item Personality Inventory (TIPI-G). *Eur J Psychol Assess.* 2007;23(3): 166-175. <https://doi.org/10.1027/1015-5759.23.3.166>.
42. McCulloch P, Kaul A, Wagstaff GF, Wheatcroft J. Tolerance of uncertainty, extroversion, neuroticism and attitudes to randomized controlled trials among surgeons and physicians. *Br J Surg.* 2005;92(10): 1293-1297. <https://doi.org/10.1002/bjs.4930>.

43. Heros RC. Neurosurgical education: the "other" competencies. The 2003 presidential address. *J Neurosurg.* 2003;99(4): 623-629. <https://doi.org/10.3171/jns.2003.99.4.0623>.
44. Yule S, Flin R, Paterson-Brown S, Maran N. Non-technical skills for surgeons in the operating room: a review of the literature. *Surgery.* 2006;139(2): 140-149. <https://doi.org/10.1016/j.surg.2005.06.017>.
45. Henaux PL, Jannin P, Riffaud L. Nontechnical Skills in Neurosurgery: A Systematic Review of the Literature. *World Neurosurg.* 2019;130: e726-e736. <https://doi.org/10.1016/j.wneu.2019.06.204>.
46. Wortman J, Lucas RE, Donnellan MB. Stability and change in the Big Five personality domains: evidence from a longitudinal study of Australians. *Psychol Aging.* 2012;27(4): 867-874. <https://doi.org/10.1037/a0029322>.
47. Costa PT, Jr., McCrae RR, Zonderman AB, Barbano HE, Lebowitz B, Larson DM. Cross-sectional studies of personality in a national sample: 2. Stability in neuroticism, extraversion, and openness. *Psychol Aging.* 1986;1(2): 144-149.
48. McCrae RR, Costa PT, Jr., Pedroso de Lima M, et al. Age differences in personality across the adult life span: parallels in five cultures. *Dev Psychol.* 1999;35(2): 466-477. <https://doi.org/10.1037//0012-1649.35.2.466>.
49. Whitaker M. The surgical personality: does it exist? *Ann R Coll Surg Engl.* 2018;100(1): 72-77. <https://doi.org/10.1308/rcsann.2017.0200>.
50. Siddaiah-Subramanya M, To H, Haigh C. The psychosocial impact of surgical complications on the operating surgeon: A scoping review. *Ann Med Surg (Lond).* 2021;67: 102530. <https://doi.org/10.1016/j.amsu.2021.102530>.
51. Gawande AA, Zinner MJ, Studdert DM, Brennan TA. Analysis of errors reported by surgeons at three teaching hospitals. *Surgery.* 2003;133(6): 614-621. <https://doi.org/10.1067/msy.2003.169>.
52. Shanafelt TD, Balch CM, Bechamps G, et al. Burnout and medical errors among American surgeons. *Ann Surg.* 2010;251(6): 995-1000. <https://doi.org/10.1097/SLA.0b013e3181bfdab3>.
53. Balch CM, Shanafelt T. Combating stress and burnout in surgical practice: a review. *Adv Surg.* 2010;44: 29-47. <https://doi.org/10.1016/j.yasu.2010.05.018>.
54. Han K, Bohnen JD, Peponis T, et al. The Surgeon as the Second Victim? Results of the Boston Intraoperative Adverse Events Surgeons' Attitude (BISA) Study. *J Am Coll Surg.* 2017;224(6): 1048-1056. <https://doi.org/10.1016/j.jamcollsurg.2016.12.039>.
55. Patel AM, Ingalls NK, Mansour MA, Sherman S, Davis AT, Chung MH. Collateral damage: the effect of patient complications on the surgeon's psyche. *Surgery.* 2010;148(4): 824-828; discussion 828-830. <https://doi.org/10.1016/j.surg.2010.07.024>.
56. Arnold-Forster A. 'A small cemetery': death and dying in the contemporary British operating theatre. *Med Humanit.* 2020;46(3): 278-287. <https://doi.org/10.1136/medhum-2019-011668>.
57. Cuijpers P, Smit F, Penninx BW, de Graaf R, ten Have M, Beekman AT. Economic costs of neuroticism: a population-based study. *Arch Gen Psychiatry.* 2010;67(10): 1086-1093. <https://doi.org/10.1001/archgenpsychiatry.2010.130>.
58. Anglim J, Horwood S, Smillie LD, Marrero RJ, Wood JK. Predicting psychological and subjective well-being from personality: A meta-analysis. *Psychol Bull.* 2020;146(4): 279-323. <https://doi.org/10.1037/bul0000226>.
59. Zell E, Lesick TL. Big five personality traits and performance: A quantitative synthesis of 50+ meta-analyses. *J Pers.* 2021. <https://doi.org/10.1111/jopy.12683>.
60. Ullstrom S, Andreen Sachs M, Hansson J, Ovreteit J, Brommels M. Suffering in silence: a qualitative study of second victims of adverse events. *BMJ Qual Saf.* 2014;23(4): 325-331. <https://doi.org/10.1136/bmjqs-2013-002035>.
61. Pani JR, Chariker JH. The psychology of error in relation to medical practice. *J Surg Oncol.* 2004;88(3): 130-142. <https://doi.org/10.1002/jso.20123>.

62. Engel KG, Rosenthal M, Sutcliffe KM. Residents' responses to medical error: coping, learning, and change. *Acad Med*. 2006;81(1): 86-93. <https://doi.org/10.1097/00001888-200601000-00021>.
63. Rosenthal R, Hoffmann H, Dwan K, Clavien PA, Bucher HC. Reporting of adverse events in surgical trials: critical appraisal of current practice. *World J Surg*. 2015;39(1): 80-87. <https://doi.org/10.1007/s00268-014-2776-8>.
64. Allemand M, Schaffhuser K, Martin M. Long-term correlated change between personality traits and perceived social support in middle adulthood. *Pers Soc Psychol Bull*. 2015;41(3): 420-432. <https://doi.org/10.1177/0146167215569492>.
65. Roberts BW, Luo J, Briley DA, Chow PI, Su R, Hill PL. A systematic review of personality trait change through intervention. *Psychol Bull*. 2017;143(2): 117-141. <https://doi.org/10.1037/bul0000088>.
66. Kennair LEO, Solem S, Hagen R, Havnen A, Nysaeter TE, Hjemdal O. Change in personality traits and facets (Revised NEO Personality Inventory) following metacognitive therapy or cognitive behaviour therapy for generalized anxiety disorder: Results from a randomized controlled trial. *Clin Psychol Psychother*. 2021;28(4): 872-881. <https://doi.org/10.1002/cpp.2541>.

Table 1. Predicting levels of neuroticism for non-surgeons, surgeons, and occasionals.

	Model 1	Model 2	Model 3	Model 4
Specialty group (ref: Non-surgeons)				
Surgeons	-0.330*** (0.0470)	-0.501*** (0.151)	-0.651** (0.242)	0.694 (0.618)
Occasionals	-0.00707 (0.0420)	-0.155 (0.128)	-0.140 (0.177)	0.944 (0.554)
Age/100	-0.446** (0.155)	-0.688** (0.228)	-1.607*** (0.345)	-5.950** (1.987)
Male (ref: Female)	-0.428*** (0.0380)	-0.429*** (0.0380)	-1.118*** (0.173)	-1.056* (0.515)
Surgeons × Age/100		0.486 (0.411)	0.643 (0.729)	-7.102* (3.354)
Occasionals × Age/100		0.411 (0.335)	0.290 (0.504)	-5.735 (3.025)
Surgeons × Male			0.494 (0.319)	-0.662 (0.875)
Occasionals × Male			-0.0493 (0.269)	-0.990 (0.836)
Male × Age/100			1.772*** (0.464)	1.883 (2.691)
Surgeons × Male × Age/100			-0.851 (0.899)	5.806 (4.575)
Occasionals × Male × Age/100			0.244 (0.695)	5.616 (4.310)
Age/100 × Age/100				5.673* (2.557)
Surgeons × Age/100 × Age/100				10.10* (4.275)
Occasionals × Age/100 × Age/100				7.418 (3.824)
Male × Age/100 × Age/100				-0.738 (3.299)
Surgeons × Male × Age/100 × Age/100				-8.643 (5.622)
Occasionals × Male × Age/100 × Age/100				-6.802 (5.168)
Constant	2.262*** (0.0592)	2.347*** (0.0838)	2.676*** (0.119)	3.432*** (0.361)
Observations	4,678	4,678	4,678	4,678
Adjusted R ²	0.049	0.049	0.055	0.066

Standard errors in parentheses

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$




Figure 1. Linear predictions of levels of neuroticism across age for non-surgeons, surgeons, and occasionals. Shaded areas are 95% confidence intervals. Estimates based on Model 4 (see Table 1), restricted to respondents aged 20 to 70 and averaged over gender.

Abbreviations list

Analysis of variance (ANOVA)

Five-factor model (FFM)

Institutional review board of the Canton St.Gallen, Switzerland (Ethikkommission des Kantons St. Gallen (EKSG))

Standard deviation (SD)

Ten-Item Personality Inventory (TIPI)