

CATEGORICAL TORELLI THEOREMS FOR GUSHEL–MUKAI THREEFOLDS

AUGUSTINAS JACOVSKIS, XUN LIN, ZHIYU LIU AND SHIZHUO ZHANG

ABSTRACT. We show that a general ordinary Gushel–Mukai (GM) threefold X can be reconstructed from its Kuznetsov component $\mathcal{Ku}(X)$ together with an extra piece of data coming from tautological subbundle of the Grassmannian $\mathrm{Gr}(2, 5)$. We also prove that $\mathcal{Ku}(X)$ determines the birational isomorphism class of X , while $\mathcal{Ku}(X')$ determines the isomorphism class of a special GM threefold X' if it is general. As an application, we prove a conjecture of Kuznetsov–Perry in dimension three under a mild assumption. Finally, we use $\mathcal{Ku}(X)$ to restate a conjecture of Debarre–Iliev–Manivel regarding fibers of the period map for ordinary GM threefolds.

CONTENTS

1. Introduction	1
2. Semiorthogonal decompositions	6
3. Gushel–Mukai threefolds and their derived categories	7
4. Bridgeland stability conditions	8
5. Projection of \mathcal{E} into $\mathcal{Ku}(X)$	11
6. Conics on GM threefolds	13
7. Conics and Bridgeland moduli spaces	15
8. The moduli space $M_G(2, 1, 5)$ for GM threefolds	22
9. Refined and birational categorical Torelli theorems for GM threefolds	27
10. The Debarre–Iliev–Manivel conjecture	29
Appendix A. Uniqueness of Serre-invariant stability conditions	31
References	37

1. INTRODUCTION

In recent times, derived categories have played an important role in algebraic geometry; in many cases, much of the geometric information of a variety/scheme X is encoded by its bounded derived category of coherent sheaves $\mathrm{D}^b(X)$. In this setting, one of the most fundamental questions that can be asked is whether $\mathrm{D}^b(X)$ recovers X up to isomorphism, in other words, whether a *derived Torelli theorem* holds for X . For varieties with ample or anti-ample canonical bundle (which include Fano varieties and varieties of general type), this question was answered affirmatively by Bondal–Orlov in [BO01].

1.1. Kuznetsov components and categorical Torelli theorems. Therefore, for the class of varieties above, it is natural to ask whether they are also determined up to isomorphism by *less* information than the whole derived category $\mathrm{D}^b(X)$. A natural candidate for this is a subcategory $\mathcal{Ku}(X)$ of $\mathrm{D}^b(X)$ called the *Kuznetsov component*. This subcategory has been studied extensively by Kuznetsov and others (e.g. [Kuz04, Kuz09, KP18]) for many Fano varieties, including Gushel–Mukai (GM) varieties.

The question of whether $\mathcal{Ku}(X)$ determines X up to isomorphism has been studied for certain cases in the setting of Fano threefolds. In [BMMS12], the authors show that the Kuznetsov component completely determines cubic threefolds up to isomorphism, in other words, a *categorical Torelli theorem* holds for

Date: November 10, 2023.

2010 Mathematics Subject Classification. Primary 14F05; secondary 14J45, 14D20, 14D23.

Key words and phrases. Derived categories, Bridgeland moduli spaces, Kuznetsov components, Gushel–Mukai threefolds, Categorical Torelli theorem.

cubic threefolds Y . The same result was also verified in [PY22]. On the other hand, for many Fano varieties, the Kuznetsov component $Ku(X)$ does not determine the isomorphism class, but only the birational isomorphism class of X . This is known as a *birational categorical Torelli theorem*. For instance, Kuznetsov components determine the birational isomorphism class of every index 1 prime Fano threefolds of even genus $g \geq 8$. For GM threefolds – the focus of our paper – by [KP23] it is known that there are birational GM threefolds with equivalent Kuznetsov components. So there are two natural questions to ask in this setting:

Question 1.1.

- (1) Does $Ku(X)$ determine the birational equivalence class of X ?
- (2) What extra data along with $Ku(X)$ do we need to identify a particular GM threefold X from its birational equivalence class?

1.2. Main Results.

1.2.1. (Refined) categorical Torelli for Gushel–Mukai threefolds. In the present paper, we deal with the case of index 1 prime Fano threefolds of degree 10 and genus 6, also known as Gushel–Mukai threefolds (GM threefolds for short), which are split into two types: ordinary GM threefolds which arise as a quadric section of a linear section of the Grassmannian $\mathrm{Gr}(2, 5)$, and special GM threefolds which arise as double covers of a codimension three linear section of $\mathrm{Gr}(2, 5)$, branched over a degree ten K3 surface. By [KP18], we have a semiorthogonal decomposition

$$D^b(X) = \langle Ku(X), \mathcal{E}, \mathcal{O}_X \rangle,$$

where \mathcal{E} is the pull-back of the tautological subbundle on $\mathrm{Gr}(2, 5)$ along the natural map $X \rightarrow \mathrm{Gr}(2, 5)$.

Our first main theorem is concerned with ordinary GM threefolds and answers Question 1.1 (2):

Theorem 1.2 (Theorem 9.2). *Let X be a general ordinary GM threefold and $\pi: D^b(X) \rightarrow Ku(X)$ be the right adjoint to the inclusion $Ku(X) \subset D^b(X)$. Then the data of $Ku(X)$ along with the object $\pi(\mathcal{E})$ is enough to determine X up to isomorphism.*

On the other hand, for special GM threefolds which are general (“general special” for short), we show that a categorical Torelli theorem holds:

Theorem 1.3 (Theorem 9.9). *Let X and X' be general special GM threefolds, and assume that there is an equivalence of categories $Ku(X) \simeq Ku(X')$. Then X and X' are isomorphic.*

1.2.2. Birational categorical Torelli for Gushel–Mukai threefolds. Next, returning to the setting of ordinary GM threefolds, we show that a birational categorical Torelli theorem holds for general ordinary GM threefolds, which answers Question 1.1 (1).

Theorem 1.4 (Theorem 9.3). *Let X and X' be general ordinary GM threefolds, and suppose that there is an equivalence of categories $Ku(X) \simeq Ku(X')$. Then X is birationally equivalent to X' .*

In [KP23], the authors studied GM varieties of arbitrary dimension and proved the Duality Conjecture [KP18, Conjecture 3.7] for them, i.e. they showed that the period partner or period dual of a GM variety X shares the same Kuznetsov component $Ku(X)$ as X . Combining earlier results [DK18, Theorem 4.20] on the birational equivalence of these varieties, this gives strong evidence for the following conjecture:

Conjecture 1.5 ([KP23, Conjecture 1.7]). *If X and X' are GM varieties of the same dimension such that there is an equivalence $Ku(X) \simeq Ku(X')$, then X and X' are birationally equivalent.*

Thus our result Theorem 1.4 actually proves Conjecture 1.5 under the assumption that X and X' are both of dimension 3, ordinary and general.

Moreover, by a careful study of Bridgeland moduli spaces of stable objects in the Kuznetsov components \mathcal{A}_X for not only smooth ordinary GM threefolds but also special GM threefolds X , we can prove that the Kuznetsov component of a general ordinary GM threefold can not be equivalent to the one of a general special GM threefold. Therefore, combined with Theorem 1.4 and 1.3, we have the following improved version of Theorem 1.4, which allows threefolds to be either ordinary or special:

Theorem 1.6 (Theorem 9.7 and Corollary 9.8). *If X and X' are general ordinary or general special GM threefolds such that there is an equivalence $Ku(X) \simeq Ku(X')$, then X and X' are birationally equivalent.*

1.2.3. *The Debarre–Iliev–Manivel Conjecture.* In [DIM12], the authors conjecture that the general fiber of the classical period map from the moduli space of ordinary GM threefolds to the moduli space of 10 dimensional principally polarised abelian varieties is birational to the disjoint union of the minimal model $\mathcal{C}_m(X)$ of the Fano surface of conics and a moduli space of stable sheaves $M_G^X(2, 1, 5)$, both quotiented by involutions, which we call the *Debarre–Iliev–Manivel Conjecture* (cf. Conjecture 10.1). Within the moduli space of smooth ordinary GM threefolds, we define the fiber of the “categorical period map” through $[X]$ as the isomorphism classes of all ordinary GM threefolds X' whose Kuznetsov components satisfy $Ku(X') \simeq Ku(X)$. Then the following categorical analogue of the *Debarre–Iliev–Manivel conjecture* follows from Theorem 1.4 and results on Bridgeland moduli spaces with respect to the two (-1) -classes in the numerical Grothendieck group of \mathcal{A}_X .

Theorem 1.7 (Theorem 10.3). *A general fiber of the “categorical period map” through an ordinary GM threefold X is the union of $\mathcal{C}_m(X)/\iota$ and $M_G^X(2, 1, 5)/\iota'$ where ι, ι' are geometrically meaningful involutions.*

As an application, the *Debarre–Iliev–Manivel Conjecture* 10.1 can be restated in an equivalent form as follows:

Conjecture 1.8. *Let X be a general ordinary GM threefold. The intermediate Jacobian $J(X)$ determines the Kuznetsov component $Ku(X)$.*

Remark 1.9. In [DIM12], the authors actually conjecture that a general fiber of the period map is *birational* to the disjoint union of two surfaces, parametrizing conic transforms and conic transforms of a line transform of X , which is birational to the disjoint union of $\mathcal{C}_m(X)$ and $M_G^X(2, 1, 5)$, both quotiented by involutions. In Corollary 9.5 we show that this birational equivalence is indeed an *isomorphism*.

1.2.4. *Uniqueness of Serre-invariant stability conditions.* One of the key steps when we identify Bridgeland moduli spaces via an equivalence of Kuznetsov components in the proofs of Theorems 1.2 and 1.4. A stability condition σ on the Kuznetsov component $Ku(X)$ of a prime Fano threefold X is *Serre-invariant* if $S_{Ku(X)} \cdot \sigma = \sigma \cdot g$ for some $g \in \widetilde{GL}^+(2, \mathbb{R})$ (see Section 4.4). Serre-invariance is one of the fundamental tools in studying relationship of classical Gieseker moduli spaces and Bridgeland moduli spaces for Kuznetsov components (cf. [APR22, PY22, Zha20, LZ22, FP23]). A natural question is whether any two Serre-invariant stability conditions are in the same $\widetilde{GL}^+(2, \mathbb{R})$ -orbit. In the present paper, we answer this question affirmatively.

Theorem 1.10 (Theorem A.10). *Let X be a prime Fano threefold of index 1 of genus $g \geq 6$, or a del Pezzo threefold of degree $d \geq 2$. Then all Serre-invariant stability conditions on $Ku(X)$ are in the same $\widetilde{GL}^+(2, \mathbb{R})$ -orbit.*

1.3. **Methods.** For convenience, we work with the alternative Kuznetsov component \mathcal{A}_X , defined by the semiorthogonal decomposition $D^b(X) = \langle \mathcal{A}_X, \mathcal{O}_X, \mathcal{E}^\vee \rangle$ and there is an equivalence $\Xi: Ku(X) \simeq \mathcal{A}_X$. We prove the above theorems 1.2, 1.4, 1.6 and 1.7 by considering the moduli spaces of Bridgeland stable objects in the alternative Kuznetsov component \mathcal{A}_X with respect to (-1) -classes in the numerical Grothendieck group of \mathcal{A}_X , i.e. a vector v with $\chi(v, v) = -1$ where χ is the Euler form. Up to sign, there are two (-1) -classes in the numerical Grothendieck group of \mathcal{A}_X , call them $-x$ and $y - 2x$.

First, we show that the moduli space with the class $-x$ is isomorphic to the minimal model $\mathcal{C}_m(X)$ of the Fano surface of conics (Theorem 7.12). Indeed, we first show that the unique exceptional curve contracted in $\mathcal{C}(X)$ is the rational curve of conics whose ideal sheaf I_C is not in \mathcal{A}_X and that the image is the smooth point represented by $\pi(\mathcal{E})$ (Proposition 7.1), so $\mathcal{C}_m(X)$ forms an irreducible component of the moduli space $\mathcal{M}_\sigma(\mathcal{A}_X, -x)$ of stable objects in \mathcal{A}_X with respect to $-x$.

Then we show this component actually occupies the whole moduli space (Proposition 7.11), which is the most difficult and technical part of the article and we only briefly sketch the argument here. We start

with a stable object $F \in \mathcal{A}_X$ of the class $-x$. It suffices to show that F is isomorphic to the projection of ideal sheaf I_C of a conic $C \subset X$. First, we assume that F is semistable in the double tilted heart $\text{Coh}_{\alpha,\beta}^0(X)$ (cf. Section 4.4). Then by a wall-crossing argument, we prove that $F[-1]$ is a slope-semistable sheaf of rank one. Since its class is $[F] = -[I_C]$, we get $F \cong I_C[1]$. Next, we assume that F is not semistable in the double tilted heart $\text{Coh}_{\alpha,\beta}^0(X)$. Our main tools are inequalities in [Li19] and Theorem 4.7, which allow us to bound the rank and first two Chern characters ch_1, ch_2 of the destabilizing objects and their cohomology objects. Since $F \in \mathcal{A}_X$, by using the Euler characteristics $\chi(\mathcal{O}_X, -)$ and $\chi(\mathcal{E}^\vee, -)$ we can obtain a bound on ch_3 . Then we deduce that the Harder–Narasimhan factors of F are the expected ones (Proposition 7.10). As a result, $\mathcal{M}_\sigma(\mathcal{A}_X, -x) \cong \mathcal{C}_m(X)$. Similarly, we identify the moduli space $M_G^X(2, 1, 5)$ of Gieseker semistable sheaves of rank 2, $c_1 = 1, c_2 = 5$ and $c_3 = 0$ on X with the Bridgeland moduli space $\mathcal{M}_\sigma(\mathcal{A}_X, y - 2x)$ in Theorem 8.9.

As we have seen, $\mathcal{C}(X)$ is exactly the blow-up of $\mathcal{C}_m(X) \cong \mathcal{M}_\sigma(\mathcal{A}_X, -x)$ at the point $\Xi(\pi(\mathcal{E}))$, hence the data $(\mathcal{K}u(X), \pi(\mathcal{E}))$ determines $\mathcal{C}(X)$. A classical result of Logachev [Log12] states that X can be determined up to isomorphism from $\mathcal{C}(X)$. Thus Theorem 1.2 is proved.

We prove Theorem 1.3 via another method. By considering the equivariant Kuznetsov components $\mathcal{K}u(X)^{\mu_2}$, first discussed in [KP17], and exploiting the fact that X is the double cover of a degree 5 index 2 prime Fano threefold Y , branched over a quadric hypersurface $\mathcal{B} \subset Y$. In this case, the equivariant Kuznetsov component is equivalent to $D^b(\mathcal{B})$ where \mathcal{B} is a K3 surface. Therefore, a number of results concerning the Fourier–Mukai partners of K3 surfaces can be used to deduce that $\mathcal{K}u(X)^{\mu_2} \simeq \mathcal{K}u(X')^{\mu_2}$ implies $\mathcal{B} \cong \mathcal{B}'$. Then the fact that the del Pezzo threefold Y of degree 5 is rigid can be used to deduce that indeed, $X \cong X'$.

To prove Theorem 1.4, we invoke a few more results from [DIM12]. More precisely, an equivalence of categories $\Phi: \mathcal{A}_X \simeq \mathcal{A}_{X'}$ identifies the moduli space $\mathcal{M}_\sigma(\mathcal{A}_X, -x)$ with either $\mathcal{M}_\sigma(\mathcal{A}_X, -x)$ or $\mathcal{M}_\sigma(\mathcal{A}_{X'}, y - 2x)$. The former case gives an isomorphism of minimal surfaces $\mathcal{C}_m(X) \cong \mathcal{C}_m(X')$. Blowing $\mathcal{C}_m(X)$ up at the smooth point associated to $\pi(\mathcal{E})$ gives $\mathcal{C}(X)$, and blowing up $\mathcal{C}_m(X')$ at the image of $\pi(\mathcal{E})$ under Φ gives $\mathcal{C}(X'_C)$, where X'_C is certain birational transformation of X' , associated with a conic $C \subset X'$. Then by Logachev’s Reconstruction Theorem for $\mathcal{C}(X)$, X is isomorphic to X'_C which is birational to X' . For the latter case, we start with the isomorphism $\mathcal{C}_m(X) \cong M_G^{X'}(2, 1, 5)$. In fact, $M_G^{X'}(2, 1, 5)$ is birational to $\mathcal{C}(X'_L)$, where X'_L is another birational transformation of X' , associated with a line $L \subset X'$. Since $\mathcal{C}(X'_L)$ is a surface of general type, we get $\mathcal{C}_m(X) \cong \mathcal{C}_m(X'_L)$. Then by the same argument as in the previous case, X is isomorphic to some birational transformation of X' .

Finally, the proof of Theorem 1.6 is similar to that of Theorem 1.4. Firstly, we identify the Bridgeland moduli spaces $\mathcal{M}_\sigma(\mathcal{A}_{X'}, -x)$ and $\mathcal{M}_\sigma(\mathcal{A}_{X'}, y - 2x)$ on a special GM threefold X' with $\mathcal{C}_m(X')$ and $M_G^{X'}(2, 1, 5)$ respectively (Theorem 7.12 and Theorem 8.9), where $\mathcal{C}_m(X')$ is the contraction of the Fano surface $\mathcal{C}(X')$ of conics on X' along one of the components to a singular point. Then if X is ordinary, the equivalence $\Phi: \mathcal{A}_X \simeq \mathcal{A}_{X'}$ would identify those moduli spaces on a general ordinary GM threefold X with those on a special GM threefold X' ; we show that this is impossible by analyzing their singularities. Then Theorem 1.6 reduces to Theorem 1.4 and Theorem 1.3.

1.4. Related Work.

1.4.1. *Categorical Torelli theorems.* There is a very nice survey article [PS23] on recent results and remaining open questions on this topic. In [BMMS12] and [PY22], the authors prove categorical Torelli theorems for cubic threefolds. In [APR22] and [BT16], the authors prove categorical Torelli theorems for general quartic double solids. In [LNSZ21] and [LSZ22], the authors prove a refined categorical Torelli theorem for Enriques surfaces. In [JLZ22], the authors generalize Theorem 9.2 to all prime Fano threefolds of genus $g \geq 6$. In [GLZ22], the authors prove a birational categorical Torelli theorem for general non-Hodge-special Gushel–Mukai fourfolds.

1.4.2. *Identifying classical moduli spaces as Bridgeland moduli spaces for Kuznetsov components.* In the present article, we realize the Fano surface of conics and a certain Gieseker moduli space of semistable sheaves as Bridgeland moduli spaces of stable objects in Kuznetsov components of GM threefolds. In

[PY22], the authors realize the Fano surface of lines $\Sigma(Y_d)$ (for $d \geq 2$) as a Bridgeland moduli space of stable objects in the Kuznetsov component $\mathcal{Ku}(Y_d)$. In [LZ22], the authors realize the moduli space of rank two instanton sheaves on a del Pezzo threefold Y_d (for $d \geq 3$) and the compactification of the moduli space of ACM sheaves on X_{4d+2} (for $d \geq 3$) as Bridgeland moduli spaces of stable objects in $\mathcal{Ku}(Y_d)$ and $\mathcal{Ku}(X_{4d+2})$, respectively. In [FP23], the authors realize the moduli space of Ulrich bundles of arbitrary rank on a cubic threefold Y_3 as an open locus of a Bridgeland moduli space of stable objects in $\mathcal{Ku}(Y_3)$.

1.4.3. Serre-invariant stability conditions. In [PR23a] and [PY22], the authors prove that stability conditions on Kuznetsov components of every del Pezzo threefold Y_d of degree $d \geq 1$ and every index 1 prime Fano threefold of genus $g \geq 6$ are Serre-invariant. In [FP23], the authors prove the uniqueness of Serre-invariant stability conditions for a general triangulated category satisfying a list of very natural assumptions, which include Kuznetsov components of a series of prime Fano threefolds.

1.5. Notation and conventions.

- We work over the field $k = \mathbb{C}$. All triangulated categories and abelian categories are assumed to be k -linear.
- We use hom and ext^i to represent the dimension of the vector spaces Hom and Ext^i .
- The numerical K group of a triangulated category \mathcal{D} is denoted by $\mathcal{N}(\mathcal{D})$, which is the Grothendieck group $K_0(\mathcal{D})$ modulo the kernel of the Euler form $\chi(E, F) = \sum_i (-1)^i \mathrm{ext}^i(E, F)$.
- We denote the bounded derived category of a smooth projective variety X by $\mathrm{D}^b(X)$. The derived dual functor is denoted by $\mathbb{D} := \mathrm{RHom}_X(-, \mathcal{O}_X)$.
- We denote the phase and slope with respect to a weak stability condition σ by ϕ_σ and μ_σ , respectively. The maximal and minimal slopes (phases) of the Harder–Narasimhan factors of a given object F will be denoted by $\mu_\sigma^+(F)$ ($\phi_\sigma^+(F)$) and $\mu_\sigma^-(F)$ ($\phi_\sigma^-(F)$), respectively.
- $\mathcal{H}_\mathcal{A}^i$ means the i -th cohomology with respect to the heart \mathcal{A} . When the \mathcal{A} subscript is dropped, we take the heart to be $\mathrm{Coh}(X)$.
- The symbol \simeq denotes an equivalence of categories and a birational equivalence of varieties. The symbol \cong denotes an isomorphism of varieties.
- Let X be a GM threefold. Then a conic means a closed subscheme $C \subset X$ with Hilbert polynomial $p_C(t) = 1 + 2t$, and a line means a closed subscheme $L \subset X$ with Hilbert polynomial $p_L(t) = 1 + t$.

1.6. Organization of the paper. In Section 2, we collect basic facts about semiorthogonal decompositions. In Section 3, we introduce Gushel–Mukai threefolds and their Kuznetsov components. In Section 4, we introduce the definition of weak stability conditions on $\mathrm{D}^b(X)$, and the induced stability conditions on the alternative Kuznetsov components \mathcal{A}_X of GM threefolds. In Section 5, we introduce a distinguished object $\pi(\mathcal{E}) \in \mathcal{Ku}(X)$ and its alternative Kuznetsov component analogue $\Xi(\pi(\mathcal{E})) \in \mathcal{A}_X$ and prove its stability. In Section 6 we discuss the geometry of the Fano surface of conics of a GM threefold. In Section 7, we construct the Bridgeland moduli space of σ -stable objects with class $-x$ in \mathcal{A}_X . In Section 8, we construct the Bridgeland moduli space of σ -stable objects with respect to the other (-1) -class $y - 2x$ in \mathcal{A}_X . In Section 9, we prove several birational/refined categorical Torelli theorems (Theorems 1.2, 1.3 and 1.4) and Conjecture 1.5 in dimension three with mild assumptions. In Section 10, we describe the general fiber of the “categorical period map” for ordinary GM threefolds 1.7, and restate the *Debarre–Iliev–Manivel conjecture* in terms of Conjecture 10.6. Finally, we study Serre-invariant stability conditions on Kuznetsov components and show that they are contained in one $\widetilde{\mathrm{GL}}^+(2, \mathbb{R})$ orbit in Appendix A.

Acknowledgements. Firstly, it is our pleasure to thank Arend Bayer for very useful discussions on the topics of this project. We would like to thank Sasha Kuznetsov for answering many of our questions on Gushel–Mukai threefolds. We thank Atanas Iliev, Laurent Manivel, Daniele Faenzi, Dmitry Logachev, Will Donovan, Bernhard Keller, Alexey Elagin, Xiaolei Zhao, Chunyi Li, Laura Pertusi, Song Yang, Alex Perry, Pieter Belmans, Qingyuan Jiang, Enrico Fatighenti, Naoki Koseki, Bingyu Xia, Yong Hu and Luigi Martinelli for helpful conversations on several related topics. We would like to thank Daniele Faenzi for sending us the preprint [FV21] and Soheyla Feyzbakhsh for sending us the preprint [FP23]. We thank

Pieter Belmans for useful comments on the first draft of our article. The fourth author would like to thank Tingyu Sun for constant support and encouragement. We would like to thank the anonymous referee for their careful reading of our paper, and for their very useful and insightful comments.

The first and last authors are supported by ERC Consolidator Grant WallCrossAG, no. 819864. The first author was also supported by the Luxembourg National Research Fund (FNR–17113194).

2. SEMIORTHOGONAL DECOMPOSITIONS

In this section, we collect some useful facts about semiorthogonal decompositions. Background on triangulated categories and derived categories of coherent sheaves can be found in [Huy06], for example. From now on, let $D^b(X)$ denote the bounded derived category of coherent sheaves on a smooth projective variety X , and for $E, F \in D^b(X)$, define

$$\mathrm{RHom}^\bullet(E, F) = \bigoplus_{i \in \mathbb{Z}} \mathrm{Hom}(E, F[i])[-i].$$

2.1. Exceptional collections and semiorthogonal decompositions.

Definition 2.1. Let \mathcal{D} be a triangulated category and $E \in \mathcal{D}$. We say that E is an *exceptional object* if $\mathrm{RHom}^\bullet(E, E) = k$. Now let (E_1, \dots, E_m) be a collection of exceptional objects in \mathcal{D} . We say it is an *exceptional collection* if $\mathrm{RHom}^\bullet(E_i, E_j) = 0$ for $i > j$.

Definition 2.2. Let \mathcal{D} be a triangulated category and \mathcal{C} be a triangulated subcategory of \mathcal{D} . We define the *right orthogonal complement* of \mathcal{C} in \mathcal{D} as the full triangulated subcategory

$$\mathcal{C}^\perp = \{X \in \mathcal{D} \mid \mathrm{Hom}(Y, X) = 0 \text{ for all } Y \in \mathcal{C}\}.$$

The *left orthogonal complement* is defined similarly, as

$${}^\perp\mathcal{C} = \{X \in \mathcal{D} \mid \mathrm{Hom}(X, Y) = 0 \text{ for all } Y \in \mathcal{C}\}.$$

Definition 2.3. Let \mathcal{D} be a triangulated category. We say a triangulated subcategory $\mathcal{C} \subset \mathcal{D}$ is *admissible* if the inclusion functor $i: \mathcal{C} \hookrightarrow \mathcal{D}$ has left adjoint i^* and right adjoint $i^!$.

Definition 2.4. Let \mathcal{D} be a triangulated category, and $(\mathcal{C}_1, \dots, \mathcal{C}_m)$ be a collection of full admissible subcategories of \mathcal{D} . We say that $\mathcal{D} = \langle \mathcal{C}_1, \dots, \mathcal{C}_m \rangle$ is a *semiorthogonal decomposition* of \mathcal{D} if $\mathcal{C}_j \subset \mathcal{C}_i^\perp$ for all $i > j$, and the subcategories $(\mathcal{C}_1, \dots, \mathcal{C}_m)$ generate \mathcal{D} , i.e. the category resulting from taking all shifts and cones of objects in the categories $(\mathcal{C}_1, \dots, \mathcal{C}_m)$ is equivalent to \mathcal{D} .

Let $S_{\mathcal{D}}$ be the Serre functor of \mathcal{D} , then we have the following standard result, see e.g. [BLMS23, Section 3]:

Proposition 2.5 ([BLMS23, Section 3]). *If $\mathcal{D} = \langle \mathcal{D}_1, \mathcal{D}_2 \rangle$ is a semiorthogonal decomposition, then $\mathcal{D} = \langle S_{\mathcal{D}}(\mathcal{D}_2), \mathcal{D}_1 \rangle = \langle \mathcal{D}_2, S_{\mathcal{D}}^{-1}(\mathcal{D}_1) \rangle$ are also semiorthogonal decompositions.*

2.2. Mutations. Let $\mathcal{C} \subset \mathcal{D}$ be an admissible triangulated subcategory. Then the *left mutation functor* $\mathbf{L}_{\mathcal{C}}$ through \mathcal{C} is defined as the functor lying in the canonical functorial exact triangle

$$ii^! \rightarrow \mathrm{id} \rightarrow \mathbf{L}_{\mathcal{C}}$$

and the *right mutation functor* $\mathbf{R}_{\mathcal{C}}$ through \mathcal{C} is defined similarly, by the triangle

$$\mathbf{R}_{\mathcal{C}} \rightarrow \mathrm{id} \rightarrow ii^*.$$

When $E \in D^b(X)$ is an exceptional object, and $F \in D^b(X)$ is any object, the left mutation $\mathbf{L}_E F$ fits into the triangle

$$E \otimes \mathrm{RHom}^\bullet(E, F) \rightarrow F \rightarrow \mathbf{L}_E F,$$

and the right mutation $\mathbf{R}_E F$ fits into the triangle

$$\mathbf{R}_E F \rightarrow F \rightarrow E \otimes \mathrm{RHom}^\bullet(F, E)^\vee.$$

Proposition 2.6 ([Kuz19, Lemma 2.6]). *Let $\mathcal{D} = \langle \mathcal{A}, \mathcal{B} \rangle$ be a semiorthogonal decomposition. Then*

$$S_{\mathcal{B}} = \mathbf{R}_{\mathcal{A}} \circ S_{\mathcal{D}} \quad \text{and} \quad S_{\mathcal{A}}^{-1} = \mathbf{L}_{\mathcal{B}} \circ S_{\mathcal{D}}^{-1}.$$

Lemma 2.7 ([Kuz10, Lemma 2.7]). *Let $\mathcal{D} = \langle \mathcal{C}_1, \mathcal{C}_2, \dots, \mathcal{C}_n \rangle$ be a semiorthogonal decomposition with all components being admissible. Then for each $1 \leq k \leq n-1$, there is a semiorthogonal decomposition*

$$\mathcal{D} = \langle \mathcal{C}_1, \dots, \mathcal{C}_{k-1}, \mathbf{L}_{\mathcal{C}_k} \mathcal{C}_{k+1}, \mathcal{C}_k, \mathcal{C}_{k+2}, \dots, \mathcal{C}_n \rangle$$

and for each $2 \leq k \leq n$ there is a semiorthogonal decomposition

$$\mathcal{D} = \langle \mathcal{C}_1, \dots, \mathcal{C}_{k-2}, \mathcal{C}_k, \mathbf{L}_{\mathcal{C}_k} \mathcal{C}_{k-1}, \mathcal{C}_{k+1}, \dots, \mathcal{C}_n \rangle.$$

3. GUSHEL–MUKAI THREEFOLDS AND THEIR DERIVED CATEGORIES

Let X be a prime Fano threefold of index 1 and degree $H^3 = 10$, where H is the ample generator of $\text{CaCl}(X)$. Then X is either a quadric section of a linear section of codimension 2 of the Grassmannian $\text{Gr}(2, 5)$, in which case it is called an ordinary Gushel–Mukai (GM) threefold, or X is a double cover of a degree 5 and index 2 Fano threefold Y ramified in a quadric hypersurface, in which case it is called a special GM threefold. In the latter case, it has a natural involution $\tau: X \rightarrow X$ induced by the double cover $\pi: X \rightarrow Y$. By [Muk92, BMK23], there exists a stable vector bundle \mathcal{E} of rank 2 with $c_1(\mathcal{E}) = -H$, $c_2(\mathcal{E}) = 4L$ and $c_3(\mathcal{E}) = 0$, where L is the class of a line on X . In addition, \mathcal{E} is exceptional and $H^\bullet(X, \mathcal{E}) = 0$. In fact, \mathcal{E} is the pullback of the tautological bundle on the Grassmannian $\text{Gr}(2, 5)$. By [DIM12, Proposition 4.1], \mathcal{E} is the unique stable sheaf with $c_1(\mathcal{E}) = -H$, $c_2(\mathcal{E}) = 4L$ and $c_3(\mathcal{E}) = 0$.

Furthermore, there is a standard short exact sequence

$$0 \rightarrow \mathcal{E} \rightarrow \mathcal{O}_X^{\oplus 5} \rightarrow \mathcal{Q} \rightarrow 0 \quad (1)$$

where \mathcal{Q} is the pull-back of the tautological quotient bundle on $\text{Gr}(2, 5)$ along the natural map $X \rightarrow \text{Gr}(2, 5)$. Since $\text{rk}(\mathcal{E}) = 2$, we have $\mathcal{E}(H) \cong \mathcal{E}^\vee$.

Definition 3.1. Let X be a GM threefold.

- The *Kuznetsov component* of X is defined as $\mathcal{Ku}(X) := \langle \mathcal{E}, \mathcal{O}_X \rangle^\perp$. In particular, it fits into the semiorthogonal decomposition $\text{D}^b(X) = \langle \mathcal{Ku}(X), \mathcal{E}, \mathcal{O}_X \rangle$;
- The *alternative Kuznetsov component* of X is defined as $\mathcal{A}_X := \langle \mathcal{O}_X, \mathcal{E}^\vee \rangle^\perp$. In particular, it fits into the semiorthogonal decomposition $\text{D}^b(X) = \langle \mathcal{A}_X, \mathcal{O}_X, \mathcal{E}^\vee \rangle$.

Remark 3.2. By [KP18, Proposition 2.6], there is a natural involutive autoequivalence functor $\tau_{\mathcal{A}} := S_{\mathcal{A}_X}[-2]$ of \mathcal{A}_X . When X is special, it is induced by the natural involution τ on X as $\tau_{\mathcal{A}} = \tau^*|_{\mathcal{A}_X}$.

Definition 3.3. The left adjoint to the inclusion $\mathcal{A}_X \hookrightarrow \text{D}^b(X)$ is given by $\text{pr} := \mathbf{L}_{\mathcal{O}_X} \mathbf{L}_{\mathcal{E}^\vee}: \text{D}^b(X) \rightarrow \mathcal{A}_X$. We call this the *projection functor*.

The analogous natural projection functor can be defined for $\mathcal{Ku}(X)$, and we denote it by $\text{pr}' := \mathbf{L}_{\mathcal{E}} \mathbf{L}_{\mathcal{O}_X}$.

3.1. Kuznetsov components. Let $K_0(\mathcal{D})$ denote the Grothendieck group of a triangulated category \mathcal{D} . We have the bilinear Euler form

$$\chi(E, F) = \sum_{i \in \mathbb{Z}} (-1)^i \text{ext}^i(E, F)$$

for $[E], [F] \in K_0(\mathcal{D})$. By the Hirzebruch–Riemann–Roch formula, it takes the following form on GM threefolds. We have [Kuz09, p. 5] $\chi(u, v) = \chi_0(u^* \cap v)$ where $u \mapsto u^*$ is an involution of $\oplus_{i=0}^3 H^i(X, \mathbb{Q})$ given by multiplication with $(-1)^i$ on $H^{2i}(X, \mathbb{Q})$, and χ_0 is given by

$$\chi_0(x + yH + zL + wP) = x + \frac{17}{6}y + \frac{1}{2}z + w,$$

where L is the class of lines and P is the class of points. The *numerical Grothendieck group* of \mathcal{D} is $\mathcal{N}(\mathcal{D}) = K_0(\mathcal{D}) / \ker \chi$.

Lemma 3.4 ([Kuz09, p. 5]). *The numerical Grothendieck group $\mathcal{N}(\mathcal{K}u(X))$ of the Kuznetsov component is a rank 2 integral lattice generated by the basis elements $v = 1 - 3L + \frac{1}{2}P$ and $w = H - 6L + \frac{1}{6}P$. Using this basis, χ is given by the matrix*

$$\begin{pmatrix} -2 & -3 \\ -3 & -5 \end{pmatrix}.$$

3.2. Alternative Kuznetsov components. As in [Kuz09, Proposition 3.9], the following lemma follows from a straightforward computation.

Lemma 3.5. *The numerical Grothendieck group of \mathcal{A}_X is a rank 2 integral lattice with basis vectors $x = 1 - 2L$ and $y = H - 4L - \frac{5}{6}P$, and the Euler form with respect to the basis is*

$$\begin{pmatrix} -1 & -2 \\ -2 & -5 \end{pmatrix}.$$

Remark 3.6. It is straightforward to check that the (-1) -classes of $\mathcal{N}(\mathcal{A}_X)$ are $x = 1 - 2L$ and $2x - y = 2 - H + \frac{5}{6}P$, up to sign.

Indeed, the Kuznetsov components from Subsection 3.1 and the alternative Kuznetsov components from this section are equivalent:

Lemma 3.7. *The original and alternative Kuznetsov components are equivalent. More precisely, there is an equivalence of categories $\Xi: \mathcal{K}u(X) \xrightarrow{\sim} \mathcal{A}_X$ given by $E \mapsto \mathbf{L}_{\mathcal{O}_X}(E \otimes \mathcal{O}_X(H))$, with inverse given by $F \mapsto (\mathbf{R}_{\mathcal{O}_X}F) \otimes \mathcal{O}_X(-H)$.*

Proof. Using Lemma 2.7 and noting that $\mathcal{E} \otimes \mathcal{O}_X(H) \cong \mathcal{E}^\vee$, we manipulate the semiorthogonal decomposition as follows:

$$\begin{aligned} \mathrm{D}^b(X) &= \langle \mathcal{K}u(X), \mathcal{E}, \mathcal{O}_X \rangle \\ &\simeq \langle \mathcal{K}u(X) \otimes \mathcal{O}_X(H), \mathcal{E}^\vee, \mathcal{O}_X(H) \rangle \\ &\simeq \langle \mathcal{O}_X, \mathcal{K}u(X) \otimes \mathcal{O}_X(H), \mathcal{E}^\vee \rangle \\ &\simeq \langle \mathbf{L}_{\mathcal{O}_X}(\mathcal{K}u(X) \otimes \mathcal{O}_X(H)), \mathcal{O}_X, \mathcal{E}^\vee \rangle. \end{aligned}$$

Now comparing with the definition of \mathcal{A}_X , we get $\mathcal{A}_X \simeq \mathbf{L}_{\mathcal{O}_X}(\mathcal{K}u(X) \otimes \mathcal{O}_X(H))$ and the desired result follows. The reverse direction is similar. \square

4. BRIDGELAND STABILITY CONDITIONS

In this section, we recall (weak) Bridgeland stability conditions on $\mathrm{D}^b(X)$, and the notions of tilt stability, double-tilt stability, and stability conditions induced on Kuznetsov components from weak stability conditions on $\mathrm{D}^b(X)$. We follow [BLMS23, § 2].

4.1. Weak stability conditions. Let \mathcal{D} be a triangulated category, and $K_0(\mathcal{D})$ its Grothendieck group. Fix a surjective morphism $v: K_0(\mathcal{D}) \rightarrow \Lambda$ to a finite rank lattice.

Definition 4.1. A *stability condition* (resp. *weak stability condition*) on \mathcal{D} is a pair $\sigma = (\mathcal{A}, Z)$ where \mathcal{A} is the heart of a bounded t-structure on \mathcal{D} , and $Z: \Lambda \rightarrow \mathbb{C}$ is a group homomorphism such that the following conditions hold:

- (1) The composition $Z \circ v: K_0(\mathcal{A}) \cong K_0(\mathcal{D}) \rightarrow \mathbb{C}$ satisfies: for any $E \neq 0 \in \mathcal{D}$ we have $\mathrm{Im} Z(E) \geq 0$ and if $\mathrm{Im} Z(E) = 0$ then $\mathrm{Re} Z(E) < 0$ (resp. $\mathrm{Re} Z(E) \leq 0$). From now on, we write $Z(E)$ rather than $Z(v(E))$.

We define a *slope function* μ_σ for σ using Z . For any $E \in \mathcal{A}$, set

$$\mu_\sigma(E) := \begin{cases} -\frac{\mathrm{Re} Z(E)}{\mathrm{Im} Z(E)}, & \mathrm{Im} Z(E) > 0 \\ +\infty, & \text{else.} \end{cases}$$

We say an object $0 \neq E \in \mathcal{A}$ is σ -(semi)stable if $\mu_\sigma(F) < \mu_\sigma(E/F)$ (respectively $\mu_\sigma(F) \leq \mu_\sigma(E/F)$) for all proper subobjects $F \subset E$.

- (2) Any object $E \in \mathcal{A}$ has a Harder–Narasimhan filtration in terms of σ -semistability defined above.
- (3) There exists a quadratic form Q on $\Lambda \otimes \mathbb{R}$ such that $Q|_{\ker Z}$ is negative definite, and $Q(E) \geq 0$ for all σ -semistable objects $E \in \mathcal{A}$. This is known as the *support property*.

Definition 4.2. Let $\sigma = (\mathcal{A}, Z)$ be a stability condition on \mathcal{D} . The *phase* of a σ -semistable object $E \in \mathcal{A}$ is

$$\phi(E) := \frac{1}{\pi} \arg(Z(E)) \in (0, 1].$$

Specially, if $Z(E) = 0$ then $\phi(E) = 1$. If $F = E[n]$, then we define

$$\phi(F) := \phi(E) + n$$

A *slicing* \mathcal{P} of \mathcal{D} consists of full additive subcategories $\mathcal{P}(\phi) \subset \mathcal{D}$ for each $\phi \in \mathbb{R}$ satisfying

- (1) for $\phi \in (0, 1]$, the subcategory $\mathcal{P}(\phi)$ is given by the zero object and all σ -semistable objects whose phase is ϕ ;
- (2) for $\phi + n$ with $\phi \in (0, 1]$ and $n \in \mathbb{Z}$, we set $\mathcal{P}(\phi + n) := \mathcal{P}(\phi)[n]$.

We will use both notations $\sigma = (\mathcal{A}, Z)$ and $\sigma = (\mathcal{P}, Z)$ for a stability condition σ with heart $\mathcal{A} = \mathcal{P}((0, 1])$ where \mathcal{P} is the slicing of σ .

We say σ is a *numerical stability condition* on \mathcal{D} if the surjective morphism $v: K_0(\mathcal{D}) \rightarrow \Lambda$ factors through the natural surjection $K_0(\mathcal{D}) \rightarrow \mathcal{N}(\mathcal{D})$ (assuming $\mathcal{N}(\mathcal{D})$ is well-defined).

Next, we recall two natural group actions on the set of stability conditions $\text{Stab}(\mathcal{D})$.

- (1) An element $\tilde{g} = (g, G)$ in the universal covering $\widetilde{\text{GL}}^+(2, \mathbb{R})$ of the group $\text{GL}^+(2, \mathbb{R})$ consists of an increasing function $g: \mathbb{R} \rightarrow \mathbb{R}$ such that $g(\phi + 1) = g(\phi) + 1$ and a matrix $G \in \text{GL}^+(2, \mathbb{R})$ with $\det(G) > 0$. It acts on the right on the stability manifold by $\sigma \cdot \tilde{g} := (G^{-1} \circ Z, \mathcal{P}(g(\phi)))$ for any $\sigma = (\mathcal{P}, Z) \in \text{Stab}(\mathcal{D})$ (see [Bri07, Lemma 8.2]).
- (2) Let $\text{Aut}_\Lambda(\mathcal{D})$ be the group of exact autoequivalences of \mathcal{D} , whose action Φ_* on $K_0(\mathcal{D})$ is compatible with $v: K_0(\mathcal{D}) \rightarrow \Lambda$. For $\Phi \in \text{Aut}_\Lambda(\mathcal{D})$ and $\sigma = (\mathcal{P}, Z) \in \text{Stab}(\mathcal{D})$, we define a left action of the group of linear exact autoequivalences $\text{Aut}_\Lambda(\mathcal{D})$ by $\Phi \cdot \sigma = (\Phi(\mathcal{P}), Z \circ \Phi_*^{-1})$, where Φ_* is the automorphism of $K_0(\mathcal{D})$ induced by Φ .

4.2. Tilt-stability. Let (X, H) be a polarised smooth projective variety of dimension n and $\sigma_H = (\text{Coh}(X), Z_H)$ be the standard weak stability condition on $\text{Coh}(X)$ defined as

$$Z_H(E) := -H^{n-1} \text{ch}_1(E) + iH^n \text{rk}(E).$$

Its σ_H -stability coincides with classical μ_H -stability (slope stability). Now for a fixed real number β , consider the following subcategories¹ of $\text{Coh}(X)$:

$$\begin{aligned} \mathcal{T}^\beta &= \langle E \in \text{Coh}(X) \mid E \text{ is } \sigma_H\text{-semistable with } \mu_{\sigma_H}(E) > \beta \rangle \\ \mathcal{F}^\beta &= \langle E \in \text{Coh}(X) \mid E \text{ is } \sigma_H\text{-semistable with } \mu_{\sigma_H}(E) \leq \beta \rangle. \end{aligned}$$

Then it is a result of [HRS96] that the tilted heart $\text{Coh}^\beta(X) := \langle \mathcal{T}^\beta, \mathcal{F}^\beta[1] \rangle$ is the heart of a bounded t-structure on $\text{Coh}(X)$.

Proposition 4.3 ([BMT14, BMS16]). *Let $\alpha > 0$ and $\beta \in \mathbb{R}$. Then the pair $\sigma_{\alpha, \beta} = (\text{Coh}^\beta(X), Z_{\alpha, \beta})$ defines a weak stability condition on $\text{D}^b(X)$, where*

$$Z_{\alpha, \beta}(E) = \frac{1}{2} \alpha^2 H^n \text{ch}_0^\beta(E) - H^{n-2} \text{ch}_2^\beta(E) + iH^{n-1} \text{ch}_1^\beta(E).$$

The quadratic form Q is given by the discriminant

$$\Delta_H(E) = (H^{n-1} \text{ch}_1(E))^2 - 2H^n \text{ch}_0(E) H^{n-2} \text{ch}_2(E).$$

We denote the slope function by $\mu_{\alpha, \beta} := \mu_{\sigma_{\alpha, \beta}}$.

¹The angle brackets here mean extension closure.

The weak stability conditions $\sigma_{\alpha,\beta}$ constructed above are also known as *tilt-stability* and the heart $\text{Coh}^\beta(X)$ are called the *tilted heart*.

Now pick a weak stability condition $\sigma_{\alpha,\beta}$. We define

$$\begin{aligned}\mathcal{T}_{\alpha,\beta}^0 &= \langle E \in \text{Coh}^\beta(X) \mid E \text{ is } \sigma_{\alpha,\beta}\text{-semistable with } \mu_{\alpha,\beta}(E) > 0 \rangle \\ \mathcal{F}_{\alpha,\beta}^0 &= \langle E \in \text{Coh}^\beta(X) \mid E \text{ is } \sigma_{\alpha,\beta}\text{-semistable with } \mu_{\alpha,\beta}(E) \leq 0 \rangle.\end{aligned}$$

Moreover, we “rotate” the stability function $Z_{\alpha,\beta}$ by setting

$$Z_{\alpha,\beta}^0 := \frac{1}{i} Z_{\alpha,\beta}.$$

Then we have the following result:

Proposition 4.4 ([BLMS23, Proposition 2.15]). *The pair $\sigma_{\alpha,\beta}^0 = (\text{Coh}_{\alpha,\beta}^0(X) = \langle \mathcal{T}_{\alpha,\beta}^0, \mathcal{F}_{\alpha,\beta}^0[1] \rangle, Z_{\alpha,\beta}^0)$ defines a weak stability condition on $\text{D}^b(X)$. We denote the slope function by $\mu_{\alpha,\beta}^0 := \mu_{\sigma_{\alpha,\beta}^0}$.*

We now state a useful lemma that relates 2-Gieseker-stability (see [BBF⁺20, Definition 4.3]) and tilt-stability.

Lemma 4.5 ([BMS16, Lemma 2.7], [BBF⁺20, Proposition 4.8, 4.9]). *Let $E \in \text{D}^b(X)$.*

- (1) *Let $\beta < \mu(E)$. Then $E \in \text{Coh}^\beta(X)$ is $\sigma_{\alpha,\beta}$ -(semi)stable for $\alpha \gg 0$ if and only if $E \in \text{Coh}(X)$ and E is 2-Gieseker-(semi)stable.*
- (2) *If $E \in \text{Coh}^\beta(X)$ is $\sigma_{\alpha,\beta}$ -semistable for $\beta \geq \mu(E)$ and $\alpha \gg 0$, then $\mathcal{H}^{-1}(E)$ is a torsion free μ -semistable sheaf and $\mathcal{H}^0(E)$ is supported in dimension not greater than one. If $\beta > \mu(E)$ and $\alpha > 0$, then $\mathcal{H}^{-1}(E)$ is also reflexive.*

4.3. Stronger BG inequalities. In this subsection, we state stronger Bogomolov–Gieseker (BG) style inequalities, which hold for tilt-semistable objects. These will be useful later on for ruling out potential walls for tilt-stability of objects in $\text{D}^b(X)$. The first is a stronger version of Proposition 4.3, which was proved by Chunyi Li in [Li19, Proposition 3.2] for Fano threefolds of Picard number one.

Lemma 4.6 (Stronger BG I). *Let X be an index 1 prime Fano threefold with degree d , and $E \in \text{D}^b(X)$ a $\sigma_{\alpha,\beta}$ -stable object where $\alpha > 0$. Let $k := \lfloor \mu(E) \rfloor$. Then we have:*

$$\frac{H \cdot \text{ch}_2(E)}{H^3 \cdot \text{ch}_0(E)} \leq \max \left\{ k\mu_H(E) - \frac{k^2}{2}, \frac{1}{2}\mu_H(E)^2 - \frac{3}{4d}, (k+1)\mu_H(E) - \frac{(k+1)^2}{2} \right\}.$$

Moreover, if the equality holds, then E has rank one or two.

The second is due to Naoki Koseki and Chunyi Li. It is based on [Kos20, Lemma 4.2, Theorem 4.3]. Chunyi Li also sent us a similar inequality from his upcoming paper [Li21].

Theorem 4.7 (Stronger BG II). *Let X be an index 1 Fano threefold of degree d , and $E \in \text{Coh}^0(X)$ be a $\sigma_{\alpha,0}$ -semistable object for some $\alpha > 0$ with $|\mu_H(E)| \in [0, 1]$ and $\text{rk}(E) \geq 2$. Then*

$$\frac{H \cdot \text{ch}_2(E)}{H^3 \cdot \text{ch}_0(E)} \leq \max \left\{ \frac{1}{2}\mu_H(E)^2 - \frac{3}{4d}, \mu_H(E)^2 - \frac{1}{2}|\mu_H(E)| \right\}.$$

Before we prove Theorem 4.7, we first state an easy lemma.

Lemma 4.8. *Let S be a K3 surface of degree d and H_S the ample polarisation. Let E be a μ_{H_S} -semistable sheaf in $\text{D}^b(S)$ with $\text{rk}(E) \geq 2$. Then*

$$\frac{\text{ch}_2(E)}{H_S^2 \cdot \text{rk}(E)} \leq \frac{1}{2}\mu_{H_S}(E)^2 - \frac{3}{4d}.$$

Proof. Let $v(E)$ be the Mukai vector of E . We have

$$\begin{aligned}v(E)^2 &= H_S^2 \cdot \text{ch}_1(E)^2 - 2\text{rk}(E)^2 - 2\text{rk}(E) \cdot \text{ch}_2(E) \\ &\geq -2 \geq -\frac{1}{2}\text{rk}(E)^2.\end{aligned}$$

Dividing through by $\text{rk}(E)^2$ and rearranging, we get

$$\frac{\text{ch}_2(E)}{\text{rk}(E)} \leq \frac{1}{2}\mu_{H_S}(E)^2 H_S^2 - \frac{3}{4}$$

as required. \square

Proof of Theorem 4.7. Let $f: [0, 1] \rightarrow \mathbb{R}$ be defined as

$$f(t) := \max \left\{ \frac{1}{2}t^2 - \frac{3}{4d}, t^2 - \frac{1}{2}t \right\}$$

Note that f is star-shaped [Kos20, Definition 3.2] and satisfies $f(0) = 0$ and $f(1) = 1/2$ as well as

$$t^2 - \frac{1}{2}t \leq f(t) \leq \frac{1}{2}t^2$$

for all $t \in [0, 1]$. We now follow the strategy of proof in [Kos20, Theorem 4.3]. Assume for a contradiction that there is an $E \in \mathcal{D}^b(X)$ such that the inequality in the statement of Theorem 4.7 is not true. Then conditions (a) and (b) in [Kos20, Lemma 3.3] are satisfied for f . Then by *loc. cit.*, the restriction $E|_{S_d}$ where S_d is a general hyperplane section of X_d is $\mu_{H_{S_d}}$ -semistable. Also note that $\mu_{H_{S_d}}(E|_{S_d}) = \mu_H(E)$ and S_d is a smooth K3 surface. But then by assumption

$$\frac{\text{ch}_2(E|_{S_d})}{H_{S_d}^2 \cdot \text{rk}(E|_{S_d})} > \frac{1}{2}\mu_{H_{S_d}}(E)^2 - \frac{3}{4d}$$

which contradicts Proposition 4.8, so the assumption is false and the result follows. \square

4.4. Stability conditions on the Kuznetsov component of a GM threefold. Proposition 5.1 in [BLMS23] gives a criterion for checking when weak stability conditions on a triangulated category can be used to induce stability conditions on a subcategory. Each of the criteria of this proposition can be checked for $\mathcal{A}_X \subset \mathcal{D}^b(X)$ to give stability conditions on \mathcal{A}_X .

More precisely, let $\mathcal{A}(\alpha, \beta) = \text{Coh}_{\alpha, \beta}^0(X) \cap \mathcal{A}_X$ and $Z(\alpha, \beta) = Z_{\alpha, \beta}^0|_{\mathcal{A}_X}$. Furthermore, if we take suitable (α, β) , by [BLMS23, Theorem 6.9] and [PR23a, Proposition 3.2] we have:

Theorem 4.9. *Let X be a GM threefold. Then $\sigma(\alpha, \beta)$ is a stability condition on \mathcal{A}_X for all $(\alpha, \beta) \in V$, where*

$$V := \{(\alpha, \beta) : -\frac{1}{10} < \beta < 0, 0 < \alpha < -\beta\}.$$

Now we introduce a special class of stability condition, which will play a central role in our paper.

Definition 4.10. Let σ be a stability condition on a triangulated category \mathcal{D} . It is called *Serre-invariant* if $S_{\mathcal{D}} \cdot \sigma = \sigma \cdot g$ for some $g \in \widetilde{\text{GL}}^+(2, \mathbb{R})$, where $S_{\mathcal{D}}$ is the Serre functor of \mathcal{D} .

We recall a recent result proved in [PR23a].

Theorem 4.11. *Let X be a GM threefold and σ (resp. σ') be a stability condition on $\mathcal{K}u(X)$ (resp. \mathcal{A}_X) defined by [BLMS23]. Then σ (resp. σ') is Serre-invariant.*

Proposition 4.12. *Let X be a GM threefold and E a non-zero object in \mathcal{A}_X such that $\text{ext}^1(E, E) \leq 3$ and $-\chi(E, E)$ is not a perfect square. Then E is σ -stable for every Serre-invariant stability condition σ on \mathcal{A}_X .*

Proof. The proof is the same as in [Zha20, Lemma 9.12]. We omit the details. \square

5. PROJECTION OF \mathcal{E} INTO $\mathcal{K}u(X)$

In this section, we consider the object that results from projecting the vector bundle \mathcal{E} into $\mathcal{K}u(X)$, and its stability in $\mathcal{K}u(X)$. We start with a lemma.

Lemma 5.1. *Let X be a GM threefold.*

- (1) $\text{RHom}^\bullet(\mathcal{Q}(-H), \mathcal{E}) = \text{RHom}^\bullet(\mathcal{E}, \mathcal{Q}^\vee) = k^2$ when X is ordinary.
- (2) $\text{RHom}^\bullet(\mathcal{Q}(-H), \mathcal{E}) = \text{RHom}^\bullet(\mathcal{E}, \mathcal{Q}^\vee) = k^3 \oplus k[-1]$ when X is special.

- (3) $\mathrm{RHom}^\bullet(\mathcal{E}, \mathcal{Q}(-H)) = k[-2]$.
 (4) $\mathrm{RHom}^\bullet(\mathcal{Q}^\vee, \mathcal{E}) = \mathrm{RHom}^\bullet(\mathcal{E}^\vee, \mathcal{Q}) = k[-2]$.

Proof. When X is ordinary, (1) and (2) follow from the Koszul resolution of $X \subset \mathrm{Gr}(2, 5)$ and the Borel–Weil–Bott Theorem. When X is special with the double cover $\pi: X \rightarrow Y$, note that $\pi_* \mathcal{O}_X = \mathcal{O}_Y \oplus \mathcal{O}_Y(-1)$. Then the (1) and (2) follow from the projection formula and [San14, Lemma 2.14, Proposition 2.15]. And applying $\mathrm{Hom}(-, \mathcal{E})$ to (1) and using Serre duality, we get $\mathrm{RHom}^\bullet(\mathcal{E}, \mathcal{Q}(-H)) = \mathrm{RHom}^\bullet(\mathcal{Q}, \mathcal{E})^\vee[-3] = k[-2]$, which proves (3). Finally, (4) follows from applying $\mathrm{Hom}(-, \mathcal{E})$ to (1) and using Serre duality and $\mathrm{RHom}^\bullet(\mathcal{E}, \mathcal{E}) = k$. \square

5.1. The projection of \mathcal{E} into $\mathcal{K}u(X)$. Let $\pi := \mathbf{R}_{\mathcal{O}_X(-H)} \mathbf{R}_{\mathcal{E}(-H)}: \mathrm{D}^b(X) \rightarrow \mathcal{K}u(X)$ be the right adjoint to the inclusion $\mathcal{K}u(X) \hookrightarrow \mathrm{D}^b(X)$. Here $\mathcal{K}u(X) = \langle \mathcal{E}, \mathcal{O}_X \rangle^\perp$ is the original Kuznetsov component.

Lemma 5.2. *The projection object $\pi(\mathcal{E})$ is the unique object that fits into a non-trivial exact triangle*

$$\mathcal{Q}(-H)[1] \rightarrow \pi(\mathcal{E}) \rightarrow \mathcal{E}. \quad (2)$$

Proof. By Serre duality, we have $\mathrm{RHom}^\bullet(\mathcal{E}, \mathcal{E}(-H)) = \mathrm{RHom}^\bullet(\mathcal{E}, \mathcal{E})^\vee[-3] = k[-3]$. Then we have an exact triangle $\mathbf{R}_{\mathcal{E}(-H)} \mathcal{E} \rightarrow \mathcal{E} \rightarrow \mathcal{E}(-H)[3]$. And by (1), we see $\mathbf{R}_{\mathcal{O}_X(-H)} \mathcal{E}(-H) = \mathcal{Q}(-H)[-1]$. Thus, from $\mathbf{R}_{\mathcal{O}_X(-H)} \mathcal{E} = \mathcal{E}$ we obtain the triangle (2). It is non-trivial since $\pi(\mathcal{E}) \in \mathcal{K}u(X)$, so \mathcal{E} cannot be a direct summand of $\pi(\mathcal{E})$. Finally, the uniqueness follows from Lemma 5.1 (4). \square

Lemma 5.3. *Let X be a GM threefold. Then we have*

- $\mathrm{RHom}^\bullet(\pi(\mathcal{E}), \pi(\mathcal{E})) = k \oplus k^2[-1]$ when X is ordinary.
- $\mathrm{RHom}^\bullet(\pi(\mathcal{E}), \pi(\mathcal{E})) = k \oplus k^3[-1] \oplus k[-2]$ when X is special.

Hence $\pi(\mathcal{E})$ is stable with respect to every Serre-invariant stability condition on $\mathcal{K}u(X)$.

Proof. The first statement follows from applying $\mathrm{Hom}(-, \mathcal{E})$ to triangle (2) and Lemma 5.1, and also the fact that $\mathrm{RHom}^\bullet(\pi(\mathcal{E}), \pi(\mathcal{E})) = \mathrm{RHom}^\bullet(\pi(\mathcal{E}), \mathcal{E})$ which is by adjunction. The last statement follows from Lemma 4.12. \square

5.2. The analogous projection object for \mathcal{A}_X . In this subsection, we state and prove the analogous results as in Subsection 5.1, except for \mathcal{A}_X instead of $\mathcal{K}u(X)$. Let $\pi' := \mathbf{R}_{\mathcal{E}} \mathbf{R}_{\mathcal{O}_X(-H)}: \mathrm{D}^b(X) \rightarrow \mathcal{A}_X$ be the right adjoint to the inclusion $\mathcal{A}_X \hookrightarrow \mathrm{D}^b(X)$.

Lemma 5.4. *The projection object $\pi'(\mathcal{Q}^\vee)$ is the unique object fits into a non-trivial exact triangle*

$$\mathcal{E}[1] \rightarrow \pi'(\mathcal{Q}^\vee) \rightarrow \mathcal{Q}^\vee. \quad (3)$$

Proof. The proof is completely analogous to the proof of Lemma 5.2. By Serre duality, we have the vanishing $\mathrm{RHom}^\bullet(\mathcal{Q}^\vee, \mathcal{O}_X(-H)) = \mathrm{RHom}^\bullet(\mathcal{O}_X, \mathcal{Q}^\vee)^\vee = 0$. Thus $\pi'(\mathcal{Q}^\vee) = \mathbf{R}_{\mathcal{E}} \mathcal{Q}^\vee$. Then the result follows from Lemma 5.1 (4). Finally, the uniqueness also follows from Lemma 5.1 (4) and (3) is non-trivial since $\mathrm{RHom}^\bullet(\mathcal{Q}^\vee, \pi'(\mathcal{Q}^\vee)) = 0$. \square

Remark 5.5. Later in Section 7, we will see that we have $\pi'(\mathcal{Q}^\vee) \cong \mathrm{pr}(I_C)[1]$ where $C \subset X$ is a conic such that $I_C \notin \mathcal{A}_X$.

Lemma 5.6. *Let X be a GM threefold. Then*

- $\mathrm{RHom}^\bullet(\pi'(\mathcal{Q}^\vee), \pi'(\mathcal{Q}^\vee)) = k \oplus k^2[-1]$ when X is ordinary.
- $\mathrm{RHom}^\bullet(\pi'(\mathcal{Q}^\vee), \pi'(\mathcal{Q}^\vee)) = k \oplus k^3[-1] \oplus k[-2]$ when X is special.

Hence $\pi'(\mathcal{Q}^\vee)$ is stable with respect to every Serre-invariant stability condition on \mathcal{A}_X .

Proof. It is not hard to check that $\Xi(\pi(\mathcal{E})) \cong \pi'(\mathcal{Q}^\vee)[1]$, where Ξ is the equivalence $\mathcal{K}u(X) \simeq \mathcal{A}_X$ in Lemma 3.7. Then the result follows from Lemma 5.3. \square

6. CONICS ON GM THREEFOLDS

In this section, we collect some useful results regarding the birational geometry of GM threefolds and their Hilbert schemes of conics. The results in this section are all from [DIM12], [Log12], and [Ili94a].

Recall that a conic means a closed subscheme $C \subset X$ with Hilbert polynomial $p_C(t) = 1 + 2t$, and a line means a closed subscheme $L \subset X$ with Hilbert polynomial $p_L(t) = 1 + t$. Denote their Hilbert schemes by $\mathcal{C}(X)$ and $\Gamma(X)$, respectively.

6.1. Conics on ordinary GM threefolds. Let X be an ordinary GM threefold. Recall that it is a quadric section of a linear section of codimension 2 of the Grassmannian $\mathrm{Gr}(2, 5) = \mathrm{Gr}(2, V_5)$, where V_5 is a 5-dimensional complex vector space. Let V_i be an i -dimensional vector subspace of V_5 . There are two types of 2-planes in $\mathrm{Gr}(2, 5)$; σ -planes are given set-theoretically as $\{[V_2] \mid V_1 \subset V_2 \subset V_4\}$, and ρ -planes are given by $\{[V_2] \mid V_2 \subset V_3\}$.

Remark 6.1. In [DIM12, Section 3.1], the σ -planes and ρ -planes are called α -planes and β -planes, respectively.

By [DIM12, Section 3.1] and [IM11, Section 3.1], we have the following classification of conics on X .

Definition 6.2 ([DIM12, p. 5]).

- A conic $C \subset X$ is called a τ -conic if the 2-plane $\langle C \rangle$ is not contained in $\mathrm{Gr}(2, V_5)$, there is a unique $V_4 \subset V_5$ such that $C \subset \mathrm{Gr}(2, V_4)$, the conic C is reduced and if it is smooth, the union of corresponding lines in $\mathbb{P}(V_5)$ is a smooth quadric surface in $\mathbb{P}(V_4)$.
- A conic $C \subset X$ is called a σ -conic if the 2-plane $\langle C \rangle$ spanned by C is an σ -plane, and if there is a unique hyperplane $V_4 \subset V_5$ such that $C \subset \mathrm{Gr}(2, V_4)$ and the union of the corresponding lines in $\mathbb{P}(V_5)$ is a quadric cone in $\mathbb{P}(V_4)$.
- A conic $C \subset X$ is called a ρ -conic if the 2-plane $\langle C \rangle$ spanned by C is a ρ -plane, and the union of corresponding lines in $\mathbb{P}(V_5)$ is this 2-plane.

The following lemma is very useful for computations:

Lemma 6.3. *Let X be an ordinary GM threefold and C be a conic on X .*

- (1) *If C is a τ -conic, then we have $\mathrm{RHom}^\bullet(\mathcal{E}, I_C) = k$ and $\mathrm{RHom}^\bullet(\mathcal{E}^\vee, I_C) = 0$.*
- (2) *If C is a ρ -conic, then we have $\mathrm{RHom}^\bullet(\mathcal{E}, I_C) = k^2 \oplus k[-1]$ and $\mathrm{RHom}^\bullet(\mathcal{E}^\vee, I_C) = 0$.*
- (3) *If C is a σ -conic, then we have $\mathrm{RHom}^\bullet(\mathcal{E}, I_C) = k$ and $\mathrm{RHom}^\bullet(\mathcal{E}^\vee, I_C) = k[-1] \oplus k[-2]$.*

Proof. Note that if $\mathrm{Hom}(\mathcal{E}, I_C) = k^a$, then $C \subset \mathrm{Gr}(2, 5 - a) \cap X$. Since for any conic C , there is some V_4 such that $C \subset \mathrm{Gr}(2, V_4)$, then we have $\mathrm{hom}(\mathcal{E}, I_C) \geq 1$ for any conic C . Now if $\mathrm{hom}(\mathcal{E}, I_C) \geq 2$, we know that C is contained in a ρ -plane $\mathrm{Gr}(2, 3)$. Since $\langle C \rangle$ is not in $\mathrm{Gr}(2, 5)$ for a τ -conic C , and $\langle C \rangle$ is a σ -plane $\{[V_2] \mid V_1 \subset V_2 \subset V_4\}$ for a σ -conic, for these two types of conics we have $\mathrm{Hom}(\mathcal{E}, I_C) = k$. Also, for a ρ -conic C , since $\langle C \rangle = \mathrm{Gr}(2, 3)$, we have $\mathrm{hom}(\mathcal{E}, I_C) \geq 2$. But if $\mathrm{hom}(\mathcal{E}, I_C) \geq 3$, we know that $C \subset \mathrm{Gr}(2, 2)$ which is impossible. Hence for a ρ -conic C we have $\mathrm{Hom}(\mathcal{E}, I_C) = k^2$. Now the result for Ext groups follows from applying $\mathrm{Hom}(\mathcal{E}, -)$ to the short exact sequence $0 \rightarrow I_C \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_C \rightarrow 0$ and $\chi(\mathcal{E}, I_C) = 1$.

First by stability and Serre duality, we have $\mathrm{Hom}(\mathcal{E}^\vee, I_C) = \mathrm{Ext}^3(\mathcal{E}^\vee, I_C) = 0$. From $\chi(\mathcal{E}^\vee, I_C) = 0$, we only need to compute $\mathrm{Ext}^1(\mathcal{E}^\vee, I_C)$. Since $\mathrm{RHom}^\bullet(\mathcal{O}_X, I_C) = 0$, applying $\mathrm{Hom}(-, I_C)$ to the tautological sequence, we have $\mathrm{Hom}(\mathcal{Q}^\vee, I_C) = \mathrm{Ext}^1(\mathcal{E}^\vee, I_C)$. Note that if $\mathrm{Hom}(\mathcal{Q}^\vee, I_C) = k^a$, then $C \subset \mathrm{Gr}(2 - a, 5 - a) \cap X$. Thus we have $\mathrm{hom}(\mathcal{Q}^\vee, I_C) \leq 1$ for any conic C . And since $\mathrm{hom}(\mathcal{Q}^\vee, I_C) = 1$ if and only if C is contained in the zero locus of a global section of \mathcal{Q} , which is a σ -3-plane in $\mathrm{Gr}(2, 5)$, we know that $\mathrm{Hom}(\mathcal{Q}^\vee, I_C) = 0$ for C of type τ or ρ , and $\mathrm{Hom}(\mathcal{Q}^\vee, I_C) = k$ for a σ -conic. Then the result follows. \square

Now we recall some properties of the Fano surface of conics $\mathcal{C}(X)$.

Theorem 6.4 ([Log12], [DIM12]). *Let X be an ordinary GM threefold. Then $\mathcal{C}(X)$ is an irreducible projective surface. If X is furthermore general, then $\mathcal{C}(X)$ is smooth.*

It is a fact that there is a unique ρ -conic on X , and there is a curve $L_\sigma \subset \mathcal{C}(X)$ parameterise all σ -conics on X (cf. [DIM12, Section 5.1]), and we denote it by c_X . Furthermore, we have the following result which is a corollary of Logachev's Tangent Bundle Theorem ([Log12, Section 4]).

Lemma 6.5 ([DIM12, p. 16]). *The only rational curve in $\mathcal{C}(X)$ is L_σ . Furthermore, there exists a surface $\mathcal{C}_m(X)$ and a map $\mathcal{C}(X) \rightarrow \mathcal{C}_m(X)$ which contracts L_σ to a point $[\pi]$. If X is general, then $\mathcal{C}_m(X)$ is the minimal surface of $\mathcal{C}(X)$.*

Theorem 6.6 ([DIM12, Section 5.2]). *Let X be a general ordinary GM threefold. Then there is a natural involution ι on $\mathcal{C}_m(X)$, switching the points $[c_X]$ and $[\pi]$.*

Another important result that we require is Logachev's Reconstruction Theorem. This was originally proved in [Log12, Theorem 7.7], and then reproved later in [DIM12, Theorem 9.1].

Theorem 6.7 (Logachev's Reconstruction Theorem). *Let X and X' be general ordinary GM threefolds. If $\mathcal{C}(X) \cong \mathcal{C}(X')$, then $X \cong X'$.*

6.2. Conic and line transforms. For this section, we follow [DIM12, Section 6.1]. Let X be a general ordinary GM threefold, and let C be a conic. Then in [DIM12, § 6.1, Theorem 6.4], the authors construct a new GM threefold X_C and a birational map $\psi_C: X \dashrightarrow X_C$, called the *conic transform*. Similarly, for any line $L \subset X$, a new GM threefold X_L and a birational morphism $\psi_L: X \rightarrow X_L$ are constructed in [DIM12, Section 6.2], called the *line transform*.

Note that in [DK18], such an X_C is called the *period partner* of X , and the line transforms are called the *period duals*. We now list some important results about conic and line transforms below.

Theorem 6.8 ([DIM12, Theorem 6.4]). *Let X be a general ordinary GM threefold, and let $C \subset X$ be a conic. Then $\mathcal{C}(X_C)$ is isomorphic to $\mathcal{C}_m(X)$ blown up at the point $[C] \in \mathcal{C}_m(X)$, where $\mathcal{C}_m(X)$ is the minimal surface of $\mathcal{C}(X)$.*

Proposition 6.9 ([DIM12, Theorem 6.4, Remark 7.2]). *Let X be a general ordinary GM threefold. Then the isomorphism classes of conic transforms of X are parametrized by the surface $\mathcal{C}_m(X)/\iota$.*

Theorem 6.10 ([KP23, Theorem 1.6]). *Let X be a general ordinary GM threefold. Then the Kuznetsov components of all conic transforms and line transforms of X are equivalent to \mathcal{A}_X .*

6.3. Conics on special GM threefolds. Let X be a special GM threefold. Recall that X is a double cover $X \rightarrow Y$ of a degree five del Pezzo threefold Y with branch locus a quadric hypersurface $\mathcal{B} \subset Y$. When X is general, \mathcal{B} is a smooth K3 surface of Picard number 1 and degree 10. Recall that Y is a codimension 3 linear section of $\text{Gr}(2, 5)$. Let \mathcal{V} be the tautological quotient bundle on Y . We recall some properties of $\mathcal{C}(X)$ from [Ili94a].

Theorem 6.11 ([Ili94a]). *Let X be a special GM threefold. Then $\mathcal{C}(X)$ has two components \mathcal{C}_1 and \mathcal{C}_2 . One of the components $\mathcal{C}_2 \cong \Sigma(Y) \cong \mathbb{P}^2$ parametrizes the preimage of lines on Y . Moreover, when X is general, $\mathcal{C}(X)$ is smooth away from $\mathcal{C}_1 \cap \mathcal{C}_2$.*

The following lemma will be useful in computations; it is similar to Lemma 6.3.

Lemma 6.12. *Let X be a special GM threefold and C a conic on X . Then $\text{RHom}^\bullet(\mathcal{E}^\vee, I_C) \neq 0$ if and only if C is the preimage of a line on Y . In this case $\text{RHom}^\bullet(\mathcal{E}^\vee, I_C) = k[-1] \oplus k[-2]$, and such a family of conics is parametrized by the Hilbert scheme of lines $\Sigma(Y) \cong \mathbb{P}^2$ on Y .*

Proof. The proof is almost the same as the second part of the proof of Lemma 6.3. The same argument shows that $\text{RHom}^\bullet(\mathcal{E}^\vee, I_C) \neq 0$ if and only if $\text{Hom}(\mathcal{Q}^\vee, I_C) \neq 0$. The image of a non-trivial map $\mathcal{Q}^\vee \rightarrow I_C$ is the ideal sheaf of the zero locus of a section s of \mathcal{Q} , which is the preimage of the zero locus of a section of \mathcal{V} . By [San14, Lemma 2.18], the zero locus of a section of \mathcal{V} is either a line or a point. Thus the zero locus of a section of \mathcal{Q} is either the preimage of a line on Y which is a conic on X , or a zero-dimensional closed subscheme of length two. But this zero locus contains a conic $C \subset X$, so $C = Z(s)$ is the preimage of a line on Y and the map $\mathcal{Q}^\vee \rightarrow I_C$ is surjective. In particular, such conics are exactly the preimages of lines on Y and are parametrized by $\Sigma(Y) \cong \mathbb{P}^2$. \square

7. CONICS AND BRIDGELAND MODULI SPACES

In this section, we study the moduli space of σ -stable objects of the (-1) -class $-x$ in the alternative Kuznetsov component \mathcal{A}_X of a GM threefold X and its relation to $\mathcal{C}(X)$. Our main result in this section is Theorem 7.12, which realizes the Bridgeland moduli space as a contraction of $\mathcal{C}(X)$.

First, we study those conics C such that $I_C \notin \mathcal{A}_X$.

Proposition 7.1. *Let $C \subset X$ be a conic on a GM threefold X . Then $I_C \notin \mathcal{A}_X$ if and only if*

- (1) *C is a σ -conic when X is ordinary. In particular, such a family of conics is parametrized by the line L_σ .*
- (2) *C is the preimage of a line on Y when X is special. In particular, such a family of conics is parametrized by the Hilbert scheme of lines $\Sigma(Y) \cong \mathbb{P}^2$ on Y .*

Moreover, we have an exact sequence

$$0 \rightarrow \mathcal{E} \rightarrow \mathcal{Q}^\vee \rightarrow I_C \rightarrow 0.$$

Proof. Note that $I_C \notin \mathcal{A}_X$ if and only if $\mathrm{RHom}^\bullet(\mathcal{E}^\vee, I_C) \neq 0$. When X is ordinary, (1) follows from Lemma 6.3. When X is special, we deduce (2) from Lemma 6.12. Note that since $I_C \notin \mathcal{A}_X$, we have $\mathrm{Hom}(\mathcal{Q}^\vee, I_C) \neq 0$. The non-trivial map $\mathcal{Q}^\vee \rightarrow I_C$ is surjective by the arguments in Lemma 6.3 and 6.12. Note that by the stability of \mathcal{Q}^\vee , the kernel of $\mathcal{Q}^\vee \rightarrow I_C$ is μ -stable with the same Chern character as \mathcal{E} , hence we have $\ker(\mathcal{Q}^\vee \rightarrow I_C) \cong \mathcal{E}$ by [DIM12, Proposition 4.1]. \square

Proposition 7.2. *Let X be a GM threefold and $C \subset X$ a conic on X . If $I_C \notin \mathcal{A}_X$, then we have the exact triangle*

$$\mathcal{E}[1] \rightarrow \mathrm{pr}(I_C) \rightarrow \mathcal{Q}^\vee$$

and $\mathrm{pr}(I_C) \cong \pi'(\mathcal{Q}^\vee)$

Proof. By Proposition 7.1, I_C fits into the short exact sequence

$$0 \rightarrow \mathcal{E} \rightarrow \mathcal{Q}^\vee \rightarrow I_C \rightarrow 0.$$

Applying the projection functor to this exact sequence, and note that applying the functor pr to the dual exact sequence of (1) gives $\mathrm{pr}(\mathcal{Q}^\vee) = 0$. Then we have $\mathrm{pr}(I_C) \cong \mathrm{pr}(\mathcal{E})[1]$. Now we compute the projection $\mathrm{pr}(\mathcal{E})$. Since $\mathrm{RHom}^\bullet(\mathcal{E}^\vee, \mathcal{E}) \cong k[-3]$, we get an exact triangle $\mathcal{E}^\vee[-3] \rightarrow \mathcal{E} \rightarrow \mathbf{L}_{\mathcal{E}^\vee} \mathcal{E}$. Now applying $\mathbf{L}_{\mathcal{O}_X}$ to this triangle and using $\mathbf{L}_{\mathcal{O}_X} \mathcal{E}^\vee = \mathcal{Q}^\vee[1]$, we get

$$\mathcal{Q}^\vee[-2] \rightarrow \mathcal{E} \rightarrow \mathrm{pr}(\mathcal{E}).$$

Therefore we obtain the triangle

$$\mathcal{E}[1] \rightarrow \mathrm{pr}(\mathcal{E})[1] \rightarrow \mathcal{Q}^\vee$$

and the desired result follows from Lemma 5.4. \square

Now the following two results follow from Proposition 7.2 and Lemma 5.6.

Lemma 7.3. *Let X be a GM threefold. If $C \subset X$ is a conic such that $I_C \notin \mathcal{A}_X$, then*

- $\mathrm{RHom}^\bullet(\mathrm{pr}(I_C), \mathrm{pr}(I_C)) = k \oplus k^2[-1]$ when X is ordinary.
- $\mathrm{RHom}^\bullet(\mathrm{pr}(I_C), \mathrm{pr}(I_C)) = k \oplus k^3[-1] \oplus k[-2]$ when X is special.

Lemma 7.4. *Let X be a GM threefold. If $I_C \notin \mathcal{A}_X$, the projection $\mathrm{pr}(I_C)[1]$ is stable with respect to every Serre-invariant stability condition on \mathcal{A}_X .*

When $I_C \in \mathcal{A}_X$, we cannot use Proposition 4.12 to prove the Bridgeland stability of I_C , since $\mathcal{C}(X)$ can be singular and $\mathrm{Ext}^1(I_C, I_C)$ may have large dimension. Instead, we use a wall-crossing argument and the uniqueness of Serre-invariant stability conditions (Theorem A.10).

Lemma 7.5. *Let X be a GM threefold. Let F be an object with $\mathrm{ch}_{\leq 2}(F) = (1, 0, -2L)$. Then there are no walls for F in the range $-\frac{1}{2} \leq \beta < 0$ and $\alpha > 0$.*

Proof. Recall that by [BBF⁺20, Theorem 4.13], $\beta = 0$ is the unique vertical wall of F . Any other wall is a semicircle centered along the β -axis, and its apex lies on the hyperbola $\mu_{\alpha,\beta}(F) = 0$. Moreover, no two walls intersect.

Note that when $\mu_{\alpha,\beta}(F) = 0$ holds, we have $\beta < -\sqrt{\frac{2}{5}} < -\frac{1}{2}$, thus we know that there is no semicircular wall centered in the interval $-\frac{1}{2} \leq \beta < 0$. Therefore, any semicircular wall in the range $-\frac{1}{2} \leq \beta < 0$ will intersect $\beta = -\frac{1}{2}$. To prove the statement, we only need to show that there are no walls when $\beta = -\frac{1}{2}$. This follows from the fact that $\text{ch}_1^{-\frac{1}{2}}(F)$ is minimal. \square

Lemma 7.6. *Let $C \subset X$ be a conic on a GM threefold X such that $I_C \in \mathcal{A}_X$. Then $I_C[1]$ is stable with respect to every Serre-invariant stability condition on \mathcal{A}_X .*

Proof. By Lemma 4.5 and Lemma 7.5, we know that I_C is $\sigma_{\alpha,\beta}$ -semistable for every $(\alpha, \beta) \in V$. Since I_C is torsion-free, $I_C[1] \in \text{Coh}_{\alpha,\beta}^0(X)$ is $\sigma_{\alpha,\beta}^0$ -semistable. Thus $I_C[1] \in \mathcal{A}(\alpha, \beta)$ is $\sigma(\alpha, \beta)$ -semistable. Then stability with respect to every Serre-invariant stability condition follows from Theorem 4.11 and Theorem A.10. \square

7.1. The Bridgeland moduli space of class $-x$. In this subsection, we are going to describe the Bridgeland moduli space $\mathcal{M}_\sigma(\mathcal{A}_X, -x)$ in Theorem 7.12.

The proofs in this section seem technical. However, the only results in this section that will be used in other sections are Proposition 7.11 in the proof of Theorem 7.12, so there is no harm for readers in skipping this whole section and assuming Proposition 7.11 and Theorem 7.12.

We start with two lemmas.

Lemma 7.7. *Let X be a GM threefold and E a μ -semistable sheaf on X with truncated Chern character $\text{ch}_{\leq 2}(E) = (2, -H, aL)$. If $a \geq 1$ and $c_3(E) \geq 0$ and then we have $E \cong \mathcal{E}$.*

Proof. By Lemma 4.6, we have $a \leq 1$ which means $a = 1$ by our assumption. Then $c_1(E) = -1$ and $c_2(E) = 4$. Since $c_3(E) \geq 0$, by [BF14, Proposition 3.5] we have $\chi(E) = 0$. This implies $c_3(E) = 0$. Moreover, $E^{\vee\vee}$ also satisfies the assumptions above. Hence by the previous argument, we have $c_1(E^{\vee\vee}) = -1$, $c_2(E^{\vee\vee}) = 4$ and $c_3(E^{\vee\vee}) = 0$ as well. In other words, $E = E^{\vee\vee}$. Since $c_3(E) = 0$ and E is reflexive of rank two, it is a vector bundle. Moreover, E is a globally generated bundle by [BF14, Proposition 3.5]. Thus $E \cong \mathcal{E}$ by [DIM12, Proposition 4.1]. \square

Lemma 7.8. *Let X be a GM threefold and E a μ -semistable sheaf on X with $\text{ch}(E) = \text{ch}(\mathcal{Q})$. Then we have $E \cong \mathcal{Q}$.*

Proof. First we show that $h^2(E) = 0$; then from $\chi(E) = 5$ we have $h^0(E) \geq 5$. Indeed, if $h^2(E) \neq 0$, then $\text{Hom}(E, \mathcal{O}_X(-H)[1]) \neq 0$ by Serre duality. Therefore, we have a non-trivial extension

$$0 \rightarrow \mathcal{O}_X(-H) \rightarrow F \rightarrow E \rightarrow 0.$$

If F is not μ -semistable, then by the stability of $\mathcal{O}_X(-H)$ and E , the minimal destabilizing quotient sheaf F' of F has $\text{ch}_{\leq 1}(F') = (1, -H)$. Thus $F'^{\vee\vee} \cong \mathcal{O}_X(-H)$. But if we apply $\text{Hom}(-, \mathcal{O}_X(-H))$ to the exact sequence above, we obtain $\text{Hom}(F, \mathcal{O}_X(-H)) = 0$ since this extension is non-trivial, which gives a contradiction. Then F is μ -semistable with $\text{ch}_{\leq 2}(F) = (4, 0, 4L)$, which is impossible since $\Delta(F) < 0$.

Now we can take five linearly independent elements in $H^0(E)$ and obtain a map $t: \mathcal{O}_X^{\oplus 5} \rightarrow E$. From the stability of \mathcal{O}_X and E , we have $\mu(\text{Im}(t)) = 0$ or $\mu(\text{Im}(t)) = \frac{1}{3}$. But the first case cannot happen, since then $\text{Im}(t)$ is the direct sum of a number of copies of \mathcal{O}_X , and this contradicts the construction of t . Thus $\mu(\text{Im}(t)) = \frac{1}{3}$ and $\text{ch}_{\leq 1}(\text{Im}(t)) = (3, H)$. Also $\text{ch}_{\leq 2}(\ker(t)) = (2, -H, xL)$, where $x \geq 1$. Note that $\ker(t)$ is reflexive, thus we have $c_3(\ker(t)) \geq 0$ since $\ker(t)$ has rank two. Then by stability of \mathcal{O}_X and $\text{Hom}(\mathcal{O}_X, \ker(t)) = 0$, it is not hard to see that $\ker(t)$ is μ -semistable. Thus by Lemma 7.7 we have $\ker(t) \cong \mathcal{E}$. Therefore $\text{ch}(\text{Im}(t)) = \text{ch}(E)$ and thus t is surjective.

Now applying $\text{Hom}(\mathcal{Q}, -)$ to the exact sequence

$$0 \rightarrow \mathcal{E} \rightarrow \mathcal{O}_X^{\oplus 5} \rightarrow E \rightarrow 0,$$

from $\mathrm{RHom}^\bullet(\mathcal{Q}, \mathcal{O}_X) = 0$ and $\mathrm{Ext}^1(\mathcal{Q}, \mathcal{E}) = k$ we have $\mathrm{Hom}(\mathcal{Q}, E) = k$. Thus from the stability of E and \mathcal{Q} , we have $E \cong \mathcal{Q}$ and the result follows. \square

Now we introduce some notations. Let $\alpha > 0$ and $\beta < 0$. For an object $E \in \mathrm{D}^b(X)$, the limit central charge $Z_{0,0}^0(E)$ is defined as the limit of $Z_{\alpha,\beta}^0(E)$ when $(\alpha, \beta) \rightarrow (0, 0)$. Note that $Z_{\alpha,\beta}^0(E)$ is given by \mathbb{Q} -linear combinations of $\alpha, \beta, \alpha^2, \beta^2$, thus such a limit $Z_{0,0}^0(E)$ always exists. For $Z_{0,0}^0(E) \neq 0$, we can also define the limit slope $\mu_{0,0}^0(E)$ as follows:

- If $\mathrm{Im}(Z_{0,0}^0(E)) \neq 0$, then we define $\mu_{0,0}^0(E) := -\frac{\mathrm{Re}(Z_{0,0}^0(E))}{\mathrm{Im}(Z_{0,0}^0(E))}$.
- If $\mathrm{Im}(Z_{0,0}^0(E)) = 0$ and $\mathrm{Re}(Z_{0,0}^0(E)) > 0$, then we define $\mu_{0,0}^0(E) := -\infty$.
- If $\mathrm{Im}(Z_{0,0}^0(E)) = 0$ and $\mathrm{Re}(Z_{0,0}^0(E)) < 0$, then we define $\mu_{0,0}^0(E) := +\infty$.

Note that $Z_{0,0}^0(E) = 0$ if and only if $\mathrm{ch}_{\leq 2}(E)$ is a multiple of $\mathrm{ch}_{\leq 2}(\mathcal{O}_X)$.

Let $E \in \mathrm{Coh}_{\alpha,\beta}^0(X)$. By continuity, we can find a neighborhood U_E of the origin such that for any $(\alpha, \beta) \in U_E$, the slopes $\mu_{\alpha,\beta}^0(E)$ and $\mu_{0,0}^0(E)$ are both negative or positive. Let $F \in \mathrm{Coh}_{\alpha,\beta}^0(X)$ be another object such that E, F are both $\sigma_{\alpha,\beta}^0$ -semistable in a neighborhood $U_{E,F}$ of the origin. If $\mu_{0,0}^0(E) > \mu_{0,0}^0(F)$, then by continuity, we can find a smaller neighborhood $U'_{E,F}$ such that $\mu_{\alpha,\beta}^0(E) > \mu_{\alpha,\beta}^0(F)$ holds for every $(\alpha, \beta) \in U'_{E,F}$. Thus we have $\mathrm{Hom}(E, F) = 0$. We will use these two elementary facts repeatedly.

Proposition 7.9. *If $F \in \mathcal{A}(\alpha, \beta)$ is $\sigma(\alpha, \beta)$ -stable such that $[F] = -x$ and F is $\sigma_{\alpha,\beta}^0$ -semistable for some $(\alpha, \beta) \in V$, then $F \cong I_C[1]$ for a conic C on X .*

Proof. Since F is $\sigma_{\alpha,\beta}^0$ -semistable and $\mu_{\alpha,\beta}^0(F) > 0$, as in [PY22, Proposition 4.6] there is a triangle

$$F_1[1] \rightarrow F \rightarrow F_2$$

where $F_1 \in \mathrm{Coh}^\beta(X)$ with $\mu_{\alpha,\beta}^+(F_1) < 0$ and F_2 is supported on points. Thus $\mathrm{ch}(F_1) = (1, 0, -2L, mP)$, where m is the length of F_2 . By Lemmas 7.5 and 4.5, F_1 is a rank one torsion-free sheaf, hence it is the ideal sheaf of a closed subscheme. Thus by [San14, Corollary 1.38], we have $m \leq 0$, which means $F_2 = 0$ and $F_1 \cong F[-1]$. Thus by Lemma 7.5 again, $F[-1]$ is a μ -semistable torsion free sheaf, which is of the form $F[-1] \cong I_C$ for a conic C on X since $\mathrm{Pic}(X) = \mathbb{Z} \cdot H$. \square

When F is not $\sigma_{\alpha,\beta}^0$ -semistable for $(\alpha, \beta) \in V$, the argument is more complicated. Our main tools are the inequalities in [PR23b], [PY22, Proposition 4.1], Lemma 4.6 and Theorem 4.7, which allow us to bound the rank and first two Chern characters $\mathrm{ch}_1, \mathrm{ch}_2$ of the destabilizing objects and their cohomology objects. Since $F \in \mathcal{A}_X$, by using the Euler characteristics $\chi(\mathcal{O}_X, -)$ and $\chi(\mathcal{E}^\vee, -)$ we can obtain a bound on ch_3 . Finally, via a similar argument as in Lemma 7.7 we deduce that the Harder–Narasimhan factors of F are the ones we expect.

Proposition 7.10. *If $F \in \mathcal{A}(\alpha, \beta)$ is $\sigma(\alpha, \beta)$ -stable such that $[F] = -x$ and F is not $\sigma_{\alpha,\beta}^0$ -semistable for every $(\alpha, \beta) \in V$, then F fits into a triangle*

$$\mathcal{E}[2] \rightarrow F \rightarrow \mathcal{Q}^\vee[1].$$

Proof. Since there are no walls for F tangent to the wall $\beta = 0$, by the local finiteness of walls and [BMT14, Proposition 2.2.2] we can find an open neighborhood U' of the origin such that the Harder–Narasimhan filtration with respect to $\sigma_{\alpha,\beta}^0$ is constant for every $(\alpha, \beta) \in U := U' \cap V$. In the following we will only consider $\sigma_{\alpha,\beta}^0$ for $(\alpha, \beta) \in U$.

Let B be the minimal destabilizing quotient object of F and $0 \rightarrow A \rightarrow F \rightarrow B \rightarrow 0$ be the destabilizing short exact sequence of F in $\mathrm{Coh}_{\alpha,\beta}^0(X)$. Hence we know that $A, B \in \mathrm{Coh}_{\alpha,\beta}^0(X)$ and B is $\sigma_{\alpha,\beta}^0$ -semistable with $\mu_{\alpha,\beta}^{0,-}(A) > \mu_{\alpha,\beta}^0(F) > \mu_{\alpha,\beta}^0(B)$ for all $(\alpha, \beta) \in U$. By [BLMS23, Remark 5.12], we have $\mu_{\alpha,\beta}^0(B) \geq \min\{\mu_{\alpha,\beta}^0(F), \mu_{\alpha,\beta}^0(\mathcal{O}_X), \mu_{\alpha,\beta}^0(\mathcal{E}^\vee)\}$. Hence the following relations hold for all $(\alpha, \beta) \in U$:

- $\mu_{\alpha,\beta}^0(A) > \mu_{\alpha,\beta}^0(F) > \mu_{\alpha,\beta}^0(B)$,
- $\mathrm{Im}(Z_{\alpha,\beta}^0(A)) \geq 0, \mathrm{Im}(Z_{\alpha,\beta}^0(B)) > 0$,
- $\mu_{\alpha,\beta}^0(B) \geq \min\{\mu_{\alpha,\beta}^0(F), \mu_{\alpha,\beta}^0(\mathcal{O}_X), \mu_{\alpha,\beta}^0(\mathcal{E}^\vee)\}$,
- $\Delta(B) \geq 0$.

By continuity and taking $(\alpha, \beta) \rightarrow (0, 0)$, we have:

- (1) $\mu_{0,0}^0(A) \geq \mu_{0,0}^0(F) = 0 \geq \mu_{0,0}^0(B)$,
- (2) $\text{Im}(Z_{0,0}^0(A)) \geq 0$, $\text{Im}(Z_{0,0}^0(B)) \geq 0$,
- (3) $\mu_{0,0}^0(B) \geq \min\{\mu_{0,0}^0(F), \mu_{0,0}^0(\mathcal{O}_X), \mu_{0,0}^0(\mathcal{E}^\vee)\}$,
- (4) $\Delta(B) \geq 0$.

Assume that $[A] = a[\mathcal{O}_X] + b[\mathcal{O}_H] + c[\mathcal{O}_L] + d[\mathcal{O}_P]$ for integers $a, b, c, d \in \mathbb{Z}$. Then we have $[B] = (-1 - a)[\mathcal{O}_X] - b[\mathcal{O}_H] + (2 - c)[\mathcal{O}_L] - (1 + d)[\mathcal{O}_P]$. Then we see

- $\text{ch}(A) = (a, bH, \frac{c-5b}{10}H^2, \frac{5b+\frac{c}{5}+d}{10}H^3)$
- $Z_{0,0}^0(A) = bH^3 + (\frac{c-5b}{10}H^3)i$, $Z_{0,0}^0(B) = -bH^3 + (\frac{2-c+5b}{10}H^3)i$
- $\mu_{0,0}^0(A) = \frac{10b}{5b-c}$, $\mu_{0,0}^0(B) = \frac{-10b}{c-5b-2}$.

Note that $[F] = -[\mathcal{O}_X] + 2[\mathcal{O}_L] - [\mathcal{O}_P]$. From (2) we know $c - 5b = 0, 1$ or 2 . But when $c - 5b = 2$, it is not hard to see that (c) fails near the origin. Thus $c - 5b = 0$ or 1 .

We begin with two claims.

Claim 1: *We have $\text{RHom}^\bullet(\mathcal{O}_X, B) = \text{Hom}(\mathcal{O}_X, B)$ and $\text{RHom}^\bullet(\mathcal{O}_X, A) = \text{Ext}^1(\mathcal{O}_X, A)[-1]$.*

Since $F \in \mathcal{A}_X$, we only need to prove that $\text{Ext}^i(\mathcal{O}_X, A) = 0$ for $i \neq 1$. Indeed, since $\mathcal{O}_X \in \text{Coh}_{\alpha,\beta}^0(X)$ and $F \in \mathcal{A}_X$, we have $\text{Ext}^i(\mathcal{O}_X, A) = 0$ for all $i \leq 0$. Also, by Serre duality we have $\text{Ext}^i(\mathcal{O}_X, A) = \text{Hom}(A, \mathcal{O}_X(-H)[3-i])$. Thus from the fact that $\mathcal{O}_X(-H) \in \text{Coh}_{\alpha,\beta}^0(X)$, we obtain $\text{Hom}(A, \mathcal{O}_X(-H)[3-i]) = 0$ for $i \geq 2$. Therefore we have $\text{Ext}^i(\mathcal{O}_X, A) = 0$ for $i \neq 1$.

Claim 2: *We have $\text{RHom}^\bullet(\mathcal{E}^\vee, B) = \text{Hom}(\mathcal{E}^\vee, B)$ and $\text{RHom}^\bullet(\mathcal{E}^\vee, A) = \text{Ext}^1(\mathcal{E}^\vee, A)[-1]$.*

Since \mathcal{E}^\vee and $\mathcal{E}[2] \in \text{Coh}_{\alpha,\beta}^0(X)$, the argument is the same as Claim 1.

Now we deal with the cases $c - 5b = 0$ and $c - 5b = 1$ separately.

Case 1 ($c - 5b = 0$):

First, we assume that $c - 5b = 0$. By 7.1, we have:

- (1) $-2 \leq b \leq 0$,
- (2) $b^2 + \frac{2a+2}{5} \geq 0$.

case 1.1 ($b = 0$): If $b = 0$, then $c = 0$ and $a \geq -1$. In this case we have $\text{ch}_{\leq 2}(B) = (-1 - a, 0, 2L)$. If $a = -1$, then $\text{ch}_{\leq 2}(A) = \text{ch}_{\leq 2}(\mathcal{O}_X[1]) = (-1, 0, 0)$, which is impossible since $\text{Im}(Z_{\alpha,\beta}^0(A)) < 0$ for $(\alpha, \beta) \in V$. Thus $a \geq 0$, and $a \neq 0$ otherwise $\mu_{\alpha,\beta}^0(F) = \mu_{\alpha,\beta}^0(B)$ for any $(\alpha, \beta) \in V$. But then we have $\mu_{\alpha,\beta}^0(F) < \mu_{\alpha,\beta}^0(B)$ when $(\alpha, \beta) \in U$ is sufficiently close to the origin. This contradicts our assumption on B .

case 1.2 ($b = -1$): If $b = -1$, we have $c = -5$. In this case $\text{ch}_{\leq 2}(A) = (a, -H, 0)$. Since $A \in \text{Coh}_{\alpha,\beta}^0(X)$, we have $\text{Im}(Z_{\alpha,\beta}^0(A)) \geq 0$ for every $(\alpha, \beta) \in U$. Note that $\text{Im}(Z_{\alpha,\beta}^0(A)) = (\beta + \frac{a(\beta^2 - \alpha^2)}{2})H^3$ and $0 < \alpha < -\beta$, and we have $a \geq \frac{-2\beta}{\beta^2 - \alpha^2}$. But note that when $\alpha = \frac{-\beta}{2}$ and $\beta \rightarrow 0$, we have $\frac{-2\beta}{\beta^2 - \alpha^2} \rightarrow +\infty$, thus we get a contradiction since a is a finite number.

case 1.3 ($b = -2$): If $b = -2$, we have $c = -10$. In this case we have $\text{ch}_{\leq 2}(A) = (a, -2H, 0)$. Similarly to case 1.2, we have $\text{Im}(Z_{\alpha,\beta}^0(A)) \geq 0$ for every $(\alpha, \beta) \in U$. Note that $\text{Im}(Z_{\alpha,\beta}^0(A)) = (2\beta + \frac{a(\beta^2 - \alpha^2)}{2})H^3$ and $\alpha < -\beta$, and we have $a \geq \frac{-4\beta}{\beta^2 - \alpha^2}$. Then as in case 1.2, we get a contradiction.

Case 2 ($c - 5b = 1$): Now we assume that $c - 5b = 1$. Then by 7.1, we have:

- (1) $-1 \leq b \leq 0$,
- (2) $b^2 + \frac{a+1}{5} \geq 0$.

case 2.1 ($b = 0$): If $b = 0$, then $c = 1$. Therefore $-1 \leq a$. If $a = -1$, since B is $\sigma_{\alpha,\beta}^0$ -semistable, we know $\mathcal{H}_{\text{Coh}^\beta(X)}^0(B)$ is either 0 or supported on points. Thus $\text{ch}_{\leq 2}(\mathcal{H}_{\text{Coh}^\beta(X)}^{-1}(B)) = (0, 0, -L)$. But $\text{Re}(Z_{\alpha,\beta}(\mathcal{H}_{\text{Coh}^\beta(X)}^{-1}(B))) > 0$ which is impossible since $\mathcal{H}_{\text{Coh}^\beta(X)}^{-1}(B) \in \text{Coh}^\beta(X)$ with $\text{Im}(Z_{\alpha,\beta}(\mathcal{H}_{\text{Coh}^\beta(X)}^{-1}(B))) = 0$.

Therefore we have $a \geq 0$. Hence $\text{ch}_{\leq 2}(B) = -(a+1, 0, -L)$, where $a+1 \geq 1$. This is also impossible since when $(\alpha, \beta) \in U$ is sufficiently close to the origin, we have $\mu_{\alpha,\beta}^0(B) > \mu_{\alpha,\beta}^0(F)$.

case 2.2 ($b = -1$): We have $b = -1$ and $c = -4$. Hence $-6 \leq a$. In this case $\text{ch}_{\leq 2}(B) = (-1 - a, H, L)$ and we have $\mu_{\alpha,\beta}^0(B) < 0$ and $\text{ch}_1^\beta(B) > 0$ for when $(\alpha, \beta) \in U$ is sufficiently close to the origin. Thus $B \in \text{Coh}^\beta(X)$ is $\sigma_{\alpha,\beta}$ -semistable. Applying Lemma 4.6 to B , we have $a \geq -3$.

We first prove a claim.

Claim 3: *In the situation of case 2.2, we have A is $\sigma_{\alpha,\beta}^0$ -semistable. Hence $\text{RHom}^\bullet(\mathcal{O}_X, A) = 0$, $\text{ch}(A) = (a, -H, L, (\frac{7}{3} - a)P)$ and $\chi(\mathcal{E}^\vee, A) = 3 - 2a$.*

Assume A is not $\sigma_{\alpha,\beta}^0$ -semistable for some $(\alpha, \beta) \in U$. Then we can take a neighborhood U'_A of the origin such that A has constant Harder–Narasimhan factors for any $(\alpha, \beta) \in U_A := U \cap U'_A \cap V$. Let C be the minimal destabilizing quotient object of A with respect to $\sigma_{\alpha,\beta}^0$ for $(\alpha, \beta) \in U_A$. In this case we have $\text{ch}_{\leq 2}(A) = (a, -H, L)$. Since $\text{Im}(Z_{0,0}^0(A)) = \frac{1}{10}H^3$, we know that $\text{Im}(Z_{0,0}^0(C)) = 0$ or $\frac{1}{10}H^3$. If $\text{Im}(Z_{0,0}^0(C)) = 0$, then $\mu_{0,0}^0(C) = +\infty$ or $-\infty$. But the previous case contradicts $\mu_{\alpha,\beta}^0(A) > \mu_{\alpha,\beta}^0(C)$ and the latter case contradicts $\mu_{\alpha,\beta}^0(C) > \mu_{\alpha,\beta}^0(F)$. Therefore we have $\text{Im}(Z_{0,0}^0(C)) = \frac{1}{10}H^3$ and we can assume that $\text{ch}_{\leq 2}(C) = (e, fH, L)$ where $e, f \in \mathbb{Z}$. Since $\mu_{0,0}^0(A) \geq \mu_{0,0}^0(C) \geq \mu_{0,0}^0(F) = 0$, we have $10 \geq -10f \geq 0$. If $f = 0$, then $\text{ch}_{\leq 2}(C) = (e, 0, L)$ and $\text{ch}_{\leq 2}(D) = (a - e, -H, 0)$, where $D = \text{cone}(A \rightarrow C)[-1]$. Then $\mu_{\alpha,\beta}^0(D) > \mu_{\alpha,\beta}^0(A)$ for any $(\alpha, \beta) \in U_A$. Hence

$$\mu_{\alpha,\beta}^0(D) = \frac{1 + (a - e)\beta}{\beta + \frac{a-e}{2}(\beta^2 - \alpha^2)} > \mu_{\alpha,\beta}^0(A) > \mu_{\alpha,\beta}^0(F).$$

But note that $\mu_{\alpha,\beta}^0(D) < 0$ for $(\alpha, \beta) \in U_A$ that sufficiently closed to the origin, which gives a contradiction since $\mu_{\alpha,\beta}^0(D) > \mu_{\alpha_0,\beta_0}^0(F)$ holds for any $(\alpha, \beta) \in U_A$.

Therefore the only possible case is $f = -1$, and hence $\mu_{0,0}^0(C) = 10$. Since $\mu_{\alpha,\beta}^0(A) > \mu_{\alpha,\beta}^0(C)$ for $(\alpha, \beta) \in U_A$, we have $\text{rk } C > a$. But this is impossible since $D, \mathcal{O}_X \in \text{Coh}_{\alpha,\beta}^0(X)$ but $\text{ch}_{\leq 2}(D) = (s, 0, 0) = s \cdot \text{ch}_{\leq 2}(\mathcal{O}_X)$ where $s = a - \text{rk } C < 0$. Now for the last statement, note that $\mathcal{O}_X(-H)[2] \in \text{Coh}_{\alpha,\beta}^0(X)$ is $\sigma_{\alpha,\beta}^0$ -semistable with $\mu_{0,0}^0(\mathcal{O}_X(-H)[2]) = 2$, hence we have $\text{Hom}(A, \mathcal{O}_X(-H)[2]) = \text{Hom}(\mathcal{O}_X, A[1]) = 0$. Now combined with Claim 1, this proves our claim.

Now we deal with the three cases $a = -3$, $-2 \leq a \leq 1$ and $a \geq 2$ separately.

When $a = -3$, we have $\text{ch}_{\leq 2}(B) = \text{ch}_{\leq 2}(\mathcal{E}^\vee)$. Then since $\text{ch}_{\leq 2}(B)$ is on the boundary of Lemma 4.6, by a standard argument we know that B is $\sigma_{\alpha,\beta}$ -semistable for every $\alpha > 0$ and $\beta < 0$, as explained in [PR23a, Proposition 3.2]. Thus by Lemma 4.5, B is a μ -semistable sheaf. From Claim 3 we have $\chi(\mathcal{O}_X, B) = 0$, hence $\text{ch}(B) = \text{ch}(\mathcal{E}^\vee)$ and by Lemma 7.7 we have $B \cong \mathcal{E}^\vee$. But this implies $\text{Hom}(\mathcal{O}_X, A[1]) = k^5$ since $F \in \mathcal{A}_X$, which contradicts Claim 3.

When $-2 \leq a \leq 1$, we have $\mu_{\alpha,\beta}^0(A) > \mu_{\alpha,\beta}^0(\mathcal{E}[2])$. Since A is $\sigma_{\alpha,\beta}^0$ -semistable, we have $\text{Hom}(A, \mathcal{E}[2]) = \text{Hom}(\mathcal{E}^\vee, A[1]) = 0$. Thus $\text{RHom}^\bullet(\mathcal{E}^\vee, A) = 0$ by Claim 2. But this contradicts Claim 3 since $\chi(\mathcal{E}^\vee, A) = 3 - 2a$.

When $a \geq 2$, applying Theorem 4.7 to B , we have $a = 2$. Thus $\text{ch}_{\leq 2}(B) = \text{ch}_{\leq 2}(\mathcal{Q}^\vee[1])$. By Claim 3, we know that $\text{RHom}^\bullet(\mathcal{O}_X, B) = 0$ and we get $\text{ch}(B) = \text{ch}(\mathcal{Q}^\vee[1])$. Thus $\chi(\mathcal{E}^\vee, B) = \text{hom}(\mathcal{E}^\vee, B) > 0$. Therefore, if we apply $\text{Hom}(-, B)$ to the exact sequence $0 \rightarrow \mathcal{Q}^\vee \rightarrow \mathcal{O}_X^{\oplus 5} \rightarrow \mathcal{E}^\vee \rightarrow 0$, we obtain $\text{hom}(\mathcal{Q}^\vee[1], B) > 0$. Now by stability, we have $B \cong \mathcal{Q}^\vee[1]$. Now $\text{ch}(A) = \text{ch}(\mathcal{E}[2])$. By Claim 2 and Claim 3, we have $\text{ext}^1(\mathcal{E}^\vee, A) = \text{hom}(A, \mathcal{E}[2]) = 1$. Since A is $\sigma_{\alpha,\beta}^0$ -semistable and $\mathcal{E}[2]$ is $\sigma_{\alpha,\beta}^0$ -stable, we have $A \cong \mathcal{E}[2]$. \square

Proposition 7.11. *Let X be a GM threefold. Then every object in the moduli space $\mathcal{M}_\sigma(\mathcal{A}_X, -x)$ is of form $\text{pr}(I_C)[1]$ for a conic $C \subset X$.*

Proof. Note that $\text{hom}(\mathcal{Q}^\vee[1], \mathcal{E}[2]) = 1$. Then the result follows from Proposition 7.9 and Proposition 7.10. \square

Now we are ready to realize the Bridgeland moduli space $\mathcal{M}_\sigma(\mathcal{A}_X, -x)$ as the contraction $\mathcal{C}_m(X)$ of the Fano surface $\mathcal{C}(X)$:

Theorem 7.12. *Let X be a GM threefold and σ a Serre-invariant stability condition on \mathcal{A}_X . The projection functor $\text{pr}: \text{D}^b(X) \rightarrow \mathcal{A}_X$ induces a surjective morphism $p: \mathcal{C}(X) \rightarrow \mathcal{M}_\sigma(\mathcal{A}_X, -x)$, where p is*

- a blow-down morphism to a smooth point when X is ordinary;
- a contraction of the component \mathbb{P}^2 to a singular point when X is special.

In particular, when X is general and ordinary, $\mathcal{M}_\sigma(\mathcal{A}_X, -x)$ is isomorphic to the minimal model $\mathcal{C}_m(X)$ of the Fano surface of conics on X . When X is general and special, the moduli space $\mathcal{M}_\sigma(\mathcal{A}_X, -x)$ has only one singular point.

Proof. By Lemma 7.4 and Lemma 7.6, $\mathrm{pr}(I_C)[1]$ is σ -stable for any conic $C \subset X$. Then we obtain a morphism $p: \mathcal{C}(X) \rightarrow \mathcal{M}_\sigma(\mathcal{A}_X, -x)$. Moreover, Proposition 7.11 implies that p is surjective.

Now according to Proposition 7.1, the family of conics $C \subset X$ with the property that $I_C \notin \mathcal{A}_X$ is parametrized by the line L_σ when X is ordinary, and the component \mathbb{P}^2 when X is special. Since $\mathrm{pr}(I_C)[1] \cong \pi'(\mathcal{Q}^\vee)[1]$ for $I_C \notin \mathcal{A}_X$ by Proposition 7.2, we know that $p(L_\sigma) = [\pi'(\mathcal{Q}^\vee)[1]]$ when X is ordinary, and $p(\mathbb{P}^2) = [\pi'(\mathcal{Q}^\vee)[1]]$ when X is special, where $[\pi'(\mathcal{Q}^\vee)[1]] \in \mathcal{M}_\sigma(\mathcal{A}_X, -x)$ is the point represented by the object $\pi'(\mathcal{Q}^\vee)[1]$. Thus p is a blow-down morphism to a smooth point when X is ordinary and a contraction of the component \mathbb{P}^2 to a singular point when X is special by Lemma 5.6.

When X is general and ordinary, the Fano surface $\mathcal{C}(X)$ is smooth by Theorem 6.4. Thus $\mathcal{M}_\sigma(\mathcal{A}_X, -x)$ is a smooth surface obtained by blowing down a smooth rational curve L_σ on the smooth irreducible projective surface $\mathcal{C}(X)$. This implies that $\mathcal{M}_\sigma(\mathcal{A}_X, -x)$ is also a smooth irreducible projective surface. On the other hand, it is known that there is a unique rational curve $L_\sigma \subset \mathcal{C}(X)$ and it is the unique exceptional curve by Lemma 6.5. Thus $\mathcal{M}_\sigma(\mathcal{A}_X, -x)$ is isomorphic to the minimal model $\mathcal{C}_m(X)$ of Fano surface of conics on X .

When X is general and special, the last statement follows from Theorem 6.11 and Lemma 7.3. \square

7.2. Involutions on $\mathcal{C}_m(X)$. In this section, we are going to describe the involution ι on $\mathcal{C}_m(X)$ in Theorem 6.6, described in [DIM12, Section 5.2] using the involution on \mathcal{A}_X . Recall that there is a natural involutive autoequivalence functor of \mathcal{A}_X , denoted by $\tau_{\mathcal{A}}$ (cf. Remark 3.2). When X is special, it is induced by the natural involution τ on X , which comes from the double cover $X \rightarrow Y$. In this case it is easy to see that $\tau_{\mathcal{A}}(\mathrm{pr}(I_C)) \cong \mathrm{pr}(I_{\tau(C)})$.

When X is ordinary, the situation is more subtle. In the following, we describe the action of $\tau_{\mathcal{A}}$ on the projection into \mathcal{A}_X of an ideal sheaf of a conic $\mathrm{pr}(I_C)$ in this case, and compare with the involution ι on $\mathcal{C}_m(X)$ described in [DIM12, Section 5.2].

Proposition 7.13. *Let X be an ordinary GM threefold and C a conic on X .*

- (1) *If $I_C \in \mathcal{A}_X$, then $\tau_{\mathcal{A}}(I_C)$ is either*
 - (a). *$I_{C'}$ such that $C \cup C' = Z(s)$ for $s \in H^0(\mathcal{E}^\vee)$, where $Z(s)$ is the zero locus of the section s ;*
 - (b). *or $\pi'(\mathcal{Q}^\vee)$, and in this case C is the ρ -conic*
- (2) *If $I_C \notin \mathcal{A}_X$, then $\tau_{\mathcal{A}}(\mathrm{pr}(I_C)) \cong I_{C''}$ for the ρ -conic $C'' \subset X$.*

Therefore, the involution induced by $\tau_{\mathcal{A}}$ on $\mathcal{C}_m(X)$ is the same as ι in Theorem 6.6.

Remark 7.14. We can define a birational involution on $\mathcal{C}(X)$ for any GM threefold X as in Proposition 7.13 (1)(a), which is regular on the locus of conics C with $\mathrm{hom}(\mathcal{E}, I_C) = 1$.

We first state some lemmas which we require for the proof of the proposition above.

Lemma 7.15. *Let X be an ordinary GM threefold and C be the ρ -conic on X . Then the natural morphism $s': \mathcal{E}^{\oplus 2} \rightarrow I_C$ is surjective and there is a short exact sequence*

$$0 \rightarrow \mathcal{Q}(-H) \rightarrow \mathcal{E}^{\oplus 2} \xrightarrow{s'} I_C \rightarrow 0.$$

Proof. By Lemma 6.3, we have $\mathrm{Hom}(\mathcal{E}, I_C) = k^2$. Thus, taking two linearly independent elements in $\mathrm{Hom}(\mathcal{E}, I_C)$, we have a natural map $s': \mathcal{E}^{\oplus 2} \rightarrow I_C$. Moreover, since $\langle C \rangle = \mathrm{Gr}(2, 3)$ and $\langle C \rangle \cap X = C$, we know that s' is surjective. Let $K := \ker(s')$. Then it is not hard to see that $\mathrm{ch}(K) = \mathrm{ch}(\mathcal{Q}(-H))$. Note that $\mathrm{Hom}(\mathcal{E}, K) = 0$ and K is reflexive.

We claim that K is μ -semistable. Indeed, suppose K is not μ -semistable and let K' be its maximal destabilizing subsheaf. Then K' is also reflexive. Since $\mathrm{Hom}(\mathcal{E}, K) = 0$, we have $K' \neq \mathcal{E}$. By the stability

of \mathcal{E} and the fact that $K \subset \mathcal{E}^{\oplus 2}$, we know that $\mu(K') = -\frac{1}{2}$. Since $\text{Hom}(K', \mathcal{E}) \neq 0$, by the stability of K' and \mathcal{E} we have $K' \subset \mathcal{E}$. Thus from $\text{ch}_{\leq 1}(K') = \text{ch}_{\leq 1}(\mathcal{E})$ we know that \mathcal{E}/K' is supported in codimension ≥ 2 , which gives a contradiction since \mathcal{E} and K' are both reflexive.

Now the result follows from Lemma 7.8, since $K(H)$ is μ -semistable with $\text{ch}(K(H)) = \text{ch}(\mathcal{Q})$. \square

Lemma 7.16. *Let X be an ordinary GM threefold. Let C be a conic on X . Then*

$$\mathbf{L}_{\mathcal{E}}(I_C) = \begin{cases} \mathbb{D}(I_{C'}) \otimes \mathcal{O}_X(-H)[1], & \text{RHom}^{\bullet}(\mathcal{E}, I_C) = k \\ \pi(\mathcal{E}), & \text{RHom}^{\bullet}(\mathcal{E}, I_C) = k^2 \oplus k[-1] \end{cases}$$

such that $C \cup C' = Z(s)$ for $s \in H^0(\mathcal{E}^{\vee})$, where $Z(s)$ is the zero locus of the section s

Proof. By Lemma 6.3, we have that $\text{RHom}^{\bullet}(\mathcal{E}, I_C)$ is either k or $k^2 \oplus k[-1]$. If $\text{RHom}^{\bullet}(\mathcal{E}, I_C) = k$, then we have the triangle

$$\mathcal{E} \rightarrow I_C \rightarrow \mathbf{L}_{\mathcal{E}}(I_C).$$

Taking cohomology with respect to the standard heart we get

$$0 \rightarrow \mathcal{H}^{-1}(\mathbf{L}_{\mathcal{E}}(I_C)) \rightarrow \mathcal{E} \xrightarrow{s} I_C \rightarrow \mathcal{H}^0(\mathbf{L}_{\mathcal{E}}(I_C)) \rightarrow 0.$$

The image of the map s is the ideal sheaf of an elliptic quartic $D = Z(s)$ for $s \in H^0(\mathcal{E}^{\vee})$, thus we have following two short exact sequences: $0 \rightarrow \mathcal{H}^{-1}(\mathbf{L}_{\mathcal{E}}(I_C)) \rightarrow \mathcal{E} \rightarrow I_D \rightarrow 0$ and $0 \rightarrow I_D \rightarrow I_C \rightarrow \mathcal{H}^0(\mathbf{L}_{\mathcal{E}}(I_C)) \rightarrow 0$. Then $\mathcal{H}^{-1}(\mathbf{L}_{\mathcal{E}}(I_C))$ is a torsion-free sheaf of rank 1 with the same Chern character as $\mathcal{O}_X(-H)$. It is easy to show that it must be $\mathcal{O}_X(-H)$. On the other hand $\mathcal{H}^0(\mathbf{L}_{\mathcal{E}}(I_C))$ is supported on the residual curve C' of C in D and $\mathcal{H}^0(\mathbf{L}_{\mathcal{E}}(I_C)) \cong \mathcal{O}_{C'}(-H)$. Thus we have the triangle

$$\mathcal{O}_X(-H)[1] \rightarrow \mathbf{L}_{\mathcal{E}}(I_C) \rightarrow \mathcal{O}_{C'}(-H)$$

and we observe that $\mathbf{L}_{\mathcal{E}}(I_C)$ is exactly the twisted derived dual of the ideal sheaf $I_{C'}$ of a conic $C' \subset X$, i.e. $\mathbf{L}_{\mathcal{E}}(I_C) \cong \mathbb{D}(I_{C'}) \otimes \mathcal{O}_X(-H)[1]$.

If $\text{RHom}^{\bullet}(\mathcal{E}, I_C) = k^2 \oplus k[-1]$, then we have the triangle

$$\mathcal{E}^2 \oplus \mathcal{E}[-1] \rightarrow I_C \rightarrow \mathbf{L}_{\mathcal{E}}(I_C).$$

Taking the long exact sequence in cohomology with respect to the standard heart, we get

$$0 \rightarrow \mathcal{H}^{-1}(\mathbf{L}_{\mathcal{E}}(I_C)) \rightarrow \mathcal{E}^2 \xrightarrow{s'} I_C \rightarrow \mathcal{H}^0(\mathbf{L}_{\mathcal{E}}(I_C)) \rightarrow \mathcal{E} \rightarrow 0.$$

Now by Lemma 7.15, s' is surjective and the cohomology objects are given by $\mathcal{H}^{-1}(\mathbf{L}_{\mathcal{E}}(I_C)) \cong \mathcal{Q}(-H)$ and $\mathcal{H}^0(\mathbf{L}_{\mathcal{E}}(I_C)) \cong \mathcal{E}$, which implies that $\mathbf{L}_{\mathcal{E}}(I_C) \cong \pi(\mathcal{E})$. \square

Proof of Proposition 7.13. Since $\tau_{\mathcal{A}} \circ \tau_{\mathcal{A}} \cong \text{id}$, we have $\tau_{\mathcal{A}} \cong \tau_{\mathcal{A}}^{-1}$. By Proposition 2.6, we have $\tau_{\mathcal{A}} \cong \tau_{\mathcal{A}}^{-1} \cong \mathbf{L}_{\mathcal{O}_X} \circ \mathbf{L}_{\mathcal{E}^{\vee}}(- \otimes \mathcal{O}_X(H))[-1]$. Then

$$\begin{aligned} \tau_{\mathcal{A}}(I_C) &\cong \mathbf{L}_{\mathcal{O}_X} \circ \mathbf{L}_{\mathcal{E}^{\vee}}(I_C \otimes \mathcal{O}_X(H))[-1] \\ &\cong \mathbf{L}_{\mathcal{O}_X}(\mathbf{L}_{\mathcal{E}}(I_C) \otimes \mathcal{O}_X(H))[-1]. \end{aligned}$$

The left mutation $\mathbf{L}_{\mathcal{E}}(I_C)$ is given by

$$\text{RHom}^{\bullet}(\mathcal{E}, I_C) \otimes \mathcal{E} \rightarrow I_C \rightarrow \mathbf{L}_{\mathcal{E}}(I_C).$$

Note that by Lemma 6.3, $\text{RHom}^{\bullet}(\mathcal{E}, I_C)$ is either k or $k^2 \oplus k[-1]$, and in the latter case C is the unique ρ -conic. Then by Lemma 7.16,

$$\mathbf{L}_{\mathcal{E}}(I_C) = \begin{cases} \mathbb{D}(I_{C'}) \otimes \mathcal{O}_X(-H)[1], & \text{RHom}^{\bullet}(\mathcal{E}, I_C) = k \\ \pi(\mathcal{E}), & \text{RHom}^{\bullet}(\mathcal{E}, I_C) = k^2 \oplus k[-1] \end{cases}$$

If $\text{RHom}^{\bullet}(\mathcal{E}, I_C) = k$, then $\tau_{\mathcal{A}}(I_C) \cong \mathbf{L}_{\mathcal{O}_X}(\mathbb{D}(I_{C'}))$. We have the triangle

$$\text{RHom}^{\bullet}(\mathcal{O}_X, \mathbb{D}(I_{C'})) \otimes \mathcal{O}_X \rightarrow \mathbb{D}(I_{C'}) \rightarrow \mathbf{L}_{\mathcal{O}_X}(\mathbb{D}(I_{C'})).$$

Note that $\text{RHom}^{\bullet}(\mathcal{O}_X, \mathbb{D}(I_{C'})) \cong \text{RHom}^{\bullet}(I_{C'}, \mathcal{O}_X) = k \oplus k[-1]$. Then we have the triangle

$$\mathcal{O}_X \oplus \mathcal{O}_X[-1] \rightarrow \mathbb{D}(I_{C'}) \rightarrow \mathbf{L}_{\mathcal{O}_X}(\mathbb{D}(I_{C'})). \quad (4)$$

The derived dual $\mathbb{D}(I_{C'})$ is given by the triangle $\mathcal{O}_X \rightarrow \mathbb{D}(I_{C'}) \rightarrow \mathcal{O}_{C'}[-1]$. Then taking cohomology with respect to the standard heart of triangle (4) we have the long exact sequence

$$\begin{aligned} 0 &= \mathcal{H}^{-1}(\mathbb{D}(I_{C'})) \rightarrow \mathcal{H}^{-1}(\mathbf{L}_{\mathcal{O}_X}(\mathbb{D}(I_{C'}))) \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_X \\ &\rightarrow \mathcal{H}^0(\mathbf{L}_{\mathcal{O}_X}(\mathbb{D}(I_{C'}))) \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_{C'} \rightarrow \mathcal{H}^1(\mathbf{L}_{\mathcal{O}_X}(\mathbb{D}(I_{C'}))) \rightarrow 0. \end{aligned}$$

Thus we have $\mathcal{H}^{-1}(\mathbf{L}_{\mathcal{O}_X}(\mathbb{D}(I_{C'}))) = 0$, $\mathcal{H}^1(\mathbf{L}_{\mathcal{O}_X}(\mathbb{D}(I_{C'}))) = 0$ and $\mathcal{H}^0(\mathbf{L}_{\mathcal{O}_X}(\mathbb{D}(I_{C'}))) \cong I_{C'}$. Hence $\tau_{\mathcal{A}}(I_C) \cong \mathbf{L}_{\mathcal{O}_X}(\mathbb{D}(I_{C'})) \cong I_{C'}$.

If $\mathrm{RHom}^\bullet(\mathcal{E}, I_C) = k^2 \oplus k[-1]$, then $\tau_{\mathcal{A}}(I_C) \cong \mathbf{L}_{\mathcal{O}_X} \circ \mathbf{L}_{\mathcal{E}^\vee}(I_C \otimes \mathcal{O}_X(H))[-1] \cong \mathbf{L}_{\mathcal{O}_X}(\pi(\mathcal{E}) \otimes \mathcal{O}_X(H)[-1])$ by Lemma 7.16. Then using the triangle (2), we have $\tau_{\mathcal{A}}(I_C) \cong \pi'(\mathcal{Q}^\vee)$. Then (2) follows from $\tau_{\mathcal{A}} \cong \tau_{\mathcal{A}}^{-1}$.

Now since $\tau_{\mathcal{A}} = S_{\mathcal{A}_X}[-2]$ and $\tau_{\mathcal{A}}$ acts trivially on $\mathcal{N}(\mathcal{A}_X)$, it induces an involution on the Bridgeland moduli space of any class with respect to any Serre-invariant stability condition. In particular, $\tau_{\mathcal{A}}$ induces an involution on $\mathcal{C}_m(X) \cong \mathcal{M}_\sigma(\mathcal{A}_X, -x)$ by Theorem 7.12. By (1) and (2), this induced involution coincides with ι in Theorem 6.6, described in [DIM12, Section 5.2]. \square

Remark 7.17. Smooth τ -conics form an open subscheme U of $\mathcal{C}(X)$. Therefore, the open subscheme $U \cap \iota(U)$ parameterizes smooth τ -conics C such that their involutive conics in Proposition 7.13 are smooth as well. The same also works for special GM threefolds, but replace τ -conics with conics with $\mathrm{hom}(\mathcal{E}, I_C) = 1$ and $I_C \in \mathcal{A}_X$, which are parametrized by $\mathcal{C}(X) \setminus \mathbb{P}^2$. In other words, for any GM threefold X , there is a two-dimensional open subscheme $\mathcal{C}_1 \subset \mathcal{C}(X)$ parameterizing smooth conics C with $\mathrm{hom}(\mathcal{E}, I_C) = 1$ such that their involutive conics are smooth.

8. THE MODULI SPACE $M_G(2, 1, 5)$ FOR GM THREEFOLDS

In this section, we investigate the moduli space of rank 2 Gieseker-semistable sheaves on a GM threefold X with Chern classes $c_1 = H, c_2 = 5L$ and $c_3 = 0$, denoted $M_G^X(2, 1, 5)$. We drop X from the notation when it is clear from context on which threefold we work. Note that if $F \in M_G(2, 1, 5)$, then

$$\mathrm{ch}(F) = (2, H, 0, -\frac{5}{6}P).$$

We are interested in $M_G(2, 1, 5)$ since it naturally appears in the description of the period fiber in [DIM12]. Our main theorem in Section is Theorem 8.9, which realizes $M_G(2, 1, 5)$ as the Bridgeland moduli space $\mathcal{M}_\sigma(\mathcal{A}_X, y - 2x)$.

First, we prove a classification result of sheaves in $M_G(2, 1, 5)$.

Proposition 8.1. *Let X be a GM threefold and $F \in M_G(2, 1, 5)$. Then we have $\mathrm{RHom}^\bullet(\mathcal{O}_X, F) = k^4$ and $\mathrm{RHom}^\bullet(\mathcal{O}_X, F(-H)) = 0$. Moreover, F is either a*

- (1) *globally generated bundle which fits into a short exact sequence*

$$0 \rightarrow \mathcal{O}_X \rightarrow F \rightarrow I_Z(H) \rightarrow 0$$

where Z is a projective normal smooth elliptic quintic curve;

- (2) *non-locally free sheaf with a short exact sequence*

$$0 \rightarrow F \rightarrow \mathcal{E}^\vee \rightarrow \mathcal{O}_L \rightarrow 0$$

where L is a line on X . Moreover, F is uniquely determined by L .

Remark 8.2. In [DIM12, Section 8], they also did computations for non-globally generated bundles in $M_G^X(2, 1, 5)$. However, in the following proof, we will show such sheaves do not exist.

Proof. The first statement follows from [BF14, Proposition 3.5 (1)] and the fact $\chi(F) = 4$. (1) and (2) also follow from [BF14, Proposition 3.5] or the argument in [DIM12, Section 8]. So we only need to prove the non-existence of non-globally generated bundles in $M_G^X(2, 1, 5)$. If $F \in M_G^X(2, 1, 5)$ is a non-globally generated bundle, then as showed in [DIM12, Section 8], we have an exact sequence

$$0 \rightarrow F^\vee \rightarrow \mathcal{O}_X^{\oplus 4} \rightarrow \mathcal{E}^\vee \xrightarrow{a} \mathcal{O}_L \rightarrow 0.$$

By (2), we know that $E := \ker(a)$ is a non-locally free stable sheaf and $E \in M_G^X(2, 1, 5)$. Thus we have an exact sequence $0 \rightarrow F^\vee \rightarrow \mathcal{O}_X^{\oplus 4} \rightarrow E \rightarrow 0$. In particular, E is generated by global sections.

However, we also have the following commutative diagram of exact sequences:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{O}_X^{\oplus 4} & \longrightarrow & \mathcal{O}_X^{\oplus 5} & \longrightarrow & \mathcal{O}_X \longrightarrow 0 \\ & & \downarrow \text{ev} & & \downarrow & & \downarrow \\ 0 & \longrightarrow & E & \longrightarrow & \mathcal{E}^\vee & \longrightarrow & \mathcal{O}_L \longrightarrow 0 \end{array}$$

where $\text{ev}: \mathcal{O}_X^{\oplus 4} \rightarrow E$ is the evaluation map. Then using Snake Lemma, we have an exact sequence

$$0 \rightarrow \ker(\text{ev}) \rightarrow \mathcal{Q}^\vee \xrightarrow{s} I_L \rightarrow \text{cok}(\text{ev}) \rightarrow 0.$$

As shown in Lemma 6.3 and Proposition 7.1, the image of s is the zero locus of a non-zero section of \mathcal{Q} . It is a σ -conic when X is ordinary, and a preimage of a line on Y when X is special. Hence in both cases, $\text{im}(s)$ is an ideal sheaf of a conic, and s is not surjective. Therefore, ev is not surjective as well and we get a contradiction. \square

A natural question to ask is what Bridgeland moduli space we get after projecting a sheaf in $M_G(2, 1, 5)$ into the Kuznetsov component. Since it is easier in this setting, we will work with the alternative Kuznetsov component \mathcal{A}_X in this section. Our analysis of the projections of objects in $M_G(2, 1, 5)$ is based on the three cases listed in Proposition 8.1. We begin with a Hom-vanishing result.

Lemma 8.3. *Let X be a GM threefold and $F \in M_G^X(2, 1, 5)$. Then we have $\text{RHom}^\bullet(\mathcal{E}^\vee, F) = 0$.*

Proof. By Serre duality and the stability of \mathcal{E}^\vee and F , we have $\text{Hom}(\mathcal{E}^\vee, F) = \text{Ext}^3(\mathcal{E}^\vee, F) = 0$. Since $\chi(\mathcal{E}^\vee, F) = 0$, we only need to show that $\text{Ext}^2(\mathcal{E}^\vee, F) = 0$ or $\text{Ext}^1(\mathcal{E}^\vee, F) = 0$. By Serre duality, we have $\text{Ext}^2(\mathcal{E}^\vee, F) = \text{Hom}(F, \mathcal{E}[1])$. Since $\text{ch}_1^0(F) = \text{ch}_1^0(\mathcal{E}[1]) = 1$, by Lemma 4.5 we know that F and $\mathcal{E}[1]$ are both $\sigma_{\alpha, 0}$ -stable for any $\alpha > 0$. Then $\text{Hom}(F, \mathcal{E}[1]) = 0$ since $\mu_{\alpha, 0}(F) > \mu_{\alpha, 0}(\mathcal{E}[1])$ when $0 < \alpha$ is sufficiently small. \square

We are now ready to give an explicit description of $\text{pr}(F)$, for all objects $F \in M_G(2, 1, 5)$. Recall that for any line $L \subset X$, we have $\mathcal{Q}|_L \cong \mathcal{O}_L^{\oplus 2} \oplus \mathcal{O}_L(1)$. Hence L is contained in a unique σ -conic C . We define the *residue line* of L to be the support of $\mathcal{O}_C \twoheadrightarrow \mathcal{O}_L$. Note that when C is a double line, we have $L' = L$.

Lemma 8.4. *Let X be a GM threefold and $F \in M_G(2, 1, 5)$.*

- *If F is a globally generated bundle, then*

$$\text{pr}(F) \cong \ker(\text{ev})[1],$$

where $\text{ev}: \mathcal{O}_X^{\oplus 4} \twoheadrightarrow F$ is the evaluation map.

- *If F is a non-locally free sheaf determined by a line $L \subset X$, then $\text{pr}(F)$ is the unique object fits into a non-trivial exact triangle*

$$\mathcal{E}[1] \rightarrow \text{pr}(F) \rightarrow \mathcal{O}_{L'}(-1),$$

where L' is the residue line of L .

Proof. As a result of Lemma 8.3, $\mathbf{L}_{\mathcal{E}^\vee} F = F$, so $\text{pr}(F) = \mathbf{L}_{\mathcal{O}_X} F$. By Proposition 8.1 we have $\text{RHom}^\bullet(\mathcal{O}_X, F) = k^4$, and the triangle defining the left mutation is

$$\mathcal{O}_X^{\oplus 4} \xrightarrow{\text{ev}} F \rightarrow \text{pr}(F). \quad (5)$$

In the cases where F is globally generated, the evaluation map ev is surjective, so $\text{pr}(F) = \ker(\text{ev})[1]$.

When F is non-locally free, as in Proposition 8.1, we have an exact sequence

$$0 \rightarrow \ker(\text{ev}) \rightarrow \mathcal{Q}^\vee \xrightarrow{s} I_L \rightarrow \text{cok}(\text{ev}) \rightarrow 0.$$

As shown in Lemma 6.3 and Proposition 7.1, the image of s is the zero locus of a non-zero section of \mathcal{Q} , which is a σ -conic. Hence by Proposition 7.1, we obtain $\ker(\text{ev}) = \mathcal{E}$ and $\text{cok}(\text{ev}) = \mathcal{O}_{L'}(-1)$. Since $\text{pr}(F) \in \mathcal{A}_X$, by Serre duality we have $\text{RHom}^\bullet(\mathcal{E}^\vee, \text{pr}(F)) = \text{RHom}^\bullet(\text{pr}(F), \mathcal{E})^\vee[-3] = 0$, which implies such triangle is non-trivial. And the uniqueness follows from $\text{Ext}^2(\mathcal{O}_{L'}(-1), \mathcal{E}) = H^1(\mathcal{E}(-1)|_L) = k$. \square

8.1. Stability of projection objects. In the following, we prove the stability of $\mathrm{pr}(F)$ for any $F \in M_G^X(2, 1, 5)$.

Lemma 8.5. *The functor $\mathrm{pr}: D^b(X) \rightarrow \mathcal{A}_X$ induces isomorphisms of $\mathrm{Ext}^k(\mathrm{pr}(F_1), \mathrm{pr}(F_2))$ and $\mathrm{Ext}^k(F_1, F_2)$ for all k and for all $F_1, F_2 \in M_G(2, 1, 5)$.*

Proof. We apply $\mathrm{Hom}(F_1, -)$ to the exact triangle $\mathcal{O}_X^{\oplus 4} \rightarrow F_2 \rightarrow \mathrm{pr}(F_2)$. By adjunction of pr and the inclusion $\mathcal{A}_X \hookrightarrow D^b(X)$, we have $\mathrm{Ext}^k(F_1, \mathrm{pr}(F_2)) = \mathrm{Ext}^k(\mathrm{pr}(F_1), \mathrm{pr}(F_2))$ for all k . Thus we get a long exact sequence

$$\cdots \rightarrow \mathrm{Ext}^k(F_1, \mathcal{O}_X)^{\oplus 4} \rightarrow \mathrm{Ext}^k(F_1, F_2) \rightarrow \mathrm{Ext}^k(\mathrm{pr}(F_1), \mathrm{pr}(F_2)) \rightarrow \mathrm{Ext}^{k+1}(F_1, \mathcal{O}_X)^{\oplus 4} \rightarrow \cdots$$

Note that $\mathrm{Ext}^k(F_1, \mathcal{O}_X) = \mathrm{Ext}^{3-k}(\mathcal{O}_X, F_1(-H)) = 0$ for all k by Proposition 8.1. Thus the desired result follows. \square

Before we show the stability of projection objects, let us recall a classical result:

Proposition 8.6. *Let X be an ordinary GM threefold and $L \subset X$ be a line. Then $\mathrm{RHom}^\bullet(\mathcal{O}_L, \mathcal{O}_L) = k \oplus k[-1]$ or $k \oplus k^2[-1] \oplus k[-2]$. Moreover, when X is general, we always have $\mathrm{RHom}^\bullet(\mathcal{O}_L, \mathcal{O}_L) = k \oplus k[-1]$.*

Proof. The first statement follows from [PS99, Lemma 4.2.1] and the second one follows from [PS99, Theorem 4.2.7]. \square

Now we are ready to prove the stability of $\mathrm{pr}(F)$.

Proposition 8.7. *Let X be a GM threefold and $F \in M_G^X(2, 1, 5)$. Then we have $\mathrm{RHom}^\bullet(F, F) = k \oplus k^2[-1]$ or $\mathrm{RHom}^\bullet(F, F) = k \oplus k^3[-1] \oplus k[-2]$. Hence $\mathrm{pr}(F)$ is stable with respect to every Serre-invariant stability condition on \mathcal{A}_X .*

Moreover,

- (1) *when X is ordinary, if $\mathrm{RHom}^\bullet(F, F) = k \oplus k^3[-1] \oplus k[-2]$ then F is a non-globally generated bundle or a non-locally free sheaf determined by a line L , and $[L] \in \Gamma(X)$ is a singular point. In particular, we always have $\mathrm{RHom}^\bullet(F, F) = k \oplus k^2[-1]$ when X is general;*
- (2) *when X is special, $\mathrm{RHom}^\bullet(F, F) = k \oplus k^3[-1] \oplus k[-2]$ if and only if $\tau^*F \cong F$, where τ is the natural involution on X .*

Proof. First, we assume that X is ordinary. We have $\mathrm{hom}(F, F) = 1$ and $\mathrm{ext}^3(F, F) = 0$ by Serre duality and the stability of F . Since $\chi(F, F) = -1$, we need to prove $\mathrm{ext}^2(F, F) = 0$ or 1.

When F is a globally generated bundle, by the proof of [DIM12, Theorem 8.2], we have $\mathrm{ext}^1(F, F) = 2$ and $\mathrm{ext}^2(F, F) = 0$. When F is non-locally free, there is a mistake made in the proof of [DIM12, Theorem 8.2] and we fix it here. From Proposition 8.1, we have an exact sequence $0 \rightarrow F \rightarrow \mathcal{E}^\vee \rightarrow \mathcal{O}_L \rightarrow 0$. Since $\mathrm{RHom}^\bullet(\mathcal{E}^\vee, F) = 0$ by Lemma 8.3, applying $\mathrm{Hom}(-, F)$ to this exact sequence, we get $\mathrm{Ext}^k(F, F) = \mathrm{Ext}^{k+1}(\mathcal{O}_L, F)$ for any k . Now applying $\mathrm{Hom}(\mathcal{O}_L, -)$ to this exact sequence, we get a long exact sequence

$$\cdots \rightarrow \mathrm{Ext}^2(\mathcal{O}_L, \mathcal{O}_L) \rightarrow \mathrm{Ext}^3(\mathcal{O}_L, F) \rightarrow \mathrm{Ext}^3(\mathcal{O}_L, \mathcal{E}^\vee) \rightarrow 0.$$

By Serre duality, we have $\mathrm{Ext}^3(\mathcal{O}_L, \mathcal{E}^\vee) = H^0(\mathcal{E}(-1)|_L) = 0$. Then from Proposition 8.6, we have $\mathrm{ext}^2(F, F) = \mathrm{ext}^3(\mathcal{O}_L, F) \leq \mathrm{ext}^2(\mathcal{O}_L, \mathcal{O}_L) \leq 1$. Moreover, if $\mathrm{ext}^2(F, F) = 1$, then $\mathrm{ext}^2(\mathcal{O}_L, \mathcal{O}_L) = 1$. In other words, $[L] \in \Sigma(X)$ is a singular point. This proves (1).

Now we assume that X is special. Then by Lemma 8.5 and Serre duality in $\mathcal{K}u(X)$, we have

$$\begin{aligned} \mathrm{Ext}^2(F, F) &\cong \mathrm{Ext}^2(\mathrm{pr}(F), \mathrm{pr}(F)) \\ &\cong \mathrm{Hom}(\mathrm{pr}(F), \tau_{\mathcal{A}}(\mathrm{pr}(F))) \\ &\cong \mathrm{Hom}(\mathrm{pr}(F), \mathrm{pr}(\tau^*F)) \cong \mathrm{Hom}(F, \tau^*F), \end{aligned}$$

where τ is the involution on X induced by the double cover. Thus when $F \cong \tau^*F$, we have $\mathrm{Ext}^2(F, F) = k$, and $\mathrm{Ext}^2(F, F) = 0$ otherwise. Since $\mathrm{Ext}^3(F, F) = 0$ and $\mathrm{Hom}(F, F) = k$, the result follows from $\chi(F, F) = -1$.

Finally, the stability of $\mathrm{pr}(F)$ follows from Lemma 8.5 and Proposition 4.12. \square

8.2. Involutions on $M_G(2, 1, 5)$. In this subsection, we briefly recall the involutions that exist on $M_G(2, 1, 5)$ and compare it with the one induced by $\tau_{\mathcal{A}}$. Let F be a globally generated vector bundle, and consider the short exact sequence

$$0 \rightarrow \ker(\text{ev}) \rightarrow H^0(X, F) \otimes \mathcal{O}_X \xrightarrow{\text{ev}} F \rightarrow 0.$$

Note that $\ker(\text{ev})$ is a rank 2 vector bundle with $c_1 = -H$ and $c_2 = 5L$ and no global sections, hence $\ker(\text{ev})^\vee \in M_G(2, 1, 5)$. Define $\iota F := \ker(\text{ev})^\vee$. This bundle ιF is globally generated, and we have $H^0(X, \iota F) \cong H^0(X, F)^\vee$ [DIM12, p. 29]. This defines a birational involution on $M_G^X(2, 1, 5)$.

Note that there is no non-globally generated bundle in $M_G^X(2, 1, 5)$ by Proposition 8.1, then the definition of ι on the non-locally free locus in [DIM12, Theorem 8.2] does not work. However, we can fix this issue as follows: for any non-locally free $F \in M_G^X(2, 1, 5)$ determined by a line L , we define $\iota(F) := F'$, where $F' := \ker(\mathcal{E}^\vee \rightarrow \mathcal{O}_{L'})$ is a non-locally free stable sheaf determined by the residue line L' of L . This extends ι to be a regular involution on $M_G^X(2, 1, 5)$.

Note that for a special GM threefold, there is another involution on $M_G(2, 1, 5)$ induced by the involution τ on X ,

$$\tau^*: M_G(2, 1, 5) \rightarrow M_G(2, 1, 5), \quad F \mapsto \tau^* F.$$

And it is clear that $\tau_{\mathcal{A}}(\text{pr}(F)) \cong \text{pr}(\tau^* F)$.

Now let X be an ordinary GM threefold, $\tau_{\mathcal{A}}$ be the involution of \mathcal{A}_X , and ι be the geometric involution of $M_G(2, 1, 5)$ defined above. Then $\tau_{\mathcal{A}}$ induces involutions of the Bridgeland moduli spaces of σ -stable objects $\mathcal{M}_\sigma(\mathcal{A}_X, -x)$ and $\mathcal{M}_\sigma(\mathcal{A}_X, y - 2x)$. In Proposition 7.13, we already showed that the action of $\tau_{\mathcal{A}}$ on $\mathcal{M}_\sigma(\mathcal{A}_X, -x)$ induces a geometric involution on $\mathcal{C}_m(X)$. In this section, we show that the involution induced by $\tau_{\mathcal{A}}$ is also compatible with ι on $M_G(2, 1, 5)$.

Proposition 8.8. *Let X be an ordinary GM threefold and $F \in M_G^X(2, 1, 5)$. Then $\tau_{\mathcal{A}}\text{pr}(F) \cong \text{pr}(\iota(F))$.*

Proof.

- (1) If F is a non-locally free sheaf determined by a line L , then by Lemma 8.4 we have the triangle

$$\mathcal{E}[1] \rightarrow \text{pr}(F) \rightarrow \mathcal{O}_{L'}(-1).$$

Then since $\tau_{\mathcal{A}} \cong \tau_{\mathcal{A}}^{-1} \cong \mathbf{L}_{\mathcal{O}_X} \circ \mathbf{L}_{\mathcal{E}^\vee}(- \otimes \mathcal{O}_X(H))[-1]$, $\tau_{\mathcal{A}}(\text{pr}(F))$ is given by a triangle

$$\mathbf{L}_{\mathcal{O}_X} \mathbf{L}_{\mathcal{E}^\vee}(\mathcal{E}^\vee) \rightarrow \tau_{\mathcal{A}}(\text{pr}(F)) \rightarrow \mathbf{L}_{\mathcal{O}_X} \mathbf{L}_{\mathcal{E}^\vee}(\mathcal{O}_{L'})[-1].$$

Note that $\mathbf{L}_{\mathcal{E}^\vee}(\mathcal{E}^\vee) = 0$, hence $\tau_{\mathcal{A}}(\text{pr}(F)) \cong \mathbf{L}_{\mathcal{O}_X} \mathbf{L}_{\mathcal{E}^\vee}(\mathcal{O}_{L'})[-1]$. It is easy to see $\text{RHom}^\bullet(\mathcal{E}^\vee, \mathcal{O}_{L'}) = k$, therefore we have $\mathcal{E}^\vee \rightarrow \mathcal{O}_{L'} \rightarrow \mathbf{L}_{\mathcal{E}^\vee} \mathcal{O}_{L'}$. Also, since $\mathcal{E}^\vee \rightarrow \mathcal{O}_{L'}$ is surjective, we have $\mathbf{L}_{\mathcal{E}^\vee} \mathcal{O}_{L'} \cong F'[1]$, where $F' := \ker(\mathcal{E}^\vee \rightarrow \mathcal{O}_{L'})$ is a non-locally free sheaf in $M_G(2, 1, 5)$ determined by L' as in Proposition 8.1. Thus $\tau_{\mathcal{A}}(\text{pr}(F)) \cong \mathbf{L}_{\mathcal{O}_X} F' \cong \text{pr}(F') = \text{pr}(\iota(F))$.

- (2) If F is a globally generated vector bundle, consider the standard short exact sequence

$$0 \rightarrow \ker(\text{ev}) \rightarrow H^0(X, F) \otimes \mathcal{O}_X \xrightarrow{\text{ev}} F \rightarrow 0.$$

Dualizing the sequence and applying pr , we get the triangle

$$\text{pr}(F^\vee) \rightarrow \text{pr}(\mathcal{O}_X^{\oplus 4}) \rightarrow \text{pr}(\ker(\text{ev})^\vee) \cong \text{pr}(\iota F).$$

Note that $F^\vee \in \mathcal{A}_X$ and $\text{pr}(\mathcal{O}_X) = 0$, thus we get $\text{pr}(\iota F) \cong F^\vee[1]$. Since $F \in M_G(2, 1, 5)$ is a globally generated vector bundle, we have $F \cong \iota E$ for some globally generated vector bundle E . Then $\text{pr}(F) = \text{pr}(\iota E) \cong E^\vee[1] \cong E \otimes \mathcal{O}_X(-H)[1]$, hence $\tau_{\mathcal{A}}(\text{pr}(F)) \cong \tau_{\mathcal{A}}(E \otimes \mathcal{O}_X(-H))[1] \cong \text{pr}(E) \cong \text{pr}(\iota F)$.

□

8.3. The Bridgeland moduli space of class $y - 2x$. In this subsection, we show that $M_G(2, 1, 5) \cong \mathcal{M}_\sigma(\mathcal{A}_X, y - 2x)$.

Theorem 8.9. *Let X be a GM threefold and σ be a Serre-invariant stability condition on \mathcal{A}_X . Then the projection functor $\mathrm{pr}: \mathrm{D}^b(X) \rightarrow \mathcal{A}_X$ induces an isomorphism $M_G^X(2, 1, 5) \cong \mathcal{M}_\sigma(\mathcal{A}_X, y - 2x)$.*

We split the proof of this theorem into a series of lemmas and propositions. Recall that in 4.9 we defined

$$V := \{(\alpha, \beta): -\frac{1}{10} < \beta < 0, 0 < \alpha < -\beta\}.$$

Proposition 8.10. *Let $F \in \mathcal{A}(\alpha, \beta)$ be a $\sigma(\alpha, \beta)$ -stable object with numerical class $y - 2x$ for every $(\alpha, \beta) \in V$. Then $F = \mathrm{pr}(E)$ for some $E \in M_G(2, 1, 5)$.*

Proof. First, we argue as in [PY22, Proposition 4.6]. When $(\alpha_0, \beta_0) = (0, 0)$, we have $\mu_{\alpha_0, \beta_0}^0(F) = -\infty$. Since there are no walls intersecting with $\beta = 0$ as in [PY22, Proposition 4.6], we know that F is $\sigma_{\alpha, 0}^0$ -semistable for all $\alpha > 0$. By the definition of the double-tilted heart, we have a triangle

$$A[1] \rightarrow F \rightarrow B$$

such that A (respectively B) is in $\mathrm{Coh}^0(X)$ with its $\sigma_{\alpha, 0}$ -semistable factors having slope $\mu_{\alpha, 0} \leq 0$ (respectively $\mu_{\alpha, 0} > 0$). Since F is $\sigma_{\alpha, 0}^0$ -semistable and $\mu_{\alpha, 0}^0(F) < 0$, we have that $A[1] = 0$ and $B \cong F$. Since $\mathrm{ch}_1^0(F)$ is minimal, there are no walls on $\beta = 0$, and we know that F is $\sigma_{\alpha, 0}$ -semistable for every $\alpha > 0$. Thus by Lemma 4.5, $\mathcal{H}^{-1}(F)$ is a μ -semistable reflexive sheaf and $\mathcal{H}^0(F)$ is 0 or supported in dimension ≤ 1 .

If $\mathcal{H}^0(F)$ is supported in dimension 0, then $\mathrm{ch}(\mathcal{H}^0(F)) = bP$ for $b \geq 1$. But this is impossible since then $c_3(\mathcal{H}^{-1}(F)) > 0$ and by [BF14, Proposition 3.5] we have $\chi(\mathcal{H}^{-1}(F)) = 0$, which implies $b = 0$.

If $\mathcal{H}^0(F)$ is supported in dimension 1, we can assume $\mathrm{ch}(\mathcal{H}^0(F)) = aL + \frac{b}{2}P$ where $a \geq 1$ and b are integers. Thus $\mathrm{ch}(\mathcal{H}^{-1}(F)) = 2 - H + aL + (\frac{5}{6} + \frac{b}{2})P$. Now from Lemma 7.7, we know $\mathcal{H}^{-1}(F) \cong \mathcal{E}$ and $\mathrm{ch}(\mathcal{H}^0(F)) = L - \frac{P}{2}$. Thus $\mathcal{H}^0(F) \cong \mathcal{O}_L(-1)$ for some line L on X . Therefore we have a triangle

$$\mathcal{E}[1] \rightarrow F \rightarrow \mathcal{O}_L(-1).$$

In this case we have $\mathrm{Hom}(\mathcal{O}_L(-1), \mathcal{E}[2]) = \mathrm{Hom}(\mathcal{E}^\vee(1), \mathcal{O}_L[1]) = H^1(L, \mathcal{E}(-1)|_L) = H^1(L, \mathcal{O}_L(-1) \oplus \mathcal{O}_L(-2)) = k$. Hence by Lemma 8.4, $F \cong \mathrm{pr}(E)$ for some $E \in M_G(2, 1, 5)$ such that E is locally free but not globally generated.

If $\mathcal{H}^0(F) = 0$, we have $F[-1] \cong \mathcal{H}^{-1}(F)$. Then $F[-1]$ is a μ -semistable sheaf. Since $F[-1]$ is reflexive and $c_3(F[-1]) = 0$, $F[-1] \in M_G(2, -1, 5)$ is a stable vector bundle. Thus by Lemma 8.4, we know $F[-1] = \mathrm{pr}(E)$ for some $E \in M_G(2, 1, 5)$ such that E is a globally generated vector bundle. \square

Lemma 8.11. *The functor $\mathrm{pr}: \mathrm{D}^b(X) \rightarrow \mathcal{A}_X$ is injective on all objects in $M_G(2, 1, 5)$, i.e. if $\mathrm{pr}(F_1) \cong \mathrm{pr}(F_2)$, then $F_1 \cong F_2$.*

Proof. For the case of globally generated vector bundles, by Corollary 8.4, $\mathrm{pr}(F_1) \cong \mathrm{pr}(F_2)$ implies that

$$(\iota F_1)^\vee \cong (\iota F_2)^\vee.$$

Note that $(\iota F_i)^\vee \cong \iota F_i \otimes \mathcal{O}_X(-H)$ for $i = 1, 2$. Then we get $\iota F_1 \cong \iota F_2$. Finally, we apply ι to both sides. Since it is an involution $\iota^2 = \mathrm{id}$, so $F_1 \cong F_2$ as required.

For the case of non-locally free sheaves F , recall that from Lemma 8.4 we have $\mathcal{H}^{-1}(\mathrm{pr}(F)) = \mathcal{E}$ and $\mathcal{H}^0(\mathrm{pr}(F)) = \mathcal{O}_L(-H)$. Since F is uniquely determined by the line L , and $\mathrm{Hom}(\mathcal{O}_L(-H), \mathcal{E}[2]) = k$, the object $\mathrm{pr}(F)$ is also uniquely determined by the line L . Thus $\mathrm{pr}(F_1) \cong \mathrm{pr}(F_2)$ implies $F_1 \cong F_2$, as required. \square

Proof of Theorem 8.9. Using Proposition 8.7, we know that the projection functor pr induces a morphism

$$p: M_G(2, 1, 5) \rightarrow \mathcal{M}_\sigma(\mathcal{A}_X, y - 2x)$$

which is bijective on points by Proposition 8.10 and Lemma 8.11, and bijective on tangent spaces by Lemma 8.5. Hence it is an isomorphism. \square

9. REFINED AND BIRATIONAL CATEGORICAL TORELLI THEOREMS FOR GM THREEFOLDS

In this section, we will prove several refined/birational categorical Torelli theorems for GM threefolds, using results from the previous sections.

9.1. The universal family for $\mathcal{C}_m(X)$. In this subsection, we show that $\mathcal{C}_m(X) \cong \mathcal{M}_\sigma(\mathcal{A}_X, -x)$ admits a universal family, which thus gives a fine moduli space. Let \mathcal{I} be the universal ideal sheaf of conics on $X \times \mathcal{C}(X)$ and \mathcal{I}_{L_σ} be the universal ideal sheaf of conics restricted to $X \times L_\sigma$. Let $q: X \times \mathcal{C}(X) \rightarrow X$ and $\pi: X \times \mathcal{C}(X) \rightarrow \mathcal{C}(X)$ be the projection maps on the first and second factors, respectively. Let $\mathcal{G}' := \text{pr}(\mathcal{I}_{L_\sigma})$ be the projected family in $\mathcal{A}_{X \times L_\sigma}$. Let $t \in L_\sigma \cong \mathbb{P}^1$ be any point. Then $j_t^* \text{pr}(\mathcal{I}_{L_\sigma}) \cong A$, where $j_t: X_t \rightarrow X_t \times L_\sigma$ and $A \in \mathcal{A}_X$ is $A \cong \text{pr}(I_C)$ for $I_C \notin \mathcal{A}_X$ by Proposition 7.2. Then $\mathcal{G}' \cong q^*(A) \otimes \pi^* \mathcal{O}_{L_\sigma}(k)$ for some $k \in \mathbb{Z}$. Now let $\mathcal{G} := \text{pr}(\mathcal{I}) \otimes \pi^* \mathcal{O}_{\mathcal{C}(X)}(kE)$, where $E \cong L_\sigma \cong \mathbb{P}^1$ is the unique exceptional curve on $\mathcal{C}(X)$.

Proposition 9.1. *The object $(p_X)_* \mathcal{G}$ is the universal family of $\mathcal{C}_m(X)$, where $p_X = \text{id}_X \times p: X \times \mathcal{C}(X) \rightarrow X \times \mathcal{C}_m(X)$.*

Proof.

- (1) If $s = [A] = \pi \in \mathcal{C}_m(X)$, s is contracted from the unique rational curve $L_\sigma \cong \mathbb{P}^1 \subset \mathcal{C}(X)$. Note that in this case $p_X|_{L_\sigma} = q$. Then

$$\begin{aligned} i_s^*(p_X)_* \mathcal{G} &\cong i_s^*(p_X)_*(\mathcal{G}' \otimes \pi^* \mathcal{O}_{\mathcal{C}(X)}(kE)) \\ &\cong i_s^* q_*(q^*(A) \otimes \pi^* \mathcal{O}_{L_\sigma}(k) \otimes \pi^* \mathcal{O}_{\mathcal{C}(X)}(kE)) \\ &\cong i_s^* q_*(q^*(A) \otimes (\pi^* \mathcal{O}_{L_\sigma}(k) \otimes \mathcal{O}_{L_\sigma}(kE))) \\ &\cong i_s^* q_*(q^*(A) \otimes \pi^*(\mathcal{O}_{L_\sigma}(k) \otimes \mathcal{O}_{L_\sigma}(-k))) \\ &\cong i_s^* q_*(q^*(A)) \cong i_s^*(A) \cong A. \end{aligned}$$

- (2) If $s = [I_C]$, then $\mathcal{C}_m(X)$ and $\mathcal{C}(X)$ are isomorphic outside L_σ . Note that p restricts to id on $\mathcal{C}(X) \setminus L_\sigma$. Then

$$\begin{aligned} i_s^*(p_X)_* \mathcal{G} &\cong i_s^*(p_X)_*(\text{pr}(\mathcal{I}) \otimes \pi^* \mathcal{O}_{\mathcal{C}(X)}(kE)) \\ &\cong j_s^*(\text{pr}(\mathcal{I})) \otimes j_s^* \pi^* \mathcal{O}_{\mathcal{C}(X)}(kE) \\ &\cong I_C \otimes (\pi \circ j_s)^* \mathcal{O}_{\mathcal{C}(X)}(kE) \\ &\cong I_C \otimes (i_s \circ \pi_s)^* \mathcal{O}_{\mathcal{C}(X)}(kE) \cong I_C. \end{aligned}$$

See below for the commutative diagrams which summarise the maps in the proof:

$$\begin{array}{ccccc} X_s & \xrightarrow{\cong} & X_s & \xrightarrow{\pi_s} & \{s\} \\ \downarrow j_s & & \downarrow i_s & & \downarrow \\ X \times \mathcal{C}(X) & \xrightarrow{p_X} & X \times \mathcal{C}_m(X) & \longrightarrow & \mathcal{C}_m(X) \end{array}$$

$$\begin{array}{ccc} X_s & \xrightarrow{j_s} & X \times \mathcal{C}(X) \\ \downarrow \pi_s & & \downarrow \pi \\ \{s\} & \xrightarrow{i_s} & \mathcal{C}(X) \end{array}$$

□

9.2. A refined categorical Torelli theorem for ordinary GM threefolds. We now prove a refined categorical Torelli theorem for ordinary GM threefolds.

Theorem 9.2. *Let X and X' be general ordinary GM threefolds such that $\Phi: Ku(X) \simeq Ku(X')$ is an equivalence and $\Phi(\pi(\mathcal{E})) \cong \pi(\mathcal{E}')$. Then $X \cong X'$.*

Proof. Since Φ commutes with Serre functors, it preserves the stability of an object with respect to any Serre-invariant stability condition. Then the existence of the universal family on $\mathcal{C}_m(X) \cong \mathcal{M}_\sigma(\mathcal{A}_X, -x)$ guarantees a morphism from $\mathcal{C}_m(X)$ to $\mathcal{C}_m(X')$, denoted by ψ , which is induced by Ψ (for more details on the construction of the morphism ψ , see [BMMS12, APR22]). Since Φ is an equivalence, ψ is an isomorphism. On the other hand, we have $\psi([\pi_X]) = [\pi_{X'}]$ by the assumption, where $\pi_X := \pi'(\mathcal{Q}^\vee)$ and $\pi_{X'} := \pi'(\mathcal{Q}^\vee)$. Then ψ induces an isomorphism $\phi: \mathcal{C}(X) \cong \mathcal{C}(X')$ by blowing up $[\pi_X] \in \mathcal{C}_m(X)$ and $[\pi_{X'}] \in \mathcal{C}_m(X')$, respectively. Then we have $X \cong X'$ by Logachev's Reconstruction Theorem 6.7. \square

9.3. Birational categorical Torelli theorem for ordinary GM threefolds. In this subsection, we show a birational categorical Torelli theorem for ordinary GM threefolds, i.e. assuming the Kuznetsov components are equivalent leads to a birational equivalence of the ordinary GM threefolds.

Theorem 9.3. *Let X and X' be general ordinary GM threefolds such that $\mathcal{A}_X \simeq \mathcal{A}_{X'}$. Then X' is a conic transform or a conic transform of a line transform of X . In particular, we have $X \simeq X'$.*

Proof. The equivalence $\Phi: \mathcal{A}_X \xrightarrow{\sim} \mathcal{A}_{X'}$ sends $-x$ to either itself or $y - 2x$ in $\mathcal{N}(\mathcal{A}_{X'})$ up to sign, since they are only (-1) -class. By the same argument as in Theorem 9.2 and [BMMS12, APR22], we thus get two possible induced isomorphisms between Bridgeland moduli spaces

$$\begin{array}{ccc} \mathcal{C}_m(X) \cong \mathcal{M}_\sigma(\mathcal{A}_X, -x) & \xrightarrow{\gamma} & \mathcal{C}_m(X') \cong \mathcal{M}_{\Phi(\sigma)}(\mathcal{A}_{X'}, -x) \\ & \searrow \gamma' & \\ & & M_G^{X'}(2, 1, 5) \cong \mathcal{M}_{\Phi(\sigma)}(\mathcal{A}_{X'}, y - 2x) \end{array}$$

If we have the isomorphism γ , then we blow up $\mathcal{C}_m(X)$ at the distinguished point $[\pi_X] := [\Xi(\pi(\mathcal{E}))]$, and blow up $\mathcal{C}_m(X')$ at the point $[C] := [\Phi(\pi_X)] = \gamma([\pi_X])$. We have

$$\mathcal{C}(X) \cong \text{Bl}_{[\pi_X]} \mathcal{C}_m(X) \cong \text{Bl}_{[C]} \mathcal{C}_m(X'),$$

and $\text{Bl}_{[C]} \mathcal{C}_m(X') \cong \mathcal{C}(X'_C)$ by Theorem 6.8, so $\mathcal{C}(X) \cong \mathcal{C}(X'_C)$. Therefore by Logachev's Reconstruction Theorem 6.7 we have $X \cong X'_C$.

For the second case, we get $\mathcal{C}_m(X) \cong M_G^{X'}(2, 1, 5)$. And by [DIM12, Proposition 8.1] we have a birational equivalence $M_G^{X'}(2, 1, 5) \simeq \mathcal{C}(X'_L)$ of surfaces, where $L \subset X'$ is a line. Then we see $\mathcal{C}_m(X)$ is birationally equivalent to $\mathcal{C}(X'_L)$. Let $\mathcal{C}_m(X'_L)$ be the minimal surface of $\mathcal{C}(X'_L)$. Note that the surfaces here are all smooth surfaces of general type. By the uniqueness of minimal models of surfaces of general type, we get $\mathcal{C}_m(X) \cong \mathcal{C}_m(X'_L)$, which implies $X \cong (X'_L)_C \simeq X'$ for a conic $C \subset X'_L$ as in the first case. \square

Remark 9.4. Theorem 9.3 proves a conjecture [KP23, Conjecture 1.7] of Kuznetsov–Perry for *general ordinary* GM varieties of dimension 3.

In [DIM12], the authors proved that $\mathcal{C}_m(X_L)$ is birational to $M_G^X(2, 1, 5)$. The following corollary shows that they are indeed isomorphic.

Corollary 9.5. *Let X be a general ordinary GM threefold, and X_L be a line transform of X . Then we have $\mathcal{C}_m(X_L) \cong M_G^X(2, 1, 5)$. Moreover, this isomorphism commutes with involutions ι and ι' on both sides, thus giving an isomorphism $\mathcal{C}_m(X_L)/\iota \cong M_G^X(2, 1, 5)/\iota'$.*

Proof. By the same argument as in the proof of Theorem 9.3, we have $\mathcal{C}_m(X_L) \cong \mathcal{C}_m(X)$ or $\mathcal{C}_m(X_L) \cong M_G^X(2, 1, 5)$. Note that $\mathcal{C}_m(X_L) \cong \mathcal{C}_m(X)$ implies that $X_L \cong X_C$ for some conic $C \subset X$ as in Theorem 9.3. But this is impossible by [DIM12, Remark 7.3]. Thus we always have $\mathcal{C}_m(X_L) \cong M_G^X(2, 1, 5)$. The last statement follows from the fact that any equivalence between Kuznetsov components commutes with Serre functors, and the involutions on $\mathcal{C}_m(X_L)$ and $M_G^X(2, 1, 5)$ can be induced by Serre functors up to shift by Propositions 7.13 and Proposition 8.8. \square

Since the intermediate Jacobian $J(X)$ is invariant under conic and line transforms, as a corollary we have

Corollary 9.6. *Let X and X' be general ordinary GM threefolds. If $Ku(X) \simeq Ku(X')$, then we have $J(X) \cong J(X')$.*

In fact, we can relax the assumptions on X by looking at the singularities of Bridgeland moduli spaces.

Theorem 9.7. *Let X and X' be general GM threefolds (they can be either general ordinary or general special) and suppose their Kuznetsov components $\mathcal{A}_X \simeq \mathcal{A}_{X'}$ are equivalent. Then X is birationally equivalent to X' .*

Proof. First, we claim that if X and X' are general GM threefolds such that $\Phi: \mathcal{A}_X \simeq \mathcal{A}_{X'}$, then both X and X' are ordinary or special simultaneously. Indeed, we may assume X' is ordinary and X is special. Then the equivalence would identify the moduli space $\mathcal{C}_m(X) \cong \mathcal{M}_\sigma(\mathcal{A}_X, -x)$ of stable objects of class $-x$ in \mathcal{A}_X with either the moduli space $\mathcal{C}_m(X') \cong \mathcal{M}_{\sigma'}(\mathcal{A}_{X'}, -x)$ or $M_G^{X'}(2, 1, 5) \cong \mathcal{M}_{\sigma'}(\mathcal{A}_{X'}, y - 2x)$. But $\mathcal{C}_m(X)$ has a unique singular point by Theorem 7.12, and both $\mathcal{C}_m(X')$ and $M_G^{X'}(2, 1, 5)$ are smooth for X' general by Theorem 7.12 and Theorem 8.9. This means that neither identification is possible, so the claim follows.

Now X and X' are both general ordinary or general special, hence the result follows from Theorem 9.3 and 9.4. \square

Corollary 9.8. *Let X and X' be general GM threefolds such that one of them is ordinary and their Kuznetsov components $\mathcal{A}_X \simeq \mathcal{A}_{X'}$ are equivalent. Then they are both general ordinary and X is birationally equivalent to X' .*

9.4. A categorical Torelli theorem for special GM threefolds. In this subsection, we show that the Kuznetsov component of a general special GM threefold X determines the isomorphism class of X .

Recall from Section 3 that every special GM threefold X is a double cover of a degree 5 index 2 prime Fano threefold Y branched over a quadric hypersurface \mathcal{B} in Y . Since X is smooth and general, (\mathcal{B}, h) is a smooth degree $h^2 = 10$ K3 surface with Picard number 1. There is a natural geometric involution τ on X induced by the double cover. The Serre functor on $Ku(X)$ is given by $S_{Ku(X)} = \tau \circ [2]$.

Theorem 9.9. *Let X and X' be general special GM threefolds with $\Phi: Ku(X) \simeq Ku(X')$. Then $X \cong X'$.*

Proof. By [KP17, Theorem 1.1, Section 8.2], the equivariant triangulated category $Ku(X)^{\mu_2}$ is equivalent to $D^b(\mathcal{B})$, where μ_2 is the group of square roots of 1 generated by the involution τ acting on $Ku(X)$. Assume there is an equivalence $\Phi: Ku(X) \simeq Ku(X')$. Since $S_{Ku(X)} \cong \tau[2]$ and $S_{Ku(X')} \cong \tau'[2]$, Φ commutes with the involutions τ and τ' on $Ku(X)$ and $Ku(X')$, respectively. Then we get an induced equivalence

$$\Psi: Ku(X)^{\mu_2} \simeq Ku(X')^{\mu'_2}$$

where $\mu_2 = \langle \tau \rangle$, $\mu'_2 = \langle \Phi \circ \tau \circ \Phi^{-1} = \tau' \rangle$ and $\mu_2 \cong \mu'_2$. Thus we have $\Psi: D^b(\mathcal{B}) \simeq D^b(\mathcal{B}')$. We know that \mathcal{B} and \mathcal{B}' are smooth projective surfaces with polarizations h and h' , respectively, so Ψ is a Fourier–Mukai functor by Orlov’s Representability Theorem [Orl97, Theorem 2.2]. Moreover, (\mathcal{B}, h) and (\mathcal{B}', h') are both Picard number 1 smooth projective K3 surfaces of degree $h^2 = h'^2 = 10 = 2 \times 5$. Then by [Ogu02, Theorem 1.10] and [HLOY03, Corollary 1.7], there is an isomorphism $\phi: \mathcal{B} \cong \mathcal{B}'$. Since they both have Picard number one, we obtain $\phi^*(h') = h$. On the other hand Y_5 is rigid [Kuz09, § 4.1], which implies $X \cong X'$. \square

Remark 9.10. Theorem 9.9 can also be proved via Bridgeland moduli spaces with respect to the Kuznetsov component \mathcal{A}_X . The details are contained in another paper of authors [JLZ22].

10. THE DEBARRE–ILIEV–MANIVEL CONJECTURE

Let \mathcal{X}_{10} be the moduli space of smooth ordinary GM threefolds and \mathcal{A}_{10} be the moduli space of 10-dimensional principal polarised abelian varieties. In [DIM12, pp. 3-4], the authors make the following conjecture regarding the general fiber of the period map:

Conjecture 10.1 ([DIM12, pp. 3-4]). *A general fiber $\mathcal{P}^{-1}([J(X)])$ of the period map $\mathcal{P}: \mathcal{X}_{10} \rightarrow \mathcal{A}_{10}$ at the intermediate Jacobian $J(X)$ of an ordinary GM threefold X is the union of $\mathcal{C}_m(X)/\iota$ and a surface birationally equivalent to $M_G^X(2, 1, 5)/\iota'$, where ι, ι' are geometrically meaningful involutions.*

Remark 10.2. Note that by Corollary 9.5, the surface birationally equivalent to $M_G(2, 1, 5)/\iota'$ in [DIM12], parametrizing conic transforms of a line transform of X , is actually isomorphic to $M_G(2, 1, 5)/\iota'$. Thus this conjecture predicts that a general fiber $\mathcal{P}^{-1}([J(X)])$ is actually the disjoint union of $\mathcal{C}_m(X)/\iota$ and $M_G(2, 1, 5)/\iota'$.

We will prove a categorical analogue of this conjecture. Consider the “categorical period map”

$$\mathcal{P}_{\text{cat}}: \mathcal{X}_{10} \rightarrow \{\mathcal{A}_X\}/\simeq, \quad X \mapsto \mathcal{A}_X$$

where \mathcal{X}_{10} is the moduli space of isomorphism classes of GM threefolds and $\{\mathcal{A}_X\}/\simeq$ is the set of equivalence classes of Kuznetsov components of GM threefolds. Note that a global description of a “moduli of Kuznetsov components” $\{\mathcal{A}_X\}/\simeq$ is not known, however, local deformations are controlled by the second Hochschild cohomology $\text{HH}^2(\mathcal{A}_X)$. The fiber of the “categorical period map” \mathcal{P}_{cat} over \mathcal{A}_X for an ordinary GM threefold is defined as the isomorphism classes of all ordinary GM threefolds X' such that $\mathcal{A}_{X'} \simeq \mathcal{A}_X$.

Theorem 10.3. *The general fiber $\mathcal{P}_{\text{cat}}^{-1}([\mathcal{A}_X])$ of the categorical period map over the alternative Kuznetsov component of an ordinary GM threefold X is the union of $\mathcal{C}_m(X)/\iota$ and $M_G^X(2, 1, 5)/\iota'$ where ι, ι' are geometrically meaningful involutions.*

Proof. The general fiber $\mathcal{P}_{\text{cat}}^{-1}([\mathcal{A}_X])$ of the categorical period map consists of GM threefolds X' such that there is an equivalence of Kuznetsov components $\mathcal{A}_{X'} \simeq \mathcal{A}_X$. Then by Theorem 9.7, X' is also a general ordinary GM threefold. Thus by Theorem 9.3 and Theorem 6.10, we know that $\mathcal{A}_{X'} \simeq \mathcal{A}_X$ if and only if X' is a conic transform of X , or a conic transform of a line transform of X . Then the result follows from Proposition 6.9 and Corollary 9.5. \square

The Kuznetsov components of prime Fano threefolds of index 1 and 2 are often regarded as categorical analogues of the intermediate Jacobians of these threefolds, and it is known that if there is a Fourier–Mukai type equivalence $\mathcal{K}u(X) \simeq \mathcal{K}u(X')$ (or $\mathcal{A}_X \simeq \mathcal{A}_{X'}$), then $J(X) \cong J(X')$ by [Per22]. For the converse, we have the following result.

Theorem 10.4. *For smooth prime Fano threefolds X , if X is one of the following:*

- Y_d , $2 \leq d \leq 5$
- X_{2g-2} , $g = 5, 7, 8, 9, 10, 12$,

then the intermediate Jacobian $J(X)$ uniquely determines the Kuznetsov component $\mathcal{K}u(X)$, i.e. for another prime Fano threefold X' of the same degree, if $J(X) \cong J(X')$, then $\mathcal{K}u(X) \simeq \mathcal{K}u(X')$.

Proof. If X is an index 2 prime Fano threefold Y_d of degree $2 \leq d \leq 5$, then the statement follows from the Torelli theorems for Y_d . Now let X_d be a degree d index one prime Fano threefold. If $X = X_8$, the statement follows from its Torelli theorem. If $X = X_{12}, X_{18}, X_{16}$, their intermediate Jacobians are Jacobians of curves: $J(X_{12}) \cong J(C_7)$, $J(X_{16}) \cong J(C_3)$, and $J(X_{18}) \cong J(C_2)$. But $\mathcal{K}u(X_{12}) \simeq \text{D}^b(C_7)$, $\mathcal{K}u(X_{16}) \simeq \text{D}^b(C_3)$ and $\mathcal{K}u(X_{18}) \simeq \text{D}^b(C_2)$. Thus the statement follows from the classical Torelli theorem for curves. If $X = X_{14}$, the statement follows from the Kuznetsov conjecture for the pair (Y_3, X_{14}) [Kuz04] and the Torelli theorem for cubic threefolds. If $X = X_{22}$, the statement is trivial since $\mathcal{K}u(X_{22}) \cong \mathcal{K}u(Y_5)$ ([KPS18]) and Y_5 is rigid, so $\mathcal{K}u(X) \simeq \mathcal{K}u(X')$ is always true. \square

Therefore, it is natural to make the following conjecture:

Conjecture 10.5. *Let X be a prime Fano threefold of index one or two. Then the intermediate Jacobian $J(X)$ uniquely determines the Kuznetsov component $\mathcal{K}u(X)$, i.e. for another prime Fano threefold X' of the same degree, if $J(X) \cong J(X')$, then $\mathcal{K}u(X) \simeq \mathcal{K}u(X')$.*

Surprisingly, in the case of general ordinary GM threefolds, we can restate the Debarre–Iliev–Manivel Conjecture 10.1 as Conjecture 10.5:

Proposition 10.6. *The Debarre–Iliev–Manivel Conjecture 10.1 is equivalent to Conjecture 10.5 for general ordinary GM threefolds.*

Proof. First, we assume that Conjecture 10.5 holds. Then by Corollary 9.6 and Theorem 10.3, the Debarre–Iliev–Manivel Conjecture 10.1 holds.

On the other hand, we assume the Debarre–Iliev–Manivel Conjecture 10.1 holds. Then for any X and X' such that $J(X) \cong J(X')$, X is either a conic transform of X' , or X is a conic transform of a line transform of X' . In both cases, we have $\mathcal{K}u(X) \simeq \mathcal{K}u(X')$ by the Duality Conjecture Theorem 6.10. Thus Conjecture 10.5 holds. \square

APPENDIX A. UNIQUENESS OF SERRE-INVARIANT STABILITY CONDITIONS

In this appendix, we aim to prove the uniqueness of Serre-invariant stability conditions on $\mathcal{K}u(X)$ for several prime Fano threefolds X (Theorem A.10). We start with a general criterion for when two numerical stability conditions with the same central charge are equal. We always assume that any triangulated category is k -linear and of finite type, i.e. $\sum_i \text{ext}^i(A, B) < +\infty$ for any two objects A, B . Therefore, the Euler form and the numerical Grothendieck group are well-defined.

Theorem A.1. *Let \mathcal{D} be a k -linear triangulated category of finite type. Assume that*

- (A). $\chi(x, x) \leq 1 - n$ for a positive integer n and any non-zero $x \in \mathcal{N}(\mathcal{D})$,
- (B). *there exists an object D satisfies*

$$n \leq \text{ext}^1(D, D) < 2n,$$

Let $\sigma_1 = (\mathcal{A}_1, Z)$ be a numerical stability condition on \mathcal{D} and $D_1, D_2 \in \mathcal{A}_1$ be two σ_1 -stable objects satisfying:

- (C). *for any two objects $A, B \in \mathcal{D}$, if $\phi_{\sigma_1}^+(B) < \phi_{\sigma_1}^-(A)$, then $\text{Hom}(B, A[2]) = 0$,*
- (D). *if E is a σ_1 -semistable object with $\chi(E, D_1) \geq 0$ and $\chi(E, D_2) \geq 0$, then there exist $k \in \{1, 2\}$ such that $\chi(E, D_k) > 0$ and $\mu_{\sigma_1}(E) < \mu_{\sigma_1}(D_k)$, and*
- (E). *if E is a σ_1 -semistable object with $\chi(D_1, E) \geq 0$ and $\chi(D_2, E) \geq 0$, then there exist $k \in \{1, 2\}$ such that $\chi(D_k, E) > 0$ and $\mu_{\sigma_1}(E) > \mu_{\sigma_1}(D_k)$.*

If $\sigma_2 = (\mathcal{A}_2, Z)$ is a numerical stability condition on \mathcal{D} satisfies (C), (D) and (E) such that D_1 and D_2 are σ_2 -stable with $\phi_{\sigma_2}(D_1) = \phi_{\sigma_1}(D_1)$ and $\phi_{\sigma_2}(D_2) = \phi_{\sigma_1}(D_2)$, then $\sigma_1 = \sigma_2$.

We first prove several lemmas. By the same proof as in [BB17, Lemma 2.5], we have the following generalized version of Weak Mukai Lemma:

Lemma A.2. *Let \mathcal{D} be a k -linear triangulated category with finite-dimensional Hom-space. Then for any exact triangle $A \rightarrow E \rightarrow B$ with $\text{Hom}(A, B) = \text{Hom}(B, A[2]) = 0$, we have*

$$\text{ext}^1(A, A) + \text{ext}^1(B, B) \leq \text{ext}^1(E, E).$$

Lemma A.3. *Let \mathcal{D} be a k -linear triangulated category of finite type satisfies (A). Assume that there is a stability condition $\sigma = (\mathcal{A}, Z)$ on \mathcal{D} satisfies (C).*

- (1) *The homological dimension of \mathcal{A} is at most 2.*
- (2) *For any exact triangle $A \rightarrow E \rightarrow B$ with $\phi_{\sigma}^-(A) > \phi_{\sigma}^+(B)$, we have*

$$\text{ext}^1(A, A) + \text{ext}^1(B, B) \leq \text{ext}^1(E, E).$$

- (3) *For any non-zero object $A \in \mathcal{D}$, we have $\text{ext}^1(A, A) \geq n$.*
- (4) *If a non-zero object E is not σ -semistable, then any Harder–Narasimhan factor A of E satisfies*

$$\text{ext}^1(A, A) < \text{ext}^1(E, E).$$

(5) Any object E with

$$n \leq \text{ext}^1(E, E) < 2n$$

is σ -semistable.

Proof. Let $A, B \in \mathcal{A}$. Then we have $\phi_\sigma^+(A) \leq 1 < \phi_\sigma^-(B[k])$ for any $k \geq 1$. Therefore, by (C) we get $\text{Hom}(A, B[k+2]) = \text{Hom}(A, B[k][2]) = 0$ for any $k \geq 0$. This proves (1).

Now for (2), note that $\text{Hom}(A, B) = 0$ and by (C) we have $\text{Hom}(B, A[2])$. Then the result follows from Lemma A.2.

Next, we prove (3). If $A \neq 0 \in \mathcal{A}$, then from (1), we get $\chi(A, A) = \text{hom}(A, A) - \text{ext}^1(A, A) + \text{ext}^2(A, A)$. Since $\chi(A, A) \leq 1 - n$ by (A), we know that $\text{ext}^1(A, A) \geq n$ in this case. Now for a general non-zero object $A \in \mathcal{D}$, if A is σ -semistable, then it is in \mathcal{A} up to shift and the result follows from the previous argument. So we assume that A is not σ -semistable. Let A' be the first Harder–Narasimhan factor of A with respect to σ , and $A'' := \text{cone}(A' \rightarrow A)$. We have $\phi_\sigma(A') > \phi_\sigma^+(A'')$. Using (2) and σ -semistability of A' , we obtain $n \leq \text{ext}^1(A', A') + \text{ext}^1(A'', A'') \leq \text{ext}^1(A, A)$ and hence (3) follows. And (4) follows from the induction on the number of Harder–Narasimhan factors of E and using (2) and (3).

Finally, if such E in (5) is not σ -semistable, then by the existence of Harder–Narasimhan filtration, we can find a triangle $A \rightarrow E \rightarrow B$ with $\phi_\sigma^-(A) > \phi_\sigma^+(B)$. By (2) and (3), this contradicts our assumption on $\text{ext}^1(E, E)$. Thus E is σ -semistable. \square

Now we are ready to prove our criterion.

Proof of Theorem A.1. Since σ_1 and σ_2 have the same central charge, it remains to show $\mathcal{A}_1 = \mathcal{A}_2$. By our assumptions, $D_1, D_2 \in \mathcal{A}_1 \cap \mathcal{A}_2$ are both σ_1 -stable and σ_2 -stable with phases in $(0, 1]$.

Step 1. First, we show that if E is a σ_i -semistable object which is also σ_j -semistable, then $\phi_{\sigma_1}(E) = \phi_{\sigma_2}(E)$, where $\{i, j\} = \{1, 2\}$. Since σ_1 and σ_2 satisfy the same assumptions, in the following, we will take $i = 2$ and $j = 1$. The other case can be deduced from the same argument but exchanges the role of σ_1 and σ_2 .

Up to shift, we can assume that $E \in \mathcal{A}_2$. Since σ_1 and σ_2 have the same central charge, we have $\phi_{\sigma_1}(E) - \phi_{\sigma_2}(E) = 2m$ for an integer m . Then we see

$$2m < \phi_{\sigma_1}(E) \leq 2m + 1. \quad (6)$$

- Assume that there exist $k, l \in \{1, 2\}$ such that $\chi(D_k, E) < 0$ and $\chi(E, D_l) < 0$. Then by Lemma A.3 (1) and the fact that $E, D_1, D_2 \in \mathcal{A}_2$, we have

$$\text{ext}^1(D_k, E) \neq 0, \quad \text{ext}^1(E, D_l) \neq 0,$$

which imply

$$-1 < \phi_{\sigma_1}(D_k) - 1 \leq \phi_{\sigma_1}(E) \leq \phi_{\sigma_1}(D_l) + 1 \leq 2.$$

Hence, by (6) we get $2m < 2$ and $-1 < 2m + 1$, which means $m = 0$ and we obtain $\phi_{\sigma_1}(E) = \phi_{\sigma_2}(E)$ as required.

- Assume that $\chi(E, D_1) \geq 0$ and $\chi(E, D_2) \geq 0$. By (D), there is an integer $s \in \{1, 2\}$ such that $\chi(E, D_s) > 0$ and $\mu_{\sigma_1}(E) < \mu_{\sigma_1}(D_s)$. Thus we have $\mu_{\sigma_2}(E) < \mu_{\sigma_2}(D_s)$ as well. Since $E, D_s \in \mathcal{A}_2$, we get $\phi_{\sigma_2}(E) < \phi_{\sigma_2}(D_s)$, which implies $\text{Hom}(E, D_s[2]) = 0$ by (C). Then from $\chi(E, D_s) > 0$ and Lemma A.3 (1), we obtain $\text{Hom}(E, D_s) \neq 0$, and hence

$$\phi_{\sigma_1}(E) \leq \phi_{\sigma_1}(D_s) \leq 1.$$

Now if one of $\chi(D_1, E)$ and $\chi(D_2, E)$ is negative, the same argument as in the first case shows that $-1 < \phi_{\sigma_1}(E)$.

If $\chi(D_1, E) \geq 0$ and $\chi(D_2, E) \geq 0$, then by (E), there is an integer $t \in \{1, 2\}$ such that $\chi(D_t, E) > 0$ and $\mu_{\sigma_1}(E) = \mu_{\sigma_2}(E) > \mu_{\sigma_1}(D_t) = \mu_{\sigma_2}(D_t)$. Since $E, D_t \in \mathcal{A}_2$, we get $\phi_{\sigma_2}(E) > \phi_{\sigma_2}(D_t)$, which by (C) implies $\text{Hom}(D_t, E[2]) = 0$. Then together with $\chi(D_t, E) > 0$ and Lemma A.3 (1), we see $\text{Hom}(D_t, E) \neq 0$. Therefore, we have $0 < \phi_{\sigma_1}(D_t) \leq \phi_{\sigma_1}(E)$. In both cases, we always have $\phi_{\sigma_1}(E) \in (-1, 2]$. By (6), we get $m = 0$ and $\phi_{\sigma_1}(E) = \phi_{\sigma_2}(E)$ as required.

- Assume that $\chi(D_1, E) \geq 0$ and $\chi(D_2, E) \geq 0$. Then using (E), by a similar argument as the second case, we obtain $\phi_{\sigma_1}(E) = \phi_{\sigma_2}(E)$. This completes the first step.

Step 2. Next, we prove that an object E is σ_1 -semistable if and only if σ_2 -semistable. We show this by induction on $\text{ext}^1(E, E)$. If $\text{ext}^1(E, E) < 2n$, then from (B) we know such E exists. By Lemma A.3 (5), E is both σ_1 -semistable and σ_2 -semistable.

Now assume that the statement holds for any object F with $\text{ext}^1(F, F) < N$ for an integer $N > 2n$. Let E be an object with $\text{ext}^1(E, E) = N$. If E is σ_i -semistable but not σ_j -semistable for $\{i, j\} = \{1, 2\}$, let A be the first Harder–Narasimhan factor of E with respect to σ_j and B be the last one. Therefore, we see

$$\phi_{\sigma_j}(A) > \phi_{\sigma_j}(B). \quad (7)$$

And from Lemma A.3 (4), we have

$$\text{ext}^1(A, A) < \text{ext}^1(E, E), \quad \text{ext}^1(B, B) < \text{ext}^1(E, E).$$

Therefore, by the induction hypothesis, A and B are σ_i -semistable as well and the first step implies

$$\phi_{\sigma_1}(A) = \phi_{\sigma_2}(A), \quad \phi_{\sigma_1}(B) = \phi_{\sigma_2}(B). \quad (8)$$

But then from $\text{Hom}(A, E) \neq 0$ and $\text{Hom}(E, B) \neq 0$, we have $\phi_{\sigma_i}(A) \leq \phi_{\sigma_i}(E) \leq \phi_{\sigma_i}(B)$, which contradicts (7) and (8). Hence E is σ_j -semistable. This completes our induction argument.

Step 3. Finally, by the previous two steps, we know that an object E is σ_1 -semistable if and only if σ_2 -semistable with $\phi_{\sigma_1}(E) = \phi_{\sigma_2}(E)$. Since every non-zero object in the heart is obtained by extensions of semistable objects with phases in $(0, 1]$, we know that $\mathcal{A}_1 = \mathcal{A}_2$. This ends the proof of our theorem. \square

A.1. Applications to Kuznetsov components of Fano threefolds. Let Y_d be smooth index 2 degree $d \geq 2$ prime Fano threefold and X_{4d+2} an index 1 degree $4d+2$ prime Fano threefold. In this section, we apply Theorem A.1 to show that all Serre-invariant stability conditions on $\mathcal{K}u(Y_d)$ and $\mathcal{K}u(X_{4d+2})$ (or $\mathcal{A}_{X_{4d+2}}$) are in the same $\widehat{\text{GL}}^+(2, \mathbb{R})$ -orbit for each $d \geq 2$ (Theorem A.10).

Recall that the Kuznetsov component of an index two prime Fano threefold Y_d of degree d is defined by $\mathcal{K}u(Y_d) := \langle \mathcal{O}_{Y_d}, \mathcal{O}_{Y_d}(H) \rangle^\perp$. The numerical Grothendieck group $\mathcal{N}(\mathcal{K}u(Y_d))$ is a rank two lattice generated by two classes

$$v = 1 - \frac{1}{d}H^2, \quad w = H - \frac{1}{2}H^2 + \left(\frac{1}{6} - \frac{1}{d}\right)H^2.$$

Moreover, under this basis, the Euler form is given by the matrix

$$\begin{pmatrix} -1 & -1 \\ 1-d & -d \end{pmatrix}.$$

For index one cases, we assume that $d \geq 2$. Then the Kuznetsov component is defined by $\mathcal{K}u(X_{4d+2}) := \langle \mathcal{E}_{X_{4d+2}}, \mathcal{O}_{X_{4d+2}} \rangle^\perp$, where $\mathcal{E}_{X_{4d+2}}$ is a certain exceptional bundle pulled back from a Grassmannian (cf. [Kuz04]).

By [BLMS23], $\sigma(\alpha, \beta)$ is a stability condition on $\mathcal{K}u(Y_d)$ and $\mathcal{K}u(X_{4d+2})$ for suitable (α, β) . Moreover, according to [PY22, PR23a], these stability conditions are all Serre-invariant.

Since for every index one prime Fano threefold X_{4d+2} with $d \geq 3$, there is an index two prime Fano threefold Y_d with $\mathcal{K}u(Y_d) \simeq \mathcal{K}u(X_{4d+2})$ by [Kuz04], hence we only need to consider Kuznetsov components of Y_d for $d \geq 2$ and X_{10} . Moreover, $\mathcal{K}u(Y_4)$ is equivalent to the derived category of a smooth curve, and $\mathcal{K}u(Y_5)$ is equivalent to the derived category of the 3-Kronecker quiver. In these two cases, the result is known by [Mac07] and [DK19]. So in the following, we mainly focus on $\mathcal{D} = \mathcal{K}u(Y_d)$ for $2 \leq d \leq 3$ or $\mathcal{K}u(X_{10})$. We first prove some properties of Serre-invariant stability conditions.

Lemma A.4. *Let $\mathcal{D} = \mathcal{K}u(Y_d)$ for $2 \leq d \leq 3$ or $\mathcal{K}u(X_{10})$ and $\sigma = (\mathcal{A}, Z)$ be a Serre-invariant stability condition on \mathcal{D} . Then \mathcal{D} satisfies (A) and (B) in Theorem A.1 and σ satisfies (C). Moreover, for any σ -semistable object $E \in \mathcal{D}$, we have :*

(1) *if $\mathcal{D} = \mathcal{K}u(Y_3)$, then*

$$\phi_\sigma(E) + 1 \leq \phi_\sigma(S_{\mathcal{D}}(E)) < \phi_\sigma(E) + 2,$$

(2) if $\mathcal{D} = \mathcal{K}u(Y_2)$ or $\mathcal{K}u(X_{10})$, then

$$\phi_\sigma(S_{\mathcal{D}}(E)) = \phi_\sigma(E) + 2.$$

Proof. It is clear that \mathcal{D} satisfies (A). And by [PY22, Lemma 5.16] and Proposition 8.7, \mathcal{D} also satisfies (B). When $\mathcal{D} = \mathcal{K}u(Y_3)$, by [PY22, Lemma 5.9] we have $\phi_\sigma(S_{\mathcal{D}}(E)) < \phi_\sigma(E) + 2$. When $\mathcal{D} = \mathcal{K}u(Y_2)$ or $\mathcal{K}u(X_{10})$, recall that $S_{\mathcal{D}}^2 \cong [4]$. Then the same argument as in [PY22, Lemma 5.9] shows that $\phi_\sigma(S_{\mathcal{D}}(E)) \leq \phi_\sigma(E) + 2$. Then for any two objects $A, B \in \mathcal{D}$ with $\phi_\sigma^+(B) < \phi_\sigma^-(A)$, using (1) and (2) we see

$$\phi_\sigma^-(A[2]) = \phi_\sigma^-(A) + 2 > \phi_\sigma^+(B) + 2 \geq \phi_\sigma^+(S_{\mathcal{D}}(B)).$$

Hence $\text{Hom}(B, A[2]) = \text{Hom}(A[2], S_{\mathcal{D}}(B)) = 0$ and the condition (C) is satisfied.

When $\mathcal{D} = \mathcal{K}u(Y_3)$, from [PY22, Lemma 5.11], we get $\text{ext}^1(E, E) \neq 0$, which implies $\phi_\sigma(E) + 1 \leq \phi_\sigma(S_{\mathcal{D}}(E))$. This proves (1).

When $\mathcal{D} = \mathcal{K}u(Y_2)$ or $\mathcal{K}u(X_{10})$, since \mathcal{D} satisfies (A) and (C), by Lemma A.3 we have $\text{ext}^1(E, E) \neq 0$, which implies $\phi_\sigma(E) + 1 \leq \phi_\sigma(S_{\mathcal{D}}(E))$. Now since $[E] = [S_{\mathcal{D}}(E)] \in \mathcal{N}(\mathcal{D})$, we have $\phi_\sigma(E) - \phi_\sigma(S_{\mathcal{D}}(E)) \in 2\mathbb{Z}$. Hence we get $\phi_\sigma(S_{\mathcal{D}}(E)) = \phi_\sigma(E) + 2$. □

Before verifying (D) and (E), we need several lemmas.

Lemma A.5. *Let X be a GM threefold.*

- (1) $\text{Hilb}_X^{3t+m} = \emptyset$ for $m < 1$. Thus for any conic $C \subset X$ and line $L \subset X$, we have $\text{Hom}(I_C, \mathcal{O}_L(-k)) = 0$ for any $k > 1$.
- (2) If a line L and a conic C satisfies $L \cap C \neq \emptyset$, then $L \cap C$ is of length one and $L \cup C$ is a twisted cubic.
- (3) Let $\mathcal{I} \subset \Gamma(X) \times \mathcal{C}(X)$ be the incidence variety, i.e.

$$\mathcal{I} = \{(L, C) : L \cap C \neq \emptyset\}.$$

Then the projection maps $\mathcal{I} \rightarrow \mathcal{C}(X)$ and $\mathcal{I} \rightarrow \Gamma(X)$ are surjective.

Proof. By [San14, Corollary 1.38], we have $\text{Hilb}_X^{3t+m} = \emptyset$ for $m < 0$. Thus to prove (1), we only need to show $\text{Hilb}_X^{3t} = \emptyset$. From [San14, Corollary 1.38], $\langle C \rangle \cong \mathbb{P}^2$ for any $[C] \in \text{Hilb}_X^{3t}$. Since X is an intersection of quadrics, such a C cannot exist on X . Hence $\text{Hilb}_X^{3t} = \emptyset$. Now note that the kernel of any non-zero map $I_C \rightarrow \mathcal{O}_L(-k)$ is the ideal sheaf of a closed subscheme with the Hilbert polynomial $3t + m$ for $m \leq 2 - k$. Therefore, $\text{Hom}(I_C, \mathcal{O}_L(-k)) = 0$ when $k > 1$. This proves (1). For (2), note that $\chi(\mathcal{O}_{L \cup C}) = 2 - \text{length}(L \cap C)$, then the result follows from (1).

Finally, we prove (3). Since $\dim \Gamma(X) = 1$, all lines on X sweep out a surface S in X . By $\text{Pic}(X) = \mathbb{Z} \mathcal{O}_X(H)$, we see $S \in |mH|$ for $m > 0$. Thus $C \cdot S \geq C \cdot H > 0$ for any conic C . In other words, $C \cap S \neq \emptyset$, hence any conic on X intersects with a line. Thus $\mathcal{I} \rightarrow \mathcal{C}(X)$ is surjective. Similarly, since X is covered by conics, any line intersects with a conic. Then $\mathcal{I} \rightarrow \Gamma(X)$ is surjective. □

Lemma A.6. *Let X be a GM threefold. Then there exists a line L and twisted cubics C and D on X such that*

- (1) $[L] \in \Gamma(X)$ is a smooth point,
- (2) $I_C \notin \mathcal{K}u(X)$ and $L \cup C = Z(s)$ for a section $s \in H^0(\mathcal{E}^\vee)$,
- (3) $L \subset D$, $I_D \in \mathcal{K}u(X)$ and $\text{ext}^1(I_D, I_D) = 3$.

Proof. Let $\mathcal{I} \subset \Gamma(X) \times \mathcal{C}(X)$ be the incidence variety. We denote by \mathcal{C}_1 the sublocus of $\mathcal{C}(X)$ parametrizing smooth conics Z such that their involutive conics are also smooth and $\text{hom}(\mathcal{E}, I_Z) = 1$. By Remark 7.17, \mathcal{C}_1 is an open subscheme of $\mathcal{C}(X)$. Let $\mathcal{I}_1 := \mathcal{I}|_{\Gamma(X) \times \mathcal{C}_1}$. From [Isk78, Theorem 3.4 (iii)] and [Ili94b, Section 3.1], $\Gamma(X)$ is generic smooth. This implies that the image of $p: \mathcal{I}_1 \rightarrow \Gamma(X)$ contains a smooth point.

Let $L \subset X$ be a line such that $[L] \in \Gamma(X)$ is smooth and contained in the image of p . Then $p^{-1}([L])$ is non-empty and there is a conic $[Z] \in \mathcal{C}_1$ such that $L \cap Z \neq \emptyset$. We set $D := L \cup Z$. And since $\text{Hom}(\mathcal{E}, I_L) \neq 0$, there is a section $s \in H^0(\mathcal{E})$ such that $L \subset Z(s)$. We define C to be the residue curve

of L in $Z(s)$. It is clear that C and D are twisted cubics by Lemma A.5. Moreover, L and Z intersect transversely at a single point. Then it remains to check $I_C \notin \mathcal{K}u(X)$, $\text{ext}^1(I_D, I_D) = 3$ and $I_D \in \mathcal{K}u(X)$.

Since $C \subset Z(s)$, it is clear that $\text{Hom}(\mathcal{E}, I_C) \neq 0$, i.e. $I_C \notin \mathcal{K}u(X)$. Moreover, by the construction, we have an exact sequence

$$0 \rightarrow I_D \rightarrow I_Z \rightarrow \mathcal{O}_L(-1) \rightarrow 0. \quad (9)$$

Note that all conics are connected, hence the smoothness implies irreducibility. Since $\text{Hom}(\mathcal{E}, I_Z) = k$ and the involutive conic Z' is smooth, we know that $Z \cup Z'$ only has two irreducible components which are both of degree 2, hence does not contain D . This means the unique non-zero map in $\text{Hom}(\mathcal{E}, I_Z) = k$ does not factor through I_D . Hence the induced map $\text{Hom}(\mathcal{E}, I_Z) \rightarrow \text{Hom}(\mathcal{E}, \mathcal{O}_L(-1))$ is injective. By $\text{RHom}^\bullet(\mathcal{E}, \mathcal{O}_L(-1)) = k$, this map is actually an isomorphism. Therefore, applying $\text{Hom}(\mathcal{E}, -)$ to (9), we obtain $\text{RHom}^\bullet(\mathcal{E}, I_D) = 0$, which implies $I_D \in \mathcal{K}u(X)$.

To show $\text{ext}^1(I_D, I_D) = 3$, since $\text{hom}(I_D, I_D) = 1$ and $\text{ext}^3(I_D, I_D) = 0$, by $\chi(I_D, I_D) = -2$ we only need to prove $\text{ext}^2(I_D, I_D) = 0$. From the construction above, we see $\text{ext}^2(I_Z, I_Z) = 0$. Moreover, $\text{ext}^2(\mathcal{O}_L(-1), \mathcal{O}_L(-1)) = 0$ since $[L] \in \Gamma(X)$ is a smooth point. And by the transversality of the intersection of L and Z , we see the derived restriction $\mathcal{O}_Z|_L \cong \mathcal{O}_{L \cap Z} \in D^b(L)$. Hence $\text{ext}^2(\mathcal{O}_Z, \mathcal{O}_L(-1)) = \text{ext}^1(I_Z, \mathcal{O}_L(-1)) = 0$. Finally, by Lemma A.5 we have $\text{hom}(I_Z, \mathcal{O}_L(-2)) = \text{ext}^3(\mathcal{O}_L(-1), I_Z) = 0$. Then $\text{ext}^2(I_D, I_D) = 0$ follows from applying [Pir23, Lemma 2.27] to (9). \square

Lemma A.7. *Let X be a GM threefold and L, C, D as in Lemma A.6. We define $F_1 := \text{pr}'(I_C)$, $F'_1 := I_D$ and $F_2 := \text{pr}'(I_L)$. Then the objects F_1, F'_1, F_2 are stable with respect to any Serre-invariant stability condition on $\mathcal{K}u(X)$. Moreover, F_1 and F'_1 have the same phase.*

Proof. From the construction, we see $\text{ext}^1(F'_1, F'_1) = 3$. By the same argument as in [Zha20, Corollary 5.4], we have $\text{ext}^1(F_1, F_1) = 3$. Finally, applying [Pir23, Lemma 2.27] to $\mathcal{E}^{\oplus 2} \rightarrow I_L \rightarrow F_2$ and using $\text{RHom}^\bullet(\mathcal{O}_L, \mathcal{O}_L) = k \oplus k[-1]$ implies $\text{ext}^1(F_2, F_2) = 3$. Then the stability of F_1, F'_1 and F_2 follows from Proposition 4.12.

As $[F_1] = [F'_1] \in \mathcal{N}(\mathcal{K}u(X))$, we have $\chi(F_1, F'_1) < 0$. Since $\text{ext}^i(F_1, F'_1) = \text{ext}^i(I_C, I_D) = 0$ for $i \notin \{1, 2\}$, we get $\text{Hom}(F_1, F'_1[1]) = \text{Hom}(F'_1[1], S_{\mathcal{K}u(X)}(F_1)) \neq 0$. Using Lemma A.4, we obtain $\phi_\sigma(F'_1) - 1 < \phi_\sigma(F_1) < \phi_\sigma(F'_1) + 1$ for any Serre-invariant stability condition σ . Thus $\phi_\sigma(F_1) = \phi_\sigma(F'_1)$ since $[F_1] = [F'_1] \in \mathcal{N}(\mathcal{K}u(X))$. \square

Lemma A.8. *Let $\mathcal{D} = \mathcal{K}u(Y_d)$ for $2 \leq d \leq 3$ or $\mathcal{K}u(X_{10})$. Then there exist two objects $F_1, F_2 \in \mathcal{D}$ such that for any Serre-invariant stability condition σ on \mathcal{D} , F_1 and F_2 are σ -stable with*

$$\phi_\sigma(F_2) - 1 < \phi_\sigma(F_1) < \phi_\sigma(F_2).$$

In particular, the image of the central charge is not contained in a line for any Serre-invariant stability condition on \mathcal{D} .

Proof. When $\mathcal{D} = \mathcal{K}u(Y_d)$, we define $F_2 := \text{RHom}(I_L, \mathcal{O}_{Y_d}(-H))[1]$ and $F_1 = I_L$, where $L \subset Y_d$ is a line. Then by [PY22, Lemma 5.13] and [PY22, Remark 4.8], F_1 and F_2 are σ -stable for any Serre-invariant stability condition σ on \mathcal{D} with $\phi_\sigma(F_2) - 1 < \phi_\sigma(F_1) < \phi_\sigma(F_2)$.

Now assume that $\mathcal{D} = \mathcal{K}u(X_{10})$. We take F_1, F'_1 and F_2 as in Lemma A.7. By [Zha20, Proposition 3.3, 5.3], we have $\text{pr}'(I_C) \cong \text{pr}'(G)$, where G fits into an exact triangle

$$\mathcal{O}_X(-H)[1] \rightarrow G \rightarrow \mathcal{O}_L(-2)$$

and is the twisted derived dual of the line L .

First, we prove that $\text{Hom}(F_2, F_1[1]) \neq 0$. By adjunction, we have $\text{Hom}(F_2, F_1[1]) = \text{Hom}(I_L, \text{pr}'(G)[1])$. And by [Zha20, Proposition 5.3], $\text{pr}'(G)$ fits into an exact triangle

$$G \rightarrow \text{pr}'(G) \rightarrow \mathcal{E}. \quad (10)$$

Since $\mathcal{E}|_L \cong \mathcal{O}_L \oplus \mathcal{O}_L(-1)$, it is easy to see $\text{Ext}^i(I_L, \mathcal{E})$ for $i \neq 2$. So applying $\text{Hom}(I_L, -)$ to (10), we get $\text{Hom}(I_L, \text{pr}'(G)[1]) = \text{Hom}(I_L, G[1]) = \text{Hom}(I_L, \mathcal{O}_L(-2)[1])$. As the normal bundle $N_{L/X_{10}}$ is either $\mathcal{O}_L \oplus \mathcal{O}_L(-1)$ or $\mathcal{O}_L(1) \oplus \mathcal{O}_L(-2)$ by [PS99, Lemma 4.2.1], we see the derived restriction $I_L|_L \cong$

$N_{L/X}^\vee \oplus \mathcal{O}_L(1)[1]$ from [Huy06, Proposition 11.8]. Then $\text{Hom}(I_L, \mathcal{O}_L(-2)[1]) \neq 0$ follows from a direct computation.

Next, we show that $\text{Hom}(F'_1, F_2) \neq 0$. By the definition of pr' , $\text{pr}'(I_L)$ fits into an exact triangle $\mathcal{E}^{\oplus 2} \rightarrow I_L \rightarrow \text{pr}'(I_L)$. Then applying $\text{Hom}(I_D, -)$ to this triangle, the result follows from $\text{Hom}(I_D, \mathcal{E}) = 0$ and $\text{Hom}(I_D, I_L) \neq 0$ since $L \subset D$.

By Lemma A.7, F_1, F'_1 and F_2 are all stable with respect to any Serre-invariant stability condition on \mathcal{D} . Therefore, combined with above results we get $\phi_\sigma(F_1) = \phi_\sigma(F'_1) < \phi_\sigma(F_2) < \phi_\sigma(F_1) + 1$ as desired. \square

Now we are ready to verify conditions (D) and (E) in Theorem A.1.

Lemma A.9. *Let $\mathcal{D} = \text{Ku}(Y_d)$ for $2 \leq d \leq 3$ or $\text{Ku}(X_{10})$. Then there exists a Serre-invariant stability condition σ_1 on \mathcal{D} with two σ_1 -stable objects D_1, D_2 satisfying (D) and (E). Moreover,*

- *we can assume that for any Serre-invariant stability condition σ on \mathcal{D} , D_1 and D_2 are σ -stable with*

$$\phi_\sigma(D_1) - 1 < \phi_\sigma(D_2) < \phi_\sigma(D_1), \text{ or } \phi_\sigma(D_1) < \phi_\sigma(D_2) < \phi_\sigma(D_1) + 1,$$

- *any Serre-invariant stability condition on \mathcal{D} with the same central charge as σ_1 satisfies (D) and (E).*

Proof. When $\mathcal{D} = \text{Ku}(Y_d)$, we define $D_1 := R\text{Hom}(I_L, \mathcal{O}_{Y_d}(-H))[1]$ and $D_2 = I_L[1]$, where $L \subset Y_d$ is a line. Then by [PY22, Lemma 5.13] and [PY22, Remark 4.8], D_1 and D_2 are σ -stable for any Serre-invariant stability condition σ on \mathcal{D} with $\phi_\sigma(D_1) < \phi_\sigma(D_2) < \phi_\sigma(D_1) + 1$. In this case, we take $\sigma_1 := \sigma(\alpha, -\frac{1}{2})$ for $\alpha > 0$ sufficiently small. Then by [PY22, Section 4], $D_1, D_2 \in \mathcal{A}(\alpha, -\frac{1}{2})$. Now a direct computation shows that, for any object E with $[E] = av + bw$, we have

- $\chi(E, D_2) = a + (d-1)b$, $\chi(D_2, E) = a + b$; and $\mu_{\alpha, -\frac{1}{2}}^0(E) > \mu_{\alpha, -\frac{1}{2}}^0(D_2) \iff b < 0$
- $\chi(E, D_1) = -b$, $\chi(D_1, E) = -[(d-2)a + (d-1)b]$; and $\mu_{\alpha, -\frac{1}{2}}^0(E) > \mu_{\alpha, -\frac{1}{2}}^0(D_1) \iff a + b < 0$.

Then it is straightforward to check (D) and (E) for σ_1 .

When $\mathcal{D} = \text{Ku}(X_{10})$, we use the equivalence Ξ in Lemma 3.7 and prove every thing on $\mathcal{A}_{X_{10}}$. Let $\sigma_1 := \sigma(\alpha, \beta)$, where $\beta < 0$ and $\alpha > 0$ with $-\beta$ and α are sufficiently small. We set $D_1 = I_C[1]$ and $D_2 = \text{pr}(F)[1]$, where $C \subset X$ is a smooth conic with $I_C \in \mathcal{A}_{X_{10}}$ and $F \in M_G(2, 1, 5)$ is non-locally free. It is clear that $D_1, D_2 \in \mathcal{A}(\alpha, \beta)$ and are stable with respect to any Serre-invariant stability condition on $\mathcal{A}_{X_{10}}$ by Lemma 7.5 and Proposition 8.7. As in the previous case, it is straightforward to check (D) and (E) for σ_1 . Now we show that for any Serre-invariant stability condition σ on $\mathcal{A}_{X_{10}}$, we have $\phi_\sigma(D_1) - 1 < \phi_\sigma(D_2) < \phi_\sigma(D_1)$. Indeed, if $\sigma = \sigma_1$, then this follows from a direct computation of the slope function of σ_1 . When $\sigma \neq \sigma_1$, by Lemma A.8, up to $\widetilde{\text{GL}}^+(2, \mathbb{R})$ -action, we can assume that σ and σ_1 has the same central charge and $\phi_\sigma(D_1) = \phi_{\sigma_1}(D_1)$. Thus $\phi_\sigma(D_2) - \phi_{\sigma_1}(D_2) \in 2\mathbb{Z}$. We claim that $\phi_\sigma(D_1) - 2 < \phi_\sigma(D_2) < \phi_\sigma(D_1)$, which implies $\phi_\sigma(D_2) = \phi_{\sigma_1}(D_2)$ and the result follows. Indeed, by Proposition 8.1 we have an exact sequence $0 \rightarrow F \rightarrow \mathcal{E} \rightarrow \mathcal{O}_L(-1) \rightarrow 0$ for a line $L \subset X_{10}$. Hence applying $\text{Hom}(-, D_1)$ to this exact sequence and use $\text{Hom}(\mathcal{E}, D_1[-1]) \neq 0$ (Lemma 6.3) and adjunction of pr , we have $\text{Hom}(F, I_C) = \text{Hom}(D_2, D_1) = \text{Hom}(D_1, S_{\mathcal{A}_{X_{10}}}(D_2)) \neq 0$. Then by Lemma A.4 we obtain $\phi_\sigma(D_1) - 2 < \phi_\sigma(D_2) < \phi_\sigma(D_1)$ as desired.

The final statement follows from the fact that (D) and (E) in this case only depend on the central charge and numerical classes $[D_1]$ and $[D_2]$, as we have seen above. \square

Applying Theorem A.1, we obtain the uniqueness of Serre-invariant stability conditions.

Theorem A.10. *Let $\mathcal{D} = \text{Ku}(Y_d)$ for $2 \leq d \leq 3$ or $\text{Ku}(X_{10})$. Then all Serre-invariant stability conditions on \mathcal{D} are in the same $\widetilde{\text{GL}}^+(2, \mathbb{R})$ -orbit.*

Proof. Let σ_1, D_1 and D_2 as in Lemma A.9. Let σ_2 be another Serre-invariant stability condition on \mathcal{D} . By Lemma A.8, up to $\widetilde{\text{GL}}^+(2, \mathbb{R})$ -action, we can assume that σ_1 and σ_2 have the same central charge. Moreover, up to shift we can assume that $\phi_{\sigma_1}(D_1) = \phi_{\sigma_2}(D_1)$. Thus $\phi_{\sigma_1}(D_2) - \phi_{\sigma_2}(D_2) \in 2\mathbb{Z}$. And by Lemma A.9, σ_2 also satisfies (D) and (E).

Now from Lemma A.9, we have $\phi_{\sigma_k}(D_1) - 1 < \phi_{\sigma_k}(D_2) < \phi_{\sigma_k}(D_1)$ or $\phi_{\sigma_k}(D_1) < \phi_{\sigma_k}(D_2) < \phi_{\sigma_k}(D_1) + 1$ for any $k \in \{1, 2\}$. This implies $\phi_{\sigma_1}(D_2) = \phi_{\sigma_2}(D_2)$. Therefore, by Lemma A.4 and Lemma A.9, we can apply Theorem A.1 and get $\sigma_1 = \sigma_2$. \square

Remark A.11. The idea of the proof of Theorem A.10 was first explained to us by Arend Bayer. In [Zha20, Proposition 4.21], one of the authors made an attempt to prove this statement but the argument is incomplete. Here, we fill the gaps and give a more general argument. Later, in [FP23, Theorem 3.1], the authors also prove the uniqueness of Serre-invariant stability conditions for a general triangulated category satisfying a list of assumptions and include Kuznetsov components of cubic threefolds and very general cubic fourfolds. The assumptions used in [FP23, Theorem 3.1] are (A), (B), and the Serre functor of \mathcal{D} satisfies $S_{\mathcal{D}}^r = [k]$ with $0 < k/r < 2$ or $r = 2$ and $k = 4$, while our Theorem A.1 also works for general triangulated categories which are not fractional Calabi–Yau but with extra assumptions (C), (D) and (E). Indeed, if we take $D_1 = D$ and $D_2 = S_{\mathcal{D}}(D)[-2]$ in (D) and (E) where D is an object in (B), then one can show that when $k/r < 2$, Theorem A.1 implies [FP23, Theorem 3.1]. Moreover, Theorem A.1 can be applied to the derived category of a smooth projective curve or a generalized Kronecker quiver as well.

REFERENCES

- [APR22] Matteo Altavilla, Marin Petković, and Franco Rota. Moduli spaces on the Kuznetsov component of Fano threefolds of index 2. *Épjournal Géom. Algébrique*, 6:Art. 13, 31, 2022.
- [BB17] Arend Bayer and Tom Bridgeland. Derived automorphism groups of K3 surfaces of Picard rank 1. *Duke Math. J.*, 166(1):75–124, 2017.
- [BBF⁺20] Arend Bayer, Sjoerd Beentjes, Soheyla Feyzbakhsh, Georg Hein, Diletta Martinelli, Fatemeh Rezaee, and Benjamin Schmidt. The desingularization of the theta divisor of a cubic threefold as a moduli space. *arXiv preprint, arXiv:2011.12240*, 2020.
- [BF14] Maria Chiara Brambilla and Daniele Faenzi. Vector bundles on Fano threefolds of genus 7 and Brill–Noether loci. *Internat. J. Math.*, 25(3):1450023, 59, 2014.
- [BLMS23] Arend Bayer, Martí Lahoz, Emanuele Macrì, and Paolo Stellari. Stability conditions on Kuznetsov components. *Ann. Sci. Éc. Norm. Supér. (4)*, 56(2):517–570, 2023. With an appendix by Bayer, Lahoz, Macrì, Stellari and X. Zhao.
- [BMK23] Arend Bayer, Emanuele Macrì, and Alexander Kuznetsov. Vector bundles Fano threefolds. *preprint*, 2023.
- [BMMS12] Marcello Bernardara, Emanuele Macrì, Sukhendu Mehrotra, and Paolo Stellari. A categorical invariant for cubic threefolds. *Adv. Math.*, 229(2):770–803, 2012.
- [BMS16] Arend Bayer, Emanuele Macrì, and Paolo Stellari. The space of stability conditions on abelian threefolds, and on some Calabi–Yau threefolds. *Invent. Math.*, 206(3):869–933, 2016.
- [BMT14] Arend Bayer, Emanuele Macrì, and Yukinobu Toda. Bridgeland stability conditions on threefolds I: Bogomolov–Gieseker type inequalities. *J. Algebraic Geom.*, 23(1):117–163, 2014.
- [BO01] Alexei Bondal and Dmitri Orlov. Reconstruction of a variety from the derived category and groups of autoequivalences. *Compositio Math.*, 125(3):327–344, 2001.
- [Bri07] Tom Bridgeland. Stability conditions on triangulated categories. *Ann. of Math. (2)*, 166(2):317–345, 2007.
- [BT16] Marcello Bernardara and Gonçalo Tabuada. From semi-orthogonal decompositions to polarized intermediate Jacobians via Jacobians of noncommutative motives. *Mosc. Math. J.*, 16(2):205–235, 2016.
- [DIM12] Olivier Debarre, Atanas Iliev, and Laurent Manivel. On the period map for prime Fano threefolds of degree 10. *J. Algebraic Geom.*, 21(1):21–59, 2012.
- [DK18] Olivier Debarre and Alexander Kuznetsov. Gushel–Mukai varieties: classification and birationalities. *Algebr. Geom.*, 5(1):15–76, 2018.
- [DK19] George Dimitrov and Ludmil Katzarkov. Bridgeland stability conditions on wild Kronecker quivers. *Adv. Math.*, 352:27–55, 2019.
- [FP23] Soheyla Feyzbakhsh and Laura Pertusi. Serre-invariant stability conditions and Ulrich bundles on cubic threefolds. *Épjournal Géom. Algébrique*, 7:Art. 1, 32, 2023.
- [FV21] Daniele Faenzi and Alessandro Verra. Fano threefolds of genus 10 and Coble cubics. *In preparation*, 2021.

- [GLZ22] Hanfei Guo, Zhiyu Liu, and Shizhuo Zhang. Conics on Gushel–Mukai fourfolds, EPW sextics and Bridgeland moduli spaces. *arXiv preprint, arXiv:2203.05442*, 2022.
- [HLOY03] Shinobu Hosono, Bong H. Lian, Keiji Oguiso, and Shing-Tung Yau. Fourier–Mukai partners of a $K3$ surface of Picard number one. In *Vector bundles and representation theory (Columbia, MO, 2002)*, volume 322 of *Contemp. Math.*, pages 43–55. Amer. Math. Soc., Providence, RI, 2003.
- [HRS96] Dieter Happel, Idun Reiten, and Sverre O. Smalø. Tilting in abelian categories and quasitilted algebras. *Mem. Amer. Math. Soc.*, 120(575):viii+ 88, 1996.
- [Huy06] Daniel Huybrechts. *Fourier–Mukai transforms in algebraic geometry*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, Oxford, 2006.
- [Ili94a] Atanas Iliev. The Fano surface of the Gushel threefold. *Compositio Math.*, 94(1):81–107, 1994.
- [Ili94b] Atanas Iliev. Lines on the Gushel’ threefold. *Indag. Math. (N.S.)*, 5(3):307–320, 1994.
- [IM11] Atanas Iliev and Laurent Manivel. Fano manifolds of degree ten and EPW sextics. *Ann. Sci. Éc. Norm. Supér. (4)*, 44(3):393–426, 2011.
- [Isk78] V. A. Iskovskih. Fano threefolds. II. *Izv. Akad. Nauk SSSR Ser. Mat.*, 42(3):506–549, 1978.
- [JLZ22] Augustinas Jacovskis, Zhiyu Liu, and Shizhuo Zhang. Brill–Noether theory and categorical Torelli theorems for Kuznetsov components of index 1 Fano threefolds. *arXiv preprint, arXiv:2207.01021*, 2022.
- [Kos20] Naoki Koseki. On the Bogomolov–Gieseker inequality for hypersurfaces in the projective spaces. *arXiv preprint, arXiv:2008.09799*, 2020.
- [KP17] Alexander Kuznetsov and Alexander Perry. Derived categories of cyclic covers and their branch divisors. *Selecta Math. (N.S.)*, 23(1):389–423, 2017.
- [KP18] Alexander Kuznetsov and Alexander Perry. Derived categories of Gushel–Mukai varieties. *Compos. Math.*, 154(7):1362–1406, 2018.
- [KP23] Alexander Kuznetsov and Alexander Perry. Categorical cones and quadratic homological projective duality. *Ann. Sci. Éc. Norm. Supér. (4)*, 56(1):1–57, 2023.
- [KPS18] Alexander G. Kuznetsov, Yuri G. Prokhorov, and Constantin A. Shramov. Hilbert schemes of lines and conics and automorphism groups of Fano threefolds. *Jpn. J. Math.*, 13(1):109–185, 2018.
- [Kuz04] Alexander Kuznetsov. Derived category of a cubic threefold and the variety V_{14} . *Tr. Mat. Inst. Steklova*, 246:183–207, 2004.
- [Kuz09] Alexander Kuznetsov. Derived categories of Fano threefolds. *Tr. Mat. Inst. Steklova*, 264:116–128, 2009.
- [Kuz10] Alexander Kuznetsov. Derived categories of cubic fourfolds. In *Cohomological and geometric approaches to rationality problems*, volume 282 of *Progr. Math.*, pages 219–243. Birkhäuser Boston, Boston, MA, 2010.
- [Kuz19] Alexander Kuznetsov. Calabi–Yau and fractional Calabi–Yau categories. *J. Reine Angew. Math.*, 753:239–267, 2019.
- [Li19] Chunyi Li. Stability conditions on Fano threefolds of Picard number 1. *J. Eur. Math. Soc. (JEMS)*, 21(3):709–726, 2019.
- [Li21] Chunyi Li. Stronger Bogomolov–Gieseker inequality for Fano varieties of Picard number one. *In preparation*, 2021.
- [LNSZ21] Chunyi Li, Howard Nuer, Paolo Stellari, and Xiaolei Zhao. A refined derived Torelli theorem for Enriques surfaces. *Math. Ann.*, 379(3-4):1475–1505, 2021.
- [Log12] Dmitry Logachev. Fano threefolds of genus 6. *Asian J. Math.*, 16(3):515–559, 2012.
- [LSZ22] Chunyi Li, Paolo Stellari, and Xiaolei Zhao. A refined derived Torelli theorem for Enriques surfaces, II: the non-generic case. *Math. Z.*, 300(4):3527–3550, 2022.
- [LZ22] Zhiyu Liu and Shizhuo Zhang. A note on Bridgeland moduli spaces and moduli spaces of sheaves on X_{14} and Y_3 . *Math. Z.*, 302(2):803–837, 2022.
- [Mac07] Emanuele Macrì. Stability conditions on curves. *Math. Res. Lett.*, 14(4):657–672, 2007.
- [Muk92] Shigeru Mukai. Fano 3-folds. In *Complex projective geometry (Trieste, 1989/Bergen, 1989)*, volume 179 of *London Math. Soc. Lecture Note Ser.*, pages 255–263. Cambridge Univ. Press, Cambridge, 1992.
- [Ogu02] Keiji Oguiso. $K3$ surfaces via almost-primes. *Math. Res. Lett.*, 9(1):47–63, 2002.
- [Orl97] D. O. Orlov. Equivalences of derived categories and $K3$ surfaces. *J. Math. Sci. (New York)*, 84(5):1361–1381, 1997. Algebraic geometry, 7.

- [Per22] Alexander Perry. The integral Hodge conjecture for two-dimensional Calabi–Yau categories. *Compos. Math.*, 158(2):287–333, 2022.
- [Pir23] Dmitrii Pirozhkov. Admissible subcategories of del Pezzo surfaces. *Adv. Math.*, 424:Paper No. 109046, 62, 2023.
- [PR23a] Laura Pertusi and Ethan Robinett. Stability conditions on Kuznetsov components of Gushel–Mukai threefolds and Serre functor. *Math. Nachr.*, 296(7):2975–3002, 2023.
- [PR23b] Marin Petković and Franco Rota. A note on the Kuznetsov component of the Veronese double cone. *J. Pure Appl. Algebra*, 227(3):Paper No. 107214, 23, 2023.
- [PS99] A. N. Parshin and I. R. Shafarevich, editors. *Algebraic geometry. V*, volume 47 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 1999. Fano varieties, A translation of *Algebraic geometry. 5* (Russian), Ross. Akad. Nauk, Vseross. Inst. Nauchn. i Tekhn. Inform., Moscow.
- [PS23] Laura Pertusi and Paolo Stellari. Categorical Torelli theorems: results and open problems. *Rend. Circ. Mat. Palermo (2)*, 72(5):2949–3011, 2023.
- [PY22] Laura Pertusi and Song Yang. Some remarks on Fano threefolds of index two and stability conditions. *Int. Math. Res. Not. IMRN*, (17):12940–12983, 2022.
- [San14] Giangiacomo Sanna. Rational curves and instantons on the Fano threefold Y_5 . *arXiv preprint, arXiv:1411.7994*, 2014.
- [Zha20] Shizhuo Zhang. Bridgeland moduli spaces and Kuznetsov’s Fano threefold conjecture. *arXiv preprint, arXiv:2012.12193*, 2020.

DEPARTMENT OF MATHEMATICS, MAISON DU NOMBRE, 6 AVENUE DE LA FONTE, L-4364 ESCH-SUR-ALZETTE, LUXEMBOURG
Email address: `augustinas.jacovskis@uni.lu`

MAX PLANCK INSTITUTE FOR MATHEMATICS, VIVATSGASSE 7, 53111 BONN, GERMANY
Email address: `xlin@mpim-bonn.mpg.de`, `lin-x18@tsinghua.org.cn`

INSTITUTE FOR ADVANCED STUDY IN MATHEMATICS, ZHEJIANG UNIVERSITY, HANGZHOU, ZHEJIANG PROVINCE 310030, P. R. CHINA
Email address: `jasonlzy0617@gmail.com`

MAX PLANCK INSTITUTE FOR MATHEMATICS, VIVATSGASSE 7, 53111 BONN, GERMANY

INSTITUT DE MATHÉMATIQUES DE TOULOUSE, UMR 5219, UNIVERSITÉ DE TOULOUSE, UNIVERSITÉ PAUL SABATIER, 118 ROUTE DE NARBONNE, 31062 TOULOUSE CEDEX 9, FRANCE
Email address: `shizhuozhang@mpim-bonn.mpg.de`, `shizhuo.zhang@math.univ-toulouse.fr`