

PAPER • OPEN ACCESS

Topological theory of phase transitions

To cite this article: Matteo Gori *et al* 2022 *J. Phys. A: Math. Theor.* **55** 375002

View the [article online](#) for updates and enhancements.

You may also like

- [Optical kerr effect measurements on supercooled water: The experimental perspective](#)
P Bartolini, A Taschin, R Eramo et al.
- [The LHCf detector at the CERN Large Hadron Collider](#)
The LHCf Collaboration, O Adriani, L Bonechi et al.
- [Effective potential, Mori product and quantum dynamics](#)
A Cuccoli, R Giachetti, V Tognetti et al.

Topological theory of phase transitions

Matteo Gori^{1,2}, Roberto Franzosi^{3,4,5},
Giulio Pettini^{6,9} and Marco Pettini^{7,8,*} 

¹ Quantum Biology Lab, Howard University, 2400 6th St NW, Washington, DC 20059, United States of America

² Department of Physics and Sciences of Materials, University of Luxembourg, Luxembourg

³ DSFTA, University of Siena, Via Roma 56, 53100 Siena, Italy

⁴ INFN Sezione di Perugia, I-06123 Perugia, Italy

⁵ QSTAR, CNR—Istituto Nazionale di Ottica, largo E. Fermi 6, 50125 Firenze, Italy

⁶ Dipartimento di Fisica, Università di Firenze, I.N.F.N., Sezione di Firenze, via G. Sansone 1, I-50019 Sesto Fiorentino, Italy

⁷ Aix-Marseille Univ, Université de Toulon, CNRS, Marseille, France

⁸ CNRS, Centre de Physique Théorique, UMR7332, 13288 Marseille, France

⁹ Sezione di Firenze, INFN, 50019 Sesto Fiorentino, Italy

E-mail: gori6matteo@gmail.com, roberto.franzosi@ino.it, pettini@fi.unifi.it and pettini@cpt.univ-mrs.fr

Received 5 December 2021, revised 27 June 2022

Accepted for publication 6 July 2022

Published 17 August 2022



CrossMark

Abstract

The investigation of the Hamiltonian dynamical counterpart of phase transitions, combined with the Riemannian geometrization of Hamiltonian dynamics, has led to a preliminary formulation of a differential-topological theory of phase transitions. In fact, in correspondence of a phase transition there are peculiar geometrical changes of the mechanical manifolds that are found to stem from changes of their topology. These findings, together with two theorems, have suggested that a topological theory of phase transitions can be formulated to go beyond the limits of the existing theories. Among other advantages, the new theory applies to phase transitions in small N systems (that is, at nanoscopic and mesoscopic scales), and in the absence of symmetry-breaking. However, the preliminary version of the theory was incomplete and still falsifiable by counterexamples. The present work provides a relevant leap forward leading to an accomplished development of the topological theory of phase transitions paving the way to further developments and applications of the theory that can be no longer hampered.

* Author to whom any correspondence should be addressed.



Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](https://creativecommons.org/licenses/by/4.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Keywords: phase transitions, topological methods, microcanonical ensemble

(Some figures may appear in colour only in the online journal)

1. Introduction

Phase transitions phenomena are ubiquitous in nature at very different scales in space and in energy. Therefore, from the theoretical viewpoint, understanding their origin, and the way of classifying them, is of central interest. In spite of a huge literature on this topic, a general theory is still lacking. In the framework of Landau's phenomenological theory, phase transitions are generally related to the mechanism of spontaneous symmetry breaking. However, Landau's theory is not all-encompassing. Indeed, many systems do not fit in this theory and undergo a phase transition lacking spontaneous symmetry breaking and lacking an order parameter. Some notable examples are: Kosterlitz–Thouless transitions after Mermin–Wagner theorem, systems with local gauge symmetries after Elitzur's theorem, liquid–gas transitions, transitions in supercooled glasses and liquids, transitions in amorphous and disordered systems, folding transitions in homopolymers and proteins. Furthermore, to account for the loss of analyticity of thermodynamic observables, the mathematical description of phase transitions requires the limit of an infinite number of particles (thermodynamic limit) as is the case of the Yang–Lee theory [1] and of the Dobrushin–Lanford–Ruelle theory [2]. However, the contemporary research on nanoscopic and mesoscopic systems, on the biophysics of polymers [4, 5], on Bose–Einstein condensation, Dicke superradiance in microlaser, superconducting transitions in small metallic objects, tackles transition phenomena in systems of finite—and often very small—number of particles.

Within all the hitherto developed theoretical frameworks, also including the monumental theory of renormalization group and critical phenomena, it is assumed that the primitive object at the grounds of a theory is a given statistical measure; schematically: the grand-canonical measure in the old Yang–Lee theory, the canonical measure in the Dobrushin–Lanford–Ruelle theory, and the microcanonical measure in a still somewhat open and more recent approach [3–5]. However, there are several general results suggesting that the possibility for a system to undergo a phase transition depends on some measure-independent properties, as is its spatial dimension, the dimensionality of its order parameter, the range of its interactions, the symmetry group (discrete or continuous) of its Hamiltonian. This hints at the possibility that the same information might be encoded already at a more fundamental level completely determined by the internal interactions of a system, interactions described by their potential function.

Therefore, looking for generalisations of the existing theories is a well motivated and timely purpose. The present paper puts forward a new starting point for a line of thought initiated several years ago and based on a variety of results which hitherto did not appear to fit in a coherent theoretical framework. The central idea of this line of thought is that the singular energy dependence of the thermodynamic observables at a phase transition is the 'shadow' of some adequate *change of topology* of the energy level sets in phase space (or of the potential level sets in configuration space, as well).

1.1. Why topology

Recently, the study of equilibrium phase transitions in the microcanonical ensemble has attracted increasing interest, being very important in presence of ensemble inequivalence, when only the microcanonical ensemble gives the correct results. Two complementary approaches

have been undertaken. One of these is of a statistical kind [3, 4], recently summarized in a very interesting, powerful and rich classification of microcanonical phase transitions by Bachmann in reference [5]. On another side, as the ergodic invariant measure of nonintegrable Hamiltonian systems is the microcanonical measure, the other approach resorts to the study of Hamiltonian dynamics of systems undergoing phase transitions. This dynamical approach brings about interesting novelties with respect to the standard studies of phase transitions, eventually leading to the topological hypothesis (TH) through the following logical chain. The dynamics of a generic system of N degrees of freedom described by a Hamiltonian $H = \frac{1}{2} \sum_{i=1}^N p_i^2 + V(q_1, \dots, q_N)$, or equivalently by the corresponding Lagrangian function $L = \frac{1}{2} \sum_{i=1}^N \dot{q}_i^2 - V(q_1, \dots, q_N)$, is chaotic. The degree of chaoticity of the dynamics is measured by the largest Lyapunov exponent, a new observable that has been proven useful to characterize phase transitions from a dynamical viewpoint [6]. Then, the explanation of the origin of Hamiltonian chaos—encompassing the computation of the largest Lyapunov exponent—proceeds by identifying a Hamiltonian flow with a geodesic flow of an appropriate Riemannian differentiable manifold. This differential geometric framework is given by configuration space endowed with the non-Euclidean metric of components [7] $g_{ij} = 2[E - V(q)]\delta_{ij}$, whence the infinitesimal arc element $ds^2 = 2[E - V(q)]^2 dq_i dq^i$; then Newton equations are retrieved from the geodesic equations

$$\frac{d^2 q^i}{ds^2} + \Gamma_{jk}^i \frac{dq^j}{ds} \frac{dq^k}{ds} = 0,$$

where Γ_{jk}^i are the Christoffel connection coefficients of the manifold. The degree of instability of the dynamics is described by means of the Jacobi–Levi-Civita equation for the geodesic spread

$$\frac{\nabla^2 J}{ds^2} + R(J, \dot{\gamma})\dot{\gamma} = 0,$$

where the vector field J locally measures the distance between nearby geodesics, $\frac{\nabla}{ds}$ is the covariant derivative along the configuration space geodesic $\dot{\gamma}$, and $R(\cdot, \cdot)$ is the Riemann curvature tensor. The largest Lyapunov exponent for high dimensional Hamiltonian flows is found to depend on the curvature ‘landscape’ of the configuration space manifold [6]. Hence, a natural consequence has been to investigate whether the occurrence of phase transitions has some peculiar counterpart in geometrical changes of the manifolds underlying the flows. And it has been discovered that this is actually the case. Moreover, the peculiar geometrical changes associated with phase transitions were discovered to be the effects of deeper topological changes of the potential level sets $\Sigma_v^{V_N} := \{V_N(q_1, \dots, q_N) = v \in \mathbb{R}\}$ in configurations space, and, equivalently, of the balls $\{M_v^{V_N} = V_N^{-1}((-\infty, v])\}_{v \in \mathbb{R}}$ bounded by the $\Sigma_v^{V_N}$.

A topological approach to the study of phase transitions has been considered for a variety of systems, ranging from those undergoing entropy driven transitions [8, 9] (having also applications to robotics), and hard spheres systems [10], to quantum phase transitions [11–13], glasses and supercooled liquids [14, 15], classical models in statistical mechanics [16–18], discrete spin models [19], DNA denaturation [20], peptide structure [21], to quote just a few of them. In fact, in many contexts, well before an explicit formulation of the TH [22, 23], topological concepts were implicitly entering the study of phase transitions while talking of energy

landscapes [24, 25] and of saddle points for disordered systems, glasses [14, 15], spin glasses: saddle points being critical points in the language of Morse theory of differential topology.

On a completely different field, more recently, handling big data—outsourcing from complex systems—through methods referred to as topological data analysis (TDA) it happens to highlight the existence of phase transition-like phenomena in the absence of a statistical measure. Here the concept of phase transition is intended as the emergence of qualitatively new properties when a control parameter crosses a critical value (the prototype can be dated back to the Erdős–Renyi giant component appearance in random graphs). To quote a fascinating example in this field, in reference [26] the discovery is reported of topological phase transitions in functional brain networks by merging concepts from TDA, topology, geometry, physics, and network theory.

The present paper is organized as follows: in section 2 the basic definitions and concepts of extrinsic geometry of hypersurfaces is recalled for the sake of self-containedness, and the definition of asymptotic diffeomorphicity is also therein introduced. In section 3 the main theorem is formulated and proved; this theorem states that a topological change of the potential level sets of a physical system is a necessary condition for the appearance of a phase transition. Finally, in section 4 the problem raised by the counterexample to a preceding formulation of the main theorem is fixed. Section 5 addresses some past controversial points. Section 6 is devoted to some concluding remarks. Two appendices contain computational details.

2. Topological origin of phase transitions

On the one side the study of the Hamiltonian dynamical counterpart of phase transitions, combined with the geometrization of Hamiltonian dynamics, has led to find out the crucial role of topology at the grounds of these transition phenomena, on the other side a mathematical relationship exists between macroscopic thermodynamics and topological properties of the manifolds M_v^{VN} , as expressed by [6]

$$\begin{aligned}
 S_N(v) &= (k_B/N) \log \left[\int_{M_v^{VN}} d^N q \right] \\
 &= \frac{k_B}{N} \log \left[\text{vol} \left[M_v^{VN} \setminus \bigcup_{i=1}^{\mathcal{N}(v)} \Gamma(x_c^{(i)}) \right] + \sum_{i=0}^N w_i \mu_i(M_v^{VN}) + \mathcal{R} \right], \quad (1)
 \end{aligned}$$

where S_N is the configurational entropy, v is the potential energy, and the $\mu_i(M_v^{VN})$ are the Morse indexes (in one-to-one correspondence with topology) of the manifolds M_v^{VN} ; in square brackets: the first term is the result of the excision of certain neighborhoods of the critical points of the interaction potential from M_v^{VN} ; the second term is a weighed sum of the Morse indexes, and the third term is a smooth function of N and v .

As a consequence, major topology changes with v of the submanifolds M_v^{VN} —bringing about sharp changes of the potential energy pattern of at least some of the $\mu_i(M_v^{VN})$ —can affect the v -dependence of $S_N(v)$ and of its derivatives (figure 1).




Figure 1. Low-dimensional pictorial representation of the transition between complex topologies as a metaphor of the origin of a phase transition. From the ground level up to the crossover level v_c , the manifolds M_v have a genus which increases with v . Above the crossover level v_c , the manifolds M_v have also a nonvanishing linking number which increases with v .

Hence, it has been surmised [6] that, at least for a broad class of physical systems, phase transitions stem from a suitable change of the topology of the potential level sets $\Sigma_v^{V_N}$ and, equivalently, of the manifolds $M_v^{V_N}$, when v , playing the role of the control parameter, takes a critical value v_c . This hypothesis has turned into the start of a new theory by putting together several studies on specific models [6, 27] and two theorems [28–30]. These theorems state that an equilibrium phase transition—is necessarily due to appropriate topological transitions in configuration space. However, a counterexample to these theorems has been found in reference [31] thus undermining this version of the topological theory of phase transitions. The counterexample is provided by the second order phase transition of the 2D lattice ϕ^4 -model that occurs at a critical value v_c of the potential energy density which belongs to a broad interval of v -values void of critical points of the potential function. The difficulty raised by this counterexample has stimulated a deeper investigation of the transition of the ϕ^4 -model which led to figure out a crucial point associated with the breaking of the \mathbb{Z}_2 symmetry (and possibly with the breaking of discrete symmetries in general), that is, the possibility of a phase transition to stem from an asymptotic loss of diffeomorphicity of the relevant manifolds [32]. In what follows this fact is formalized into a new and more consistent version of the theory. Other alleged difficulties of the theory are briefly discussed in Section 5.

2.1. Basic definitions and concepts

Consider now an open set of v -values $I \subseteq \mathbb{R}$ such that the cylindrical subset of configuration space $\Gamma_I^N = \bigcup_{v \in I} \Sigma_v^{V_N}$ contains only non-singular level sets, that is, $\nabla V_N(q) \neq 0$ for any $q \in \Gamma_I^N$, meaning that V_N has no critical points for any $v \in I$.

For any $v_0, v_1 \in I$, the two level sets $\Sigma_{v_0}^{V_N} \subset \Gamma_I^N$ and $\Sigma_{v_1}^{V_N} \subset \Gamma_I^N$ are diffeomorphic under the action of an explicitly known diffeomorphism given by the integral lines of the vector field $\xi_N = \nabla V_N / \|\nabla V_N\|^2$, that is, any initial condition $q_\alpha \in \Sigma_{v_0}^{V_N}$ is diffeomorphically mapped onto




Figure 2. Pictorial representation of the action of the vector field ξ in equation (2) diffeomorphically mapping each point of the level set Σ_a onto each point of the level set Σ_b . Pictorial representation of the action of the vector field ξ in equation (2) diffeomorphically mapping each point of the level set Σ_a onto each point of the level set Σ_b .

a point $q_\beta \in \Sigma_{v_1}^{V_N}$ by the equation [33] (figure 2)

$$\frac{dq}{dv} = \frac{\nabla V_N}{\|\nabla V_N\|^2}. \tag{2}$$

2.2. Extrinsic geometry of hypersurfaces

In this section some basic definitions and concepts are given about the extrinsic geometry of hypersurfaces of a Euclidean space. The basic tool consists in measuring the way of changing from point to point on the surface of the normal direction in order to describe how the n -surface Σ curves around in \mathbb{R}^N . The rate of change of the normal vector \mathcal{N} at a point $x \in \Sigma$ in a given direction \mathbf{u} is described by the *shape operator* (also known as Weingarten’s map) $L_x(\mathbf{u}) = -\nabla_{\mathbf{u}} \mathcal{N} = -(\mathbf{u} \cdot \nabla) \mathcal{N}$, where \mathbf{u} is a tangent vector at x and $\nabla_{\mathbf{u}}$ is the directional derivative; gradients and vectors are represented in \mathbb{R}^N .

The constant-energy hypersurfaces in the phase space of Hamiltonian systems or of the equipotential hypersurfaces in configuration space, are the level sets of regular functions and, for the level sets defined via a regular real-valued function f as $\Sigma_a := f^{-1}(a)$, the normal vector is $\mathcal{N} = \nabla f / \|\nabla f\|$. Let $\{\mathbf{e}_\mu\}_{\mu=1,\dots,N} = \{\mathbf{e}_1, \dots, \mathbf{e}_n, \mathcal{N}\}$, with $\mathbf{e}_\alpha \cdot \mathbf{e}_\beta = \delta_{\alpha,\beta}$ and denote with Greek subscripts, $\alpha = 1, \dots, N$, the components in the embedding space \mathbb{R}^N , and with Latin subscripts, $i = 1, \dots, n$, the components on a generic tangent space $T_x \Sigma_a$ at $x \in \Sigma_a$. We consider the case of codimension one, that is, $N = n + 1$ (figure 3).

From $\partial_\mu \mathcal{N}_\alpha \mathcal{N}_\alpha = 0 = 2\mathcal{N}_\alpha \partial_\mu \mathcal{N}_\alpha$ we see that for any \mathbf{u} , we have $\mathcal{N} \cdot L_x(\mathbf{u}) = -\mathcal{N}_\alpha \mathbf{u}_\mu \partial_\mu \mathcal{N}_\alpha = 0$, which means that $L_x(\mathbf{u})$ projects on the tangent space $T_x \Sigma_a$.

Now the *principal curvatures* $\kappa_1, \dots, \kappa_n$ of Σ_a at x are the eigenvalues of the shape operator restricted to $T_x \Sigma_a$. Considering the matrix \mathcal{L}_x to be the restriction of L_x to $T_x \Sigma_a$

$$\mathcal{L}_{ij}(x) = \mathbf{e}_i \cdot L_x(\mathbf{e}_j) = -(\mathbf{e}_i)_\alpha (\mathbf{e}_j)_\beta \partial_\beta \mathcal{N}_\alpha,$$

then the *mean curvature* is defined as

$$H(x) = \frac{1}{n} \text{Tr}^{(n)} \mathcal{L}_{ij}(x) = \frac{1}{n} \sum_{i=1}^n \kappa_i. \tag{3}$$




Figure 3. Illustration of the items entering the construction of the shape operator of a surface.

The computation of the analytic expression of the mean curvature H proceeds from

$$H(x) = \frac{1}{n} \text{Tr}^{(n)} \mathcal{L}_{ij}(x) = -\frac{1}{n} \sum_{i=1}^n (\mathbf{e}_i)_\alpha (\mathbf{e}_i)_\beta \partial_\beta \mathcal{N}_\alpha. \tag{4}$$

Defining $A_{\mu\nu} = (\mathbf{e}_\mu)_\nu$, so that $AA^T = \mathbb{I}$, we have

$$\sum_{i=1}^n (\mathbf{e}_i)_\alpha (\mathbf{e}_i)_\beta = \delta_{\alpha\beta} - \mathcal{N}_\alpha \mathcal{N}_\beta$$

and thus

$$H(x) = -\frac{1}{n} (\delta_{\alpha\beta} - \mathcal{N}_\alpha \mathcal{N}_\beta) \partial_\beta \mathcal{N}_\alpha = -\frac{1}{n} \partial_\alpha \mathcal{N}_\alpha = -\frac{1}{n} \nabla \cdot \left(\frac{\nabla f}{\|\nabla f\|} \right). \tag{5}$$

2.3. Asymptotic diffeomorphicity

In reference [32] it has been numerically found that the phase transition undergone by the 2D lattice ϕ^4 model actually corresponds to a major topological change of the potential level sets of the model, also in absence of critical points of the potential. This topological change corresponds to an asymptotic breaking of the topological transitivity of the potential level sets, what can be formalised as an asymptotic loss of diffeomorphicity of the same manifolds in the broken symmetry phase. Hence a crucial hint to fix the problem stemming from the counterexample given by the ϕ^4 model that has been hitherto considered fatal.

The first step to fix the problem thus consists in defining asymptotic diffeomorphicity, what is easily done by observing that a vector valued function of several variables, $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$, is of differentiability class C^l if all the partial derivatives $(\partial^l f / \partial x_{i_1}^{l_1} \dots \partial x_{i_k}^{l_k})$ exist and are continuous, where each of i_1, \dots, i_k is an integer between 1 and n and each l_1, \dots, l_k is an integer between 0 and l , and $l_1 + \dots + l_k = l$. Then, by taking advantage of the explicit analytic representation of the vector field generating the diffeomorphism $\xi_N : \Gamma_I^N \rightarrow T\Gamma_I^N$ previously given, uniform convergence in N of the sequence $\{\xi_N\}_{N \in \mathbb{N}}$ —and thus asymptotic diffeomorphicity in some class C^l —can be defined after the introduction of an appropriate norm containing all the derivatives up to $(\partial^l \xi_N / \partial q_{i_1}^{l_1} \dots \partial q_{i_k}^{l_k})$. At any fixed $N \in \mathbb{N}$, in the absence of critical points of a confining potential V_N , the level sets $\Sigma_v^{V_N}$ are non singular $(N - 1)$ -dimensional hypersurfaces in \mathbb{R}^N . Let us consider the already defined cylindrical subset of configuration space $\Gamma_I^N = \bigcup_{v \in I} \Sigma_v^{V_N}$ containing only non-singular level sets.

The lack of asymptotic breaking of diffeomorphicity is defined by introducing a norm for the ξ_N that allows to compare the diffeomorphisms at different dimensions

$$\|\xi_N\|_{C^k(\Gamma_{I_0}^N)} = \sup_{q_0 \in \Gamma_{I_0}^N} \|\xi_N\| + \sum_{l=1}^k \sum_{\{i_k\}} \sum_{j=1}^N \|\nabla_{\{i_k\}}^l \xi_j\|_{\Gamma_{I_0}^N}, \tag{6}$$

where $\{i_k\}$ stands for a multi-index and $\|\nabla_{\{i_k\}}^l \xi_j\|_{\Gamma_{I_0}^N}$ is the norm of the l th differential operator with $l_1 + \dots + l_k = l$

$$\|\nabla_{\{i_k\}}^l \xi_j\|_{\Gamma_{I_0}^N} = \sup_{q_0 \in \Gamma_{I_0}^N} \left| \frac{\partial^l \xi_j}{\partial q_{i_1}^{l_1} \dots \partial q_{i_k}^{l_k}} \right|. \tag{7}$$

The sequence of families of manifolds $\{\Gamma_{I_0}^N\}_{N \in \mathbb{N}}$ is said to asymptotically preserve the C^k -diffeomorphicity among the hypersurfaces $\Sigma_v^{V_N}$ —foliating each family—if there exists $B \in \mathbb{R}^+$ such that

$$\|\xi_N\|_{C^k(\Gamma_{I_0}^N)} \leq B < +\infty \quad \forall N \in \mathbb{N}. \tag{8}$$

As a consequence, from this condition we get $\|\nabla V_N\| = \|\xi_N\|^{-1} \geq 1/B = C > 0$ for each $q_0 \in \Gamma_{I_0}^N$ and all $N \in \mathbb{N}$, ruling out the existence of asymptotic critical points (i.e. $\|\nabla V_N\| \rightarrow 0$ for $N \rightarrow \infty$).

The analytic condition (8) entails remarkable consequences on the extrinsic geometry of the potential level sets. In fact, using $\sum_i \|X_i\| \geq \|\sum_i X_i\|$, from equation (7) at the lowest order with the aid of a normalised vector u tangent at q_0 to a $\Sigma_v^n \subset \Gamma_{I_0}^N$, that is, $u \in T_{q_0} \Sigma_v^n$, we can build the quadratic forms

$$\sum_{i,j=1}^N \|(\partial_i \xi_j) u_i u_j\| \geq \left\| \sum_{i,j=1}^N \left(\partial_i \frac{\partial_j V_N}{\|\nabla V_N\|^2} \right) u_i u_j \right\|, \tag{9}$$

where $\partial_i = \partial / \partial q^i$. With implicit summation on repeated indices the rhs of equation (9) is rewritten as

$$\begin{aligned} & \left\| \left[\frac{1}{\|\nabla V_N\|} \left(\partial_i \frac{\partial_j V_N}{\|\nabla V_N\|} \right) + \frac{\partial_j V_N}{\|\nabla V_N\|} \partial_i \left(\frac{1}{\|\nabla V_N\|} \right) \right] u_i u_j \right\| \\ & = \left\| \frac{1}{\|\nabla V_N\|} \left(\partial_i \frac{\partial_j V_N}{\|\nabla V_N\|} \right) u_i u_j \right\|, \end{aligned} \tag{10}$$

where the orthogonality, at any point q_0 , between the vectors \mathbf{u} and $\mathcal{N} = (\partial_1 V_N / \|\nabla V_N\|, \dots, \partial_N V_N / \|\nabla V_N\|)$ tangent and normal to $\Sigma_v^{V_N}$, respectively, has been used. Through the shape operator (Weingarten map) of $\Sigma_v^{V_N}$ [34] at q_0

$$L_{q_0}(\mathbf{u}) = -L_u \mathcal{N} = -(\nabla \mathcal{N}_1 \cdot \mathbf{u}, \dots, \nabla \mathcal{N}_N \cdot \mathbf{u}) \tag{11}$$

the quadratic form $\kappa(\mathbf{u}, q_0) = \langle \mathbf{u}, L_{q_0}(\mathbf{u}) \rangle$ is found to coincide with the one given in equation (10) (last term). The quantity $\kappa(\mathbf{u}, q_0)$ is known as the *normal curvature* of the level set $\Sigma_v^{V_N}$ at q_0 . Let $\{\kappa_1(q_0), \dots, \kappa_N(q_0)\}$ denote the principal curvatures of $\Sigma_v^{V_N}$ at q_0 , with the corresponding orthogonal principal curvature directions $\{\mathbf{v}_1, \dots, \mathbf{v}_N\}$, then the normal curvature in the direction $\mathbf{u} \in T_{q_0} \Sigma_v^{V_N}$ is given by

$$\kappa(\mathbf{u}, q_0) = \sum_{i=1}^N \kappa_i(q_0) \langle \mathbf{u}, \mathbf{v}_i \rangle = \sum_{i=1}^N \kappa_i(q_0) \cos^2 \theta_i. \tag{12}$$

By choosing $\tilde{\mathbf{u}} \in T_{q_0} \Sigma_v^{V_N}$ such that $\|\tilde{\mathbf{u}}\| = 1$ and all the angles θ_i between $\tilde{\mathbf{u}}$ and \mathbf{v}_i are equal to some $\tilde{\theta}$, we get

$$\kappa(\tilde{\mathbf{u}}, q_0) = (\cos^2 \tilde{\theta}) \sum_{i=1}^N \kappa_i(q_0) = (\cos^2 \tilde{\theta}) N H(q_0), \tag{13}$$

where $H(q_0)$ is the mean curvature (the trace of the Weingarten map) at q_0 . Thus from equations (8), (10) and (13)

$$\left\| \frac{1}{\|\nabla V_N\|} (\cos^2 \tilde{\theta}) N H(q) \right\| \leq B < +\infty \quad \forall N \in \mathbb{N} \tag{14}$$

everywhere on $\Sigma_v^{V_N}$. Since $\|\nabla V_N\| \sim \mathcal{O}(N^{1/2})$ it follows that $H(q) \sim \mathcal{O}(N^{-1/2})$ everywhere on $\Sigma_v^{V_N}$ and uniformly in N . Therefore, the first remarkable consequence of asymptotic diffeomorphicity among the potential level sets is that their mean curvature

$$H(q) = \frac{1}{N} \sum_{i=1}^N \kappa_i(q) \tag{15}$$

is everywhere uniformly bounded in N . However, this does not ensure the boundedness of each principal curvature (whose sign is not definite). *A priori* two or more principal curvatures of the same value but of opposite sign could diverge and mutually compensate leaving $H(q)$ finite. In order to get this missing information about the asymptotic boundedness of all the principal curvatures, let us consider the scalar curvature \mathcal{R} of a level set $V(q) = v$, embedded in an Euclidean space of arbitrary dimension, which reads [35]

$$\begin{aligned} \mathcal{R}(q) &= \frac{1}{N(N-1)} \sum_{i \leq j}^{1 \dots N} \kappa_i(q) \kappa_j(q) \\ &= \frac{1}{N(N-1)} \left\{ -\Delta \log \|\nabla V_N(q)\| + \nabla \cdot \left[\Delta V_N(q) \frac{\nabla V_N(q)}{\|\nabla V_N(q)\|^2} \right] \right\} \end{aligned} \tag{16}$$

let us notice that \mathcal{R} is singular at the critical points of the potential, where $\nabla V_N(q) = 0$, and can be arbitrarily large in their neighborhoods; then, using $\|\xi\| = \|\nabla V_N(q)\|^{-1}$, this can be




Figure 4. Sequence of diffeomorphic manifolds (of the same dimension) with a limit manifold which is not diffeomorphic to the members of the sequence. The infinitely tiny bridge between the two spheres of S_∞ has infinite mean curvature.

rewritten as

$$\mathcal{R} = \frac{1}{N(N-1)} \left\{ -\Delta \log \frac{1}{\|\xi\|} + \nabla \cdot [\Delta V_N(q) \xi] \right\}, \tag{17}$$

then, trivial computations (sketched in appendix A) of the rhs of this equation under the assumption of asymptotic diffeomorphicity (equations (6)–(8)) yield uniform boundedness also of $\mathcal{R}(q)$ entailing uniform boundedness in N of each $\kappa_i(q)$ everywhere on each potential level set.

To help intuition to get a hold of the relationship between boundedness of mean and scalar curvatures and asymptotic diffeomorphicity, we qualitatively illustrate in figure 4 the opposite situation, known as Gromov–Hausdorff limit [36], where a sequence of diffeomorphic manifolds of fixed dimension have a limit manifold which is not diffeomorphic to the other members of the sequence. The handles of these dumbbell shaped manifolds shrink to an asymptotic infinitely tiny cylinder of vanishing radius and thus of diverging transversal principal curvature, that is, of divergent mean curvature.

Remark 1. Summarizing, the assumption of asymptotic diffeomorphicity means that, for any pair of densities \bar{v} and \bar{v}' in some assigned interval $I_{\bar{v}} = [\bar{v}_0, \bar{v}_1]$ and N arbitrarily large, the corresponding manifolds $\Sigma_{N\bar{v}}^{V_N}$ are diffeomorphic under the action of the diffeomorphism-generating vector fields ξ_{N_k}

$$\begin{aligned} \Sigma_{N_1\bar{v}}^{V_{N_1}} &\xrightarrow{\xi_{N_1}} \Sigma_{N_1\bar{v}'}^{V_{N_1}} \\ \Sigma_{N_2\bar{v}}^{V_{N_2}} &\xrightarrow{\xi_{N_2}} \Sigma_{N_2\bar{v}'}^{V_{N_2}} \\ &\vdots \qquad \bar{v}, \bar{v}' \in [\bar{v}_0, \bar{v}_1], \quad k \in \mathbb{N} \\ \Sigma_{N_k\bar{v}}^{V_{N_k}} &\xrightarrow{\xi_{N_k}} \Sigma_{N_k\bar{v}'}^{V_{N_k}} \\ &\vdots \end{aligned} \tag{18}$$

provided that the norm of the vector fields ξ_{N_k} is uniformly bounded according to equation (6). Under this condition, all the principal curvatures $\kappa_i(q)$ of every manifold in the above diagram are uniformly bounded with N . Moreover, after the non-critical neck theorem [37] all the above manifolds $\Sigma_{N_k\bar{v}}^{V_{N_k}}$ for any $\bar{v} \in [\bar{v}_0, \bar{v}_1]$ are free of critical points of the potential functions V_N , that is of points where $\nabla V_N = 0$.

3. A necessity theorem

In its original formulation, given in references [28, 29], the theorem establishing the necessary topological origin of a phase transition was lacking a fundamental hypothesis that has led to the paradoxical situation of being falsified [31] through the example of a phase transition still related with a change of topology in configuration space, though asymptotic in the number of degrees of freedom [32], and in the absence of critical points of the potential.

The missing hypothesis suggested by the study of reference [32] is to require also asymptotic diffeomorphicity of the potential level sets in order to correspondingly get uniform convergence of the Helmholtz free energy in a differentiability class that rules out first and second order phase transitions.

Remark 2. The notation $N \in \mathbb{N}^\#$ means that $N \rightarrow \infty$ is included.

3.1. Main theorem

Theorem 1 (Absence of phase transitions under diffeomorphicity). *Let $V_N(q_1, \dots, q_N) : \mathbb{R}^N \rightarrow \mathbb{R}$, be a smooth, nonsingular, finite-range potential. Denote by $\Sigma_v^{V_N} := V_N^{-1}(v)$, $v \in \mathbb{R}$, its level sets, or equipotential hypersurfaces, in configuration space.*

Then let $\bar{v} = v/N$ be the potential energy per degree of freedom.

If for any pair of values \bar{v} and \bar{v}' belonging to a given interval $I_{\bar{v}} = [\bar{v}_0, \bar{v}_1]$ and for any $N > N_0$ with $N \in \mathbb{N}^\#$ we have

$$\Sigma_{N\bar{v}}^{V_N} \approx \Sigma_{N\bar{v}'}^{V_N},$$

that is, $\Sigma_{N\bar{v}}^{V_N}$ is diffeomorphic to $\Sigma_{N\bar{v}'}^{V_N}$, including asymptotically diffeomorphic, then the sequence of the Helmholtz free energies $\{F_N(\beta)\}_{N \in \mathbb{N}}$ —where $\beta = 1/T$ (T is the temperature) and $\beta \in I_\beta = (\beta(\bar{v}_0), \beta(\bar{v}_1))$ —is uniformly convergent at least in $\mathcal{C}^2(I_\beta \subset \mathbb{R})$, so that $F_\infty \in \mathcal{C}^2(I_\beta \subset \mathbb{R})$ and neither first- nor second-order phase transitions can occur in the (inverse) temperature interval $(\beta(\bar{v}_0), \beta(\bar{v}_1))$.

Remark 3. The configurational entropy $S_N(\bar{v})$ is related to the configurational canonical free energy, f_N in (20), for any $N \in \mathbb{N}$, $\bar{v} \in \mathbb{R}$, and $\beta \in \mathbb{R}$ through the Legendre transform

$$-f_N(\beta) = \beta \cdot \bar{v}_N - S_N(\bar{v}_N), \tag{19}$$

where the inverse of the configurational temperature $T(v)$ is given by $\beta_N(\bar{v}) = \partial S_N(\bar{v}) / \partial \bar{v}$. By following reference [57], let us consider the function $\phi(\bar{v}) = f_N[\beta(\bar{v})]$, from $\phi'(\bar{v}) = -\bar{v}$ [$d\beta_N(\bar{v})/d\bar{v}$] it is evident that if $\beta_N(\bar{v}) \in \mathcal{C}^k(\mathbb{R})$ then also $\phi(\bar{v}) \in \mathcal{C}^k(\mathbb{R})$ and thus $S_N(\bar{v}) \in \mathcal{C}^{k+1}(\mathbb{R})$ while $f_N(\beta) \in \mathcal{C}^k(\mathbb{R})$. First and second order phase transitions are associated with a discontinuity in the first or second derivatives of $f_\infty(\beta)$, that is with $f_\infty(\beta) \in \mathcal{C}^0(\mathbb{R})$ or $f_\infty(\beta) \in \mathcal{C}^1(\mathbb{R})$, respectively. Hence a first order phase transition corresponds to a discontinuity of the second derivative of the entropy $S_\infty(\bar{v})$, and a second order phase transition corresponds to a discontinuity of the third derivative of the entropy $S_\infty(\bar{v})$.

Remark 4. The proof of the main theorem follows the same conceptual path given in references [28, 29]: a *topological change* of the equipotential hypersurfaces $\Sigma_v^{V_N}$ of configuration space is a *necessary* condition for the occurrence of a thermodynamic phase transition if we prove the *equivalent proposition* that if any two hypersurfaces $\Sigma_v^{V_N}$ and $\Sigma_{v'}^{V_N}$ with $v(N), v'(N) \in (v_0(N), v_1(N))$ are *diffeomorphic* for all $N \in \mathbb{N}^\#$, then *no phase transition* can occur in the (inverse) temperature interval $[\lim_{N \rightarrow \infty} \beta(\bar{v}_0(N)), \lim_{N \rightarrow \infty} \beta(\bar{v}_1(N))]$.

Proof. For standard Hamiltonian systems (i.e. quadratic in the momenta) the relevant information is carried by the configurational microcanonical ensemble, where the configurational canonical free energy is

$$f_N(\beta) \equiv f_N(\beta; V_N) = \frac{1}{N} \log \int_{(\Lambda^d)^{\times n}} dq_1 \dots dq_N \exp[-\beta V_N(q_1, \dots, q_N)] \quad (20)$$

with and the configurational microcanonical entropy (in units s.t. $k_B = 1$) is

$$S_N(\bar{v}) \equiv S_N(\bar{v}; V_N) = \frac{1}{N} \log \int_{(\Lambda^d)^{\times n}} dq_1 \dots dq_N \delta[V_N(q_1, \dots, q_N) - v].$$

Then $S_N(\bar{v})$ is related to the configurational canonical free energy, f_N , for any $N \in \mathbb{N}$, $\bar{v} \in \mathbb{R}$, and $\beta \in \mathbb{R}$ through the Legendre transform in equation (19).

From lemma 1, proved after lemmas 2 to 9, we have that in the limit $N \rightarrow \infty$ and at constant particle density, $\text{vol}(\Lambda^d)^{\times n}/N = \text{const}$, in the interval $I_{\bar{v}} = [\bar{v}_0, \bar{v}_1]$ the sequence $\{S_N\}_{N \in \mathbb{N}^\#}$ is uniformly convergent in $\mathcal{C}^3(I_{\bar{v}} \subset \mathbb{R})$ so that $S_\infty \in \mathcal{C}^3(I_{\bar{v}} \subset \mathbb{R})$ that is, the thermodynamic limit of the entropy is three times differentiable, with continuous third-order derivative, in $I_{\bar{v}} = [\bar{v}_0, \bar{v}_1]$. Hence in the interval $I_\beta = [\lim_{N \rightarrow \infty} \beta(\bar{v}_0(N)), \lim_{N \rightarrow \infty} \beta(\bar{v}_1(N))]$ the sequence of configurational free energies $\{f_N(T)\}_{N \in \mathbb{N}^\#}$ is *uniformly convergent* at least in $\mathcal{C}^2(I_\beta \subset \mathbb{R})$, so that we have

$$-f_\infty(\beta) = \beta(\bar{v}) \cdot \bar{v} - S_\infty(\bar{v})$$

that is $\{f_\infty(T)\} \in \mathcal{C}^2(I_\beta \subset \mathbb{R})$.

Since a quadratic kinetic energy term of a standard Hamiltonian gives only a smooth contribution to the total Helmholtz free energy $F_N(\beta)$, also the asymptotic function $F_\infty(\beta)$ has differentiability class $\mathcal{C}^2(I_\beta \subset \mathbb{R})$ so that we conclude that the corresponding physical system does not undergo neither first- nor second-order phase transitions in the inverse-temperature interval $\beta \in I_\beta$. \square

3.2. Lemmas

Lemma 1 (Uniform upper bounds). *Let V_N be a standard, short-range, stable, and confining potential function bounded below. Let $\{\Sigma_v^{V_N}\}_{v \in \mathbb{R}}$ be the family of $(N - 1)$ -dimensional equipotential hypersurfaces $\Sigma_v^{V_N} := V_N^{-1}(v)$, $v \in \mathbb{R}$, of \mathbb{R}^N . If*

$$\forall N \in \mathbb{N}^\# \quad \text{and} \quad \bar{v}, \bar{v}' \in I_{\bar{v}} = [\bar{v}_0, \bar{v}_1], \quad \text{we have} \quad \Sigma_{N\bar{v}}^{V_N} \approx \Sigma_{N\bar{v}'}^{V_N},$$

then

$$\sup_{N, \bar{v} \in I_{\bar{v}}} |S_N(\bar{v})| < \infty \quad \text{and} \quad \sup_{N, \bar{v} \in I_{\bar{v}}} \left| \frac{\partial^k S_N}{\partial \bar{v}^k}(\bar{v}) \right| < \infty, \quad k = 1, 2, 3, 4.$$

Proof. The proof of this lemma amounts to proving the main theorem and proceeds as follows. After remark 2, the derivatives of the entropy are expressed in terms of the derivatives of the microcanonical configurational volume which, in turn, after lemma 2 can be expressed as surface integrals of functions of $\bar{\zeta}_N = \text{div}(\bar{\xi}_N)$ and its Lie derivatives, where $\bar{\xi}_N$ is the vector field generating the diffeomorphisms among the specific potential energy level sets. Then these integrals are replaced by averages along Monte Carlo Markov chains (MCMCs) that can be defined to have as invariant measure the microcanonical configurational measure (lemma 3 and remark 3). After lemmas 4 and 5, $\bar{\zeta}_N$ is proved to behave as a random Gaussian process

along the mentioned MCMCs, hence, after remark 5 and lemmas 6 to 9 the uniform bounds are derived of the derivatives of the entropy up to the fourth one. \square

Lemma 2 (Derivation of integrals over regular level sets ([38, 39])). *Let $O \subset \mathbb{R}^p$ be a bounded open set. Let $\psi \in \mathcal{C}^{n+1}(\overline{O})$ be constant on each connected component of the boundary ∂O and $f \in \mathcal{C}^n(O)$. Define $O_{t,t'} = \{x \in O \mid t < \psi(x) < t'\}$ and*

$$F(v) = \int_{\{\psi=v\}} f \, d\sigma^{p-1}, \tag{21}$$

where $d\sigma^{p-1}$ represents the Lebesgue measure of dimension $p - 1$. If $C > 0$ exists such that for any $x \in O_{t,t'}$, $\|\nabla\psi(x)\| \geq C$, then for any k such that $0 \leq k \leq n$, for any $v \in]t, t'[$, one has

$$\frac{d^k F}{dv^k}(v) = \int_{\{\psi=v\}} A_\psi^k f \, d\sigma^{p-1}. \tag{22}$$

with

$$A_\psi f = \nabla \left(\frac{\nabla\psi}{\|\nabla\psi\|} f \right) \frac{1}{\|\nabla\psi\|}. \tag{23}$$

This lemma allows to compute higher order derivatives of the microcanonical volume $\Omega_n(\bar{v})$, and thus of the entropy, at any order by identifying ψ with the potential $\bar{V}_N(\mathbf{q}) = V(\mathbf{q})/N$. Let us introduce the following notations: $\bar{\xi}_N = \text{div}(\bar{\xi}_N)$,

$$\bar{\chi}_N = \|\bar{\xi}_N\| = \frac{1}{\|\nabla\bar{V}_N\|}, \tag{24}$$

for the norm of $\bar{\xi}_N$, and

$$d\mu_{\bar{v}}^{N-1} = \bar{\chi}_N \, d\sigma_{\Sigma_{\bar{v}}^{\bar{V}_N}} \tag{25}$$

for the microcanonical area $(N - 1)$ -form of non critical energy level sets, and

$$\mathcal{L}_{\bar{\xi}}(\cdot) = (\bar{\xi} \cdot \nabla)(\cdot) = \sum_{i=1}^N \frac{\partial^i \bar{V}}{\|\nabla\bar{V}\|^2} \partial_i(\cdot) \tag{26}$$

for the Lie derivative along the flow of $\bar{\xi}_N$. Then, given the microcanonical configurational volume

$$\Omega_N(\bar{v}) = \int_{\Sigma_{\bar{v}}^{\bar{V}_N}} d\mu_{\bar{v}}^{N-1} \tag{27}$$

its derivatives are computed through the formula

$$\frac{d^k \Omega_N}{d\bar{v}^k}(\bar{v}) = \int_{\Sigma_{\bar{v}}^{\bar{V}_N}} \frac{1}{\bar{\chi}} A_V^k(\bar{\chi}) \, d\mu_{\bar{v}}^{N-1}, \tag{28}$$

where $A_V^k(\bar{\chi})$ stands for a k -times repeated application of the operator

$$A_V(f) = f \bar{\xi}_N + \mathcal{L}_{\bar{\xi}_N}(f). \tag{29}$$

Remark 5 (Derivatives of the entropy). The configurational microcanonical entropy density is given by

$$\bar{S}_N(\bar{v}) = \frac{1}{N} \log \Omega_N(\bar{v}) = \frac{1}{N} \log \int_{\Sigma_{\bar{V}_N}^{\bar{v}}} d\mu_{\bar{v}}^{N-1} \tag{30}$$

and its derivatives are

$$\begin{aligned} \frac{d\bar{S}_N}{d\bar{v}}(\bar{v}) &= \frac{1}{N} \frac{\Omega'_N(\bar{v})}{\Omega_N(\bar{v})} \\ \frac{d^2\bar{S}_N}{d\bar{v}^2}(\bar{v}) &= \frac{1}{N} \left[\frac{\Omega''_N(\bar{v})}{\Omega_N(\bar{v})} - \left(\frac{\Omega'_N(\bar{v})}{\Omega_N(\bar{v})} \right)^2 \right] \\ \frac{d^3\bar{S}_N}{d\bar{v}^3}(\bar{v}) &= \frac{1}{N} \left[\frac{\Omega'''_N(\bar{v})}{\Omega_N(\bar{v})} - 3 \frac{\Omega''_N(\bar{v}) \Omega'_N(\bar{v})}{\Omega_N(\bar{v})^2} + 2 \left(\frac{\Omega'_N(\bar{v})}{\Omega_N(\bar{v})} \right)^3 \right] \\ \frac{d^4\bar{S}_N}{d\bar{v}^4}(\bar{v}) &= \frac{1}{N} \left[\frac{\Omega^{(iv)}_N(\bar{v})}{\Omega_N(\bar{v})} - 4 \frac{\Omega'''_N(\bar{v}) \Omega'_N(\bar{v})}{\Omega_N(\bar{v})^2} + 12 \frac{\Omega''_N(\bar{v}) \Omega'_N(\bar{v})^2}{\Omega_N(\bar{v})^3} \right. \\ &\quad \left. - 3 \left(\frac{\Omega''_N(\bar{v})}{\Omega_N(\bar{v})} \right)^2 - 6 \left(\frac{\Omega'_N(\bar{v})}{\Omega_N(\bar{v})} \right)^4 \right], \end{aligned} \tag{31}$$

where, after lemma 2, the derivatives of configurational microcanonical volume $\Omega_N(\bar{v})$ up to the fourth order with respect to \bar{v} are found to be

$$\begin{aligned} \frac{d\Omega_N}{d\bar{v}}(\bar{v}) &= \int_{\Sigma_{\bar{V}_N}^{\bar{v}}} \bar{\zeta}_N d\mu_{\bar{v}}^{N-1} \\ \frac{d^2\Omega_N}{d\bar{v}^2}(\bar{v}) &= \int_{\Sigma_{\bar{V}_N}^{\bar{v}}} \left[\bar{\zeta}_N^2 + \mathcal{L}_{\bar{\xi}_N}(\bar{\zeta}_N) \right] d\mu_{\bar{v}}^{N-1} \\ \frac{d^3\Omega_N}{d\bar{v}^3}(\bar{v}) &= \int_{\Sigma_{\bar{V}_N}^{\bar{v}}} \left[\bar{\zeta}_N^3 + 3\bar{\zeta}_N \mathcal{L}_{\bar{\xi}_N}(\bar{\zeta}_N) + \mathcal{L}_{\bar{\xi}_N}^{(ii)}(\bar{\zeta}_N) \right] d\mu_{\bar{v}}^{N-1} \\ \frac{d^4\Omega_N}{d\bar{v}^4}(\bar{v}) &= \int_{\Sigma_{\bar{V}_N}^{\bar{v}}} \left[\bar{\zeta}_N^4 + 6\bar{\zeta}_N^2 \mathcal{L}_{\bar{\xi}_N}(\bar{\zeta}_N) + 4\bar{\zeta}_N \mathcal{L}_{\bar{\xi}_N}^{(ii)}(\bar{\zeta}_N) \right. \\ &\quad \left. + 3 \left(\mathcal{L}_{\bar{\xi}_N}(\bar{\zeta}_N) \right)^2 + \mathcal{L}_{\bar{\xi}_N}^{(iii)}(\bar{\zeta}_N) \right] d\mu_{\bar{v}}^{N-1}. \end{aligned} \tag{32}$$

On any $(N - 1)$ -dimensional hypersurface $\Sigma_{N\bar{v}}^V = V_N^{-1}(N\bar{v}) = \{X \in \mathbb{R}^N \mid V_N(X) = N\bar{v}\}$ of \mathbb{R}^N , we can define a homogeneous nonperiodic random Markov chain whose probability measure is the configurational microcanonical measure [6], namely $d\sigma/\|\nabla V_N\|$. We call this Markov chain a microcanonical-MCMC. In so doing, all the integrals giving configurational microcanonical averages are replaced by asymptotic averages along these MCMCs. Dropping the suffix N of V_N we have the following lemma:

Lemma 3 (MCMCs over regular level sets). *On each finite-dimensional level set $\Sigma_{N\bar{v}} = V^{-1}(N\bar{v})$ of a standard, smooth, confining, short-range potential V bounded below, and*

in the absence of critical points, there exists a random Markov chain of points $\{X_i \in \mathbb{R}^N\}_{i \in \mathbb{N}_+}$, constrained by the condition $V(X_i) = N\bar{v}$, which has

$$d\mu = \frac{d\sigma}{\|\nabla V\|} \left(\int_{\Sigma_{N\bar{v}}} \frac{d\sigma}{\|\nabla V\|} \right)^{-1} \tag{33}$$

as its probability measure, so that for a smooth function $F : \mathbb{R}^N \rightarrow \mathbb{R}$ we have

$$\left(\int_{\Sigma_{N\bar{v}}} \frac{d\sigma}{\|\nabla V\|} \right)^{-1} \int_{\Sigma_{N\bar{v}}} \frac{d\sigma}{\|\nabla V\|} F = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n F(X_i). \tag{34}$$

Proof. The level sets $\{\Sigma_{N\bar{v}}\}_{\bar{v} \in \mathbb{R}}$ are compact hypersurfaces of \mathbb{R}^N , therefore a partition of unity [34] can be defined on each hypersurface. Then, let $\{U_i\}$, $1 \leq i \leq m$ be an arbitrary finite covering of $\Sigma_{N\bar{v}}$ by means of domains of coordinates (for example open balls), at any point of $\Sigma_{N\bar{v}}$ an ensemble of smooth functions $\{\varphi_i\}$ exists, such that $1 \geq \varphi_i \geq 0$ and $\sum_i \varphi_i = 1$.

By means of the partition of unity $\{\varphi_i\}$ on $\Sigma_{N\bar{v}}$, associated to a collection $\{U_i\}$ of one-to-one local parametrizations of the compact and oriented hypersurfaces $\Sigma_{N\bar{v}}$, the integral of a given smooth $(N - 1)$ -form ω is given by:

$$\int_{\Sigma_{N\bar{v}}} \omega^{(N-1)} = \int_{\Sigma_{N\bar{v}}} \left(\sum_{i=1}^m \varphi_i(x) \right) \omega^{(N-1)}(x) = \sum_{i=1}^m \int_{U_i} \varphi_i \omega^{(N-1)}(x).$$

The existence of a MCMC of assigned probability measure (33) on a given $\Sigma_{N\bar{v}}$ is constructively proved as follows. Let us consider sequences of random values $\{x_i : i \in \Lambda\}$, where Λ is the finite set of indexes of the elements of the partition of unity on $\Sigma_{N\bar{v}}$, and where $x_i = (x_i^1, \dots, x_i^{N-1})$ are local coordinates with respect to U_i of an arbitrary representative point of the set U_i itself. The weight $\pi(i)$ of the i th element of the partition is then defined by

$$\pi(i) = \left(\sum_{k=1}^m \int_{U_k} \varphi_k \frac{d\sigma}{\|\nabla V\|} \right)^{-1} \int_{U_i} \varphi_i \frac{d\sigma}{\|\nabla V\|} \tag{35}$$

and the transition matrix elements [40] are given by

$$p_{ij} = \min \left[1, \frac{\pi(j)}{\pi(i)} \right] \tag{36}$$

satisfying the detailed balance equation $\pi(i)p_{ij} = \pi(j)p_{ji}$. A random Markov chain $\{i_0, i_1, \dots, i_k, \dots\}$ of indexes induces a random Markov chain of corresponding elements of the partition, that is of points $\{x_{i_0}, x_{i_1}, \dots, x_{i_k}, \dots\}$ on the hypersurface $\Sigma_{N\bar{v}}$. Denote by $(x_p^1, \dots, x_p^{N-1})$ the local coordinates of a point P on $\Sigma_{N\bar{v}}$ and define a local reference frame as $\{\partial/\partial x_p^1, \dots, \partial/\partial x_p^{N-1}, n(P)\}$, with $n(P)$ the outward unit normal vector at P ; by means of the matrix that operates a point-dependent change from this reference frame to the canonical basis $\{e_1, \dots, e_N\}$ of \mathbb{R}^N it is possible to associate to the Markov chain $\{x_{i_0}, x_{i_1}, \dots, x_{i_k}, \dots\}$ an equivalent chain $\{X_{i_0}, X_{i_1}, \dots, X_{i_k}, \dots\}$ of points specified through their coordinates in \mathbb{R}^N but still constrained to belong to the subset $V(X) = v$, that is, to $\Sigma_{N\bar{v}}$. Consequently, the invariant probability measure [40] of the Markov chain so constructed is the probability density (33). Moreover, in the absence of critical points, for smooth functions F and smooth potentials V , the variation on each set U_i of $F/\|\nabla V\|$ is limited. Therefore, by keeping it finite the partition of unity can be refined as needed to make Lebesgue integration convergent; hence equation (34) follows. \square

Remark 6. By introducing the following notation for the average of a generic measurable function $f : M^N \rightarrow \mathbb{R}$ over the hypersurface $\Sigma_{\bar{v}}^N$ endowed with the measure $d\mu_{\bar{v}}^{N-1}$

$$\langle f \rangle_{\bar{v},\mu} = \frac{\int_{\Sigma_{\bar{v}}^N} f d\mu_{\bar{v}}^{N-1}}{\int_{\Sigma_{\bar{v}}^N} d\mu_{\bar{v}}^{N-1}} = \frac{\int_{\Sigma_{\bar{v}}^N} f d\mu_{\bar{v}}^{N-1}}{\Omega_N(\bar{v})}, \tag{37}$$

the quantities

$$\begin{aligned} \text{Var}_{\bar{v},\mu}(f) &= \text{Cuml}_{\bar{v},\mu}^{(2)}(f) = \langle f^2 \rangle_{\bar{v},\mu} - \langle f \rangle_{\bar{v},\mu}^2 \\ \text{Cov}_{\bar{v},\mu}(f; g) &= \langle fg \rangle_{\bar{v},\mu} - \langle f \rangle_{\bar{v},\mu} \langle g \rangle_{\bar{v},\mu} \\ \text{Cuml}_{\bar{v},\mu}^{(3)}(f) &= \langle f^3 \rangle_{\bar{v},\mu} - 3\langle f \rangle_{\bar{v},\mu} \langle f^2 \rangle_{\bar{v},\mu} + 2\langle f \rangle_{\bar{v},\mu}^3 \\ \text{Cuml}_{\bar{v},\mu}^{(4)}(f) &= \langle f^4 \rangle_{\bar{v},\mu} - 4\langle f^3 \rangle_{\bar{v},\mu} \langle f \rangle_{\bar{v},\mu} + 12\langle f^2 \rangle_{\bar{v},\mu} \langle f \rangle_{\bar{v},\mu}^2 - 3\langle f^2 \rangle_{\bar{v},\mu}^2 \\ &\quad - 6\langle f \rangle_{\bar{v},\mu}^4 \end{aligned} \tag{38}$$

represent the variance, the correlation function, and the 3rd and 4th order cumulants on the hypersurface $\Sigma_{\bar{v}}^N$ with measure $d\mu_{\bar{v}}^{N-1}$, respectively.

With this notation, and substituting equation (32) in equation (31), the derivatives of the microcanonical entropy at a non critical value \bar{v} , and at fixed N are worked out as averages of functions of $\bar{\zeta}_N = \text{div}(\bar{\xi}_N)$, where the vector field $\bar{\xi}_N$ generates the diffeomorphisms among the equipotential level sets, as follows

$$\begin{aligned} \frac{d\bar{S}_N}{d\bar{v}}(\bar{v}) &= \frac{1}{N} \langle \bar{\zeta}_N \rangle_{\bar{v},\mu} \\ \frac{d^2\bar{S}_N}{d\bar{v}^2}(\bar{v}) &= \frac{1}{N} \left[\text{Var}_{\bar{v},\mu}(\bar{\zeta}_N) + \left\langle \mathcal{L}_{\bar{\xi}_N}(\bar{\zeta}_N) \right\rangle_{N\bar{v},\mu} \right] \\ \frac{d^3\bar{S}_N}{d\bar{v}^3}(\bar{v}) &= \frac{1}{N} \left[\text{Cuml}_{\bar{v},\mu}^{(3)}(\bar{\zeta}_N) + 3\text{Cov}_{\bar{v},\mu}(\bar{\zeta}_N; \mathcal{L}_{\bar{\xi}_N}(\bar{\zeta}_N)) + \left\langle \mathcal{L}_{\bar{\xi}_N}^{(ii)}(\bar{\zeta}_N) \right\rangle_{\bar{v},\mu} \right] \\ \frac{d^4\bar{S}_N}{d\bar{v}^4}(\bar{v}) &= \frac{1}{N} \left[\text{Cuml}_{\bar{v},\mu}^{(4)}(\bar{\zeta}_N) + 6\text{Cov}_{\bar{v},\mu}(\bar{\zeta}_N^2; \mathcal{L}_{\bar{\xi}_N}(\bar{\zeta}_N)) \right. \\ &\quad \left. + 3\text{Var}_{\bar{v},\mu}(\mathcal{L}_{\bar{\xi}_N}(\bar{\zeta}_N)) + 4\text{Cov}_{\bar{v},\mu}(\bar{\zeta}_N; \mathcal{L}_{\bar{\xi}_N}^{(ii)}(\bar{\zeta}_N)) \right. \\ &\quad \left. - 12\langle \bar{\zeta}_N \rangle_{\bar{v},\mu} \text{Cov}_{N\bar{v},\mu}(\bar{\zeta}_N; \mathcal{L}_{\bar{\xi}_N}(\bar{\zeta}_N)) + \left\langle \mathcal{L}_{\bar{\xi}_N}^{(iii)}(\bar{\zeta}_N) \right\rangle_{\bar{v},\mu} \right] \\ &= \frac{1}{N} \left[\text{Cuml}_{\bar{v},\mu}^{(4)}(\bar{\zeta}_N) + 4\text{Cov}_{\bar{v},\mu}(\bar{\zeta}_N; \mathcal{L}_{\bar{\xi}_N}^{(ii)}(\bar{\zeta}_N)) + 3\text{Var}_{\bar{v},\mu}(\mathcal{L}_{\bar{\xi}_N}(\bar{\zeta}_N)) \right. \\ &\quad \left. + 6\langle \bar{\zeta}_N \rangle_{\bar{v},\mu} \left(\text{Cov}_{\bar{v},\mu}(\Delta\bar{\zeta}_N; \mathcal{L}_{\bar{\xi}_N}(\bar{\zeta}_N)) \right) + \left\langle \mathcal{L}_{\bar{\xi}_N}^{(iii)}(\bar{\zeta}_N) \right\rangle_{\bar{v},\mu} \right], \end{aligned} \tag{39}$$

where for sake of simplicity we have introduced the quantity

$$\Delta\bar{\zeta}_N = \frac{\bar{\zeta}_N^2}{\langle \bar{\zeta}_N \rangle_{\bar{v},\mu}} - 2\bar{\zeta}_N. \tag{40}$$

Now the crucial step is to show that, under the hypothesis of diffeomorphicity that now includes *asymptotic diffeomorphicity*, the function $\bar{\zeta}_N$ —considered along a MCMC spanning any given Σ_v^{VN} —is a Gaussian random process. This is achieved through an intermediate step to show that the mean curvature H —also considered along the same MCMC—is a Gaussian random process. For the sake of notation in what follows we shall omit the suffix N of V_N .

Lemma 4 (Mean curvature along a MCMC on a level set). *The pointwise mean curvature of an N -dimensional manifold $\Sigma_{N\bar{v}}^V$*

$$H(\mathbf{q}) = \frac{1}{N} \sum_{i=1}^N \kappa_i(\mathbf{q}) = -\frac{1}{N} \left[\frac{\Delta V}{\|\nabla V\|} - \frac{\partial^i V \partial_{ij}^2 V \partial^j V}{\|\nabla V\|^3} \right] \tag{41}$$

computed along a MCMC $\{\mathbf{q}_k\}_{k \in \mathbb{N}} \in \Sigma_{N\bar{v}}^V$ such that the stationary invariant density of the MCMC is the microcanonical configurational measure, where $\Sigma_{N\bar{v}}^V$ is free of critical points of V , is a Gaussian random process.

Proof. Along a MCMC, the principal curvatures $\kappa_i(\mathbf{q})$ behave as independent random variables with probability densities $u_i(\kappa_i)$ which we do not need to know explicitly. Statistical independence means that $\langle \kappa_i(\mathbf{q}) \kappa_j(\mathbf{q}) \rangle_{N,v}^{\mu c} = \langle \kappa_i(\mathbf{q}) \rangle_{N,v}^{\mu c} \langle \kappa_j(\mathbf{q}) \rangle_{N,v}^{\mu c}$ and this can be understood as follows. Let (M^n, g) be an n -dimensional Riemannian manifold and m -codimensional submanifold of a Riemannian manifold (\bar{M}^{m+n}, \bar{g}) , let R and \bar{R} denote the Riemann curvature tensors of M^n and \bar{M}^{m+n} , respectively, and denote by $h(\cdot, \cdot)$ the second fundamental form, then the Gauss equation reads

$$\begin{aligned} \bar{g}(\bar{R}(X, Y)Z, W) &= g(R(X, Y)Z, W) + \bar{g}(h(X, Z), h(Y, W)) \\ &\quad - \bar{g}(h(X, W), h(Y, Z)) \end{aligned} \tag{42}$$

which, for sectional curvatures, obviously reads as

$$\begin{aligned} \bar{g}(\bar{R}(X, Y)X, Y) &= g(R(X, Y)X, Y) + \bar{g}(h(X, X), h(Y, Y)) \\ &\quad - \bar{g}(h(X, Y), h(Y, X)). \end{aligned} \tag{43}$$

Now, for any point $p \in M$ and basis $\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$ of $T_p M$, it is possible to choose coordinates (y^1, \dots, y^{n+1}) in \bar{M} such that the tangent vectors $\mathbf{Y}^1, \dots, \mathbf{Y}^n$ coincide with $\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$ and $\mathbf{n} = \mathbf{Y}^{n+1} \in N_p M$ is orthogonal to $T_p M$. Then M is locally given as a graph manifold: $y^1 = x^1, \dots, y^n = x^n, y^{n+1} = f(\mathbf{x})$ so that the second fundamental form has the components [34, 41]

$$h(\mathbf{e}_i, \mathbf{e}_j) = \frac{\partial^2 f}{\partial x^i \partial x^j} \mathbf{n}, \tag{44}$$

where $\mathbf{e}_i = \partial/\partial x^i$. Considering the potential level sets $\Sigma_{N\bar{v}}^V$ as hypersurfaces of \mathbb{R}^{N+1} , identifying $f(x^1, \dots, x^N)$ with $V(q^1, \dots, q^N)$, taking $\mathbf{n} = \nabla V / \|\nabla V\|$, from equations (42)–(44) we obtain

$$K(\mathbf{e}_i, \mathbf{e}_j) = \kappa_i \kappa_j = - \left(\frac{\partial^2 V}{\partial q_i^2} \right) \left(\frac{\partial^2 V}{\partial q_j^2} \right) \langle \mathbf{n}, \mathbf{n} \rangle + \left(\frac{\partial^2 V}{\partial q_i \partial q_j} \right)^2 \langle \mathbf{n}, \mathbf{n} \rangle \tag{45}$$

hence

$$\langle \kappa_i(\mathbf{q}) \kappa_j(\mathbf{q}) \rangle_{N,v}^{\mu c} = \left\langle \frac{1}{\|\nabla V\|} \left[\left(\frac{\partial^2 V}{\partial q_i \partial q_j} \right)^2 - \left(\frac{\partial^2 V}{\partial q_i^2} \right) \left(\frac{\partial^2 V}{\partial q_j^2} \right) \right] \right\rangle_{N,v}^{\mu c}. \tag{46}$$

For short-range interactions with coordination number n_0 , meaning that—with a suitable labelling of the variables— q_i and q_j do not interact if $|i - j| > n_0$, the entries of the Hessian of V vanish if $|i - j| > n_0$. Thus, locally, for $n > n_0$, we have

$$\langle \kappa_i(\mathbf{q})\kappa_{j=i+n}(\mathbf{q}) \rangle_{N,v}^{\mu c} = \left\langle - \left[\frac{1}{\|\nabla V\|^{1/2}} \left(\frac{\partial^2 V}{\partial q_i^2} \right) \right] \left[\frac{1}{\|\nabla V\|^{1/2}} \left(\frac{\partial^2 V}{\partial q_j^2} \right) \right] \right\rangle_{N,v}^{\mu c} \quad (47)$$

with evident notation, we can write

$$\begin{aligned} & \langle \kappa_i(\mathbf{q})\kappa_{j=i+n}(\mathbf{q}) \rangle_{N,v}^{\mu c} \\ &= \left\langle \left[\langle \kappa_i(\mathbf{q}) \rangle_{N,v}^{\mu c} + \delta \kappa_i(\mathbf{q}) \right] \left[\langle \kappa_{j=i+n}(\mathbf{q}) \rangle_{N,v}^{\mu c} + \delta \kappa_{j=i+n}(\mathbf{q}) \right] \right\rangle_{N,v}^{\mu c}. \end{aligned} \quad (48)$$

As we shall see in the following, $1/\|\nabla V\|^{1/2}$ tends to a constant value at increasing N , and along a MCMC sampling a potential level set with its configurational microcanonical measure, the fluctuations of $\partial^2 V/\partial q_i^2$ and $\partial^2 V/\partial q_j^2$ are independent and average to zero. In conclusion, under the condition $|i - j| > n_0$ we have

$$\langle \kappa_i(\mathbf{q})\kappa_j(\mathbf{q}) \rangle_{N,v}^{\mu c} = \langle \kappa_i(\mathbf{q}) \rangle_{N,v}^{\mu c} \langle \kappa_j(\mathbf{q}) \rangle_{N,v}^{\mu c}.$$

Having shown that the principal curvatures $\kappa_i(\mathbf{q})$ are everywhere uniformly bounded on any $\Sigma_{N\bar{v}}^V$ belonging to any cylindrical subset of the family $\{\Gamma_{I_0}^N\}_{N \in \mathbb{N}}$, the consequence is that the momenta of the distributions $u_i(\kappa_i)$ are finite and uniformly bounded too. Hence, the basic conditions are fulfilled to apply the central limit theorem (CLT) formulated by Khinchin [42] for sum functions of independent random variables, arbitrarily distributed, with bounded momenta up to the fifth order. Hence, along the MCMC $\{\mathbf{q}_k\}_{k \in \mathbb{N}} \in \Sigma_{N\bar{v}}^V$ the invariant measure of which is the configurational microcanonical one, the values of the mean curvature $H(\mathbf{q}_k)$ behave as Gaussian-distributed random variables. Notice that a finite range dependence is a weak dependence that does not prevent the CLT to apply [43].

Lemma 5 ($\zeta_N(\mathbf{q})$ along the MCMC on regular level sets). *The quantity*

$$\zeta_N(\mathbf{q}) = \frac{\Delta V}{\|\nabla V\|^2} - 2 \frac{\partial^i V \partial_{ij}^2 V \partial^j V}{\|\nabla V\|^4} \quad (49)$$

as well as $\bar{\zeta}_N(\mathbf{q})$, computed along a MCMC $\{\mathbf{q}_k\}_{k \in \mathbb{N}} \in \Sigma_{N\bar{v}}^V$ the invariant measure of which is the configurational microcanonical one, is a Gaussian random process.

Proof. After the preceding lemma it follows that the two quantities $\Delta V/\|\nabla V\|$ and $\partial^i V \partial_{ij}^2 V \partial^j V / \|\nabla V\|^3$ —computed along MCMC $\{\mathbf{q}_k\}_{k \in \mathbb{N}} \in \Sigma_{N\bar{v}}^V$ the invariant measure of which is the configurational microcanonical one—are Gaussian random processes because their sum is a Gaussian random process and the sum of Gaussian random processes is Gaussian. Now, if the quantity $\frac{\Delta V}{\|\nabla V\|} = \sum_i \partial_{ii}^2 V / \|\nabla V\|$ is asymptotically Gaussian, then the terms $\partial_{ii}^2 V / \|\nabla V\|$ have to be i.i.d. random variables as well the terms $\partial_{ii}^2 V$ because all of them are divided by the same number $\|\nabla V\|$ at each point of the MCMC, by the same token $\partial_{ii}^2 V / \|\nabla V\|^2$ have to be i.i.d. random variables because now all the terms $\partial_{ii}^2 V$ are divided by the same number $\|\nabla V\|^2$ at each point of the MCMC. The same argument applies to $\partial^i V \partial_{ij}^2 V \partial^j V / \|\nabla V\|^3$ so that in the end both $\Delta V / \|\nabla V\|^2$ and $\partial^i V \partial_{ij}^2 V \partial^j V / \|\nabla V\|^4$ are Gaussian distributed, and,

consequently, $\bar{\zeta}_N(\mathbf{q})$ in equation (49) is a Gaussian random variable along a MCMC the invariant measure of which is the configurational microcanonical one.

Remark 7. Let us emphasize that the quantity $\bar{\zeta}_N(\mathbf{q})$ is a random variable along all the MCMC $\{\mathbf{q}_k\}_{k \in \mathbb{N}} \in \Sigma_{N_k \bar{v}}^{V_{N_k}}$, with vanishing deviations from a Gaussian distribution at increasing N , under the hypothesis of asymptotic diffeomorphicity because the principal curvatures $\kappa_i(\mathbf{q})$ are uniformly bounded with N from above on any manifold, a crucial condition for the validity of lemma 2.

Remark 8. In the hypotheses of the main theorem, V contains only short-range interactions and its functional form does not change with N . In other words, we are tackling physically homogeneous systems, which, at any N , can be considered as the union of smaller and identical subsystems. If a system is partitioned into a number k of sufficiently large subsystems, the larger N the more accurate the factorization of its configuration space. Therefore, the averages of functions of interacting variables belonging to a given block depend neither on the subsystems where they are computed (the potential functions are the same on each block after suitable relabelling of the variables) nor on the total number N of degrees of freedom.

(a) Since the potential V is assumed smooth and bounded below, one has

$$\langle |\Delta V| \rangle_{N,v}^{\mu c} = \left\langle \left| \sum_{i=1}^N \partial_{ii}^2 V \right| \right\rangle_{N,v}^{\mu c} \leq \sum_{i=1}^N \langle |\partial_{ii}^2 V| \rangle_{N,v}^{\mu c} \leq N \max_{i=1, \dots, N} \langle |\partial_{ii}^2 V| \rangle_{N,v}^{\mu c}.$$

At large N (when the fluctuations of the averages are vanishingly small) $\max_{i=1, \dots, N} \langle |\partial_{ii}^2 V| \rangle_{N,v}^{\mu c}$ does not depend on N , and the same holds for $\langle |\partial^i V \partial_{ij}^2 V \partial^j V| \rangle_{N,v}^{\mu c}$ and $\max_{i,j=1, \dots, N} \langle |\partial^i V \partial_{ij}^2 V \partial^j V| \rangle_{N,v}^{\mu c}$.

Hence we set

$$\begin{aligned} m_1 &= \max_{i=1, \dots, N} \langle |\partial_{ii}^2 V| \rangle_{N,v}^{\mu c} \\ m_2 &= \max_{i,j=1, \dots, N} \langle |\partial^i V \partial_{ij}^2 V \partial^j V| \rangle_{N,v}^{\mu c}. \end{aligned} \tag{50}$$

(b) Moreover, the absence of critical points of V , implied by the hypothesis of diffeomorphicity of the equipotential hypersurfaces, means that $\|\nabla V\|^2 \geq C > 0$. Hence the terms $\langle \|\nabla V\|^{2n} \rangle_{N,v}^{\mu c}$ for $n = 1, \dots, 8$ we have

$$\begin{aligned} \langle \|\nabla V\|^2 \rangle_{N,v}^{\mu c} &= \left\langle \sum_{i=1}^N (\partial_i V)^2 \right\rangle_{N,v}^{\mu c} \\ &= \sum_{i=1}^N \langle (\partial_i V)^2 \rangle_{N,v}^{\mu c} \geq N \min_{i=1, \dots, N} \langle (\partial_i V)^2 \rangle_{N,v}^{\mu c}, \\ \langle \|\nabla V\|^4 \rangle_{N,v}^{\mu c} &= \left\langle \left[\sum_{i=1}^N (\partial_i V)^2 \right]^2 \right\rangle_{N,v}^{\mu c} = \sum_{i,j=1}^N \langle (\partial_i V)^2 (\partial_j V)^2 \rangle_{N,v}^{\mu c} \\ &\geq N^2 \min_{i,j=1, \dots, N} \langle (\partial_i V)^2 (\partial_j V)^2 \rangle_{N,v}^{\mu c}, \end{aligned}$$

which can be iterated up to $\langle \|\nabla V\|^{16} \rangle_{N,v}^{\mu c}$. By setting

$$\begin{aligned}
 c_1 &= \min_{i=1,\dots,N} \langle (\partial_i V)^2 \rangle_{N,v}^{\mu c} \\
 c_2 &= \min_{i,j=1,\dots,N} \langle (\partial_i V)^2 (\partial_j V)^2 \rangle_{N,v}^{\mu c} \\
 &\dots\dots\dots \\
 c_8 &= \min_{i_1,\dots,i_8=1,\dots,N} \langle (\partial_{i_1} V)^2 \dots (\partial_{i_8} V)^2 \rangle_{N,v}^{\mu c}. \tag{51}
 \end{aligned}$$

(c) By the same token put forward at the beginning of this remark, we can define the following quantities independent of N

$$\begin{aligned}
 m_3 &= \max_{i,j,k,l=1,N} \langle (\partial_i V \partial_{ij}^2 V \partial_j V) (\partial_k V \partial_{kl}^2 V \partial_l V) \rangle_{N,v}^{\mu c}, \\
 m_4 &= \max_{i,j,k=1,N} \langle \partial_i V \partial_{ij}^2 V \partial_{jk}^2 V \partial_k V \rangle_{N,v}^{\mu c}, \\
 m_5 &= \max_{i,j,k=1,N} \langle (\partial_i V \partial_{ij}^2 V \partial_j V) (\partial_{kk}^2 V) \rangle_{N,v}^{\mu c}, \\
 m_6 &= \max_{i,j=1,N} \langle \partial_i V \partial_{ijj}^3 V \rangle_{N,v}^{\mu c}, \\
 m_7 &= \max_{i,j,k=1,N} \langle (\partial_i V \partial_j V \partial_k V) \partial_{ijk}^3 V \rangle_{N,v}^{\mu c}, \\
 m_8 &= \max_{i,j=1,\dots,N} \langle |\partial_{ij}^2 V \partial^j V| \rangle_{N,v}^{\mu c}. \tag{52}
 \end{aligned}$$

Lemma 6 (Upper bound of the first derivative of the entropy). *After lemmas 2 to 5 and remarks 2 to 5 it is*

$$\sup_{N, \bar{v} \in I_{\bar{v}}} \left| \frac{\partial S_N}{\partial \bar{v}}(\bar{v}) \right| < \infty$$

Proof. This first derivative of the entropy is equal to the inverse of the configurational temperature, thus it is necessarily uniformly bounded with N . In fact, the property of $\bar{\zeta}_N(\mathbf{q})$ —and of $\zeta_N(\mathbf{q})$ —of being a Gaussian distributed random variable along any MCMC defined above entails the following uniform bound

$$\lim_{N \rightarrow +\infty} \langle \zeta_N \rangle_{N\bar{v},\mu} = \lim_{N \rightarrow +\infty} N^{-1} \langle \bar{\zeta}_N \rangle_{N\bar{v},\mu} \in \mathbb{R}. \tag{53}$$

□

Lemma 7 (Upper bound of the second derivative of the entropy). *After lemmas 2 to 5 and remarks 2 to 5 it is*

$$\sup_{N, \bar{v} \in I_{\bar{v}}} \left| \frac{\partial^2 S_N}{\partial \bar{v}^2}(\bar{v}) \right| < \infty. \tag{54}$$

Proof. Since $\bar{\zeta}_N(\mathbf{q})$ is a Gaussian random process, the quantity $\text{Var}_{\bar{v},\mu}(\bar{\zeta}_N)/N$ is uniformly bounded with N . Then the N -dependence of the average $\langle \mathcal{L}_{\bar{\xi}_N}(\bar{\zeta}_N) \rangle_{N\bar{v},\mu}$ follows from the explicit expression of the quantity $\mathcal{L}_{\bar{\xi}_N}(\bar{\zeta}_N)$ given by equation (74) in appendix B (by adapting it to quantities marked with an overline). Considering that the number of non-vanishing entries

of the Hessian of the potential is $\mathcal{O}(n_p N)$, where n_p is the number of nearest-neighbors in condensed matter systems and the average number of neighbors of a particle in a fluid system, using the above defined N -independent quantities in equations (50)–(52) a simple estimation term by term gives

$$\begin{aligned} \left\langle \mathcal{L}_{\bar{\xi}_N}(\bar{\zeta}_N) \right\rangle_{N\bar{v},\mu} &\leq \left\langle \left| \frac{\nabla\bar{V} \cdot \nabla(\Delta\bar{V})}{\|\nabla\bar{V}\|^4} \right| \right\rangle_{N\bar{v},\mu} + \left\langle \left| 8 \frac{(\nabla\bar{V} \cdot \text{Hess}\bar{V}\nabla\bar{V})^2}{\|\nabla\bar{V}\|^8} \right| \right\rangle_{N\bar{v},\mu} \\ &+ \left\langle \left| 2 \frac{(\nabla\bar{V} \cdot \text{Hess}(\bar{V})\nabla\bar{V})\Delta\bar{V} + 2\|\text{Hess}\bar{V}\nabla\bar{V}\|^2 + \text{D}^3\bar{V}(\nabla\bar{V}, \nabla\bar{V}, \nabla\bar{V})}{\|\nabla\bar{V}\|^6} \right| \right\rangle_{N\bar{v},\mu} \\ &\leq N \frac{m_6}{c_2} + 8 \frac{m_2^2 n_p^2}{c_4} + 2N \frac{m_5 n_p + 2m_3^2 n_p^2 + m_7 n_p}{c_3} \end{aligned} \quad (55)$$

that is, the upper bound of $\left\langle \mathcal{L}_{\bar{\xi}_N}(\bar{\zeta}_N) \right\rangle_{N\bar{v},\mu} / N$ of this quantity remains uniformly bounded in the $N \rightarrow \infty$ limit. \square

Lemma 8 (Upper bound of the third derivative of the entropy). *After lemmas 2 to 5 and remarks 2 to 5 it is*

$$\sup_{N, \bar{v} \in I_{\bar{v}}} \left| \frac{\partial^3 S_N}{\partial \bar{v}^3}(\bar{v}) \right| < \infty$$

Proof. Since $\bar{\zeta}_N(\mathbf{q})$ is a Gaussian random process, we have the following uniform bound

$$\lim_{N \rightarrow +\infty} N^2 \text{Cumul}_{N\bar{v},\mu}^{(3)} \bar{\zeta}_N = \lim_{N \rightarrow +\infty} N^{-1} \text{Cumul}_{N\bar{v},\mu}^{(3)} \bar{\zeta}_N = 0,$$

and by considering the explicit expression of $\mathcal{L}_{\bar{\xi}_N}^{(ii)}(\bar{\zeta}_N)$ given by equation (75) in appendix B, a tedious but trivial counting of the N -dependence term by term of $\mathcal{L}_{\bar{\xi}_N}^{(ii)}(\bar{\zeta}_N)$ —as in the previous case—shows that $\left\langle \mathcal{L}_{\bar{\xi}_N}^{(ii)}(\bar{\zeta}_N) \right\rangle_{N\bar{v},\mu}$ turns out of order $\mathcal{O}(n_p^3 N)$ and thus divided by N remains uniformly bounded in the $N \rightarrow \infty$ limit. Then, according to the definition (38) we have

$$\text{Cov}_{\bar{v},\mu}(\bar{\zeta}_N; \mathcal{L}_{\bar{\xi}_N}(\bar{\zeta}_N)) = \langle \bar{\zeta}_N \mathcal{L}_{\bar{\xi}_N}(\bar{\zeta}_N) \rangle_{\bar{v},\mu} - \langle \bar{\zeta}_N \rangle_{\bar{v},\mu} \langle \mathcal{L}_{\bar{\xi}_N}(\bar{\zeta}_N) \rangle_{\bar{v},\mu},$$

where the quantities $\bar{\zeta}_N$ and $\mathcal{L}_{\bar{\xi}_N}(\bar{\zeta}_N)$ are randomly varying along the MCMC whose probability measure is the configurational microcanonical measure, and the random variations of $\bar{\zeta}_N$ and of its directional (Lie) derivative in a random direction $\bar{\xi}_N$ can be considered *bona fide* statistically uncorrelated, thus their covariance vanishes. \square

Lemma 9 (Upper bound of the fourth derivative of the entropy). *After lemmas 2 to 5 and remarks 2 to 5 it is*

$$\sup_{N, \bar{v} \in I_{\bar{v}}} \left| \frac{\partial^4 S_N}{\partial \bar{v}^4}(\bar{v}) \right| < \infty.$$

Proof. Since $\bar{\zeta}_N(\mathbf{q})$ is a Gaussian random process, we have the following uniform bound

$$\lim_{N \rightarrow +\infty} N^3 \text{Cumul}_{N\bar{v},\mu}^{(4)} \bar{\zeta}_N = \lim_{N \rightarrow +\infty} N^{-1} \text{Cumul}_{N\bar{v},\mu}^{(4)} \bar{\zeta}_N = 0.$$

Then, by considering the expression of $\mathcal{L}_{\bar{\xi}_N}^{(iii)}(\bar{\zeta}_N)$ given by equation (76) in appendix B, a very tedious but trivial counting of the N -dependence term by term of $\mathcal{L}_{\bar{\xi}_N}^{(iii)}(\bar{\zeta}_N)$ —as in the previous

case—shows that $\left\langle \mathcal{L}_{\xi_N}^{(iii)}(\bar{\zeta}_N) \right\rangle_{N\bar{v},\mu}$ turns out of order $\mathcal{O}(n_p^4 N)$ and thus divided by N remains uniformly bounded in the $N \rightarrow \infty$ limit.

Now, the term $N^{-1} \text{Var}_{\bar{v},\mu}(\mathcal{L}_{\xi_N}(\bar{\zeta}_N))$ is also uniformly bounded, in fact along the MCMC spanning a given equipotential surface the terms $\bar{\xi}_i \partial_i \bar{\zeta}$ are random uncorrelated variables each bringing a factor N because $\bar{\zeta} \sim \mathcal{O}(N)$, thus

$$\begin{aligned} \text{Var}_{\bar{v},\mu}(\mathcal{L}_{\xi_N}(\bar{\zeta}_N)) &= \text{Var}_{\bar{v},\mu}\left(\sum_{i=1}^N \bar{\xi}_i \partial_i \bar{\zeta}\right) \\ &= \text{Var}_{\bar{v},\mu}\left(\frac{1}{N} \sum_{i=1}^N N \bar{\xi}_i \partial_i \bar{\zeta}\right) \\ &= \frac{1}{N^2} \sum_{i=1}^N \text{Var}_{\bar{v},\mu}(N \bar{\xi}_i \partial_i \bar{\zeta}) \\ &\leq \frac{1}{N^2} N \sigma_m^2 N^2 = N \sigma_m^2, \end{aligned} \tag{56}$$

where σ_m^2 is the largest value of all the standard deviations of the terms $\bar{\xi}_i \partial_i \bar{\zeta}$ along the MCMC.

For what concerns the two remaining terms in the fourth derivative of the entropy in equation (39) we have

$$\text{Cov}_{\bar{v},\mu}(\bar{\zeta}_N; \mathcal{L}_{\xi_N}^{(ii)}(\bar{\zeta}_N)) = \left\langle \bar{\zeta}_N \mathcal{L}_{\xi_N}^{(ii)}(\bar{\zeta}_N) \right\rangle_{\bar{v},\mu} - \langle \bar{\zeta}_N \rangle_{\bar{v},\mu} \left\langle \mathcal{L}_{\xi_N}^{(ii)}(\bar{\zeta}_N) \right\rangle_{\bar{v},\mu} \tag{57}$$

that vanishes when computed as microcanonical averages through ‘time’ averages along a MCMC, in fact, we take advantage of the resulting complete decorrelation between the random values taken by $\bar{\zeta}_N$ and the random values of its second order Lie derivative taken in a random direction ξ_N .

$$\text{Cov}_{\bar{v},\mu}(\Delta \bar{\zeta}_N; \mathcal{L}_{\xi_N}(\bar{\zeta}_N)) = \left\langle \Delta \bar{\zeta}_N \mathcal{L}_{\xi_N}(\bar{\zeta}_N) \right\rangle - \langle \Delta \bar{\zeta}_N \rangle \left\langle \mathcal{L}_{\xi_N}(\bar{\zeta}_N) \right\rangle \tag{58}$$

the same argument applies to the quantities $\Delta \bar{\zeta}_N$ and $\mathcal{L}_{\xi_N}(\bar{\zeta}_N)$ that are uncorrelated random variables along a MCMC, thus their covariance vanishes. \square

4. Fixing the problem raised by the lattice ϕ^4 model

A few years ago, an argument was raised [31] against the topological theory of phase transitions on the basis of the observation that the second order phase transition of the 2D lattice ϕ^4 -model occurs at a critical value v_c of the potential energy density which belongs to a broad interval of v -values void of critical points of the potential function. In other words, for any finite N the $\{\sum_{v < v_c}^{V_N}\}_{v \in \mathbb{R}}$ are diffeomorphic to the $\{\sum_{v > v_c}^{V_N}\}_{v \in \mathbb{R}}$ so that no topological change seems to correspond to the phase transition. This is a counterexample to the theorem in references [28, 29]. A first reply was given in [32] where, in spite of the absence of critical points of the potential in correspondence of the transition energy, a strong evidence has been given to relate the phase transition of this model with a change of topology of both the energy and potential level sets. But in this case the topology changes are asymptotic ($N \rightarrow \infty$).




Figure 5. Quantities entering equation (39) for lattices with different N . In particular, $N = 100$ (red full circles), $N = 400$ (blue squares), $N = 900$ (black diamonds), $N = 1600$ (green triangles), $N = 2500$ (purple reversed triangles).

Let us see how the main theorem proved in the present work definitely fixes the problem, so that the 2D lattice ϕ^4 -model is no longer a counterexample to the topological necessity theorem.

The model of interest, considered in reference [44], is defined by the Hamiltonian

$$\mathcal{H}_N(p, q) = \sum_i \frac{p_i^2}{2} + V_N(q), \tag{59}$$

where the potential function $V(q)$ is

$$V(q) = \sum_{i \in \mathbb{Z}^D} \left(-\frac{m^2}{2} q_i^2 + \frac{\lambda}{4!} q_i^4 \right) + \sum_{\langle ik \rangle \in \mathbb{Z}^D} \frac{1}{2} J(q_i - q_k)^2, \tag{60}$$

with $\langle ik \rangle$ standing for nearest-neighbor sites on a D dimensional lattice. This system has a discrete \mathbb{Z}_2 -symmetry and short-range interactions; therefore, according to the Mermin–Wagner theorem, in $D = 1$ there is no phase transition whereas in $D = 2$ there is a second order symmetry-breaking transition, with nonzero critical temperature, of the same universality class of the 2D Ising model.

In this section we present the results of Monte Carlo numerical simulations on equipotential level set of this model on a 2D-lattice with periodic boundary conditions and the following parameters: $J = 1$, $m^2 = 2$, and $\lambda = 0.6$. For these values of the parameters, the 2D system undergoes the symmetry-breaking phase transition at the critical potential energy density value is $v_c = \langle V \rangle_c / N \simeq 2.2$. This study has been performed in order to identify which terms composing the derivatives of the specific configurational microcanonical entropy with respect to the specific potential energy is not uniformly bounded in N , as is expected after the present main theorem.

The simulations have been performed for systems with different number of degrees of freedom: $N = 10 \times 10 = 100$, $N = 20 \times 20 = 400$, $N = 30 \times 30 = 900$, $N = 40 \times 40 = 1600$, $N = 50 \times 50 = 2500$ and $N = 70 \times 70 = 4900$. The computations were performed with vanishing magnetization as initial condition, for 2×10^7 steps, a number sufficient to guarantee the convergence of the reported quantities.




Figure 6. Quantities entering equation (39) for lattices with different N . In particular, $N = 100$ (red full circles), $N = 400$ (blue squares), $N = 900$ (black diamonds), $N = 1600$ (green triangles), $N = 2500$ (purple reversed triangles).




Figure 7. Quantities entering equation (39) for lattices with different N . In particular, $N = 100$ (red full circles), $N = 400$ (blue squares), $N = 900$ (black diamonds), $N = 1600$ (green triangles), $N = 2500$ (purple reversed triangles).




Figure 8. Quantities entering equation (39) for lattices with different N . Left panel: third cumulant of ζ_N for $N = 100$ (red full circles), $N = 400$ (blue squares), $N = 900$ (black diamonds), $N = 1600$ (green triangles), $N = 2500$ (purple reversed triangles). Right panel: fourth cumulant of ζ_N computed at the transition energy density $v_c = 2.2$, here a systematic deviation from the Gaussian scaling with N is well evident.

The results of numerical simulations are reported for each single term entering equation (39) (figures 5–8). When properly rescaled with N , under the hypothesis of diffeomorphism (at any N and also asymptotically) of the equipotential hypersurfaces, all these terms are expected to be uniformly bounded with N . Very interestingly, it is found that across the vertical dashed line denoting the phase transition point at the potential energy density $\bar{v}_c \simeq 2.2$ all these terms do not show any tendency to change with N , except for the case $N = 10 \times 10$ (for which 36% of the total number of sites belong to the boundary, making the finite size effects more relevant). There is only one very significant exception, the fourth cumulant of $\bar{\zeta}_N$ which, computed around the transition value, is found to systematically grow with N . This has been computed by means of the relation

$$\text{Cuml}_{N\bar{v},\mu}^{(4)}\bar{\zeta} = \frac{d}{d\bar{v}}[\text{Cuml}_{N\bar{v},\mu}^{(3)}\bar{\zeta}_N] - 3\langle\bar{\zeta}_N\rangle_{\bar{v},\mu}(\text{Corr}_{\bar{v},\mu}(\Delta\bar{\zeta}_N; \mathcal{L}_{\bar{\zeta}_N}(\bar{\zeta}_N))), \quad (61)$$

where the derivative of the third cumulant is evaluated numerically.

This means that $\bar{\zeta}_N$ is not a Gaussian random process along a MCMC the invariant measure of which is the microcanonical measure. As $\bar{\zeta}_N$ is proved to be a Gaussian random process under the constraining hypothesis of asymptotic diffeomorphicity of the level sets $\{\Sigma_{N\bar{v}}^{V_N}\}_{\bar{v} \in [\bar{v}_0, \bar{v}_1]}$, the growth with N of $N^{-1}\text{Cuml}_{N\bar{v},\mu}^{(4)}\bar{\zeta}$ entails the loss of asymptotic diffeomorphicity among the $\{\Sigma_{N\bar{v}}^{V_N}\}_{\bar{v} < \bar{v}_c}$ and the $\{\Sigma_{N\bar{v}}^{V_N}\}_{\bar{v} > \bar{v}_c}$ for some \bar{v}_c . This means that the 2D lattice ϕ^4 model does not fulfil a basic requirement of the main theorem formulated in the present work. Therefore, the 2D lattice ϕ^4 model is not a counterexample to the present version of the topological necessity theorem.

As it has been pointed out in remark 3, a second order phase transition in the microcanonical ensemble is associated with an asymptotic discontinuity of the third derivative of the entropy and hence an asymptotic divergence of the fourth derivative of the entropy.

5. Historical overview and challenges of the topological theory of phase transitions

After the investigation of specific, exactly solvable models [17, 45–48] corroborating the TH, during the last decade and a half other systems have been studied, shedding doubts on the general validity of the new theory. Even if all these studies are important to better define the validity limits of the TH, some authors have too quickly drawn fatal verdicts against it.

In fact, for the sake of clarity, let us begin by remarking that the proposed theory applies *a priori* to systems described by smooth, finite-range potentials such that the level sets Σ_v^{VN} and their associated balls M_v^{VN} qualify as *good differentiable and compact manifolds*. Moreover, as we have preliminarily discussed in reference [32], and as it is thoroughly tackled throughout the present paper, topology changes—as loss of diffeomorphicity among differentiable manifolds—can take place also in the absence of critical points of the potential function.

Thus, coming to specific examples, in [49] the author tackled a system modelling in one-dimension a localization–delocalization transition of interfaces. This so-called Burkhardt model was also considered in [50] where the pinning potential was considered in a modified version. The absence of coincidence between topological and thermodynamical transitions was reported in both references [49, 50].

However, the Burkhardt model has two bad properties with respect to the conditions of the TH, formalized by the theorems in references [6, 28–30] and by the theorem proved in the present paper. In fact, the pinning potentials considered in references [49, 50] are singular as they need infinitely steep potential barriers to constrain the coordinates of the system on a semi-infinite positive line, and the configuration-space submanifolds are noncompact.

In [20], and also in [50], another one dimensional model was considered, the so-called Peyrard–Bishop model describing DNA denaturation. In reference [50] the authors considered also a modified version of the Peyrard–Bishop model, and for both versions of the model the topological transition is found at a critical value of the potential energy which does not correspond to critical energy value of the thermodynamic transition. The configuration space submanifolds corresponding to these models are *noncompact*, and the critical manifolds—containing only one critical point of infinite coordinates—are infinitely large. Again, these models are outside the domain of validity of the theorems in references [6, 28–30] and in the present paper.

Another model, allegedly disproving the topological theory, is the mean-field Berlin–Kac spherical model. In reference [51] the two cases of zero and non-zero external field were considered. In the former case there is a continuous phase transition and in the latter case there is no phase transition. The two cases did not display much difference when considered from the topological viewpoint. However, for this model there is a strong statistical ensemble inequivalence, in fact the continuous phase transition for zero external field is only predicted in the framework of canonical ensemble, whereas it is absent in the framework of microcanonical ensemble [52, 53] which is the reference framework where the topological theory is formulated. Thus there is no contradiction. Moreover, considering that the ergodic invariant measure of Hamiltonian flows is the microcanonical measure, and considering that the objective reality of any physical system is dynamics, the microcanonical ensemble has to be considered the fundamental statistical ensemble.

By the way, this is not the only case of this kind of ensemble inequivalence, for example, though working in the opposite way, the clustering phase transition in a self-gravitating N -body system found in the microcanonical ensemble framework is completely absent in the canonical ensemble [54].

In [16] the authors tackle a modified version of the Berlin–Kac model which is constrained with the introduction of a long-range correlation among the degrees of freedom. In spite of the claim that this model is the first case of short-range, confining potential where the phase transition is not entailed by a topology change in phase space or configuration space, the spherical constraint, by limiting the freedom of mutual independent variation of all the degrees of freedom, makes this model a *long-range* interaction one. And even though the spherical constraint is, so to speak, a weak constraint, it plays a crucial role because without it the model is trivial.

Another system apparently going against the topological theory is the mean-field ϕ^4 model. The phase transition point of this model does not correspond to the presence of critical points of the potential, that is, it has no topological counterpart [18, 55, 56]. Some correspondence between the topological and thermodynamic transitions was recovered for this model in reference [55] by introducing a suitable weakening of the TH. However, the mean-field ϕ^4 model undergoes a \mathbb{Z}_2 -symmetry-breaking phase transition, as in the case of the short range ϕ^4 model [32]. As a consequence, the reason why this system is not a counterexample of the topological theory is twofold: on the one side it violates the condition of asymptotic diffeomorphicity of the level sets Σ_v^{VN} put forward in the present work, and, on the other side, in the broken symmetry phase even in the absence of critical points of the potential in the $N \rightarrow \infty$ limit the splitting of the configuration space into two disjoint submanifolds is actually a major change of topology in correspondence with the phase transition point.

All the attempts at falsifying the TH are useful to better outline the domain of validity of the theory, which, on the other hand, is not intended to apply to *all* the possible phase transitions. The theory can leave outside its validity domain models with unbound and long-range potentials without being invalidated by this kind of systems.

Summarizing, the TH is coherent and now free of counterexamples, nevertheless the above mentioned alleged counterexamples have the merit of showing that some—perhaps much—work remains to be done, mainly in the case of long-range interactions. In fact, for example, for the exactly solvable model in references [46, 47], which is a mean-field XY model, thus with long-range interactions, a sharp topological transition in configuration space is clearly at the origin of the phase transition. At variance with the mean-field ϕ^4 model described by polynomial potentials, the mean-field XY model is described by a potential bounded from above. Whether and why this fact could explain the different conformity of these models to the topological description of their transitional behaviour is still a wide open question.

6. Concluding remarks

The present work is a substantial leap forward of the topological theory of phase transitions which was seriously undermined by the counterexample mentioned in the previous sections. The theory is rooted in the study of thermodynamical phase transitions from the viewpoint of microscopic Hamiltonian dynamics. As Hamiltonian flows can be identified with geodesics flows of suitable differentiable manifolds, it turned out that across a phase transition point these manifolds undergo major geometrical changes of topological origin. This is to remark that topology is, so to speak, naturally implied by the fundamental/dynamical approach and it is not just conjectured to play a role. The first important consequence of this approach is that the occurrence of a phase transition is not the consequence of a loss of analyticity of statistical measures but it is already encoded in the potential function describing the interactions among the degrees of freedom of a system. This makes the *thermodynamic limit dogma* no longer necessary neither from the conceptual side nor from the mathematical description side. And, of course, this is interesting when tackling phase transition phenomena in mesoscopic

and nanoscopic systems. Moreover, phase transitions phenomena in the absence of symmetry-breaking—and thus in the absence of an order parameter—have been successfully tackled in the topological framework, at present, at least in the case of a model with a gauge symmetry [57], for a 2D-model with an $O(2)$ symmetry undergoing a Kosterlitz–Thouless transition [58], and for the protein folding transition [59]. It is worth mentioning that in a recent paper [60]—partly based on some results given in [61]—a purely geometric theory of phase transitions has been put forward. In this work it is proposed that Bachmann’s classification of phase transitions [4, 5] for finite-size systems can be reformulated in terms of geometric properties of the energy level sets associated to a given Hamiltonian function; here the energy-derivatives of the entropy are associated to specific combinations of geometric curvature properties of the energy level sets. There is no contradiction between the geometric theory and the topological theory of phase transitions, mainly because sharp changes of the geometry of the leaves (energy level sets) of a foliation of phase space can be generically attributed to deeper changes of topological kind. However, the precise relationship between geometry and topology is given by theorems in differential topology and, unfortunately, there is only a few number of these theorems that can be constructively used (essentially the Gauss–Bonnet–Hopf, the Chern–Lashof, and Pinkall theorems [6]). Therefore the geometric approach has some practical advantage with the respect to the topological one in what curvature properties of the energy level sets can be always explicitly computed.

It is noteworthy that, in principle, the topological approach to classical phase transitions—addressed in the present work, as well as the geometric approach in reference [60]—can be extended to the treatment of quantum transitions by means of Wick’s analytic prolongation to imaginary times of the path-integral generating functional of quantum field theory, this allows to map a quantum system onto a formally classical one described by a classical partition function written with the euclidean Lagrangian action, on lattice to have a countable number of degrees of freedom [6].

Finally, recent developments of powerful computational methods in algebraic topology, like those of *persistent homology* [62, 63], provide the topological description of phase transitions with new useful constructive tools in addition to the existing concepts and methods of differential topology.

Acknowledgments

This work has been done within the framework of the project MOLINT which has received funding from the Excellence Initiative of Aix-Marseille University—A*Midex, a French ‘Investissements d’Avenir’ programme. This work was also partially supported by the European Union’s Horizon 2020 Research and Innovation Programme under Grant Agreement No. 964203 (FET-Open LINKS project). Roberto Franzosi acknowledges support by the QuantERA ERA-NET Co-fund 731473 (Project Q-CLOCKS) and the support by the National Group of Mathematical Physics (GNFM-INdAM). Matteo Gori thanks the financial support of DARPA (USA) for his long term visit at Howard University at Washington D.C. during which part of this work was done.

Data availability statement

The data that support the findings of this study are available upon reasonable request from the authors.

Appendix A. Uniform boundedness of \mathcal{R}

Let us now show how asymptotic diffeomorphicity entails uniform boundedness with N of the Ricci scalar curvature defined in equation (17), and using $\|\xi\| = \|\nabla V(\mathbf{q})\|^{-1}$,

$$\mathcal{R} = \frac{1}{N(N-1)} \left\{ -\Delta \log \|\xi\|^{-1} + \nabla \cdot [\Delta V(\mathbf{q}) \xi] \right\}, \tag{62}$$

where the second term in the rhs is

$$\nabla \cdot [\Delta V(\mathbf{q}) \xi] = [\nabla \Delta V(\mathbf{q})] \cdot \xi + [\Delta V(\mathbf{q})] \nabla \cdot \xi \tag{63}$$

$$= \Delta V(\mathbf{q}) \partial^i \left(\frac{\partial_i V(\mathbf{q})}{\|\nabla V(\mathbf{q})\|^2} \right) + \frac{\partial_j V(\mathbf{q})}{\|\nabla V(\mathbf{q})\|^2} \partial^j \partial^k \partial_k V(\mathbf{q}) \tag{64}$$

and using

$$\Delta V(\mathbf{q}) = \nabla \cdot [\nabla V(\mathbf{q})] = \nabla \cdot [\|\nabla V(\mathbf{q})\|^2 \xi] = \partial^i \left(\frac{\xi_i}{\|\xi\|^2} \right) \tag{65}$$

$$= \frac{1}{\|\xi\|^2} (\partial^i \xi_i) - \frac{4}{\|\xi\|^2} \frac{\xi_i}{\|\xi\|} \frac{\xi^j}{\|\xi\|} (\partial^i \xi_j) \tag{66}$$

after equations (6)–(8) all these terms are uniformly bounded in N , and so does $\nabla \cdot \xi$, moreover, the denominator $\|\xi\|^{-4} \sim N^2$ is compensated by the pre-factor $1/N(N-1)$. Therefore the second term in the rhs of equation (63) is also uniformly bounded in N . Then the first term of equation (63) is obtained by applying the operator $\xi \cdot \nabla$ to equation (66), and, after trivial algebra of the same kind of that leading to equation (66), one obtains a lengthy expression—containing mixed second order derivatives of the components of ξ —which are uniformly bounded under the assumption of asymptotic diffeomorphicity. On the other hand, for smooth and regularized potentials, if n is the coordination number of the potential, and \mathfrak{m}_0 is the maximum value of $\partial^i \partial_i V$, then $\Delta V(\mathbf{q})$ is bounded by $n \mathfrak{m}_0 N/[N(N-1)]$. By the same token, if \mathfrak{m}_1 is the maximum value of $\partial_j \partial^i \partial_i V$ then $\xi \cdot \nabla \Delta V(\mathbf{q})$ is uniformly bounded by $n \mathfrak{m}_1 B$, where B is the constant of equation (8).

Now, coming to the first term in the rhs of equation (62), that is $\partial^i (\partial_i \log \|\nabla V(\mathbf{q})\|)$, we have

$$\partial^i (\partial_i \log \|\xi\|^{-1}) = \partial^i (\|\xi\| \partial_i \|\xi\|^{-1}) = \partial^i \|\xi\| \partial_i \|\xi\|^{-1} + \|\xi\| \partial^i \partial_i \|\xi\|^{-1} \tag{67}$$

$$= -\frac{1}{\|\xi\|^2} (\partial_i \|\xi\|)^2 + \left[-\frac{1}{\|\xi\|} (\partial^i \partial_i \|\xi\|) + \frac{2}{\|\xi\|^2} (\partial_i \|\xi\|)^2 \right] \tag{68}$$

$$= -\frac{1}{\|\xi\|} (\partial^i \partial_i \|\xi\|) + \frac{1}{\|\xi\|^2} (\partial_i \|\xi\|)^2 \tag{69}$$

and

$$\frac{1}{\|\xi\|^2} (\partial_i \|\xi\|)^2 = \frac{1}{\|\xi\|^2} (\partial_i \sqrt{\xi_j \xi^j})^2 = \frac{\xi_j \xi^j}{\|\xi\|^4} (\partial_i \xi_j)^2 \tag{70}$$

which is uniformly bounded after equations (6)–(8) and the pre-factor $1/N(N - 1)$. Then for the first term in equation (69) we get

$$\begin{aligned} \frac{1}{\|\xi\|}(\partial^i \partial_i \|\xi\|) &= \frac{1}{\|\xi\|} \partial^i \left(\frac{\xi^j}{\|\xi\|} \partial_i \xi_j \right) \\ &= \frac{1}{\|\xi\|} \left[\frac{\xi^j}{\|\xi\|} \partial^i \partial_i \xi_j + \frac{(\partial_i \xi_j)^2}{\|\xi\|} - \frac{\xi_j \xi^j (\partial_i \xi_j)^2}{\|\xi\|^3} \right] \end{aligned} \quad (71)$$

which, under the same conditions mentioned above, is also uniformly bounded in N . The relevant consequence is that under the assumption of asymptotic diffeomorphicity of potential energy level sets, the scalar Ricci curvature \mathcal{R} in equation (62) is uniformly bounded in N and so do all the principal curvatures of the manifolds transformed under the action of the vector field ξ .

Appendix B. Lie derivatives of the vector field ξ

In the following we derive explicit expressions of the Lie derivatives of the one-parameter diffeomorphism-generating vector field ξ for a potential V in ‘critical points-free’ regions of configuration space $(\mathcal{X}, g_{\mathbb{R}^N})$ endowed with a Riemannian metric. Let (q_1, \dots, q_N) be a set of coordinates in configuration space. In what follows we shall refer to $\partial_i = \partial/\partial q^i$ so that $(\nabla V)_i = \partial_i V$ and the Hessian $(\text{Hess}V)_{ij} = \partial_{ij}^2 V$.

With these choices the divergence of the vector field $\zeta = \text{div}_{\mathbb{R}^N} \xi$ reads:

$$\text{div}_{\mathbb{R}^N} \xi = \frac{\Delta V}{\|\nabla V\|^2} - 2 \frac{\nabla V \cdot (\text{Hess}V \nabla V)}{\|\nabla V\|^4}, \quad (72)$$

where $\Delta(\cdot) = \sum_i^N \partial^i \partial_i(\cdot)$ is the Laplacian operator in the Euclidean configuration space and $\|\mathbf{X}\|^2 = g_{\mathbb{R}^N}(\mathbf{X}, \mathbf{X})$ is the Euclidean norm. As the Lie derivative operator along the flow generated by the vector field ξ is

$$\mathcal{L}_\xi(\cdot) = (\xi \cdot \nabla)(\cdot) = \sum_{i=1}^N \frac{\partial^i V}{\|\nabla V\|^2} \partial_i(\cdot) \quad (73)$$

the first derivative reads

$$\begin{aligned} \mathcal{L}_\xi(\zeta) &= \frac{\nabla V \cdot \nabla(\Delta V)}{\|\nabla V\|^4} - 2 \frac{(\nabla V \cdot \text{Hess}(V) \nabla V) \Delta V + 2 \|\text{Hess}V \nabla V\|^2 + \text{D}^3 V(\nabla V, \nabla V, \nabla V)}{\|\nabla V\|^6} \\ &\quad + 8 \frac{(\nabla V \cdot \text{Hess}V \nabla V)^2}{\|\nabla V\|^8}, \end{aligned} \quad (74)$$

where $\text{D}^3 V(\nabla V, \nabla V, \nabla V) = \partial_{ijk}^3 V \partial_i V \partial_j V \partial_k V$.

The second order derivative (with the aid of symbolic manipulation with Mathematica) reads

$$\begin{aligned}
 \mathcal{L}_\xi^{(ii)}(\zeta) = & \frac{\nabla(\Delta V) \cdot (\text{Hess}V\nabla V) + \nabla V \cdot (\text{Hess}(\Delta V)\nabla V)}{\|\nabla V\|^6} + \\
 & - 2 \left[\frac{\Delta V D^3 V(\nabla V, \nabla V, \nabla V) + 2\Delta V \|\text{Hess}V\nabla V\|^2 + 4(\text{Hess}V\nabla V) \cdot (\text{Hess}V\text{Hess}V\nabla V)}{\|\nabla V\|^8} \right. \\
 & + \left. \frac{7D^3 V(\text{Hess}V\nabla V, \nabla V, \nabla V) + D^4 V(\nabla V, \nabla V, \nabla V, \nabla V) + 3(\nabla V \cdot \text{Hess}V\nabla V)(\nabla V \cdot \nabla(\Delta V))}{\|\nabla V\|^8} \right] \\
 & + \frac{28(\nabla V\text{Hess}V\nabla V)[2\|\text{Hess}V\nabla V\|^2 + D^3 V(\nabla V, \nabla V, \nabla V)] + 12(\nabla V\text{Hess}V\nabla V)^2 \Delta V}{\|\nabla V\|^{10}} + \\
 & - 64 \frac{(\nabla V\text{Hess}V\nabla V)^3}{\|\nabla V\|^{12}} \tag{75}
 \end{aligned}$$

and the third order derivative (with the aid of symbolic manipulation with Mathematica) reads

$$\begin{aligned}
 \mathcal{L}_\xi^{(iii)}(\zeta) = & \frac{3\nabla V \cdot \text{Hess}(\Delta V)\text{Hess}V\nabla V + D^3 \Delta V(\nabla V, \nabla V, \nabla V) + D^3 V(\nabla V, \nabla V, \nabla(\Delta V))}{\|\nabla V\|^8} \\
 & + \frac{\nabla(\Delta V) \cdot \text{Hess}V\text{Hess}V\nabla V}{\|\nabla V\|^8} - 2 \left[\frac{4D^3 V(\nabla V, \nabla V, \nabla V)(\nabla V \cdot \nabla(\Delta V))}{\|\nabla V\|^{10}} \right. \\
 & + \frac{7D^4 V(\nabla V, \nabla V, \nabla V, \text{Hess}V\nabla V)}{\|\nabla V\|^{10}} \\
 & + \frac{15D^3 V(\nabla V, \nabla V, \text{Hess}V\text{Hess}V\nabla V) + 7\|D^3 V(\nabla V, \nabla V)\|^2 + 18D^3 V(\text{Hess}V\nabla V, \text{Hess}V\nabla V, \nabla V)}{\|\nabla V\|^{10}} \\
 & + \frac{4DV(\nabla V, \nabla V, \nabla V, \text{Hess}V\nabla V) + D^5 V(\nabla V, \nabla V, \nabla V, \nabla V, \nabla V) + 8(\nabla V \cdot \nabla(\Delta V))\|\text{Hess}V\nabla V\|^2}{\|\nabla V\|^{10}} \\
 & + \frac{8\|\text{Hess}V\text{Hess}V\nabla V\|^2 + 7D^3 V(\nabla V, \nabla V, \text{Hess}V\nabla V)\Delta V}{\|\nabla V\|^{10}} \\
 & + \frac{\Delta V D^4 V(\nabla V, \nabla V, \nabla V, \nabla V) + 4\Delta V(\text{Hess}V\nabla V) \cdot \text{Hess}V\text{Hess}V\nabla V}{\|\nabla V\|^{10}} \\
 & + \left. \frac{6(\text{Hess}V \cdot \text{Hess}V\nabla V)(\nabla V \cdot \text{Hess}(\Delta V)\nabla V) + 6(\text{Hess}V \cdot \text{Hess}V\nabla V)(\nabla(\Delta V) \cdot \text{Hess}V\nabla V)}{\|\nabla V\|^{10}} \right] \\
 & + 4 \left[\frac{7(D^3 V(\nabla V, \nabla V, \nabla V))^2 + 28D^3 V(\nabla V, \nabla V, \nabla V)\|\text{Hess}V\nabla V\|^2}{\|\nabla V\|^{12}} \right. \\
 & + \frac{10\Delta V D^3 V(\nabla V, \nabla V, \nabla V)(\nabla V \cdot \text{Hess}V\nabla V)}{\|\nabla V\|^{12}} \\
 & + \frac{28\|\text{Hess}V\nabla V\|^4 + 20\Delta V\|\text{Hess}V\nabla V\|^2(\nabla V \cdot \text{Hess}V\nabla V)}{\|\nabla V\|^{12}} \\
 & + \frac{(\nabla V \cdot \text{Hess}V\nabla V)[7D^3 V(\nabla V, \nabla V, \text{Hess}V\nabla V)]}{\|\nabla V\|^{12}} \\
 & + \frac{11D^4 V(\nabla V, \nabla V, \nabla V, \nabla V) + 44(\text{Hess}V\nabla V) \cdot (\text{Hess}V\text{Hess}V\nabla V)}{\|\nabla V\|^{12}} \\
 & + \left. \frac{15(\text{Hess}V \cdot \text{Hess}V\nabla V)(\nabla V \cdot \nabla(\Delta V))}{\|\nabla V\|^{12}} \right] + -8 \left[\frac{59D^3 V(\nabla V, \nabla V, \nabla V)(\nabla V \cdot \text{Hess}V\nabla V)^2}{\|\nabla V\|^{14}} \right. \\
 & + \left. \frac{(\nabla V \cdot \text{Hess}V\nabla V)^2[118\|\text{Hess}V\nabla V\|^2 + 15\Delta V(\nabla V \cdot \text{Hess}V\nabla V)]}{\|\nabla V\|^{14}} \right] + 768 \frac{(\nabla V \cdot \text{Hess}V\nabla V)^4}{\|\nabla V\|^{16}}. \tag{76}
 \end{aligned}$$

ORCID iDs

Marco Pettini  <https://orcid.org/0000-0003-1561-4390>

References

- [1] Yang C N and Lee T D 1952 Statistical theory of equations of state and phase transitions: I. Theory of condensation *Phys. Rev.* **87** 404–9
Lee T D and Yang C N 1952 Statistical theory of equations of state and phase transitions: II. Lattice gas and Ising model *Phys. Rev.* **87** 410–9
- [2] Georgii H O 2011 A comprehensive account of the Dobrushin–Lanford–Ruelle theory and of its developments can be found *Gibbs Measures and Phase Transitions* 2nd edn (Berlin: de Gruyter and Co)
- [3] Gross D H E 2001 *Microcanonical Thermodynamics: Phase Transitions in ‘Small’ Systems* (Singapore: World Scientific)
- [4] Bachmann M 2014 *Thermodynamics and Statistical Mechanics of Macromolecular Systems* (Cambridge: Cambridge University Press)
- [5] Qi K and Bachmann M 2018 Classification of phase transitions by microcanonical inflection-point analysis *Phys. Rev. Lett.* **120** 180601
- [6] Pettini M 2007 *Geometry and Topology in Hamiltonian Dynamics and Statistical Mechanics (IAM Series vol 33)* (Berlin: Springer)
- [7] Pettini M 1993 Geometrical hints for a nonperturbative approach to Hamiltonian dynamics *Phys. Rev. E* **47** 828
- [8] Carlsson G, Gorham J, Kahle M and Mason J 2012 Computational topology for configuration spaces of hard disks *Phys. Rev. E* **85** 011303
- [9] Baryshnikov Y, Bubenik P and Kahle M 2013 Min-type Morse theory for configuration spaces of hard spheres *Int. Math. Res. Not.* **2014** 2577–92
- [10] Eriçok O B, Ganesan K and Mason J K 2021 Configuration spaces of hard spheres (arXiv:2106.10315v3 [cond-mat.stat-mech])
- [11] Brody D C, Hook D W and Hughston L P 2007 Quantum phase transitions without thermodynamic limits *Proc. R. Soc. A.* **463** 2021
- [12] Buonsante P, Franzosi R and Smerzi A 2015 Phase transitions at high energy vindicate negative microcanonical temperature (arXiv:1506.01933)
- [13] Volovik G E 2007 Quantum phase transitions from topology in momentum space *Quantum Analogues: From Phase Transitions to Black Holes and Cosmology* (Berlin: Springer) pp 31–73
- [14] Angelani L, Di Leonardo R, Parisi G and Ruocco G 2001 Topological description of the aging dynamics in simple glasses *Phys. Rev. Lett.* **87** 055502
- [15] Debenedetti P G and Stillinger F H 2001 Supercooled liquids and the glass transition *Nature* **410** 259
- [16] Risau-Gusman S, Ribeiro-Teixeira A C and Stariolo D A 2005 Topology, phase transitions, and the spherical model *Phys. Rev. Lett.* **95** 145702
- [17] Santos F A N, da Silva L C B and Coutinho-Filho M D 2017 Topological approach to microcanonical thermodynamics and phase transition of interacting classical spins *J. Stat. Mech.* **013202**
- [18] Garanin D A, Schilling R and Scala A 2004 Saddle index properties, singular topology, and its relation to thermodynamic singularities for a ϕ^4 mean-field model *Phys. Rev. E* **70** 036125
- [19] Cimasoni D and Delabays R 2019 The topological hypothesis for discrete spin models *J. Stat. Mech.* **033216**
- [20] Grinza P and Mossa A 2004 Topological origin of the phase transition in a model of DNA denaturation *Phys. Rev. Lett.* **92** 158102
- [21] Becker O M and Karplus M 1997 The topology of multidimensional potential energy surfaces: theory and application to peptide structure and kinetics *J. Chem. Phys.* **106** 1495
- [22] Caiani L, Casetti L, Clementi C and Pettini M 1997 Geometry of dynamics, Lyapunov exponents, and phase transitions *Phys. Rev. Lett.* **79** 4361
- [23] Caiani L, Casetti L, Clementi C, Pettini G, Pettini M and Gatto R 1998 Geometry of dynamics and phase transitions in classical lattice ϕ^4 theories *Phys. Rev. E* **57** 3886
- [24] Brooks C L, Onuchic J N and Wales D J 2001 Taking a walk on a landscape *Science* **293** 612–3

- [25] Wales D J 2001 A microscopic basis for the global appearance of energy landscapes *Science* **293** 2067–70
- [26] Santos F A N, Raposo E P, Coutinho-Filho M D, Copelli M, Stam C J and Douw L 2019 Topological phase transitions in functional brain networks *Phys. Rev. E* **100** 032414
- [27] Casetti L, Pettini M and Cohen E G D 2000 Geometric approach to Hamiltonian dynamics and statistical mechanics *Phys. Rep.* **337** 237–341
- [28] Franzosi R and Pettini M 2004 Theorem on the origin of phase transitions *Phys. Rev. Lett.* **92** 060601
- [29] Franzosi R, Pettini M and Spinelli L 2007 Topology and phase transitions: I Preliminary results *Nucl. Phys. B* **782** 189
- [30] Franzosi R and Pettini M 2007 Topology and phase transitions: II. Theorem on a necessary relation *Nucl. Phys. B* **782** 219
- [31] Kastner M and Mehta D 2011 Phase transitions detached from stationary points of the energy landscape *Phys. Rev. Lett.* **107** 160602
- [32] Gori M, Franzosi R and Pettini M 2018 Topological origin of phase transitions in the absence of critical points of the energy landscape *J. Stat. Mech.* **093204**
- [33] Hirsch M W 1976 *Differential Topology* (Berlin: Springer)
- [34] Thorpe J A 1979 *Elementary Topics in Differential Geometry* (Berlin: Springer) p 55
- [35] Zhou Y 2013 A simple formula for scalar curvature of level sets in euclidean spaces (arXiv:1301.2202[math.DG])
- [36] Sormani C 2012 How Riemannian manifolds converge *Metric and Differential Geometry (Progress in Mathematics vol 297)* ed X Dai and X Rong (Basel: Birkhäuser)
- [37] Palais R S and Terng C 1988 *Critical Point Theory and Submanifold Geometry* (Berlin: Springer)
- [38] Federer H 1969 *Geometric Measure Theory* (Berlin: Springer)
- [39] Laurence P 1989 On the convexity of geometric functional of level for solutions of certain elliptic partial differential equations *Z. Angew. Math. Phys.* **40** 258
- [40] Brémaud P 2001 *Markov Chains* (Berlin: Springer) ch 7
- [41] See lemma 3.1 Alvarez-Vizoso J, Kirby M and Peterson C 2018 Integral invariants from covariance analysis of embedded Riemannian manifolds (arXiv:1804.10425v1)
- [42] Khinchin A I 1949 *Mathematical Foundations of Statistical Mechanics* (New York: Dover)
- [43] Consider a family of random variables $\{X_k\}$, if X_i and X_j satisfy finite-range dependence, i.e. there is a positive integer n_0 such that if $|i - j| > n_0$ then X_i and X_j are independent. A law of large numbers for X_j can then be easily proved.
- [44] Caiani L, Casetti L and Pettini M 1998 Hamiltonian dynamics of the two-dimensional lattice model *J. Phys. A: Math. Gen.* **31** 3357
- [45] Casetti L, Cohen E G D and Pettini M 1999 Topological origin of the phase transition in a mean-field model *Phys. Rev. Lett.* **82** 4160
- [46] Casetti L, Cohen E G D and Pettini M 2002 Exact result on topology and phase transitions at any finite N *Phys. Rev. E* **65** 036112
- [47] Casetti L, Pettini M and Cohen E G D 2003 Phase transitions and topology changes in configuration space *J. Stat. Phys.* **111** 1091
- [48] Angelani L, Casetti L, Pettini M, Ruocco G and Zamponi F 2005 Topology and phase transitions: from an exactly solvable model to a relation between topology and thermodynamics *Phys. Rev. E* **71** 036152
- [49] Kastner M 2004 Unattainability of a purely topological criterion for the existence of a phase transition for nonconfining potentials *Phys. Rev. Lett.* **93** 150601
- [50] Angelani L, Ruocco G and Zamponi F 2005 Relationship between phase transitions and topological changes in one-dimensional models *Phys. Rev. E* **72** 016122
- [51] Ribeiro Teixeira A C and Stariolo D A 2004 Topological hypothesis on phase transitions: the simplest case *Phys. Rev. E* **70** 016113
- [52] Kastner M and Schnetz O 2006 On the mean-field spherical model *J. Stat. Phys.* **122** 1195
- [53] Kastner M 2006 Topological approach to phase transitions and inequivalence of statistical ensembles *Physica A* **359** 447
- [54] Cerruti-Sola M, Cipriani P and Pettini M 2001 On the clustering phase transition in self-gravitating N -body systems *Mon. Not. R. Astron. Soc.* **328** 339
- [55] Andronico A, Angelani L, Ruocco G and Zamponi F 2004 Topological properties of the mean-field ϕ^4 model *Phys. Rev. E* **70** 041101
- [56] Baroni F 2002 Phase transitions and topological changes in configuration space for mean-field models *Laurea Thesis in Physics* University of Florence

- [57] Pettini G, Gori M, Franzosi R, Clementi C and Pettini M 2018 On the origin of phase transitions in the absence of symmetry-breaking *Physica A* **516** 376
- [58] Bel-Hadj-Aissa G, Gori M, Franzosi R and Pettini M 2021 Geometrical and topological study of the Kosterlitz–Thouless phase transition in the XY model in two dimensions *J. Stat. Mech.* **023206**
- [59] Di Cairano L, Capelli R, Bel-Hadj-Aissa G and Pettini M 2022 Topological origin of protein folding transition <https://hal.archives-ouvertes.fr/hal-03504864> (Accessed 26 June 2022)
- [60] Di Cairano L 2022 The geometric theory of phase transitions *J. Phys. A: Math. Theor.* **55** 27LT01
- [61] Gori M 2022 Configurational microcanonical statistical mechanics from Riemannian geometry of equipotential level sets (arXiv:2205.14536)
- [62] Carlsson G and Zomorodian A 2005 Persistent homology—a survey *Discrete Comput. Geom.* **33** 249
- Carlsson G 2009 Topology and data *Bull. Am. Math. Soc.* **2** 255
- [63] Donato I, Gori M, Pettini M, Petri G, De Nigris S, Franzosi R and Vaccarino F 2016 Persistent homology analysis of phase transitions *Phys. Rev. E* **93** 052138 and references therein for persistent homology