

PERSPECTIVE

Operational perspectives for biodiversity indicators

Helmut Hillebrand^{1,2,3}  | Jan-Claas Dajka^{2,3}  | Maurits Halbach^{2,3} | Anika Happe¹ | Ralf Röchert³ | Ralf Seppelt^{4,5}  | Josef Settele^{4,6} | Markus Weitere⁷ | Marten Winter⁵  | Yves Zinggrebe⁴ | Dorothee Hodapp^{2,3}

¹School of Mathematics and Science, Institute for Chemistry and Biology of the Marine Environment (ICBM), Carl von Ossietzky Universität Oldenburg, Oldenburg, Germany; ²Helmholtz-Institute for Functional Marine Biodiversity at the University of Oldenburg [HIFMB], Oldenburg, Germany; ³Alfred Wegener Institute, Helmholtz-Centre for Polar and Marine Research [AWI], Bremerhaven, Germany; ⁴Helmholtz-Centre for Environmental Research—UFZ, Leipzig, Germany; ⁵Luxembourg Centre for Socio-Environmental Systems (LCSES), Luxembourg University, Esch-sur-Alzette, Luxembourg; ⁶Deutsches Zentrum für Integrative Biodiversitätsforschung [iDiv] Halle-Jena-Leipzig, Leipzig, Germany and ⁷Department River Ecology, Helmholtz-Centre for Environmental Research—UFZ, Magdeburg, Germany

Correspondence

Helmut Hillebrand

Email: helmut.hillebrand@hifmb.de**Funding information**

HORIZON EUROPE Climate, Energy and Mobility, Grant/Award Number: 101060072

Handling Editor: Max R Lambert

Abstract

1. Smart biodiversity indicators are needed not only for assessing and managing biodiversity (change) but also for informing target setting for environmental policies. This need has created a plethora of indicator frameworks, which are widely debated in terms of their design and usefulness. We propose that these discussions would benefit from more clearly separating different types of indicators and more operationally linking them to management targets and biodiversity goals.
2. Decision makers often consider the multitude of biodiversity indicators as being *complicated*, whereas scientists emphasize that they barely reflect how *complex* biodiversity is. It is therefore important to differentiate clearly between indicators for *diagnosing* (the drivers of) biodiversity change and indicators for *steering* policies. While the former must be scalable in time and space and reflect the full complexity of biodiversity, the latter need long-term visions and simplicity.
3. *Management targets* such as proportions of protected areas or mitigation of biodiversity drivers have the advantage of being operation-oriented, well quantifiable and immediately responsive to change. Achieving a management target, however, does not necessarily equal achieving a biodiversity goal. Likewise, *management indicators* that track how well management targets are achieved cannot replace diagnostic or steering indicators.
4. Solution: *Biodiversity goals* such as 'bending the curve' can only be achieved by closely linking diagnosis, steering and management, while accepting that their distinct design principles need to be addressed. Being aware and accepting such tailored types of different indicators and management targets could strongly improve the interplay between science and policy to reverse negative biodiversity trends in the future.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2025 The Author(s). *Ecological Solutions and Evidence* published by John Wiley & Sons Ltd on behalf of British Ecological Society.

KEYWORDS

assessment, biodiversity indicators, biodiversity strategy and actions plans (NBSAP), complexity, cross-realm, Kunming-Montreal global biodiversity framework (KM-GBF), management target, operational, steering

1 | INTRODUCTION

Biodiversity change is a global political concern as Earth's habitability for humans ultimately depends on biological processes, which in turn depend on the identity and variation of marine, freshwater and terrestrial life. In the past decades, many political instruments have been developed to address this concern ranging from the Kunming-Montreal Global Biodiversity Framework (KM-GBF; CBD, 2022b) over supranational regulations such as the European Union' Biodiversity Strategy (EU, 2021) and Framework Directives (EU, 2000, 2010, 2017) to national biodiversity strategies and action plans (NBSAPs). By mid-August 2025, 55 parties of the Convention on Biological Diversity (CBD) had submitted the requested NBSAP within the KM-GBF framework (Affinito et al., 2024; CBD, 2022a). In addition, also the private sector (economic and financial companies) started disclosing how they integrate biodiversity into their decision-making and risk management (TNFD, 2023).

All these efforts share a similar narrative of 'bending the curve': we need to invest in conservation to halt the current biodiversity loss and additionally in transformation to invert the negative trend into a biodiversity positive future, for example, through restoration (Leadley et al., 2022; Leclère et al., 2020). This narrative requires diagnosing 'underlying causes of biodiversity loss', managing 'determinants of transformative change' and steering 'options for achieving the 2050 Vision for Biodiversity' (IPBES, 2024).

Bending the curve requires smart and meaningful indicators that depict the status and trend of biodiversity as well as measure the progress made towards formulated biodiversity goals. As a consequence, a wide array of indicator frameworks exists (Feld et al., 2009; Pereira et al., 2013; Teixeira et al., 2016; Affinito et al., 2025; BDM (Coordination Office), 2014), where each framework consists of multiple indicators (which themselves may have multiple individual metrics) that capture different aspects of biodiversity and its change. These provide us with a plethora of information, which can cause confusion or be perceived as too complex, especially as different indicators can yield vastly different results despite the same underlying biodiversity change (Hill et al., 2016).

The KM-GBF's vision for 2050 formulates a joint goal for biodiversity, which is shared among scientists, practitioners and biodiversity-aware decision makers. However, scientists are expected to understand how and why biodiversity is changing, decision makers are expected to set realistic and useful goals and practitioners are expected to understand which management actions allow bending the curve. We base our perspective on the notion that each of these demands needs operational indicators, but these needs are so divergent that no single indicator (not even an indicator framework) can serve all at once. At

the same time, it is impossible to cover the entire information, as the Taskforce on Nature-related Financial Disclosures (TNFD, 2023) alone reported over 3000 unique biodiversity-related metrics.

In this perspective, we address this dilemma by first addressing the common notion that diagnosing biodiversity is (too) complex (Section 2). We then argue that explicitly acknowledging essential functional differences between diagnostic and steering indicators helps design assessments and strategy development (Section 3). Furthermore, we argue that management targets are derived from diagnosis and steering, but they require different indicators (management indicators), which cannot replace either diagnosis or steering indicators (Section 4). Finally, we discuss why the development of indicator frameworks for all three purposes needs to happen in an integrated manner across disciplines, user groups as well as realms (Section 5).

2 | COMPLEX, BUT NOT COMPLICATED

Biodiversity has many facets encompassing, for example, taxonomic, evolutionary and functional differences, biotic interactions, the abundance in populations and the intraspecific variability between individuals within populations, the species composition of communities and the size as well as functioning of ecosystems (Antonelli, 2022; Pereira et al., 2013). Each of these facets requires different ways of measuring their status and trends, but at the same time, each metric has its own biases and limits, both statistically and biologically (discussed elsewhere, e.g. Magurran & McGill, 2011; Chao et al., 2014). Moreover, even for a single driver (e.g. warming) in a single ecosystem type (e.g. lake), each of these facets can change independently or in concert. Higher temperatures can reduce the size of individual organisms within populations (Albini et al., 2025), alter the size and intraspecific composition of populations (van Dorst et al., 2019), lead to a temporal mismatch between interacting species (Kharouba et al., 2018), change species and trait composition by colonizations, extirpations and shifts in relative abundance (Hillebrand et al., 2018; Khaliq et al., 2024), shift the entire functioning of the system (Yvon-Durocher et al., 2010) and even alter the size of the ecosystem itself (Woolway et al., 2022). This long list is still non-exhaustive as it ignores links between ecosystems (lake-catchment) and the interaction of warming with other anthropogenic drivers (nutrients, pollution, land use).

Thus, the numerous ways in which biodiversity can change amount to a seemingly overwhelming complexity, which is often seen as an impediment to verify and report on biodiversity change in a way that mandates political action (Ekardt et al., 2023). While the

science–policy interface on climate change relies on temperature as an easily communicated, physically well-defined indicator, even the most consistent biodiversity indicator frameworks such as the essential biodiversity variables (Pereira et al., 2013) deal with multiple, in part abstract metrics.

Before we deal with the need to capture this complexity for diagnosis, but to simplify it for steering (Section 3), we would like to emphasize that the complex nature of biodiversity is not a nuisance, but rather its main asset. This complexity ensures the functioning of the biosphere and thus the provisioning of services forming the basis for human life on Earth (e.g. climate, food, oxygen, raw materials). Life on Earth has evolved into an incredible and fascinating array of solutions for the main necessities to obtain matter and energy, grow, reproduce and disperse. All of these facets are part of a ‘tangled bank’ (Darwin, 1859) that allows ecosystems to function and more diverse ecosystems to be more resilient (Levin, 1998). It is this complexity that ultimately provides the conditions for humans to live on Earth, and it will likely be humanity’s greatest ally in a rapidly changing climate, as it enables ecosystem resilience through the unprecedented rate of reorganization and adaptation required from species and ecological communities (Pörtner et al., 2021).

Any attempt to report biodiversity change in a single framework (or worse, a single number) must, therefore, fail as it ignores the multifaceted nature of biodiversity (spatial, temporal, community, species, functional, genetic). At the same time, reporting on several independent biodiversity facets often generates an ‘incoherent’ impression of biodiversity change, which has given biodiversity indicators the reputation of being too ‘complicated’ for decision-making and management (Sobkowiak, 2023). We argue that these different needs mandate two different indicator frameworks, one for diagnosing biodiversity change, which needs to encompass all complexity, and one for application in political steering and science communication, which requires a reduction of the complexity.

3 | DIAGNOSTIC AND STEERING INDICATORS

Diagnosis and steering have different aims, different audiences and therefore fundamentally different design principles (Figure 1). Diagnosis aims at system understanding. Therefore, diagnostic indicator frameworks need to detect the full dimensionality of biodiversity change, which requires measuring the ecological status and trends of multiple biodiversity facets and combining these with causal inference and attribution to drivers (Gonzalez, Chase, & O’Connor, 2023). To do so, the diagnosis of status and trends requires multidimensional indicator frameworks that reflect the multiple facets of biodiversity (Pereira et al., 2013).

Diagnostic indicators address mainly an audience conducting assessments, while steering indicators need to be communicable beyond those groups as they are defined in socio-political contexts and reflect a ‘policy-relevant phenomenon used to set environmental

goals and evaluate their fulfilment’ as their indicandum (Silva del Pozo et al., 2023). ‘Bending the curve’, for instance, is such a goal formulation, and the KM-GBF headline indicators give numerous examples of steering indicators for achieving biodiversity goals (<https://www.gbf-indicators.org/>). The first listed indicator ‘Red List of Ecosystems’ is a great example as an increase in the number of red-listed ecosystems helps to realize that our policies steer away from the accepted goal, but it does not help in diagnosing the causes. In fact, most headline indicators in the KM-GBF such as red lists of species or ecosystems, the extent of natural ecosystems or the proportion of small populations ($N < 500$) are steering indicators (Hébert et al., 2025). Some indicators try to aggregate steering and diagnostic aspects, such as the Living Planet Index, which however is heavily debated (Almond et al., 2020; Hébert & Gravel, 2023; Leung et al., 2020; Toszogyova et al., 2024).

Since steering indicators by definition have a normative component, they need to be relevant over longer time-scales and reflect the spatial scale of the goals, which tend to be regional (i.e. often bound to political borders such as nations or federal states) or global (KM-GBF). By contrast, diagnostic indicators mainly need short-term sensitivity and responsiveness to drivers and mainly operate at the local to regional scales of monitoring. In terms of responsiveness, indicators can be differentiated as leading, coincident and lagging (Stevenson et al., 2021). In this context, leading indicators respond before biodiversity change manifests (early warning), while coincident and lagging indicators react during or after changes in biodiversity, respectively.

Steering indicators should react at least simultaneously with the change of the variable of interest, but preferably they allow preventive adjustment of the steering, that is, be leading indicators in order to prevent certain developments (Figure 1). In this respect, many headline indicators fail as they show very slow responses to change. It takes years to decades to detect a deterioration of red list status for mammal and bird species (Piipponen-Doyle et al., 2021) or to monitor and report changing numbers of newly established introduced species (McGeoch et al., 2023). In the opposite direction, these indicators will also lag behind conservation success (Watts et al., 2020) and thus may trigger the counterfactual conclusion that conservation or restoration efforts are not working.

The same is true for diagnostic indicators that often de facto lag behind the changes they describe because biodiversity does not immediately respond to changes in environmental conditions. For example, extinction debt delays the local loss of a species as long as individuals persist despite declining abundances, while immigration credit delays the local gain of a species, for example, due to dispersal limitation, and both affect the number of species monitored over time (Jackson & Sax, 2010; Kuczynski et al., 2023).

As a result, while leading or coincident indicators would be preferable, many changes in biodiversity we measure today reflect past environmental or anthropogenically triggered shifts. This inertia does not only inhibit understanding biodiversity change and its attribution to drivers but also fails to deliver the




FIGURE 1 Different types of indicators for biodiversity status and trends, goals and management and their logical connections. Diagnostic indicators serve to assess biodiversity and understand its change. They deliver information to the societal and political discussion of biodiversity goals. Goals cannot be monitored by understanding, but need steering indicators, which also need feedback in the decision-making process. The steering decisions lead to management targets; the efficiency of this management needs its own indicators leading to, for example, conservation and restoration actions.

urgency needed regarding biodiversity action. The KM-GBF is a unique—and perhaps the only—opportunity to establish a global biodiversity observation system that can be designed to fulfil both diagnosis and steering needs (Affinito et al., 2025; Gonzalez, Vihervaara, et al., 2023). Affinito et al. (2025) concluded that the success of the KM-GBF will depend on the ambition of the parties to cover more than the so-called headline indicators and to embark on a joint and global learning exercise as 12% of the KM-GBF elements are still lacking indicators. For example, Target 10 ('Enhance Biodiversity and Sustainability in Agriculture, Aquaculture, Fisheries, and Forestry') so far has no fisheries or aquaculture headline indicator.

Despite the different audiences and design principles, it will be crucial that assessment experts, political and societal actors co-design biodiversity goals together to ensure that steering indicators lead to the same conclusions as the higher resolved diagnostic indicators, which they should be based on (Figure 1). The good news is that science is well prepared to take on this task. In a recent systematic review of the marine literature, Dajka et al. (2025) found that the indicators discussed in the KM-GBF context not only surprisingly well cover all major categories of essential biodiversity variables but they also align with the foci of the last decade of marine biodiversity research. Based on >250 scientific papers and 29 biodiversity indicators, species populations, community composition, ecosystem structure and ecosystem functioning received similar attention in science and policy; only genetic and trait-based biodiversity was underrepresented in the indicators. Hence, enough evidence exists in order to align diagnostics and steering.

The bad news is that monitoring for diagnosis requires significant additional efforts and money. As a result, management targets are often formulated and evaluated by their own management

indicators. However, these often start their own trajectories independently of the actual biodiversity trends.

4 | MANAGEMENT INDICATORS ARE NOT BIODIVERSITY INDICATORS

In the context of this article, we use the term management in a broad sense to comprise any human measure that is intended to bring the status of biodiversity closer to a predefined goal (Figure 1). Management is often guided by its own explicit targets ('30% area protected by 2030'), which define steps towards achieving the long-term overall goals. Management targets then themselves are tracked by indicators, for example, by recording the progress towards these targets quantitatively (km² or % area protected). Such management indicators are highly attractive because they are well-defined, often quantitative, easy to understand and their trajectories can be tracked comparably fast (Figure 1). For example, every newly established protected area brings the indicator '% area protected' closer to the predefined management target.

Management indicators are also needed for biodiversity credits, which have recently been proposed as means to fund conservation and restoration measures (Antonelli et al., 2024). This proposal has quickly led to the identification of a range of problems, one of which is the need to measure the impact of these credits on management goals and biodiversity status (Law, 2025; Wauchope et al., 2024).

It is important to acknowledge, though, that despite the 'ease' of tracking management targets, they cannot replace steering or diagnostic indicators. In fact, over-reliance on management targets can obscure the real drivers of biodiversity loss and jeopardize conservation efforts (Lemieux et al., 2019). To reduce climate change

impacts on targets, the EU has set explicit targets for the proportion of biofuel in energy consumption (EU, 2018), but given the potential negative impact of biofuel production on biodiversity (Tudge et al., 2021), focusing only on this target without diagnostic indicators of biodiversity change will have negative consequences for biodiversity. Generally, if decision makers focus solely on meeting numerical targets (e.g. increasing protected area coverage), they risk neglecting critical factors like habitat quality, ecosystem connectivity or species interactions in creating actions. A protected area, for instance, does not automatically lead to species recovery if it lacks effective management. Similarly, an increase in forest cover does not always equate to higher biodiversity if the new forests are monocultures or degraded habitats. Biodiversity goals should be formulated in outcome space (e.g. tropical coral reefs shall persist), not as a management target (x % of coral reefs protected). Mixing these two concepts may have highly negative consequences (see Fisher et al., 2024 for a more in-depth discussion).

Thus, management targets and their indicators are essential tools for conservation, but they cannot substitute diagnostic and steering biodiversity indicators. To genuinely address biodiversity loss, management actions must be closely linked to indicators that capture the complexity of ecosystems and long-term trends. A well-integrated approach that combines concrete conservation measures with robust biodiversity assessments is necessary to ensure lasting and effective progress.

5 | BENDING THE CURVES TOGETHER

We wrote this perspective based on a notion of opportunity and urgency. The KM-GBF and the coincident agreement on Biodiversity Beyond National Jurisdiction (<https://www.un.org/bbnjagreement/en>) are the most ambitious multilateral agreements to safeguard the biological basis of the Earth system. At the same time, the rate of nature destruction, climate change and biodiversity loss—in concert with the knowledge of how time-lagged biodiversity responds to these changes—creates a feeling of immediacy that stands in stark contrast to how slowly humanity is making progress towards biodiversity protection and restoration. While the complexity of biodiversity is often mentioned as an impediment to faster action, the implementation of measures is often impeded by the complexity of governance structures organizing the interactions of actors and regulations across levels and sectors (Paavola et al., 2009). Developing indicator frameworks should be a coordinated, joined and concerted effort across disciplines, realms (terrestrial, freshwater, marine) and spatial units (countries). In fact, however, we see multiple independent developments and little prospect for change once a framework has been adopted.

We argue here that biodiversity goals such as ‘bending the curve’ can only be achieved by closely linking diagnosis, steering and management. Scientists, decision makers and managers face the challenge that this interlinkage needs to happen *simultaneously* to the development of assessments, strategies and actions. The

development of indicator frameworks addressing diagnosis, steering and management needs to reflect the clear logical dependencies between them, but at the same time account for the differing functionalities and design principles of the different indicator types. Steering indicators do not enable diagnosis and attribution, just as management indicators cannot replace steering. Outcome goals ultimately require both valid steering indicators and an in-depth diagnosis to securely identify and adequately address root causes for biodiversity change (e.g. warming, acidification, eutrophication).

We agree with Affinito et al. (2025) that scientists, governments and societies embark on an unprecedented learning exercise to cope with the challenge of bending the curve. Conceptualizing the entire bandwidth from system understanding to effective management more clearly will improve the observation and monitoring we need to achieve our biodiversity goals and the indicator framework derived from these observations. Furthermore, if scientists, practitioners and decision makers can agree that all indicators are (and will remain) imperfect, we can stop binding resources in the search for a perfect solution. This ‘holy grail’ of biodiversity indications does not exist, and we need to move from more research on indicators to more attribution, steering and action. Otherwise, biodiversity research and decision-making will end up in a ‘trap’ comparable to that of climate change research. There, science continues to pile up evidence on man-made climate change, while enough evidence on causes and needed actions was established long ago, even with stakeholders that had contrary agendas (Supran et al., 2023). We consider emphasizing uncertainty, complexity and knowledge gaps as important scientific contribution, but this must not delay actions towards sustaining our biosphere.

As this biosphere is highly interconnected between terrestrial, marine or freshwater, we are negatively surprised by how often scientific and policy discussions around biodiversity still happen within single realms. This is inadequate from diagnosis, steering and management perspectives, as most changes in biodiversity and ecosystems are driven by processes across multiple realms. Neither climate change nor altered biogeochemical cycles stop at ecosystem boundaries, which are additionally linked through organisms’ life cycles and reciprocal subsidies (Hermoso et al., 2021; Scherer-Lorenzen et al., 2022). Last but not least, humans telecouple realms and regions of consuming and producing countries, leading to a shared responsibility of biodiversity outcomes (Zinggrebe et al., 2024). Societal transformation is only possible on land, where humans live, but must target sustainability together across realms.

6 | SUMMARY

We argue that the discussion on biodiversity indicators would profit from (i) allowing diagnostic indicators to capture as many different biodiversity facets as needed, (ii) formulating outcome-oriented steering indicators to guide public discussion and policy making and (iii) avoiding the tracking of management targets as a replacement for biodiversity assessments. It would improve communication

about biodiversity change if different indicators were mapped along their different purposes, and it would improve biodiversity research and management if the interrelationships between these indicator types were well understood. Any reporting of data leading to indicators across spatial scales, realms and taxa needs interoperable machinery to allow the leading or at least 'real time' diagnostics. A major but highly needed task across all countries, given recent developments of rebuilding national barriers, increasing federalism and reduced funding in global biodiversity conservation actions.

AUTHOR CONTRIBUTIONS

Dorothee Hodapp and Helmut Hillebrand planned the activity. Marten Winter, Anika Happe, Markus Weitere and Helmut Hillebrand outlined the manuscript; all authors wrote the first draft together and contributed to the editing of the manuscript.

ACKNOWLEDGEMENTS

The authors thank SynCom, the synthesis and communication platform within the Helmholtz Research Field 'Earth and Environment', for supporting the workshops on 'Operational Biodiversity Targets' that led to this manuscript. HH and AH acknowledge further funding by the European Union HORIZON EUROPE program ACTNOW: Advancing understanding of Cumulative Impacts on European marine biodiversity, ecosystem functions and services for human well-being (Grant no. 101060072). MWi acknowledges funding by the DFG (via iDiv, FZT 118, 202548816). We thank the associate editor and reviewer of the initial submission for comments that greatly improved the clarity of the manuscript. Open Access funding enabled and organized by Projekt DEAL.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

PEER REVIEW

The peer review history for this article is available at <https://www.webofscience.com/api/gateway/wos/peer-review/10.1002/2688-8319.70134>.

DATA AVAILABILITY STATEMENT

No data were used for this article.

ORCID

Helmut Hillebrand  <https://orcid.org/0000-0001-7449-1613>

Jan-Claas Dajka  <https://orcid.org/0000-0002-0797-9229>

Ralf Seppelt  <https://orcid.org/0000-0002-2723-7150>

Marten Winter  <https://orcid.org/0000-0002-9593-7300>

REFERENCES

- Affinito, F., Butchart, S. H. M., Nicholson, E., Hirsch, T., Williams, J. M., Campbell, J. E., Ferrari, M. F., Gabay, M., Gorini, L., Kalamujic Stroil, B., Kohsaka, R., Painter, B., Pinto, J. C., Scholz, A. H., Straza, T. R. A., Tshidada, N., Vallecillo, S., Widdicombe, S., & Gonzalez, A. (2025). Assessing coverage of the monitoring framework of the Kunming-Montreal global biodiversity framework and opportunities to fill gaps. *Nature Ecology & Evolution*, 9, 1280–1294.
- Affinito, F., Williams, J. M., Campbell, J. E., Londono, M. C., & Gonzalez, A. (2024). Progress in developing and operationalizing the monitoring framework of the global biodiversity framework. *Nature Ecology & Evolution*, 8, 2163–2171.
- Albini, D., Ransome, E., Dumbrell, A. J., Pawar, S., O'Gorman, E. J., Smith, T. P., Bell, T., Jackson, M. C., & Woodward, G. (2025). Warming alters plankton body-size distributions in a large field experiment. *Communications Biology*, 8, 162.
- Almond, R. E. A., Grooten, M., & Peterson, T. (2020). *Living planet report 2020—Bending the curve of biodiversity loss*. World Wildlife Foundation.
- Antonelli, A. (2022). *The hidden universe: Adventures in biodiversity*. Chicago University Press.
- Antonelli, A., Rueda, X., Calcagno, R., & Kalunda, P. N. (2024). How biodiversity credits could help to conserve and restore nature. *Nature*, 634, 1045–1049.
- BDM (Coordination Office). (2014). Swiss biodiversity monitoring BDM. Description of methods and indicators. In *Environmental studies* (Vol. 1410, p. 103). Swiss Federal Office for the Environment.
- CBD. (2022a). *Decision adopted by the Conference of the Parties to the Convention on Biological Diversity. 15/6 Mechanisms for planning, monitoring, reporting and review*. CBD/COP/DEC/15/6.
- CBD. (2022b). Kunming-Montreal Global Biodiversity Framework. *Conference of the Parties to the Convention on Biological Diversity. CBD/COP/DEC/15/4*.
- Chao, A., Chiu, C.-H., & Jost, L. (2014). Unifying species diversity, phylogenetic diversity, functional diversity, and related similarity and differentiation measures through Hill numbers. *Annual Review of Ecology, Evolution, and Systematics*, 45, 297–324.
- Dajka, J.-C., Dajka, J.-C., Eilrich, A. K., Franke, A., Halpern, B. S., Snow, B., Lombard, A. T., Jacob, U., Laakmann, S., Luhede, A., & Hillebrand, H. (2025). From science to policy: Evolving marine biodiversity targets. *Frontiers in Ecology and the Environment*, 23, e70000.
- Darwin, C. (1859). *The origin of species by means of natural selection*. Modern Library Paperback.
- Ekardt, F., Günther, P., Hagemann, K., Garske, B., Heyl, K., & Weyland, R. (2023). Legally binding and ambitious biodiversity protection under the CBD, the Global Biodiversity Framework, and human rights law. *Environmental Sciences Europe*, 35, 80.
- EU. (2000). Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for community action in the field of water policy. *Official Journal of the European Communities*, 53, 1–72.
- EU. (2010). 2010/477/EU: Commission decision of 1 September 2010 on criteria and methodological standards on good environmental status of marine waters. *Official Journal of the European Union*, 53, European Commission, 14–24.
- EU. (2017). 2017/848 of 17 May 2017 laying down criteria and methodological standards on good environmental status of marine waters and specifications and standardised methods for monitoring and assessment, and repealing decision 2010/477/EU. *Official Journal of the European Union*, 53, 14–24.
- EU. (2018). Directive (EU) 2018/2001 of the European Parliament and of the Council of 11 December 2018 on the promotion of the use of energy from renewable sources. *Official Journal of the European Communities*, 61, 82–209.
- EU. (2021). *EU biodiversity strategy for 2030 – Bringing nature back into our lives*. Publications Office of the European Union.
- Feld, C. K., Martins da Silva, P., Paulo Sousa, J., de Bello, F., Bugter, R., Grandin, U., Hering, D., Lavorel, S., Mountford, O., Pardo, I., Pärtel, M., Römbke, J., Sandin, L., Bruce Jones, K., & Harrison, P. (2009). Indicators of biodiversity and ecosystem services: A synthesis across ecosystems and spatial scales. *Oikos*, 118, 1862–1871.

- Fisher, L., Gross, T., Hillebrand, H., Sandberg, A., & Sayama, H. (2024). Sustainability: We need to focus on overall system outcomes rather than simplistic targets. *People and Nature*, 6, 391–401.
- Gonzalez, A., Chase, J. M., & O'Connor, M. I. (2023). A framework for the detection and attribution of biodiversity change. *Philosophical Transactions of the Royal Society, B: Biological Sciences*, 378, 20220182.
- Gonzalez, A., Vihervaara, P., Balvanera, P., Bates, A. E., Bayraktarov, E., Bellingham, P. J., Bruder, A., Campbell, J., Catchen, M. D., Cavender-Bares, J., Chase, J., Coops, N., Costello, M. J., Czúcz, B., Delavaud, A., Dornelas, M., Dubois, G., Duffy, E. J., Eggermont, H., ... Torrelio, C. Z. (2023). A global biodiversity observing system to unite monitoring and guide action. *Nature Ecology & Evolution*, 7, 1947–1952.
- Hébert, K., & Gravel, D. (2023). The living planet Index's ability to capture biodiversity change from uncertain data. *Ecology*, 104, e4044.
- Hébert, K., Jousse, M., Serrano, J., Karger, D. N., Blanchet, F. G., & Pollock, L. J. (2025). Five recommendations to fill the blank space in indicators at local and short-term scales. *Biological Conservation*, 302, 111007.
- Hermoso, V., Vasconcelos, R. P., Henriques, S., Filipe, A. F., & Carvalho, S. B. (2021). Conservation planning across realms: Enhancing connectivity for multi-realm species. *Journal of Applied Ecology*, 58, 644–654.
- Hill, S. L. L., Harfoot, M., Purvis, A., Purves, D. W., Collen, B., Newbold, T., Burgess, N. D., & Mace, G. M. (2016). Reconciling biodiversity indicators to guide understanding and action. *Conservation Letters*, 9, 405–412. <https://doi.org/10.1111/conl.12291>
- Hillebrand, H., Blasius, B., Borer, E. T., Chase, J. M., Downing, J. A., Eriksson, B. K., Filstrup, C. T., Harpole, W. S., Hodapp, D., Larsen, S., Lewandowska, A. M., Seabloom, E. W., van de Waal, D. B., & Ryabov, A. B. (2018). Biodiversity change is uncoupled from species richness trends: Consequences for conservation and monitoring. *Journal of Applied Ecology*, 55, 169–184.
- IPBES. (2024). In K. O'Brien, L. Garibaldi, A. Agrawal, E. Bennett, O. Biggs, R. Calderón Contreras, E. Carr, N. Frantzeskaki, H. Gosnell, J. Gurung, S. Lambertucci, J. Leventon, C. Liao, V. Reyes García, L. Shannon, S. Villasante, F. Wickson, Y. Zinggrebe, & L. Perianin (Eds.), *Summary for policymakers of the thematic assessment report on the underlying causes of biodiversity loss and the determinants of transformative change and options for achieving the 2050 vision for biodiversity of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services*. Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.
- Jackson, S. T., & Sax, D. F. (2010). Balancing biodiversity in a changing environment: Extinction debt, immigration credit and species turnover. *Trends in Ecology & Evolution*, 25, 153–160.
- Khalik, I., Rixen, C., Zellweger, F., Graham, C. H., Gossner, M. M., McFadden, I. R., Antão, L., Brodersen, J., Ghosh, S., Pomati, F., Seehausen, O., Roth, T., Sattler, T., Supp, S. R., Riaz, M., Zimmermann, N. E., Matthews, B., & Narwani, A. (2024). Warming underpins community turnover in temperate freshwater and terrestrial communities. *Nature Communications*, 15, 1921.
- Kharouba, H. M., Ehrlén, J., Gelman, A., Bolmgren, K., Allen, J. M., Travers, S. E., & Wolkovich, E. M. (2018). Global shifts in the phenological synchrony of species interactions over recent decades. *Proceedings of the National Academy of Sciences*, 115, 5211–5216.
- Kuczynski, L., Ontiveros, V. J., & Hillebrand, H. (2023). Biodiversity time series are biased towards increasing species richness in changing environments. *Nature Ecology & Evolution*, 7, 994–1001.
- Law, K. (2025). Biodiversity credits are more problematic than carbon credits. *Nature*, 637, 272.
- Leadley, P., Krug, C., Obura, D., Shannon, L., Gonzalez, A., Londoño-Murcia, M. C., Radulović, A., Millette, K., Rankovic, A., Archer, E., Bax, N., Chaudhari, K., Costello, M. J., Davalos, L. M., de Oliveira Roque, F., DeClerck, F., Dee, L., Essl, F., Ferrier, S., ... Xu, J. (2022). *Transformative actions on all drivers of biodiversity loss are urgently required to achieve the global goals by 2050*. Convention on Biological Diversity.
- Leclère, D., Obersteiner, M., Barrett, M., Butchart, S. H. M., Chaudhary, A., de Palma, A., DeClerck, F. A. J., di Marco, M., Doelman, J. C., Dürauer, M., Freeman, R., Harfoot, M., Hasegawa, T., Hellweg, S., Hilbers, J. P., Hill, S. L. L., Humpenöder, F., Jennings, N., Krisztin, T., ... Young, L. (2020). Bending the curve of terrestrial biodiversity needs an integrated strategy. *Nature*, 585, 551–556.
- Lemieux, C. J., Gray, P. A., Devillers, R., Wright, P. A., Dearden, P., Halpenny, E. A., Groulx, M., Beechey, T. J., & Beazley, K. (2019). How the race to achieve Aichi target 11 could jeopardize the effective conservation of biodiversity in Canada and beyond. *Marine Policy*, 99, 312–323.
- Leung, B., Hargreaves, A. L., Greenberg, D. A., McGill, B., Dornelas, M., & Freeman, R. (2020). Clustered versus catastrophic global vertebrate declines. *Nature*, 588, 267–271.
- Levin, S. A. (1998). Ecosystems and the biosphere as complex adaptive systems. *Ecosystems*, 1, 431–436.
- Magurran, A. E., & McGill, B. J. (Eds.). (2011). *Biological diversity—Frontiers in measurement and assessment*. Oxford University Press.
- McGeoch, M. A., Buba, Y., Arlé, E., Belmaker, J., Clarke, D. A., Jetz, W., Li, R., Seebens, H., Essl, F., Groom, Q., García-Berthou, E., Lenzner, B., Meyer, C., Vicente, J. R., Wilson, J. R. U., & Winter, M. (2023). Invasion trends: An interpretable measure of change is needed to support policy targets. *Conservation Letters*, 16, e12981.
- Paavola, J., Gouldson, A., & Kluvánková-Oravská, T. (2009). Interplay of actors, scales, frameworks and regimes in the governance of biodiversity. *Environmental Policy and Governance*, 19, 148–158.
- Pereira, H. M., Ferrier, S., Walters, M., Geller, G. N., Jongman, R. H. G., Scholes, R. J., Bruford, M. W., Brummitt, N., Butchart, S. H. M., Cardoso, A. C., Coops, N. C., Dulloo, E., Faith, D. P., Freyhof, J., Gregory, R. D., Heip, C., Höft, R., Hurr, G., Jetz, W., ... Wegmann, M. (2013). Essential biodiversity variables. *Science*, 339, 277–278.
- Piipponen-Doyle, S., Bolam, F. C., & Mair, L. (2021). Disparity between ecological and political timeframes for species conservation targets. *Biodiversity and Conservation*, 30, 1899–1912.
- Pörtner, H. O., Scholes, R. J., Agard, J., Archer, E., Arneeth, A., Bai, X., Barnes, D., Burrows, M., Chan, L., Cheung, W. L., Diamond, S., Donatti, C., Duarte, C., Eisenhauer, N., Foden, W., Gasalla, M. A., Handa, C., Hickler, T., Hoegh-Guldberg, O., ... Ngo, H. T. (2021). *IPBES-IPCC co-sponsored workshop report on biodiversity and climate change*. IPBES and IPCC.
- Scherer-Lorenzen, M., Gessner, M. O., Beisner, B. E., Messier, C., Paquette, A., Petermann, J. S., Sojininen, J., & Nock, C. A. (2022). Pathways for cross-boundary effects of biodiversity on ecosystem functioning. *Trends in Ecology & Evolution*, 37, 454–467.
- Silva del Pozo, M., Body, G., Rerig, G., & Basille, M. (2023). *Guide on harmonising biodiversity monitoring protocols across scales. Support to effectively integrate monitoring results*. <https://www.biodiversa.eu/biodiversity-monitoring/harmonisation-of-protocols/>
- Sobkowiak, M. (2023). The making of imperfect indicators for biodiversity: A case study of UK biodiversity performance measurement. *Business Strategy and the Environment*, 32, 336–352.
- Stevenson, S. L., Watermeyer, K., Caggiano, G., Fulton, E. A., Ferrier, S., & Nicholson, E. (2021). Matching biodiversity indicators to policy needs. *Conservation Biology*, 35, 522–532.
- Supran, G., Rahmstorf, S., & Oreskes, N. (2023). Assessing ExxonMobil's global warming projections. *Science*, 379, eabk0063.
- Teixeira, H., Berg, T., Uusitalo, L., FÜRhaupter, K., Heiskanen, A.-S., Mazik, K., Lynam, C. P., Neville, S., Rodríguez, J. G., Papadopoulou, N., Moncheva, S., Churilova, T., Kryvenko, O., Krause-Jensen, D., Zaiko, A., Veríssimo, H., Pantazi, M., Carvalho, S., Patrício, J., ... Borja, À. (2016). A catalogue of marine biodiversity indicators. *Frontiers in Marine Science*, 3, 207.
- TNFD. (2023). *Recommendations of the taskforce on nature-related financial disclosures (TNFD)*. Taskforce on Nature-related Financial Disclosures.
- Toszogyova, A., Smyčka, J., & Storch, D. (2024). Mathematical biases in the calculation of the living planet index lead to overestimation of vertebrate population decline. *Nature Communications*, 15, 5295.

- Tudge, S. J., Purvis, A., & De Palma, A. (2021). The impacts of biofuel crops on local biodiversity: A global synthesis. *Biodiversity and Conservation*, 30, 2863–2883.
- van Dorst, R. M., Gårdmark, A., Svanbäck, R., Beier, U., Weyhenmeyer, G. A., & Huss, M. (2019). Warmer and browner waters decrease fish biomass production. *Global Change Biology*, 25, 1395–1408.
- Watts, K., Whytock, R. C., Park, K. J., Fuentes-Montemayor, E., Macgregor, N. A., Duffield, S., & McGowan, P. J. K. (2020). Ecological time lags and the journey towards conservation success. *Nature Ecology & Evolution*, 4, 304–311.
- Wauchope, H. S., zu Ermgassen, S. O. S. E., Jones, J. P. G., Carter, H., Schulte to Bühne, H., & Milner-Gulland, E. J. (2024). What is a unit of nature? Measurement challenges in the emerging biodiversity credit market. *Proceedings of the Royal Society B: Biological Sciences*, 291, 20242353.
- Woolway, R. I., Sharma, S., & Smol, J. P. (2022). Lakes in hot water: The impacts of a changing climate on aquatic ecosystems. *Bioscience*, 72, 1050–1061.
- Yvon-Durocher, G., Jones, J. I., Trimmer, M., Woodward, G., & Montoya, J. M. (2010). Warming alters the metabolic balance of ecosystems. *Philosophical Transactions of the Royal Society, B: Biological Sciences*, 365, 2117–2126.
- Zinngrebe, Y., Berger, J., Bunn, C., Felipe-Lucia, M. R., Graßnick, N., Kastner, T., Pe'er, G., Schleyer, C., & Lakner, S. (2024). Prioritizing partners and products for the sustainability of the EU's agri-food trade. *One Earth*, 7, 674–686.

How to cite this article: Hillebrand, H., Dajka, J.-C., Halbach, M., Happe, A., Röchert, R., Seppelt, R., Settele, J., Weitere, M., Winter, M., Zinngrebe, Y., & Hodapp, D. (2025). Operational perspectives for biodiversity indicators. *Ecological Solutions and Evidence*, 6, e70134. <https://doi.org/10.1002/2688-8319.70134>