

ON SIGNATURE-BASED EXPRESSIONS OF SYSTEM RELIABILITY

JEAN-LUC MARICHAL, PIERRE MATHONET, AND TAMÁS WALDHAUSER

ABSTRACT. The concept of signature was introduced by Samaniego for systems whose components have i.i.d. lifetimes. This concept proved to be useful in the analysis of theoretical behaviors of systems. In particular, it provides an interesting signature-based representation of the system reliability in terms of reliabilities of k -out-of- n systems. In the non-i.i.d. case, we show that, at any time, this representation still holds true for every coherent system if and only if the component states are exchangeable. We also discuss conditions for obtaining an alternative representation of the system reliability in which the signature is replaced by its non-i.i.d. extension. Finally, we discuss conditions for the system reliability to have both representations.

1. INTRODUCTION

Consider a system made up of n ($n \geq 3$) components and let $\phi: \{0, 1\}^n \rightarrow \{0, 1\}$ be its *structure function*, which expresses the state of the system in terms of the states of its components. Denote the set of components by $[n] = \{1, \dots, n\}$. We assume that the system is *coherent*, which means that ϕ is nondecreasing in each variable and has only essential variables, i.e., for every $k \in [n]$, there exists $\mathbf{x} = (x_1, \dots, x_n) \in \{0, 1\}^n$ such that $\phi(\mathbf{x})|_{x_k=0} \neq \phi(\mathbf{x})|_{x_k=1}$.

Let X_1, \dots, X_n denote the component lifetimes and let $X_{1:n}, \dots, X_{n:n}$ be the order statistics obtained by rearranging the variables X_1, \dots, X_n in ascending order of magnitude; that is, $X_{1:n} \leq \dots \leq X_{n:n}$. Denote also the system lifetime by T and the system reliability at time $t > 0$ by $\bar{F}_S(t) = \Pr(T > t)$.

Assuming that the component lifetimes are independent and identically distributed (i.i.d.) according to an absolutely continuous joint c.d.f. F , one can show (see Samaniego [11]) that

$$(1) \quad \bar{F}_S(t) = \sum_{k=1}^n \Pr(T = X_{k:n}) \bar{F}_{k:n}(t)$$

for every $t > 0$, where $\bar{F}_{k:n}(t) = \Pr(X_{k:n} > t)$.

Under this i.i.d. assumption, Samaniego [11] introduced the *signature* of the system as the n -tuple $\mathbf{s} = (s_1, \dots, s_n)$, where

$$s_k = \Pr(T = X_{k:n}), \quad k \in [n],$$

is the probability that the k th component failure causes the system to fail. It turned out that the signature is a feature of the system design in the sense that it depends

Date: February 17, 2011.

2010 Mathematics Subject Classification. 62G30, 62N05, 90B25, 94C10.

Key words and phrases. system signature, system reliability, coherent system, order statistic.

only on the structure function ϕ (and not on the c.d.f. F). Boland [1] obtained the explicit formula

$$s_k = \phi_{n-k+1} - \phi_{n-k}$$

where

$$(2) \quad \phi_k = \frac{1}{\binom{n}{k}} \sum_{\substack{\mathbf{x} \in \{0,1\}^n \\ |\mathbf{x}|=k}} \phi(\mathbf{x})$$

and $|\mathbf{x}| = \sum_{i=1}^n x_i$. Thus, under the i.i.d. assumption, the system reliability can be calculated by the formula

$$(3) \quad \bar{F}_S(t) = \sum_{k=1}^n (\phi_{n-k+1} - \phi_{n-k}) \bar{F}_{k:n}(t).$$

Since formula (3) provides a simple and useful way to compute the system reliability through the concept of signature, it is natural to relax the i.i.d. assumption (as Samaniego [12, Section 8.3] rightly suggested) and search for necessary and sufficient conditions on the joint c.d.f. F for formulas (1) and/or (3) to still hold for every system design.

On this issue, Kochar et al. [4, p. 513] mentioned that (1) and (3) still hold when the continuous variables X_1, \dots, X_n are exchangeable (i.e., when F is invariant under any permutation of indexes); see also [6, 13] (and [7, Lemma 1] for a detailed proof). It is also noteworthy that Navarro et al. [8, Thm. 3.6] showed that (1) still holds when the joint c.d.f. F has no ties (i.e., $\Pr(X_i = X_j) = 0$ for every $i \neq j$) and the variables X_1, \dots, X_n are “weakly exchangeable” (see Remark 3 below). As we will show, all these conditions are not necessary.

Let Φ_n denote the family of nondecreasing functions $\phi: \{0, 1\}^n \rightarrow \{0, 1\}$ whose variables are all essential. In this paper, without any assumption on the joint c.d.f. F , we show that, for every $t > 0$, the representation in (3) of the system reliability holds for every $\phi \in \Phi_n$ if and only if the variables $\chi_1(t), \dots, \chi_n(t)$ are exchangeable, where

$$\chi_k(t) = \text{Ind}(X_k > t)$$

denotes the *random state* of the k th component at time t (i.e., $\chi_k(t)$ is the indicator variable of the event $(X_k > t)$). This result is stated in Theorem 4.

Assuming that the joint c.d.f. F has no ties, we also yield necessary and sufficient conditions on F for formula (1) to hold for every $\phi \in \Phi_n$ (Theorem 6). These conditions can be interpreted in terms of symmetry of certain conditional probabilities.

We also show (Proposition 7) that the condition¹

$$(4) \quad \Pr(T = X_{k:n}) = \phi_{n-k+1} - \phi_{n-k}, \quad k \in [n]$$

holds for every $\phi \in \Phi_n$ if and only if

$$(5) \quad \Pr\left(\max_{i \in [n] \setminus A} X_i < \min_{i \in A} X_i\right) = \frac{1}{\binom{n}{|A|}}, \quad A \subseteq [n].$$

Finally, we show that both (1) and (3) hold for every $t > 0$ and every $\phi \in \Phi_n$ if and only if (5) holds and the variables $\chi_1(t), \dots, \chi_n(t)$ are exchangeable for every $t > 0$ (Theorem 8).

¹Note that, according to the terminology used in [9], the left-hand side of (4) is the k th coordinate of the *probability signature*, while the right-hand side is the k th coordinate of the *system signature*.

Through the usual identification of the elements of $\{0,1\}^n$ with the subsets of $[n]$, a pseudo-Boolean function $f: \{0,1\}^n \rightarrow \mathbb{R}$ can be described equivalently by a set function $v_f: 2^{[n]} \rightarrow \mathbb{R}$. We simply write $v_f(A) = f(\mathbf{1}_A)$, where $\mathbf{1}_A$ denotes the n -tuple whose i th coordinate ($i \in [n]$) is 1, if $i \in A$, and 0, otherwise. To avoid cumbersome notation, we henceforth use the same symbol to denote both a given pseudo-Boolean function and its underlying set function, thus writing $f: \{0,1\}^n \rightarrow \mathbb{R}$ or $f: 2^{[n]} \rightarrow \mathbb{R}$ interchangeably.

Recall that the k th order statistic function $\mathbf{x} \mapsto x_{k:n}$ of n Boolean variables is defined by $x_{k:n} = 1$, if $|\mathbf{x}| \geq n - k + 1$, and 0, otherwise. As a matter of convenience, we also formally define $x_{0:n} \equiv 0$ and $x_{n+1:n} \equiv 1$.

2. SIGNATURE-BASED DECOMPOSITION OF THE SYSTEM RELIABILITY

In the present section, without any assumption on the joint c.d.f. F , we show that, for every $t > 0$, (3) holds true for every $\phi \in \Phi_n$ if and only if the state variables $\chi_1(t), \dots, \chi_n(t)$ are exchangeable.

The following result (see Dukhovny [2, Thm. 2]) gives a useful expression for the system reliability in terms of the underlying structure function and the component states. We provide a shorter proof here. For every $t > 0$, we set $\chi(t) = (\chi_1(t), \dots, \chi_n(t))$.

Proposition 1. *For every $t > 0$, we have*

$$(6) \quad \bar{F}_S(t) = \sum_{\mathbf{x} \in \{0,1\}^n} \phi(\mathbf{x}) \Pr(\chi(t) = \mathbf{x}).$$

Proof. We simply have

$$\bar{F}_S(t) = \Pr(\phi(\chi(t)) = 1) = \sum_{\substack{\mathbf{x} \in \{0,1\}^n \\ \phi(\mathbf{x}) = 1}} \Pr(\chi(t) = \mathbf{x}),$$

which immediately leads to (6). \square

Applying (6) to the k -out-of- n system $\phi(\mathbf{x}) = x_{k:n}$, we obtain

$$\bar{F}_{k:n}(t) = \sum_{|\mathbf{x}| \geq n-k+1} \Pr(\chi(t) = \mathbf{x})$$

from which we immediately derive (see [3, Prop. 13])

$$(7) \quad \bar{F}_{n-k+1:n}(t) - \bar{F}_{n-k:n}(t) = \sum_{|\mathbf{x}|=k} \Pr(\chi(t) = \mathbf{x}).$$

The following proposition, a key result of this paper, provides necessary and sufficient conditions on F for $\bar{F}_S(t)$ to be a certain weighted sum of the $\bar{F}_{k:n}(t)$, $k \in [n]$. We first consider a lemma.

Lemma 2. *Let $\lambda: \{0,1\}^n \rightarrow \mathbb{R}$ be a given function. We have*

$$(8) \quad \sum_{\mathbf{x} \in \{0,1\}^n} \lambda(\mathbf{x}) \phi(\mathbf{x}) = 0 \quad \text{for every } \phi \in \Phi_n$$

if and only if $\lambda(\mathbf{x}) = 0$ for all $\mathbf{x} \neq \mathbf{0}$.

Proof. Condition (8) defines a system of linear equations with the 2^n unknowns $\lambda(\mathbf{x})$, $\mathbf{x} \in \{0,1\}^n$. We observe that there exist $2^n - 1$ functions $\phi_A \in \Phi_n$, $A \neq \emptyset$, which are linearly independent when considered as real functions (for details, see

Appendix A). It follows that the vectors of their values are also linearly independent. Therefore the equations in (8) corresponding to the functions ϕ_A , $A \neq \emptyset$, are linearly independent and hence the system has a rank at least $2^n - 1$. This shows that its solutions are multiples of the immediate solution λ_0 defined by $\lambda_0(\mathbf{x}) = 0$, if $\mathbf{x} \neq \mathbf{0}$, and $\lambda_0(\mathbf{0}) = 1$. \square

Let $w: \{0, 1\}^n \rightarrow \mathbb{R}$ be a given function. For every $k \in [n]$ and every $\phi \in \Phi_n$, define

$$(9) \quad \phi_k^w = \sum_{|\mathbf{x}|=k} w(\mathbf{x}) \phi(\mathbf{x}).$$

Proposition 3. *For every $t > 0$, we have*

$$\overline{F}_S(t) = \sum_{k=1}^n (\phi_{n-k+1}^w - \phi_{n-k}^w) \overline{F}_{k:n}(t) \quad \text{for every } \phi \in \Phi_n$$

if and only if

$$(10) \quad \Pr(\chi(t) = \mathbf{x}) = w(\mathbf{x}) \sum_{|\mathbf{z}|=|\mathbf{x}|} \Pr(\chi(t) = \mathbf{z}) \quad \text{for every } \mathbf{x} \neq \mathbf{0}.$$

Proof. First observe that we have

$$(11) \quad \sum_{k=1}^n (\phi_{n-k+1}^w - \phi_{n-k}^w) \overline{F}_{k:n}(t) = \sum_{k=1}^n \phi_k^w (\overline{F}_{n-k+1:n}(t) - \overline{F}_{n-k:n}(t)).$$

This immediately follows from the elementary algebraic identity

$$\sum_{k=1}^n a_k (b_{n-k+1} - b_{n-k}) = \sum_{k=1}^n b_k (a_{n-k+1} - a_{n-k})$$

which holds for all real tuples (a_0, a_1, \dots, a_n) and (b_0, b_1, \dots, b_n) such that $a_0 = b_0 = 0$. Combining (7) with (9) and (11), we then obtain

$$\begin{aligned} \sum_{k=1}^n (\phi_{n-k+1}^w - \phi_{n-k}^w) \overline{F}_{k:n}(t) &= \sum_{k=1}^n \sum_{|\mathbf{x}|=k} w(\mathbf{x}) \phi(\mathbf{x}) \sum_{|\mathbf{z}|=k} \Pr(\chi(t) = \mathbf{z}) \\ &= \sum_{\mathbf{x} \in \{0,1\}^n} w(\mathbf{x}) \phi(\mathbf{x}) \sum_{|\mathbf{z}|=|\mathbf{x}|} \Pr(\chi(t) = \mathbf{z}). \end{aligned}$$

The result then follows from Proposition 1 and Lemma 2. \square

Remark 1. We observe that the existence of a c.d.f. F satisfying (10) with $\Pr(\chi(t) = \mathbf{x}) > 0$ for some $\mathbf{x} \neq \mathbf{0}$ is only possible when $\sum_{|\mathbf{z}|=|\mathbf{x}|} w(\mathbf{z}) = 1$. In this paper we will actually make use of (10) only when this condition holds (see (12) and (15)).

We now apply Proposition 3 to obtain necessary and sufficient conditions on F for (3) to hold for every $\phi \in \Phi_n$.

Theorem 4. *For every $t > 0$, the representation (3) holds for every $\phi \in \Phi_n$ if and only if the indicator variables $\chi_1(t), \dots, \chi_n(t)$ are exchangeable.*

Proof. Using (2) and Proposition 3, we see that condition (3) is equivalent to

$$(12) \quad \Pr(\chi(t) = \mathbf{x}) = \frac{1}{\binom{n}{|\mathbf{x}|}} \sum_{|\mathbf{z}|=|\mathbf{x}|} \Pr(\chi(t) = \mathbf{z}).$$

Equivalently, we have $\Pr(\chi(t) = \mathbf{x}) = \Pr(\chi(t) = \mathbf{x}')$ for every $\mathbf{x}, \mathbf{x}' \in \{0, 1\}^n$ such that $|\mathbf{x}| = |\mathbf{x}'|$. This condition clearly means that $\chi_1(t), \dots, \chi_n(t)$ are exchangeable. \square

The following well-known proposition (see for instance [10, Chap. 1] and [2, Section 2]) yields an interesting interpretation of the exchangeability of the component states $\chi_1(t), \dots, \chi_n(t)$. For the sake of self-containment, a proof is given here.

Proposition 5. *For every $t > 0$, the component states $\chi_1(t), \dots, \chi_n(t)$ are exchangeable if and only if the probability that a group of components survives beyond t (i.e., the reliability of this group at time t) depends only on the number of components in the group.*

Proof. Let $A \subseteq [n]$ be a group of components. The exchangeability of the component states means that, for every $B \subseteq [n]$, the probability $\Pr(\chi(t) = \mathbf{1}_B)$ depends only on $|B|$. In this case, the probability that the group A survives beyond t , that is

$$\bar{F}_A(t) = \sum_{B \supseteq A} \Pr(\chi(t) = \mathbf{1}_B),$$

depends only on $|A|$. Conversely, if $\bar{F}_B(t)$ depends only on $|B|$ for every $B \subseteq [n]$, then

$$\Pr(\chi(t) = \mathbf{1}_A) = \sum_{B \supseteq A} (-1)^{|B|-|A|} \bar{F}_B(t)$$

depends only on $|A|$. \square

Remark 2. Theorem 4 shows that the exchangeability of the component lifetimes is sufficient but not necessary for (3) to hold for every $\phi \in \Phi_n$ and every $t > 0$. Indeed, the exchangeability of the component lifetimes entails the exchangeability of the component states. This follows for instance from the identity (see [3, Eq. (6)])

$$\Pr(\chi(t) = \mathbf{1}_A) = \sum_{B \subseteq A} (-1)^{|A|-|B|} F(t\mathbf{1}_{[n] \setminus B} + \infty \mathbf{1}_B).$$

However, the converse statement is not true in general. As an example, consider the random vector (X_1, X_2) which takes each of the values $(2, 1)$, $(4, 2)$, $(1, 3)$ and $(3, 4)$ with probability $1/4$. The state variables $\chi_1(t)$ and $\chi_2(t)$ are exchangeable at any time t . Indeed, one can easily see that, for $|\mathbf{x}| = 1$,

$$\Pr(\chi(t) = \mathbf{x}) = \begin{cases} 1/4, & \text{if } t \in [1, 4), \\ 0, & \text{otherwise.} \end{cases}$$

However, the variables X_1 and X_2 are not exchangeable since, for instance,

$$0 = F(1.5, 2.5) \neq F(2.5, 1.5) = 1/4.$$

3. ALTERNATIVE DECOMPOSITION OF THE SYSTEM RELIABILITY

Assuming only that F has no ties (i.e., $\Pr(X_i = X_j) = 0$ for every $i \neq j$), we now provide necessary and sufficient conditions on F for formula (1) to hold for every $\phi \in \Phi_n$, thus answering a question raised implicitly in [8, p. 320].

Let $q: 2^{[n]} \rightarrow [0, 1]$ be the *relative quality function* (associated with F), which is defined as

$$q(A) = \Pr\left(\max_{i \in [n] \setminus A} X_i < \min_{i \in A} X_i\right)$$

with the convention that $q(\emptyset) = q([n]) = 1$ (see [5, Section 2]). By definition, $q(A)$ is the probability that the $|A|$ components having the longest lifetimes are exactly

those in A . It then immediately follows that the function q satisfies the following important property:

$$(13) \quad \sum_{|\mathbf{x}|=k} q(\mathbf{x}) = 1, \quad k \in [n].$$

Under the assumption that F has no ties, the authors [5, Thm. 3] proved that

$$(14) \quad \Pr(T = X_{k:n}) = \phi_{n-k+1}^q - \phi_{n-k}^q,$$

where ϕ_k^q is defined in (9).

Combining (14) with Proposition 3, we immediately derive the following result.

Theorem 6. *Assume that F has no ties. For every $t > 0$, the representation (1) holds for every $\phi \in \Phi_n$ if and only if*

$$(15) \quad \Pr(\chi(t) = \mathbf{x}) = q(\mathbf{x}) \sum_{|\mathbf{z}|=|\mathbf{x}|} \Pr(\chi(t) = \mathbf{z}).$$

Condition (15) has the following interpretation. We first observe that, for every $A \subseteq [n]$,

$$\chi(t) = \mathbf{1}_A \Leftrightarrow \max_{i \in [n] \setminus A} X_i \leq t < \min_{i \in A} X_i.$$

Assuming that q is a strictly positive function, condition (15) then means that the conditional probability

$$\frac{\Pr(\chi(t) = \mathbf{1}_A)}{q(A)} = \Pr\left(\max_{i \in [n] \setminus A} X_i \leq t < \min_{i \in A} X_i \mid \max_{i \in [n] \setminus A} X_i < \min_{i \in A} X_i\right)$$

depends only on $|A|$.

Remark 3. The concept of weak exchangeability was introduced in Navarro et al. [8, p. 320] as follows. A random vector (X_1, \dots, X_n) is said to be *weakly exchangeable* if

$$\Pr(X_{k:n} \leq t) = \Pr(X_{k:n} \leq t \mid X_{\sigma(1)} < \dots < X_{\sigma(n)}),$$

for every $t > 0$, every $k \in [n]$, and every permutation σ on $[n]$. Theorem 3.6 in [8] states that if F has no ties and (X_1, \dots, X_n) is weakly exchangeable, then (1) holds for every $\phi \in \Phi_n$. By Theorem 6, we see that weak exchangeability implies condition (15) whenever F has no ties. However, the converse is not true in general. Indeed, in the example of Remark 2, we can easily see that condition (15) holds, while the lifetimes X_1 and X_2 are not weakly exchangeable.

We now investigate condition (4) under the sole assumption that F has no ties. Navarro and Rychlik [7, Lemma 1] (see also [5, Rem. 4]) proved that this condition holds for every $\phi \in \Phi_n$ whenever the component lifetimes X_1, \dots, X_n are exchangeable. The following proposition gives a necessary and sufficient condition on F (in terms of the function q) for (4) to hold for every $\phi \in \Phi_n$.

The function q is said to be *symmetric* if $q(\mathbf{x}) = q(\mathbf{x}')$ whenever $|\mathbf{x}| = |\mathbf{x}'|$. By (13) it follows that q is symmetric if and only if $q(\mathbf{x}) = 1/\binom{n}{|\mathbf{x}|}$ for every $\mathbf{x} \in \{0, 1\}^n$.

Proposition 7. *Assume that F has no ties. Condition (4) holds for every $\phi \in \Phi_n$ if and only if q is symmetric.*

Proof. By (14) we have

$$\Pr(T = X_{k:n}) = \sum_{\mathbf{x} \in \{0,1\}^n} (\delta_{|\mathbf{x}|, n-k+1} - \delta_{|\mathbf{x}|, n-k}) q(\mathbf{x}) \phi(\mathbf{x}), \quad k \in [n],$$

where δ stands for the Kronecker delta. Similarly, by (2) we have

$$\phi_{n-k+1} - \phi_{n-k} = \sum_{\mathbf{x} \in \{0,1\}^n} (\delta_{|\mathbf{x}|, n-k+1} - \delta_{|\mathbf{x}|, n-k}) \frac{1}{\binom{n}{|\mathbf{x}|}} \phi(\mathbf{x}), \quad k \in [n].$$

The result then follows from Lemma 2. \square

We end this paper by studying the special case where both conditions (1) and (3) hold. We have the following result.

Theorem 8. *Assume that F has no ties. The following assertions are equivalent.*

- (i) *Conditions (1) and (3) hold for every $\phi \in \Phi_n$ and every $t > 0$.*
- (ii) *Condition (4) holds for every $\phi \in \Phi_n$ and the variables $\chi_1(t), \dots, \chi_n(t)$ are exchangeable for every $t > 0$.*
- (iii) *The function q is symmetric and the variables $\chi_1(t), \dots, \chi_n(t)$ are exchangeable for every $t > 0$.*

Proof. (ii) \Leftrightarrow (iii) Follows from Proposition 7.

(ii) \Rightarrow (i) Follows from Theorem 4.

(i) \Rightarrow (iii) By Theorem 4, we only need to prove that q is symmetric. Combining (12) with (15), we obtain

$$\left(q(\mathbf{x}) - \frac{1}{\binom{n}{|\mathbf{x}|}} \right) \sum_{|\mathbf{z}|=|\mathbf{x}|} \Pr(\chi(t) = \mathbf{z}) = 0.$$

To conclude, we only need to prove that, for every $k \in [n-1]$, there exists $t > 0$ such that

$$\sum_{|\mathbf{z}|=k} \Pr(\chi(t) = \mathbf{z}) > 0.$$

Suppose that this is not true. By (7), there exists $k \in [n-1]$ such that

$$0 = \bar{F}_{n-k+1:n}(t) - \bar{F}_{n-k:n}(t) = \Pr(X_{n-k:n} \leq t < X_{n-k+1:n})$$

for every $t > 0$. Then, denoting the set of positive rational numbers by \mathbb{Q}^+ , the sequence of events

$$E_m = (X_{n-k:n} \leq t_m < X_{n-k+1:n}), \quad m \in \mathbb{N},$$

where $\{t_m : m \in \mathbb{N}\} = \mathbb{Q}^+$, satisfies $\Pr(E_m) = 0$. Since \mathbb{Q}^+ is dense in $(0, \infty)$, we obtain

$$\Pr(X_{n-k:n} < X_{n-k+1:n}) = \Pr\left(\bigcup_{m \in \mathbb{N}} E_m\right) = 0,$$

which contradicts the assumption that F has no ties. \square

The following two examples show that neither of the conditions (1) and (3) implies the other.

Example 9. Let (X_1, X_2, X_3) be the random vector which takes the values $(1, 2, 3)$, $(1, 3, 2)$, $(2, 1, 3)$, $(2, 3, 1)$, $(3, 2, 1)$, $(3, 1, 2)$, with probabilities p_1, \dots, p_6 , respectively. It was shown in [8, Example 3.7] that (1) holds for every $\phi \in \Phi_n$ and every $t > 0$. However, we can easily see that $\chi_1(t), \chi_2(t), \chi_3(t)$ are exchangeable for every $t > 0$ if and only if (p_1, \dots, p_6) is a convex combination of $(0, 1/3, 1/3, 0, 1/3, 0)$ and $(1/3, 0, 0, 1/3, 0, 1/3)$. Hence, when the latter condition is not satisfied, (3) does not hold for every $\phi \in \Phi_n$ by Theorem 4.

Example 10. Let (X_1, X_2, X_3) be the random vector which takes the values $(1, 2, 4), (2, 4, 5), (3, 1, 2), (4, 2, 3), (5, 3, 4), (2, 3, 1), (3, 4, 2), (4, 5, 3)$ with probabilities $p_1 = \dots = p_8 = 1/8$. We have

$$q(\{1\}) = q(\{2\}) = q(\{1, 2\}) = q(\{1, 3\}) = 3/8 \quad \text{and} \quad q(\{3\}) = q(\{2, 3\}) = 2/8,$$

which shows that q is not symmetric. However, we can easily see that $\chi_1(t), \chi_2(t), \chi_3(t)$ are exchangeable for every $t > 0$. Indeed, we have

$$\Pr(\chi(t) = \mathbf{x}) = \begin{cases} 1/8, & \text{if } t \in [\alpha, \beta), \\ 0, & \text{otherwise,} \end{cases}$$

where $(\alpha, \beta) = (2, 5)$ whenever $|\mathbf{x}| = 1$ and $(\alpha, \beta) = (1, 4)$ whenever $|\mathbf{x}| = 2$. Thus (3) holds for every $\phi \in \Phi_n$ and every $t > 0$ by Theorem 4. However, (1) does not hold for every $\phi \in \Phi_n$ and every $t > 0$ by Theorem 8.

Remark 4. Let Φ'_n be the class of structure functions of n -component *semicoherent* systems, that is, the class of nondecreasing functions $\phi: \{0, 1\}^n \rightarrow \{0, 1\}$ satisfying the boundary conditions $\phi(\mathbf{0}) = 0$ and $\phi(\mathbf{1}) = 1$. It is clear that Proposition 1 and Lemma 2 still hold, even for $n = 2$, if we extend the set Φ_n to Φ'_n (in the proof of Lemma 2 it is then sufficient to consider the $2^n - 1$ functions $\phi_A(\mathbf{x}) = \prod_{i \in A} x_i$, $A \neq \emptyset$). We then observe that Propositions 3 and 7 and Theorems 4, 6, and 8 (which use Proposition 1 and Lemma 2 to provide conditions on F for certain identities to hold for every $\phi \in \Phi_n$) are still valid for $n \geq 2$ if we replace Φ_n with Φ'_n (that is, if we consider semicoherent systems instead of coherent systems only). This observation actually strengthens these results. For instance, from Theorem 4 we can state that, for every fixed $t > 0$, if (3) holds for every $\phi \in \Phi_n$, then the variables $\chi_1(t), \dots, \chi_n(t)$ are exchangeable; conversely, for every $n \geq 2$ and every $t > 0$, the latter condition implies that (3) holds for every $\phi \in \Phi'_n$. We also observe that the “semicoherent” version of Theorem 4 (i.e., where Φ_n is replaced with Φ'_n) was proved by Dukhovny [2, Thm. 4].

ACKNOWLEDGMENTS

The authors wish to thank M. Couceiro, G. Peccati, and F. Spizzichino for fruitful discussions. Jean-Luc Marichal and Pierre Mathonet are supported by the internal research project F1R-MTH-PUL-09MRDO of the University of Luxembourg. Tamás Waldhauser is supported by the National Research Fund of Luxembourg, the Marie Curie Actions of the European Commission (FP7-COFUND), and the Hungarian National Foundation for Scientific Research under grant no. K77409.

APPENDIX A

In this appendix we construct $2^n - 1$ functions in Φ_n which are linearly independent when considered as real functions. Here the assumption $n \geq 3$ is crucial.

Assume first that $n \neq 4$ and let π be the permutation on $[n]$ defined by the following cycles

$$\pi = \begin{cases} (1, 2, \dots, n), & \text{if } n \text{ is odd,} \\ (1, 2, 3) \circ (4, 5, \dots, n), & \text{if } n \text{ is even.} \end{cases}$$

With every $A \subsetneq [n]$, $A \neq \emptyset$, we associate $A^* \subseteq [n]$ in the following way:

- if $|A| \leq n-2$, then we choose any set A^* such that $|A^*| = n-1$ and $A \cup A^* = [n]$;
- if $A = [n] \setminus \{k\}$ for some $k \in [n]$, then we take $A^* = [n] \setminus \{\pi(k)\}$.

We now show that the $2^n - 1$ functions $\phi_A \in \Phi_n$, $A \subseteq [n]$, $A \neq \emptyset$, defined by

$$\phi_A(\mathbf{x}) = \begin{cases} \left(\prod_{i \in A} x_i \right) \sqcup \left(\prod_{i \in A^*} x_i \right), & \text{if } A \neq [n], \\ \prod_{i \in [n]} x_i, & \text{if } A = [n], \end{cases}$$

where \sqcup denotes the coproduct (i.e., $x \sqcup y = x + y - xy$), are linearly independent when considered as real functions.

Suppose there exist real numbers c_A , $A \subseteq [n]$, $A \neq \emptyset$, such that

$$\sum_{A \neq \emptyset} c_A \phi_A = 0.$$

Expanding the left-hand side of this equation as a linear combination of the functions $\prod_{i \in B} x_i$, $B \subseteq [n]$, $B \neq \emptyset$, we first see that, if $|A| \leq n - 2$, the coefficient of $\prod_{i \in A} x_i$ is c_A and hence $c_A = 0$ whenever $0 < |A| \leq n - 2$. Next, considering the coefficient of $\prod_{i \in A} x_i$ for $A = [n] \setminus \{k\}$, $k \in [n]$, we obtain

$$c_{[n] \setminus \{k\}} + c_{[n] \setminus \{\pi^{-1}(k)\}} = 0.$$

Since π is made up of odd-length cycles only, it follows that $c_A = 0$ whenever $|A| = n - 1$.

For $n = 4$ we consider the function $\pi: [4] \rightarrow [4]$ defined by $\pi(1) = \pi(4) = 2$, $\pi(2) = 3$, and $\pi(3) = 4$, and choose the functions ϕ_A as above. We then easily check that these functions are linearly independent.

REFERENCES

- [1] P. J. Boland. Signatures of indirect majority systems. *J. Appl. Prob.*, 38:597–603, 2001.
- [2] A. Dukhovny. Lattice polynomials of random variables. *Statistics & Probability Letters*, 77(10):989–994, 2007.
- [3] A. Dukhovny and J.-L. Marichal. System reliability and weighted lattice polynomials. *Probability in the Engineering and Informational Sciences*, 22(3):373–388, 2008.
- [4] S. Kochar, H. Mukerjee, and F. J. Samaniego. The “signature” of a coherent system and its application to comparisons among systems. *Naval Res. Logist.*, 46(5):507–523, 1999.
- [5] J.-L. Marichal and P. Mathonet. Extensions of system signatures to dependent lifetimes: explicit expressions and interpretations. *J. of Multivariate Analysis*, 102(5):931–936, 2011.
- [6] J. Navarro, J. M. Ruiz, and C. J. Sandoval. A note on comparisons among coherent systems with dependent components using signatures. *Stat. & Prob. Letters*, 72:179–185, 2005.
- [7] J. Navarro and T. Rychlik. Reliability and expectation bounds for coherent systems with exchangeable components. *J. Multivariate Anal.*, 98(1):102–113, 2007.
- [8] J. Navarro, F. J. Samaniego, N. Balakrishnan, and D. Bhattacharya. On the application and extension of system signatures in engineering reliability. *Naval Research Logistics*, 55:313–327, 2008.
- [9] J. Navarro, F. Spizzichino, and N. Balakrishnan. Applications of average and projected systems to the study of coherent systems. *J. Multivariate Anal.*, 101(6):1471–1482, 2010.
- [10] F. Spizzichino. *Subjective probability models for lifetimes*. Monographs on Statistics and Applied Probability, 91. Boca Raton, FL: Chapman & Hall/CRC, 2001.
- [11] F. J. Samaniego. On closure of the IFR class under formation of coherent systems. *IEEE Trans. Reliability Theory*, 34:69–72, 1985.
- [12] F. J. Samaniego. *System signatures and their applications in engineering reliability*. Int. Series in Operations Research & Management Science, 110. New York: Springer, 2007.
- [13] Z. Zhang. Ordering conditional general coherent systems with exchangeable components. *J. of Stat. Planning and Inference*, 140:454–460, 2010.

MATHEMATICS RESEARCH UNIT, FSTC, UNIVERSITY OF LUXEMBOURG, 6, RUE COUDENHOVE-KALERGI, L-1359 LUXEMBOURG, LUXEMBOURG

E-mail address: `jean-luc.marichal[at]uni.lu`

MATHEMATICS RESEARCH UNIT, FSTC, UNIVERSITY OF LUXEMBOURG, 6, RUE COUDENHOVE-KALERGI, L-1359 LUXEMBOURG, LUXEMBOURG

E-mail address: `pierre.mathonet[at]uni.lu`, `p.mathonet[at]ulg.ac.be`

MATHEMATICS RESEARCH UNIT, FSTC, UNIVERSITY OF LUXEMBOURG, 6, RUE COUDENHOVE-KALERGI, L-1359 LUXEMBOURG, LUXEMBOURG AND BOLYAI INSTITUTE, UNIVERSITY OF SZEGED, ARADI VÉRTANÚK TERE 1, H-6720 SZEGED, HUNGARY

E-mail address: `twaldha[at]math.u-szeged.hu`