

Editorial Board

Chairman: Imre J. Rudas

E-mail: rudas@bmf.hu

Editor-in-Chief: András Bakó

E-mail: bako@bmf.hu

Executive Editor: Péter Tóth

E-mail: toth.peter@tmpk.bmf.hu

Thomson Reuters Information

Acta Polytechnica Hungarica has been selected for coverage in Thomson Reuters products and custom information services. Beginning with V. 5 (1) 2008, this publication will be indexed and abstracted in the following:

- ◆ Science Citation Index Expanded (also known as SciSearch®),
- ◆ Journal Citation Reports/Science Edition.

Acta Polytechnica Hungarica

is published by Budapest Tech. Address: Bécsi út 96/B, H-1034 Budapest, Hungary.

Responsibility for the content rests upon the authors and not upon Budapest Tech.

Editors

Scientific Advisors

Péter Arató, Hungary
János Demetrovics, Hungary
László Pap, Hungary
András Prekopa, Hungary

Associate Editors

Shamsudin Amin, Malaysia
Marcelo Ang, Singapore
Miroslav Badida, Slovakia
András Benczúr, Hungary
Balázs Benyó, Hungary
János F. Bitó, Hungary
Gerard Capolino, France
Paul Drews, Germany
Wilfried Eimenreich, Austria
Ferenc Friedler, Hungary
Toshio Fukuda, Japan
Tom D. Gedeon, Australia
Ahmed Abo Ismail, Egypt
Karel Jezernik, Slovenia
Okyay M. Kaynak, Turkey
Jozef Kelemen, Czech Republic
Jim Keller, USA
Oussama Khatib, USA
Tim King, England
Dusan Kocur, Slovakia
László T. Kóczy, Hungary
Ladislav Madarász, Slovakia
András Medve, Hungary
Endre Pap, Serbia
Stefan Preitl, Romania
Tamás Réti, Hungary
J. A. Tenreiro Machado, Portugal
Spyros Tzafestas, Greece
Annamária R. Várkonyi-Kóczy, Hungary
Péter Váriaki, Hungary
Liberios Vokorokos, Slovakia
Miomir Vukobratović, Serbia

Assistant Editor

Anikó Szakál
E-mail: szakal@bmf.hu

Monograph: Aggregation Functions

Michel Grabisch^a, Jean-Luc Marichal^b, Radko Mesiar^c,
Endre Pap^d

^aUniversity of Panthéon-Sorbonne, Paris, France
e-mail: michel.grabisch@univ-paris1.fr

^bUniversity of Luxembourg, Luxembourg
e-mail: jean-luc.marichal@uni.lu

^cSlovak University of Technology
Bratislava, Slovakia
e-mail: mesiar@math.sk

^dDepartment of Mathematics and Informatics, University of Novi Sad
Trg Dositeja Obradovića 4, 21000 Novi Sad, Serbia
e-mail: pape@eunet.yu

Abstract: There is given a short overview of the monograph "Aggregation Functions" (M. Grabisch, J. L. Marichal, R. Mesiar, E. Pap), Cambridge University Press (in press) with more details from introductory Chapters 1 and 2.

Keywords: aggregation function, continuity, absolute continuity, Lipschitz condition, conjunction, disjunction, internality

1 Introduction

The central problem we are investigated in this book is the process of combining several numerical values into a single representative, one is called *aggregation*, and the numerical function performing this process is called *aggregation function*, asking also some natural conditions as monotonicity and boundary conditions. There is large field of application of aggregation: applied mathematics (e.g., probability, statistics, decision theory), computer sciences (e.g., artificial intelligence, operations research), as well as many applied fields (economy and finance, pattern recognition and image processing, data fusion, multicriteria decision aid, automated reasoning, etc.). The rapid growing of the above mentioned application fields, largely due to the arrival of computers, has made necessary the establishment of a sound theoretical basis for aggregation functions. Most of the results were disseminated in various journals or specialized

books, where usually only one specific class of aggregation functions devoted to one specific domain is discussed. Today, an important amount of literature is already available, many significant results have been found (such as characterizations of various families of aggregation functions), and many connections have been done with either related fields or former works (such as triangular norms in probabilistic metric spaces, theory of means and averages, etc.), see [1, 2, 3, 4, 5, 6, 7, 8, 9, 10]. For the researcher as well as for the practitioner, this proficiency and abundance of literature, but scattered in many domains, are more a handicap than an advantage, and there is a real lack of a unified and complete view of aggregation functions, where one could find the most important concepts and results, presented in a clear and rigorous way. This book has been written in the intention of filling this gap: it offers a full, comprehensive, rigorous and unified treatment of aggregation functions, and is intended to serve as a reference book in the field. Our main motivation has been to bring a unified viewpoint of the aggregation problem, and to provide an abstract mathematical presentation and analysis of aggregation functions used in various disciplines, without referring explicitly to a given domain. The book also provides to the field a unified terminology and notation.

After giving the contents of the book, we present only the basic definition of the aggregation function with few common examples. As an illustration, we mention only two additional elementary mathematical properties: continuity and position with respect to Min and Max. After that we mention very briefly some topics of the book.

2 Contents of the book

Preface

Chapter 1. Introduction

- 1.1 Main motivations and scope
- 1.2 Basic definitions and examples
- 1.3 Conventional notation

Chapter 2. Properties for aggregation

- 2.1 Introduction
- 2.2 Elementary mathematical properties
 - 2.2.1 Monotonicity
 - 2.2.2 Continuity
 - Standard continuity
 - A stronger form: Uniform continuity
 - A stronger form: Absolute continuity
 - A stronger form: Lipschitz condition
 - Weaker forms: left and right continuities
 - 2.2.3 Symmetry
 - Going from the symmetric case to the nonsymmetric case
- 2.2.4 Idempotency
- Idempotent functions

- Idempotent elements
- Range-idempotent functions
- Endpoint-preservation
- Strong idempotency
- Idempotizable functions
 - 2.2.5 Conjunction, disjunction, and internality
- k -conjunctive and k -disjunctive functions
- 2.3 Grouping based properties
 - 2.3.1 Associativity
 - 2.3.2 Decomposability and strong decomposability
- Decomposability
- Strong decomposability
 - 2.3.3 Autodistributivity
 - 2.3.4 Bisymmetry
- 2.4 Invariance properties
 - 2.4.1 Ratio, difference, interval, and log-ratio scales
 - 2.4.2 Ordinal scales
 - 2.4.3 Inversion of scales
- 2.5 Further properties
 - 2.5.1 Neutral and annihilator elements
 - 2.5.2 Additivity and related properties
- Chapter 3. Conjunctive and disjunctive aggregation functions**
 - 3.1 Preliminaries and general notes
 - 3.2 Generated conjunctive aggregation functions
 - 3.3 Triangular norms and related conjunctive aggregation functions
 - 3.3.1 Conjunctors and bwt-norms
 - 3.3.2 Basic t-norms
 - 3.3.3 Families of t-norms
 - 3.3.4 Different kinds of continuities for t-norms
 - 3.3.5 Continuous Archimedean t-norms
 - 3.3.6 Additive generators of t-norms
 - 3.3.7 Ordinal sums of t-norms and representation of continuous t-norms
 - 3.3.8 T-subnorms
 - 3.3.9 Convergence of t-norms
 - 3.4 Copulas and quasi-copulas
 - 3.4.1 General copulas
 - 3.4.2 Representation of copulas and Archimedean copulas
 - 3.4.3 Some additional results on copulas
 - 3.4.4 Quasi-copulas
 - 3.5 Disjunctive aggregation functions
 - 3.6 Uninorms
 - 3.6.1 Basic properties
 - 3.6.2 Characterization of uninorms
 - 3.6.3 Idempotent uninorms
 - 3.6.4 Generated uninorms
 - 3.7 Nullnorms

3.8 More aggregation functions related to t-norms

(i) Gamma operators

(ii) Exponential convex T-S-operators

(iii) Linear convex T-S-operators

(iv) Symmetric sums related to t-norms and t-conorms

3.9 Restricted distributivity

Chapter 4. Means and averages

4.1 Introduction and definitions

4.2 Quasi-arithmetic means

4.2.1 Axiomatization of quasi-arithmetic means

4.2.2 Subclasses of quasi-arithmetic means

4.3 Generalizations of quasi-arithmetic means

4.3.1 Quasi-linear means and quasi-linear functions

4.3.2 Quasi-arithmetic means with weight function

4.3.3 Nonstrict quasi-arithmetic means

4.4 Associative means

4.5 Means constructed from a mean value property

4.5.1 Lagrangian means

4.5.2 Cauchy means

4.6 Constructing means

4.6.1 Idempotization

4.6.2 Symmetrization

4.6.3 Minimization

4.7 Further extended means

4.7.1 Sierpiński carpet

4.7.2 Generated weight triangles

Chapter 5. Aggregation functions based on nonadditive integrals

5.1 Introduction

5.2 Set functions, capacities, and games

5.2.1 Starting from weight vectors

5.2.2 Capacities and related set functions

5.3 Some linear transformations of set functions

5.4 The Choquet integral

5.4.1 Definitions

5.4.2 Relation to interpolation and lattice polynomials

5.4.3 Expressions of the Choquet integral

5.4.4 Properties of the Choquet integral

5.4.5 Characterization of the Choquet integral

5.4.6 Particular Choquet integrals

5.4.7 Multilevel Choquet integral

5.5 The Sugeno integral

5.5.1 Weighted lattice polynomial functions

5.5.2 Definition and expressions of the Sugeno integral

5.5.3 Properties of the Sugeno integral

5.5.4 Characterization of the Sugeno integral

5.5.5 Particular Sugeno integrals

5.6 Other integrals

The (S, U)-integral

The Murofushi integral

The Benvenuti integral

The universal integral

Chapter 6. Construction methods

6.1 Introduction

6.2 Transformed aggregation functions

6.2.1 φ -invariant aggregation functions6.2.2 φ -duality

6.2.3 Self-dual aggregation functions

6.2.4 More general transformations

6.3 Composed aggregation

6.3.1 Composition based on associativity

6.3.2 Composition of functions

6.4 Weighted aggregation functions

6.4.1 Integer weights

6.4.2 Nonnegative real weights

6.4.3 Qualitative approach

6.5 Some other aggregation-based construction methods

6.5.1 Idempotization

6.5.2 Augmenting

6.5.3 Method of flying parameter

6.5.4 Constraining

6.6 Aggregation functions based on minimal dissimilarity

6.7 Ordinal sums of aggregation functions

6.8 Extensions to aggregation functions

Chapter 7. Aggregation on specific scale types

7.1 Introduction

7.2 Ratio scales

7.2.1 General descriptions

7.2.2 Aggregation functions on ratio scales

7.3 Difference scales

7.3.1 General descriptions

7.3.2 Aggregation functions on difference scales

7.4 Interval scales

7.4.1 General descriptions

7.4.2 Aggregation functions on interval scales

7.5 Log-ratio scales

Chapter 8. Aggregation on ordinal scales

8.1 Introduction

8.2 Order invariant subsets

8.3 Lattice polynomial functions and some of their properties

8.3.1 Lattice polynomial functions

8.3.2 Special lattice polynomial functions

8.3.3 Properties of lattice polynomial functions

- 8.4 Ordinal scale invariant functions
 - 8.4.1 General descriptions
 - 8.4.2 The nondecreasing case
 - 8.4.3 The continuous case
- 8.5 Comparison meaningful functions on a single ordinal scale
 - 8.5.1 General descriptions
 - 8.5.2 The nondecreasing case
 - 8.5.3 The continuous case
- 8.6 Comparison meaningful functions on independent ordinal scales
 - 8.6.1 General descriptions
 - 8.6.2 The nondecreasing case
 - 8.6.3 The continuous case
- 8.7 Aggregation on finite chains by chain independent functions
 - 8.7.1 Introduction
 - 8.7.2 Aggregation by ordinal scale invariant functions
 - 8.7.3 Aggregation by comparison meaningful functions on a single ordinal scale
 - 8.7.4 Aggregation by comparison meaningful functions on independent ordinal scales
- Chapter 9. Aggregation on bipolar scales**
 - 9.1 Introduction
 - 9.2 Associative bipolar operators
 - 9.2.1 Pseudo-additions
 - S is a strict t-conorm
 - S is a nilpotent t-conorm
 - S is the maximum operator
 - S is an ordinal sum of continuous Archimedean t-conorms
 - 9.2.2 Pseudo-multiplications
 - 9.3 Minimum and maximum on symmetrized linearly ordered sets
 - 9.3.1 The symmetric maximum
 - 9.3.2 Symmetric minimum
 - 9.3.3 Nonassociative calculus
 - 9.4 Separable aggregation functions
 - $A^+ = A^-$ is a strict t-conorm S
 - $A^+ = A^-$ is a continuous t-conorm S
 - A^+, A^- are integral-based aggregation functions
 - 9.5 Integral-based aggregation functions
 - 9.5.1 Bicapacities
 - 9.5.2 The Choquet integral for bicapacities
 - 9.5.3 The Sugeno integral for bicapacities
- Chapter 10. Behavioral analysis of aggregation functions**
 - 10.1 Introduction
 - 10.2 Expected values and distribution functions
 - 10.2.1 Expected values
 - 10.2.2 Andness and orness degrees of internal functions
 - 10.2.3 Idempotency degrees of conjunctive and disjunctive functions
 - 10.2.4 Distribution functions and moments

- 10.3 Importance indices
- 10.4 Interaction indices
- 10.5 Maximum improving index
- 10.6 Tolerance indices
 - 10.6.1 Veto and favor indices
 - 10.6.2 k -conjunction and k -disjunction indices
- 10.7 Measures of arguments contribution and involvement
 - 10.7.1 Index of arguments contribution
 - 10.7.2 Index of arguments involvement
- Chapter 11. Identification of aggregation functions**
 - 11.1 Introduction
 - 11.2 General formulation
 - 11.2.1 The identification problem
 - 11.2.2 Least squares regression
 - 11.3 The case of parameterized families of aggregation functions
 - Linear convex T-S-operators
 - Exponential convex T-S-operators
 - OWA operators
 - 11.4 The case of generated aggregation functions
 - 11.5 The case of integral-based aggregation functions
 - 11.5.1 Constraints
 - 11.5.2 Least mean squares criterion
 - 11.5.3 Case of ordinal information
 - Maximum separation criterion
 - Minimum distance and variance approaches
 - 11.5.4 The case of Sugeno integral
 - 11.6 Available software
- Appendix A. Aggregation of infinitely many arguments**
 - A.1 Introduction
 - A.2 Infinitary aggregation functions on sequences
 - Additive aggregation functions
 - Comonotonic additive aggregation function
 - Symmetric aggregation function
 - Countable extendable aggregation functions
 - Countable extension
 - Means and averages
 - A.3 General aggregation of infinite number of inputs
- Appendix B. Examples and applications**
 - B.1 Main domains of applications
 - Decision making under multiple criteria or attributes
 - Multiobjective optimization
 - Multiperson decision making
 - Pattern recognition and classification
 - Image analysis
 - Rule-based systems
 - B.2 A specific application: mixture of uncertainty measures

- S-measures and independence
- Hybrid mixtures
- Hybrid possibilistic-probabilistic utility function
- List of symbols
- Bibliography
- Index

3 Definition and examples

A first natural requirement comes from the fact that the output should be a synthetic value. Then, if inputs are supposed to lie in interval $[a, b]$, the output should also lie in this interval. Moreover, if all input values are identical to the lower bound a , then the output should be also a , and similarly for the case of the upper bound b . This defines a boundary condition. A second natural requirement is nondecreasing monotonicity. It means that if some of the input values increase, the representative output value should reflect this increase, or at worst, stay constant. These two requirements are commonly accepted in the field, and we adopt them as the basic definition of an aggregation function.

We assume throughout that the variables of any aggregation function have a common domain \mathbb{I} which is a nonempty real interval, bounded or not. In some exceptional cases, \mathbb{I} will represent a nonempty interval of the extended real number system $\overline{\mathbb{R}} := [-\infty, \infty]$. This latter case will always be mentioned explicitly and then we will often consider the convention that $0 \cdot \infty = 0$ and $\infty + (-\infty) = -\infty$. Denote by \mathbb{N} the set of strictly positive integers and $[n] := \{1, \dots, n\}$. Further, $|K|$ is the cardinal number of the set K .

Definition 3.1. An aggregation function in \mathbb{I}^n is a function $A^{(n)} : \mathbb{I}^n \rightarrow \mathbb{I}$ that

- (i) is nondecreasing (in each variable)
- (ii) fulfills the boundary conditions

$$\inf_{\mathbf{x} \in \mathbb{I}^n} A^{(n)}(\mathbf{x}) = \inf \mathbb{I} \quad \text{and} \quad \sup_{\mathbf{x} \in \mathbb{I}^n} A^{(n)}(\mathbf{x}) = \sup \mathbb{I}.$$

To give a first instance, the arithmetic mean, defined by

$$AM^{(n)}(\mathbf{x}) := \frac{1}{n} \sum_{i=1}^n x_i \tag{1}$$

is clearly an aggregation function in any domain \mathbb{I}^n .

The aggregation of a singleton is not an actual aggregation, the following convention is often considered: $A^{(1)}(x) = x \quad (x \in \mathbb{I})$.

We now introduce the concept of extended aggregation function.

Definition 3.2. An extended aggregation function in $\cup_{n \in \mathbb{N}} \mathbb{I}^n$ is a mapping

$$A : \bigcup_{n \in \mathbb{N}} \mathbb{I}^n \rightarrow \overline{\mathbb{R}}.$$

whose restriction $A^{(n)} := A|_{\mathbb{I}^n}$ to \mathbb{I}^n is an aggregation function in \mathbb{I}^n for any $n \in \mathbb{N}$.

For example, the arithmetic mean as an extended aggregation function is the sequence $(AM^{(n)})_{n \in \mathbb{N}}$, where $AM^{(n)}$ is defined by (1) for all $n \in \mathbb{N}$.

For illustration we now give a small list of well-known aggregation functions.

- (i) The arithmetic mean function $AM : \mathbb{I}^n \rightarrow \mathbb{I}$ and the geometric mean function $GM : \mathbb{I}^n \rightarrow \mathbb{I}$ are respectively defined by

$$AM(\mathbf{x}) := \frac{1}{n} \sum_{i=1}^n x_i, \quad GM(\mathbf{x}) := \left(\prod_{i=1}^n x_i \right)^{1/n}.$$

Note that, when $n > 1$, the geometric mean is not an aggregation function in any domain. We must consider a domain \mathbb{I}^n such that $\mathbb{I} \subseteq [0, \infty]$.

- (ii) For any $k \in [n]$, the projection function $P_k : \mathbb{I}^n \rightarrow \mathbb{I}$ and the order statistic function $OS_k : \mathbb{I}^n \rightarrow \mathbb{I}$ associated with the k th argument, are respectively defined by

$$P_k(\mathbf{x}) := x_k, \quad OS_k(\mathbf{x}) := x_{(k)},$$

where $x_{(k)}$ is the k th lowest coordinate of \mathbf{x} , that is,

$$x_{(1)} \leq \dots \leq x_{(k)} \leq \dots \leq x_{(n)}.$$

The projection onto the first and the last coordinates are defined as

$$P_F(\mathbf{x}) := P_1(\mathbf{x}) = x_1, \quad P_L(\mathbf{x}) := P_n(\mathbf{x}) = x_n.$$

Similarly, the extreme order statistics $x_{(1)}$ and $x_{(n)}$ are respectively the minimum and maximum functions

$$\text{Min}(\mathbf{x}) := \min(x_1, \dots, x_n),$$

$$\text{Max}(\mathbf{x}) := \max(x_1, \dots, x_n),$$

which will sometimes be written by means of the lattice operations \wedge and \vee , respectively, that is,

$$\text{Min}(\mathbf{x}) = \bigwedge_{i=1}^n x_i \quad \text{and} \quad \text{Max}(\mathbf{x}) = \bigvee_{i=1}^n x_i.$$

The median of an odd number of values (x_1, \dots, x_{2k-1}) is simply defined by

$$\text{Med}(x_1, \dots, x_{2k-1}) := x_{(k)},$$

which can be rewritten as

$$\text{Med}(x_1, \dots, x_{2k-1}) = \bigwedge_{\substack{K \subseteq [2k-1] \\ |K|=k}} \bigvee_{i \in K} x_i.$$

For instance, we have

$$\text{Med}(x_1, x_2, x_3) = (x_1 \vee x_2) \wedge (x_1 \vee x_3) \wedge (x_2 \vee x_3).$$

For any $\alpha \in \mathbb{I}$, we also define the α -median, $\text{Med}_\alpha : \mathbb{I}^n \rightarrow \mathbb{I}$, by

$$\text{Med}_\alpha(\mathbf{x}) = \text{Med}(x_1, \dots, x_n, \underbrace{\alpha, \dots, \alpha}_{n-1}).$$

- (iii) For any nonempty subset $K \subseteq [n]$, the *partial minimum* $\text{Min}_K : \mathbb{I}^n \rightarrow \mathbb{I}$ and the *partial maximum* $\text{Max}_K : \mathbb{I}^n \rightarrow \mathbb{I}$, associated with K , are respectively defined by

$$\text{Min}_K(\mathbf{x}) := \bigwedge_{i \in K} x_i, \quad \text{Max}_K(\mathbf{x}) := \bigvee_{i \in K} x_i.$$

- (iv) For any weight vector $\mathbf{w} = (w_1, \dots, w_n) \in [0, 1]^n$ such that $\sum_{i=1}^n w_i = 1$, the *weighted arithmetic mean function* $\text{WAM}_\mathbf{w} : \mathbb{I}^n \rightarrow \mathbb{I}$ and the *ordered weighted averaging function* $\text{OWA}_\mathbf{w} : \mathbb{I}^n \rightarrow \mathbb{I}$ associated with \mathbf{w} , are respectively defined by

$$\text{WAM}_\mathbf{w}(\mathbf{x}) := \sum_{i=1}^n w_i x_i, \quad \text{OWA}_\mathbf{w}(\mathbf{x}) := \sum_{i=1}^n w_i x_{(i)}.$$

- (v) The *sum* $\Sigma : \mathbb{R}^n \rightarrow \mathbb{R}$ and *product* $\Pi : \mathbb{R}^n \rightarrow \mathbb{R}$ functions are respectively defined by

$$\Sigma(\mathbf{x}) := \sum_{i=1}^n x_i, \quad \Pi(\mathbf{x}) := \prod_{i=1}^n x_i.$$

Note that, when $n > 1$, these latter two functions are not aggregation functions in any domain. For the sum, we need to consider the domains $[-\infty, \infty]^n, [-\infty, \infty]^{[n]}, [-\infty, \infty]^{[n]}, [-\infty, \infty]^{[n]}, [0, \infty]^n,]0, \infty]^n, [0, \infty]^{[n]},]0, \infty]^{[n]}, [-\infty, 0]^n, [-\infty, 0]^{[n]}, [-\infty, 0]^{[n]},]-\infty, 0]^n,]-\infty, 0]^n$. For the product, when restricted to $[0, \infty]^n$, we have to consider the domains $[0, 1]^n,]0, 1]^n, [0, 1]^{[n]},]0, 1]^{[n}, [1, \infty]^n,]1, \infty]^n, [1, \infty]^{[n]},]1, \infty]^{[n}, [0, \infty]^n,]0, \infty]^n, [0, \infty]^{[n]},]0, \infty]^{[n}$.

- (vi) Assume $\mathbb{I} = [a, b]$ is a closed interval. The *smallest* and the *greatest* aggregation functions in $[a, b]^n$ are respectively defined by

$$A_\perp(\mathbf{x}) := \begin{cases} b, & \text{if } x_i = b \text{ for all } i \in [n], \\ a, & \text{otherwise.} \end{cases}$$

$$A_\top(\mathbf{x}) := \begin{cases} a, & \text{if } x_i = a \text{ for all } i \in [n], \\ b, & \text{otherwise.} \end{cases}$$

By definition, we have $A_\perp \leq A \leq A_\top$ for any aggregation function $A : [a, b]^n \rightarrow [a, b]$.

4 Elementary mathematical properties

The basic properties for aggregation functions are divided into elementary mathematical properties (monotonicity, continuity, symmetry, etc.), grouping based properties (associativity, decomposability, etc.), invariance to a change of scale (ratio, difference, interval, ordinal scales), and various other properties (neutral and annihilator elements, additivity, etc.). We also introduce invariance properties related to the scale types of the variables being aggregated.

We restrict here only on an analytical property: continuity, and the position on real axe.

4.1 Continuity

Standard continuity

Definition 4.1. $F : \mathbb{I}^n \rightarrow \mathbb{R}$ is a continuous function if

$$\lim_{\substack{\mathbf{x} \rightarrow \mathbf{x}^* \\ \mathbf{x} \in \mathbb{I}^n}} F(\mathbf{x}) = F(\mathbf{x}^*) \quad (\mathbf{x}^* \in \mathbb{I}^n).$$

The continuity property essentially means that any small changes in the arguments (possible minor errors) should not entail a big change in the aggregated value (output error).

Figure 1. provides a Venn diagram showing the relationship between some classes of functions defined on bounded closed intervals.




Figure 1: Relations between some classes of functions

A stronger form: Uniform continuity

Definition 4.2. Let $\|\cdot\| : \mathbb{R}^n \rightarrow [0, \infty[$ be a norm and let $D \subseteq \mathbb{I}^n$. A function $F : \mathbb{I}^n \rightarrow \mathbb{R}$ is uniformly continuous in D (with respect to $\|\cdot\|$) if for every $\varepsilon > 0$ there exists $\delta > 0$ such that $|F(x) - F(y)| < \varepsilon$ whenever $\|x - y\| < \delta$ and $x, y \in D$.

A function $F : [a, b]^n \rightarrow \mathbb{R}$ is uniformly continuous in $[a, b]^n$ if and only if it is continuous in $[a, b]^n$.

A stronger form: Absolute continuity

Definition 4.3. We say that the unary function $f : [a, b] \rightarrow \mathbb{R}$ is absolutely continuous if, for every $\varepsilon > 0$, there exists $\delta > 0$ such that for any finite system of pairwise nonintersecting intervals $]a_i, b_i[\subset]a, b[$, $i \in [n]$, for which $\sum_{i=1}^n (b_i - a_i) < \delta$ the inequality $\sum_{i=1}^n |f(b_i) - f(a_i)| < \varepsilon$ holds.

Every absolutely continuous function on a closed interval is continuous on this interval.

We extend the notion of absolute continuity to functions of two variables. For that purpose we introduce, for a subrectangle $R = [a', b'] \times [c', d']$ of $[a, b] \times [c, d]$ and a function $F : [a, b] \times [c, d] \rightarrow \mathbb{R}$, the following notation

$$\Delta_R(F) = F(a', c') - F(b', c') - F(a', d') + F(b', d').$$

Definition 4.4. We say that $F : [a, b] \times [c, d] \rightarrow \mathbb{R}$ is absolutely continuous if the following two conditions are satisfied

(i) given $\varepsilon > 0$, there exists $\delta > 0$ such that

$$\sum_{R \in \mathcal{R}} |\Delta_R(F)| < \varepsilon$$

whenever \mathcal{R} is a finite collection of pairwise nonoverlapping subrectangles $R = [a', b'] \times [c', d']$ of $[a, b] \times [c, d]$ with

$$\sum_{R \in \mathcal{R}} \lambda(R) < \delta,$$

where λ is the Lebesgue measure;

(ii) the marginal functions $F(\cdot, d)$ and $F(b, \cdot)$ are absolutely continuous functions of a single variable on $[a, b]$ and $[c, d]$, respectively.

A stronger form: Lipschitz condition The continuity property can be strengthened into the well-known Lipschitz condition.

Definition 4.5. Let $\|\cdot\| : \mathbb{R}^n \rightarrow [0, \infty[$ be a norm. If a function $F : \mathbb{I}^n \rightarrow \mathbb{R}$ satisfies the inequality

$$|F(x) - F(y)| \leq c \|x - y\| \quad (x, y \in \mathbb{I}^n), \tag{2}$$

for some constant $c \in]0, \infty[$, then we say that F satisfies the Lipschitz condition or is Lipschitzian (with respect to $\|\cdot\|$). More precisely, any function $F : \mathbb{I}^n \rightarrow \mathbb{R}$ satisfying (2) is said to be c -Lipschitzian. The greatest lower bound of constants $c > 0$ in (2) is called the Lipschitz constant.

Important examples of norms are given by the Minkowski norm of order $p \in [1, \infty[$, namely

$$\|x\|_p := \left(\sum_{i=1}^n |x_i|^p \right)^{1/p},$$

also called the L_p -norm, and its limiting case $\|x\|_\infty := \max_i |x_i|$, which is the Chebyshev norm.

The c -Lipschitz condition has an interesting interpretation when applied in aggregation. It allows us to estimate the relative output error in comparison with input errors

$$|F(x) - F(y)| \leq c\varepsilon$$

whenever $\|x - y\| \leq \varepsilon$ for some $\varepsilon > 0$.

The Lipschitz property of functions is defined standardly on domains where the norm cannot achieve ∞ . Formally it can be defined also on \mathbb{I}^n for an unbounded interval \mathbb{I} . However, if \mathbb{I} contains $-\infty$ or ∞ , then the Lipschitz property does not imply continuity, in general. For example, the smallest aggregation function A_{\perp} on $[0, \infty]^n$ is Lipschitzian for any norm but it is not continuous.

In all cases when $\mathbb{I} \subseteq \mathbb{R}$, the Lipschitz property implies the absolute continuity. However, the converse does not hold. For instance, $\log x$ on $]0, 1]$ is absolutely continuous but not Lipschitzian.

Example 4.6. (i) The product $\Pi : [0, 1]^2 \rightarrow [0, 1]$ is 1-Lipschitzian. Therefore it is absolutely continuous.

(ii) The geometric mean $GM : [0, 1]^2 \rightarrow [0, 1]$ is not Lipschitzian. However, it is continuous and therefore uniformly continuous.

Proposition 4.7. The smallest and the greatest aggregation functions defined in $[a, b]^n$ being 1-Lipschitzian with respect to the norm $\|\cdot\|_p$ are respectively given by $A_*^{(n)} : [a, b]^n \rightarrow [a, b]$, with

$$A_*^{(n)}(x) := \text{Max}(b - \|n \cdot b - x\|_p, a),$$

and $A^{*(n)} : [a, b]^n \rightarrow [a, b]$, with

$$A^{*(n)}(x) := \text{Min}(a + \|x - n \cdot a\|_p, b).$$

4.2 Conjunction, disjunction, and internality

Considering the functions Min and Max as dominating or dominated functions gives rise to three main classes of aggregation functions: *conjunctive* functions, *disjunctive* functions and *internal* functions.

Definition 4.8. $A : \mathbb{I}^n \rightarrow \overline{\mathbb{R}}$ is conjunctive if

$$\inf \mathbb{I} \leq A \leq \text{Min}.$$

Conjunctive functions combine values as if they were related by a logical “and” operator. That is, the result of combination can be high only if all the values are high. Thus, a low value can never be compensated by a high value. The most common conjunctive functions defined on $[0, 1]^n$ are t-norms.

Definition 4.9. $A : \mathbb{I}^n \rightarrow \overline{\mathbb{R}}$ is disjunctive if

$$\text{Max} \leq A \leq \sup \mathbb{I}.$$

Disjunctive functions combine values as an “or” operator, so that the result of combination is high if at least one value is high. Thus, a high value cannot be compensated by a low value. Such functions are, in this sense, dual of conjunctive functions. The most common disjunctive functions defined on $[0, 1]^n$ are t-conorms.

Definition 4.10. A function $A : \mathbb{I}^n \rightarrow \overline{\mathbb{R}}$ is internal if

$$\text{Min} \leq A \leq \text{Max}.$$

Internality is a property shared by all the means and averaging functions, which are the most often encountered functions in the literature on aggregation. In multicriteria decision making, these functions are also called *compensative* functions. In fact, in this kind of functions, a bad (respectively good) score on one criterion can generally be compensated by a good (respectively bad) one on another criterion, so that the result of the aggregation will be medium.

5 Some topics of the book

The presentation is mathematical and rigorous, avoiding jargon and inherent imprecision from the various applied domains where aggregation functions are used (often under different names such aggregation operators, merging functions, connectives, etc.), but keeping as far as possible the standard terminology of mathematics. As far as possible, every result is given with its proof, unless the proof is long and requires extra material. In this case, a reference to the proof is always given.

A large section is devoted in chapter on disjunctive aggregation functions to triangular norms (t-norms for short): different families, continuous Archimedean t-norms, additively generated t-norms, ordinal sums, etc. There are presented

important class of copulas, well known in probability theory, as well as uninorms and nullnorms (as hybrid of t-norms and t-conorms). The concepts of means and averages functions, as well as their relationships, are presented in full generality. Then main subclasses of means, such as quasi-arithmetic ones, some of their special cases (e.g., root-mean-power functions, exponential means), and some of their generalizations (e.g., quasi-linear means, quasi-arithmetic means with weight functions, non-strict quasi-arithmetic means) are presented. Considering nonadditive integrals in the discrete finite case defines a new important class of aggregation functions. Nonadditive integrals are defined with respect to capacities (nonadditive monotone measures), and in particular generalize the notion of expected value. Special care is taken on the Choquet integral, since this is the most representative of nonadditive integrals, and the Sugeno integral, and finally other families of nonadditive integrals. There are given some operations to create new aggregation functions from existing ones, as transformation, composition, introduction of weights on variables, ordinal sums, and various other means (idempotization, symmetrization, etc.). We describe the aggregation functions that are meaningful when considering ratio, difference, and interval scales. We analyze how to extend aggregation functions to the interval $[-1, 1]$ (bipolar scale), that is, to perform a kind a symmetrization with respect to 0 while keeping properties of the aggregation function. Behavioral analysis of aggregation functions is done through various indices and values (like the expected value), which in some sense constitutes the identity card of the aggregation function.

An important topic in practice is how to choose a suitable aggregation function. There are given various ways to identify aggregation functions from data. There are presented the rather unexpected consequences of defining an aggregation function with an infinite (either countable or uncountable) number of arguments. There is given a short description with references of the main fields of applications of aggregation functions, essentially in decision making, data fusion, and artificial intelligence, and with more details on an application to the mixture of uncertainty measures.

6 Conclusion

We present here only few of basic definitions contained in Chapters 1 and 2 of the book. The book is intended primarily to researchers and graduate students in applied mathematics and computer sciences, secondarily to practitioners in, e.g., decision making, optimization, economy and finance, artificial intelligence, data fusion, computer vision, etc. It could also be used as a textbook for graduate students in applied mathematics and computer sciences.

Acknowledgment

The authors gratefully acknowledge the support of various projects which facilitated communication and collaboration among us, in particular the bilateral project between France and Serbia *Pavle Savić* N° 11092SF supported by EGIDE, Paris, and its Serbian part Action SK-SRB-19, the internal research project *Decision Mathematics and Operations Research* supported by the University of Luxembourg, projects APVV-0375-06, APVV-0012-07, and VEGA 1/4209/07, supported by the Slovak Grant Agency for Sciences, the national grants MNTRS (Serbia, Project 144012), Provincial Secretariat for Science and Technological Development of Vojvodina, and MTA HMTA (Hungary).

References

- [1] Aczél, J.: *Lectures on functional equations and their applications*, Mathematics in Science and Engineering, Vol. 19, Academic Press, New York, 1966.
- [2] Aczél, J. and Dhombres, J.: *Functional equations in several variables*, Encyclopedia of Mathematics and its Applications 31, Cambridge University Press, Cambridge, 1989.
- [3] Alsina, C. and Frank, M. J. and Schweizer, B.: *Associative functions, Triangular norms and copulas*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2006.
- [4] Beliakov, G. and Pradera, A. and Calvo, T.: *Aggregation Functions: A Guide for Practitioners*, Studies in Fuziness and Soft Computing, Springer, Berlin, 2007.
- [5] Bouchon-Meunier, B. (ed.): *Aggregation and fusion of imperfect information*, Studies in Fuzziness and Soft Computing. 12. Heidelberg: Physica-Verlag, 1998.
- [6] Bullen, P. S.: *Handbook of means and their inequalities*, Mathematics and its Applications 560, Kluwer Academic Publishers Group, Dordrecht, 2003.
- [7] Calvo, T. and Mayor, G. and Mesiar, R. (eds.): *Aggregation operators*, Studies Fuzziness Soft Computing 97, Physica, Heidelberg, 2002.
- [8] Klement, E. P. and Mesiar, R. and Pap, E.: *Triangular norms*, Trends in Logic—Studia Logica Library 8, Kluwer Academic Publishers, Dordrecht, 2000.
- [9] Nelsen, R. B.: *An introduction to copulas*, Lecture Notes in Statistics 139, Springer-Verlag, New York, 1999.
- [10] Torra, V. and Narukawa, Y.: *Modeling decisions: Information Fusion and Aggregation Operators*, Cognitive Technologies, Springer, 2007.